STUDIES OF CALCIUM AND PHOSPHORUS METABOLISM

XVIII. ON TEMPORARY FLUCTUATIONS IN THE LEVEL OF CALCIUM AND INORGANIC PHOSPHORUS IN BLOOD SERUM OF NORMAL INDIVIDUALS

RAY F. FARQUHARSON AND DOROTHY M. TIBBETTS

(From the Medical Clinic of the Massachusetts General Hospital, Boston)

(Received for publication November 26, 1930)

In the investigation of various problems of calcium and phosphorus metabolism, it has been found that great changes in the volume and direction of the calcium and phosphorus stream may occur without any notable difference in the serum levels of these elements. In hyperthyroidism the calcium excretion in the urine is relatively huge and there are large negative balances of calcium and phosphorus, yet the serum calcium and serum phosphorus¹ remain within normal limits (1). Moreover, there is no appreciable change in the serum levels of healthy people when, by changing the calcium intake, a positive balance is shifted to a negative one or vice versa.

Yet the actual and relative amounts of calcium and phosphorus in the serum is of the utmost importance in certain abnormal conditions. MacCallum and Voegtlin (2) described the low blood and tissue calcium associated with tetany after removal of the parathyroids. Greenwald (3) later demonstrated the associated rise in serum phosphorus and since that time the high phosphorus, low calcium of the serum in many types of tetany has come to be recognized. Marriot and Howland (4), DeWesselow (5) and others have found a similar relationship of the serum values in terminal nephritis. On injection of large amounts of inorganic phosphate into dogs Binger (6) produced tetany with a low serum calcium and high serum phosphorus. Similar results have been found to follow injection or ingestion of very large quantities of phosphate by numerous investigators (7, 8, 9). The

¹ Throughout this paper the term "serum phosphorus" is used in referring to serum inorganic phosphorus.

feeding of much smaller amounts of phosphate to rachitic rats has been found to produce tetany with the typical blood changes by Karelitz and Shohl (10). That in rachitic animals even the high phosphorus metabolism of fasting may have similar effects has been shown by Cavins, (11), Wilder, (12), and Shohl and Brown (13). The ingestion of sodium oxalate in dogs was found to result in a low blood calcium, high phosphorus, and tetany (14).

In states of hyperparathyroidism, on the other hand, a high serum calcium is well known to be associated with a low serum phosphorus. This has been discussed in other papers (15, 16).

It is quite clear, therefore, that gross changes in the serum phosphorus value in many conditions are associated with equally marked variation of the serum calcium in the opposite direction. That the level of serum calcium is also affected by the protein content of the serum was shown by Salvesen and Linder (17). Hastings, Murray and Sendroy (18) showed that there was a linear relationship between the protein content of serum or transudate and the calcium level. Their data, taken from human sera and transudates as drawn from the body, indicate that 0.014 millimol of calcium are bound per gram of protein at the pH value of the blood. Peters and Eiserson (19) also found in a group of cases, mostly nephritics, that the concentration of calcium in the serum varies directly with the concentration of protein and inversely with the concentration of inorganic phosphorus.

Other investigators have described temporary fluctuations in either serum calcium or phosphorus of normal individuals in response to various factors. Thus, Stewart and Haldane (20) found that an increase of from 1.5 to 2 mgm. in serum calcium might occur after ingestion of 30 grams CaCl_2 , after forced breathing for 90 minutes, and after breathing 6 to 7 per cent CO_2 for 90 minutes; and a decrease of about 2 mgm. after ingestion of 60 grams NaHCO_3 . Although values for serum phosphorus were not given in this paper it was reported separately (21) that forced breathing for 90 minutes resulted in every instance in a fall in serum inorganic phosphorus to below 1.25 mgm. Salvesen, Hastings, and McIntosh (22) found that intravenous injection of CaCl_2 resulted in a moderate rise in serum phosphorus as well as a temporary rise in serum calcium.

The object of our investigation was to determine the extent of

temporary variation in the serum levels of calcium and phosphorus in normal adults and to what degree the inverse relationship noted above would prevail when changes in one or another component could be induced at will. Accordingly, the serum levels were determined at various times of the day in subjects on ordinary diets and after ingestion of food rich respectively in carbohydrates, fat, and protein, as well as when insulin was given preceding a carbohydrate meal. The effect of ingestion of large amounts of salts of calcium and phosphorus was also followed.

METHODS

Venous blood was taken from the arm into a clean syringe, previously washed with liquid paraffin, care being taken to avoid any unnecessary manipulation and to prevent hemolysis. As soon as clotting was complete the serum was separated by centrifugalization, and removed. In some cases the CO_2 content or capacity of the plasma was determined by the Van Slyke method, the blood having been taken under oil and coagulation prevented by the use of heparin. For these determinations we are indebted to Dr. A. V. Bock. Serum protein was determined by macro Kjeldahl, serum phosphorus by the method of Fiske and Subbarow (23), and serum calcium by Fiske's (24) method. The latter method has proven most satisfactory under all conditions. Determinations were done in duplicate, usually with identical results, the variation seldom being greater than 2 per cent.

In all cases, blood was first taken during fasting and then at varying intervals after different types of meals or after ingestion of calcium salts or phosphates.

EXPERIMENTS AND RESULTS

The level during fasting

It was important to find whether the basal fasting level would be appreciably affected by changes in type of diet or by ingestion of excess acid, alkali, or phosphate as well as to note any temporary change occurring immediately after such ingestion. Also, it was interesting to observe the relative constancy of the fasting levels of the serum calcium and phosphorus over comparatively long periods. In table 1 are presented data for such observations on two healthy individuals.

274 NORMAL FLUCTUATIONS IN BLOOD CALCIUM AND PHOSPHORUS

TABLE 1
Serum values (fasting) in two healthy subjects

Date	Calcium	Phosphorus	Protein	Plasma CO ₂ content	Total fixed base	Remarks
Subject R. F. F.						
1928	mgm. per 100 cc.	mgm. per 100 cc.	per cent	volumes per cent	m.Eq. per liter	
January 27	9.6	3.4	6.9	70.7	148	Ordinary diet
February 2	9.7	4.2		71.2	152	6 grams NaHCO ₃ taken daily, January 27 to February 2
February 6	9.6	3.6	6.3	69.5		12 grams NaHCO ₃ + 12 grams sodium citrate February 2-6
February 9	9.1	3.2	6.4	71.8	168	52 grams NaHCO ₃ on February 7 56 grams NaHCO ₃ on February 8
February 11	10.0	3.7		58.6		4 grams NH ₄ Cl on February 9 12 grams NH ₄ Cl on February 10
February 13	10.0	3.4	8.0	29.1*	162	12 grams NH ₄ Cl on February 11 15 grams NH ₄ Cl on February 12
February 20	9.6	3.9				Ordinary diet
February 27	10.1	3.2				Ordinary diet
April 10	10.1	2.5	5.9			Low calcium diet April 5-14
April 19	9.6	3.2	6.4			High protein diet April 14-27
April 23	10.1	3.0		63.8		
April 26	10.2	3.6	7.1	61.7		
June 16	9.6	3.8				High protein diet June 11-16
June 19	9.5	3.4	6.6			Ordinary diet
June 22	9.6	3.6				Ordinary diet
June 26	9.6	3.2	6.6			Ordinary diet
1930						
May 26	10.0	3.1				
Subject D. M. T.						
1928						
February 27	10.1	3.3				Ordinary diet throughout
March 5	9.8	3.6				
June 27	9.2	3.3	6.9			
November 24	9.5	3.7				
December 10	10.0	3.5				
1930						
May 27	9.6	3.2				

* Total CO₂ oxygenated whole blood.

In the case of R. F. F. the fasting serum calcium was remarkably constant at about 9.6 to 10.0 mgm. per 100 cc. Under the influence of ingestion of very large amounts of alkali, it fell to 9.1 mgm., a result similar, although not as great, as that obtained by Stewart and Haldane (20). Apart from this single lower value all other determinations, including those taken during periods on very high protein diets and on ingestion of large amounts of NH_4Cl , lay between 9.5 and 10.2 mgm. Variation in the serum phosphorus level was a little greater than that of calcium, ranging as it did from 2.5 to 4.2 mgm. In eight instances, when the fasting serum calcium was 9.6 or 9.7 mgm., the phosphorus varied between 4.2 and 3.2 mgm.; when the serum calcium was 10.0 to 10.2 serum phosphorus values ranged from 2.5 to 3.7 mgm. The effect of high phosphorus intake in the form of a very high protein diet (200 grams protein daily) had no appreciable effect on the fasting values nor did the ingestion of large doses of NH_4Cl . The serum protein varied from 5.9 to 8.0 per cent.

In subject D. M. T. a smaller number of determinations showed a slightly greater variation in the calcium and smaller variation in the phosphorus level.

That ingestion of large amounts of inorganic phosphate in patients on a low calcium diet produced no appreciable effect on the fasting serum calcium and phosphorus levels has been mentioned in another paper (25). After several weeks on a low calcium diet, also, as described elsewhere (26), ingestion of large amounts of NH_4Cl was associated with slightly decreased levels for calcium and phosphorus in some cases, and ingestion of NaHCO_3 with increased levels for serum calcium. In all experiments, however, on ingestion of acid or alkali on high or low calcium diets, the total range of variation of many determinations for any given individual was never greater than 1 mgm. calcium except in one instance. In patient D. A. (Case II, period XV), described elsewhere (26), the ingestion of large amounts of NH_4Cl after the subject had been on a low calcium diet for several weeks, resulted in a fall of slightly more than 1 mgm. in serum calcium.

Absolute rest and mild exercise

In one instance, it was found that there was a definite slight increase in serum calcium and protein and fall in serum phosphorus in blood

taken from a fasting, healthy subject after lying at complete rest for one and one-quarter hours. This was repeated in a number of healthy individuals; but no material change in levels of calcium or phosphorus was found, although the serum protein did change considerably. The results are presented in table 2.

TABLE 2
Effect of absolute rest and mild exercise during fasting

Subject	Date	Calcium	Phos- phorus	Protein	Remarks
	1928	<i>mgm. per 100 cc.</i>	<i>mgm. per 100 cc.</i>	<i>per cent</i>	
R. F. F.	April 19	{ 9.6 10.1	{ 3.2 2.8	{ 6.4 7.6	{ After walking $\frac{1}{2}$ mile After 1 $\frac{1}{4}$ hours complete rest
C. F.	June 28	{ 9.5 9.5	{ 3.8 3.5	{ 5.9 6.6	{ At rest, before rising After working one hour
F. A.	June 28	{ 9.8 9.6	{ 3.6 3.6	{ 7.1 5.9*	{ After working one hour After lying down one hour
D. M. T.	November 24	{ 9.5 9.7	{ 3.7		{ After working three hours After lying down one hour
	December 10	{ 10.0 10.0	{ 3.5 3.2		{ After working three hours After lying down one hour
G. P. R.	November 24	{ 10.4 10.4	{ 4.5		{ Before rising Moving about for one hour
	December 10	{ 10.2 10.4	{ 3.9 3.5		{ Before rising Moving about for one hour

* Some hemolysis.

Fluctuations in serum calcium during the day when on ordinary diet

The serum levels of two healthy subjects were determined at different times during a day when an ordinary diet was taken and ordinary work done. Results are presented in table 3. In subject R. F. F. the calcium remained constant at 9.9 to 10.0 mgm., while the phosphorus varied between 3.0 to 3.8 mgm. In D. M. T. the serum

calcium varied between 9.4 and 10.4 mgm., a range as great as that of the fasting values over many months; and the serum phosphorus between 3.1 and 3.8 mgm. It is notable that the highest value for calcium was obtained at the same time as the highest value for phosphorus.

TABLE 3
Serum values during the day on ordinary diet

Subject	Date	Time	Calcium	Phosphorus	Protein	Remarks
	1930		<i>mgm. per 100 cc.</i>	<i>mgm. per 100 cc.</i>	<i>per cent</i>	
R. F. F.	May 26	9:00	10.0	3.1	6.8	Eggs, toast, and coffee
		9:30	Breakfast			
		11:30	10.0	3.0	7.5	
		12:45	9.9	3.2	7.5	Ham, carrots, potatoes, custard
		1:00	Lunch			
		2:15	10.0	3.8	7.2	
		5:00	10.0	3.8	7.2	
D. M. T.	May 26	9:00	9.6	3.2	7.8	Toast, marmalade, coffee
		9:30	Breakfast			
		11:30	9.7	3.1	8.0	
		12:15	Lunch			Ham, potatoes, carrots, custard
		1:30	9.4	3.7	7.8	
		4:45	10.4	3.8	8.3	

Ingestion of carbohydrate, or carbohydrate with administration of insulin

It has been clearly shown that the administration of insulin or the ingestion of large amounts of carbohydrate with or without insulin results in a prompt fall in the serum inorganic phosphorus (27, 28, 29). It was, therefore, interesting to find to what extent the serum calcium would vary when rapid changes in serum phosphorus were induced in this way. Accordingly, large amounts of carbohydrate in the form of ordinary foods or 100 grams of glucose were given to a number of healthy subjects and to some patients, including a case of tetany with high serum phosphorus. The changes in serum phosphorus, calcium, and blood sugar were followed for several hours. Results are given in table 4.

TABLE 4
Effect of ingestion of large amounts of carbohydrates

Subject	Date	Time	Cal- cium	Phos- phorus	Blood sugar	Pro- tein	Remarks
	1928		mgm. per 100 cc.	mgm. per 100 cc.	mgm. per 100 cc.	per cent	
R. F. F.	Febru- ary 2	Fasting 2½ hours	9.7 9.6	4.2 3.2			High carbohydrate break- fast
	Febru- ary 27	Fasting ½ hour 1 hour 2 hours 4 hours	10.1 10.2 10.3 9.9 10.0	3.3 3.0 3.0 3.1 3.4	81 95 68 72 68		Sugar tolerance. 100 grams glucose
D. M. T.	March 5	Fasting ¼ hour 1 hour 2 hours 4 hours	9.8 9.8 9.8 9.7 9.7	3.6 3.0 2.9 2.7 3.0	86 94 97 91 83		High carbohydrate meal Carbohydrate, 122 grams Protein, 4 grams Fat, 10 grams Calcium, 64 mgm. Phosphorus, 77 mgm.
E. R.	March 5	Fasting ¾ hour 2 hours 3½ hours	9.1 9.1 9.4 9.1	3.8 2.8 2.7 3.1	93 93 101 84		High carbohydrate meal Carbohydrate, 122 grams Protein, 4 grams Fat, 10 grams Calcium, 64 mgm. Phosphorus, 77 mgm.
B. W. (Tetany)	Febru- ary 9	Fasting ½ hour 1½ hours 2 hours 3½ hours	4.7 4.8 4.4 4.7 4.2	7.5 7.1 7.1* 6.5 6.6*	90 124 130 110 123	7.1	High carbohydrate break- fast Carbohydrate, 171 grams Calcium, 23 mgm. Phosphorus, 105 mgm.
D. S. (Acromegaly)	Febru- ary 26	Fasting ¾ hour 2½ hours	9.5 9.7 9.7	5.7 4.7 4.7	114 147 162	7.1	Sugar tolerance. 100 grams glucose
Bln (Acromegaly)	March 5	Fasting ½ hour 1 hour 2 hours 4 hours	9.8 9.8 9.6 9.8 9.6	4.3 4.2 3.8 3.7 3.8	95 143 170 154 87		Sugar tolerance. 100 grams glucose
S. A. (Hypertrichosis)	Janu- ary 9	Fasting ½ hour 1½ hours 3 hours	9.5 9.5 10.3 9.7	2.9 2.5 2.6 2.6	92 173 150 114	6.2	Sugar tolerance. 100 grams glucose

* Some hemolysis.

The effects of ingestion of large amounts of carbohydrate immediately after the administration of a large dose of insulin in three normal subjects are shown by data collected in table 5.

TABLE 5
Effect of ingestion of large amounts of carbohydrate and administration of insulin

Subject	Date	Time	Calcium	Phosphorus	Blood sugar	Protein	Remarks
	1928		mgm. per 100 cc.	mgm. per 100 cc.	mgm. per 100 cc.	per cent	
R. F. F.	June 26	Fasting	9.6	3.2	98	6.6	30 units insulin
		50 minutes	9.4	2.4	105	6.9	Breakfast
		1½ hours	10.0	2.1	64	6.7	Carbohydrate, 126 grams
		3 hours	10.4	2.1	46	6.8	Protein, 8 grams
		10 grams cane sugar with temporary relief					Fat, 3 grams
		4 hours	9.5	2.4	46	7.1	Calcium, 67 mgm.
		30 grams glucose					Phosphorus, 170 mgm.
		5½ hours	9.6	2.9	46	7.2	
		30 grams glucose plus 25 grams bread					
		8 hours	9.7	4.2	91		
D. M. T.	June 27	Fasting	9.2	3.3	84	6.9	20 units insulin
		½ hour	9.6	2.8	77	7.4	Breakfast
		2 hours	9.6	2.5	73	6.8	Carbohydrate, 109 grams
		3½ hours	9.8	3.0	41	6.1	Protein, 8 grams
		Lunch					Fat, 7 grams
		7½ hours	9.2	3.3	105	6.6	Calcium, 70 mgm.
E. R.	June 27						Phosphorus, 118 mgm.
		Fasting	9.3	4.3	93	6.2	20 units insulin
		½ hour	9.5	3.5	104	6.1	Breakfast
		2 hours	9.4	3.7	53	6.1	Carbohydrate, 109 grams
		3½ hours	9.3	3.6	60	6.1	Protein, 8 grams
		Lunch					Fat, 7 grams
		7½ hours	9.0	3.6	200		Calcium, 70 mgm.
							Phosphorus, 118 mgm.

The results, in so far as the serum calcium and phosphorus levels are concerned, were essentially the same for ingestion of carbohydrate alone as when it followed the administration of insulin. In the latter cases, however, the blood sugar fell to reaction levels.

In all subjects there was an early fall in serum phosphorus levels, averaging in eleven instances 0.8 mgm. and varying from slightly less than 0.5 mgm. to slightly more than 1.0 mgm. The initial height of the serum phosphorus seemed to have no influence on the magnitude of the change, nor did there seem to be any definite relation between the extent of the decrease in serum phosphorus and the change in the blood sugar level. When the phosphorus fell there was some tendency for the serum calcium to rise but this was by no means constant nor did the increase in serum calcium vary directly with the extent of decrease in serum phosphorus. With a maximum fall of about 1 mgm. in serum phosphorus the serum calcium remained essentially constant in R. F. F., February 2; D. M. T., March 5; E. R., March 5; and in DS (acromegaly). In the case B. W. (tetany), when the phosphorus fell from 7.5 to 6.5 the calcium remained essentially constant until the last determination, when, without further appreciable change in serum phosphorus it fell from 4.7 to 4.2. On the other hand, in R. F. F., June 26, the serum calcium rose from 9.6 to 10.4 mgm., while the phosphorus fell from 3.2 to 2.1. In eight hours, however, when the serum calcium had returned to 9.7 the serum phosphorus had risen to 4.2, 1 mgm. higher than the fasting level. In D. M. T., June 27, when the serum phosphorus was decreased by 0.8 mgm. the calcium rose from 9.2 to 9.6, reaching 9.8 one and one-half hours later when phosphorus had returned nearly to normal. In the case of hypertrichosis, the maximum rise in calcium was 0.9 mgm. and maximum fall in phosphorus 0.4 mgm. The serum calcium was essentially unchanged in at least six of the eleven cases, and only in three cases was there a variation of 0.5 mgm. or more, the greatest increase above fasting level being 1 mgm. in R. F. F. after carbohydrate and insulin. In the cases in which serum protein was determined there seemed to be no constant relation between the temporary variations in protein and those of calcium.

Thus, our results are in agreement with the well established fact that the ingestion of large amounts of carbohydrate with or without administration of insulin is constantly followed by a definite fall in serum phosphorus. They show on the whole little associated change in serum calcium, the magnitude of which was no greater than that found to occur on ordinary diet or after ingestion of other types of food

as described below. Similar values for serum calcium in dogs, after administration of insulin, were obtained by Briggs, Koechig, Doisy and Weber (30). Brougher, however, reported tremendous changes in blood calcium of rabbits (rising even to more than 20 mgm.) following injection of insulin (31). Davies, Dickens, and Dodds (32) found increases of from 2 to 4 mgm. in serum calcium of rabbits in hypoglycemic convulsions. Recently, Ellsworth (33) has reported increases in serum calcium in man similar to those that occurred in some of our cases. The maximum change in calcium did not occur always in those cases with the greatest fall in phosphorus, nor was the reciprocal inverse relationship persistent in the last determinations of a given experiment. Yet in his cases there was a more constant rise in serum calcium than in ours.

Ingestion of large amounts of protein and fat

Data showing the effect of ingestion of large amounts of protein and fat are presented in table 6.

When 127 grams of fat were taken, much of it in the form of cream, there followed an increase of calcium from 9.6 to 10.8 mgm. and a fall in phosphorus from 3.9 to 3.2 mgm. A week later the effect of 100 grams of olive oil was observed; the serum phosphorus remained constant but serum calcium increased by 0.9 mgm. in 4 hours. Similar experiments on ingestion of large amounts of protein, when the subject took 200 grams of protein daily for several days, showed in one instance a distinct rise in calcium from fasting value of 9.6 mgm. to 11.4 mgm. a few hours after breakfast, associated with a slight increase in serum phosphorus, but no material difference in serum protein. In this instance NaHCO_3 was added to neutralize the acid effect of ingested protein. When the experiment was subsequently repeated, leaving out the NaHCO_3 , the serum calcium remained stationary and serum phosphorus rose slightly. On each occasion the subject was carrying on with his ordinary work.

Data showing the effect of ingestion of single doses of phosphate and of calcium lactate are presented in table 7. The phosphate caused purgation, which persisted for several hours. The serum phosphorus, however, rose from 3.5 to 5.0 mgm., without an appreciable change in the serum calcium level. A similar rise in serum phosphorus after

ingestion of large amounts of phosphate without any associated change in the serum calcium level has been described by Schulz (34). The ingestion of calcium lactate was followed by an increase in serum

TABLE 6
Effect of ingestion of large amounts of protein and of fat

Date	Time	Cal- cium	Phos- phorus	Pro- tein	Non- protein nitrogen	Remarks
1928		<i>mgm. per 100 cc.</i>	<i>mgm. per 100 cc.</i>	<i>per cent</i>	<i>mgm. per 100 cc.</i>	
February 20	Fasting	9.6	3.9			Breakfast
	1 hour	10.5	3.2			Fat, 127 grams
	2½ hours	10.6	3.3			Carbohydrate, 15 grams
	5½ hours	10.8	3.6			Protein, 6 grams
	Lunch					Calcium, 185 mgm.
	9 hours	10.0	3.7			Phosphorus, 181 mgm.
February 27	Fasting	10.1	3.2			100 grams olive oil
	1 hour	10.8	3.2			
	2 hours	10.3	3.2			
	4 hours	11.0	3.1			
February 6	Fasting	9.6	3.6	6.3	31	Breakfast
	2½ hours	10.0	3.6	6.9	39	Protein, 61 grams Fat, 29 grams Calcium, 30 mgm. Phosphorus, 461 mgm.
April 19	8:30 a.m.	9.6	3.2	6.4	45	Total diet for day
	9:45 a.m.	10.1	2.8	7.6	46	Protein, 200 grams
	10:00 a.m.	Breakfast, plus 2 grams soda				Carbohydrate, 161 grams
	12:45 p.m.	11.4	3.7	6.9	51	Fat, 100 grams
	1:30 p.m.	Lunch, plus 3 grams soda				Calcium, 119 mgm.
	2:30 p.m.	10.2	3.8	6.8	50	Phosphorus, 1,586 mgm.
	6:30 p.m.	Dinner, plus 3 grams soda				
	9:15 p.m.	10.2	4.2		59	
April 26	8:50 a.m.	10.2	3.6	7.1	42	Same diet as on April 19
	9:00 a.m.	Breakfast				
	10:15 a.m.	10.0	3.7	6.8	50	
	12:15 p.m.	10.3	3.5	7.2	47	
	1:20 p.m.	10.2	3.3	6.8	50	
	1:30 p.m.	Lunch				
	6:15 p.m.	10.0	4.1	6.9	51	

calcium from 9.6 to 11.0 mgm. but the inorganic phosphorus remained relatively constant.

TABLE 7
Effect of ingestion of sodium phosphate and of calcium lactate

Date	Time	Calcium	Phosphorus	Remarks
		mgm. per 100 cc.	mgm. per 100 cc.	
1928				
June 19	Fasting	9.5	3.5	3 grams phosphorus in form of equimolecular solution of NaH_2PO_4 and Na_2HPO_4 . Pur-gation followed in about one hour
	1 hour	9.4	5.0	
	3½ hours	9.5	4.9	
	5 hours	9.2	5.0	
June 22	Fasting	9.6	3.6	10 grams calcium lactate = 1.3 grams calcium
	2 hours	11.0	3.6	
	3 hours	10.5	3.6	
	5 hours	10.5	4.2*	
	Lunch			
	9 hours	10.3	3.8	

* Some hemolysis.

COMMENT

It is seen that there may be greater changes in the serum calcium and phosphorus within a few hours after meals of ordinary foodstuffs than occur in the fasting levels over periods of many months. The fasting serum calcium, however, and to a lesser extent the serum phosphorus, remains remarkably constant in spite of great changes in diet and in the intake of calcium and phosphate.

Although the inverse relationship between the levels of calcium and phosphorus of the serum seems to hold in some instances during temporary fluctuations, it is quite clear that in many cases the calcium was constant while the phosphorus level changed, in others the phosphorus remained constant during changes in calcium, and in some instances both rose or fell together. Concomitant variation in the serum protein was not great enough to overshadow a specific effect of variation in phosphorus on the calcium level or vice versa, nor were temporary fluctuations in protein always paralleled by corresponding changes in serum calcium when the serum phosphorus remained constant.

These observations do not affect the truth of the statement that the serum calcium level varies directly with the protein content and inversely with the concentration of inorganic phosphorus; for the changes described occurred during periods of flux, were of short duration and did not represent equilibrium values. Their magnitude, moreover was much less than that of corresponding changes in terminal nephritis, in cases with very low serum protein, and in tetany. It is possible, however, that they represent influences of other factors that are difficult of control, factors that might affect the level of either serum calcium or serum phosphorus independently.

SUMMARY

Fasting values for serum calcium and phosphorus of healthy individuals show only slight variations over long periods, whereas greater changes may occur within a few hours after ingestion of ordinary food-stuffs. It is interesting how small are the variations.

In such temporary fluctuations the serum calcium does not always vary inversely with the serum inorganic phosphorus, although there is some tendency for it to do so.

Temporary changes in serum protein are not always associated with corresponding variations in the serum calcium.

Ingestion of large amounts of carbohydrate or carbohydrate after administration of insulin is constantly followed by a fall in the serum inorganic phosphorus and sometimes but not always by a slight increase in serum calcium.

The magnitude of the diurnal variations in the serum calcium and phosphorus was small when compared with the great changes found in states of tetany and terminal nephritis.

BIBLIOGRAPHY

1. Aub, J. C., Bauer, W., Heath C., and Ropes, M., *J. Clin. Invest.*, 1929, vii, 97. *Studies of Calcium and Phosphorus Metabolism. III. The Effects of the Thyroid Hormone and Thyroid Disease.*
2. MacCallum, W. G., and Voegtlin, C., *J. Exp. Med.*, 1909, xi, 118. *On the Relation of Tetany to the Parathyroid Glands and to Calcium Metabolism.*
3. Greenwald, I., *J. Biol. Chem.*, 1913, xiv, 369. *On the Phosphorus Content of the Blood of Normal and Parathyroidectomized Dogs.*

4. Marriott, W. M., and Howland, John, *Arch. Int. Med.*, 1916, xviii, 708. Phosphate Retention as a Factor in the Production of Acidosis in Nephritis.
5. DeWesselow, O. L. V., *Quart. J. Med.*, 1922-23, xvi, 341. On the Phosphorus and Calcium of the Blood in Renal Disease.
6. Binger, C., *J. Pharm. and Exp. Therap.*, 1917, x, 105. Toxicity of Phosphates in Relation to Blood Calcium and Tetany.
7. Tisdall, F. F., *J. Biol. Chem.*, 1922, liv, 35. The Influence of the Sodium Ion in the Production of Tetany.
8. Salvesen, H. A., Hastings, A. B., and McIntosh, J. F., *J. Biol. Chem.*, 1924, lx, 311. Blood Changes and Clinical Symptoms Following Oral Administration of Phosphates.
9. Gates, F. L., and Grant, J. H. B., *J. Exp. Med.*, 1927, xlv, 125. Experimental Observations on Irradiated, Normal, and Partially Parathyroidectomized Rabbits. II. The Effects of Injections of Calcium Chloride or of Disodium Hydrogen Phosphate.
10. Karelitz, S., and Shohl, A. T., *J. Biol. Chem.*, 1927, lxxiii, 665. Rickets in Rats. II. The Effect of Phosphorus Added to the Diet of Ricketic Rats.
11. Cavins, A. W., *J. Biol. Chem.*, 1924, lix, 237. The Effect of Fasting (and Refeeding) on the Calcium and Inorganic Phosphorus in Blood Serum of Normal and Rachitic Rats.
12. Wilder, T. S., *J. Biol. Chem.*, 1929, lxxxi, 65. The Tetany of Fasting in Experimental Rickets.
13. Shohl, A. T., and Brown, H. B., *J. Biol. Chem.*, 1929, lxxxiv, 501. Rickets in Rats. X. Fasting Tetany and Phosphate Tetany.
14. Gross, E. G., *J. Biol. Chem.*, 1923, lv, 729. Inorganic Salt Metabolism. II. Inorganic Ion Ratio After Administration of Oxalates and Citrates.
15. Albright, F., Bauer, W., Ropes, M., and Aub, J. C., *J. Clin. Invest.*, 1929, vii, 139. Studies of Calcium and Phosphorus Metabolism. IV. The Effect of the Parathyroid Hormone.
16. Bauer, W., Albright, F., and Aub, J. C., *J. Clin. Invest.*, 1930, viii, 229. A Case of Osteitis Fibrosa Cystica (Osteomalacia) with Evidence of Hyperactivity of the Parathyroid Bodies. Metabolic Study II.
17. Salvesen, H. A., and Linder, G. C., *J. Biol. Chem.*, 1923-24, lviii, 617. Observations on the Inorganic Bases and Phosphates in Relation to the Protein of Blood and Other Body Fluids in Bright's Disease, and in Heart Failure.
18. Hastings, A. B., Murray, C. D., and Sendroy, J., Jr., *J. Biol. Chem.*, 1926-27, lxxi, 723. Studies of the Solubility of Calcium Salts. I. The Solubility of Calcium Carbonate in Salt Solutions and Biological Fluids.
19. Peters, J. P., and Eiserson, L., *J. Biol. Chem.*, 1929, lxxxiv, 155. The Influence of Protein and Inorganic Phosphorus on Serum Calcium.
20. Stewart, C. P., and Haldane, J. B. S., *Biochem. J.*, 1924, xviii, 855. Experimental Alterations in the Calcium Content of Human Serum and Urine.

21. Haldane, J. B. S., Wigglesworth, V. B., and Woodrow, C. E., *Proc. Roy. Soc., Series B.*, 1924, xcvi, 1. The Effect of Reaction Changes on Human Inorganic Metabolism.
22. Salvesen, H. A., Hastings, A. B., and McIntosh, J. F., *J. Biol. Chem.*, 1924, lx, 327. The Effect of the Administration of Calcium Salts on the Inorganic Composition of the Blood.
23. Fiske, C.H., and Subbarow, Y., *J. Biol. Chem.*, 1925, lxvi, 375. The Colorimetric Determination of Phosphorus.
24. Fiske, C. H., Unpublished Method for Calcium, first described in paper by Hamilton, B., *J. Biol. Chem.* 1925, lxv, 101.
25. Farquharson, R. F., Salter, W., and Aub, J. C., *J. Clin. Invest.*, 1931, x, 251. Studies of Calcium and Phosphorus Metabolism. XIII. The Effect of Ingestion of Phosphorus on the Excretion of Calcium.
26. Farquharson, R. F., Salter, W. T., Tibbetts, D. M., and Aub, J. C., *J. Clin. Invest.*, 1931, x, 221. Studies of Calcium and Phosphorus Metabolism. XII. The Effect of the Ingestion of Acid-Producing Substances.
27. Wigglesworth, V. B., Woodrow, C. E., Smith, W., and Winter, L. B., *J. Physiol.*, 1922-23, lvii, 447. On the Effect of Insulin on Blood Phosphate.
28. Harrop, G. A., and Benedict, E. M., *J. Biol. Chem.*, 1924, lix, 683. The Participation of Inorganic Substances in Carbohydrate Metabolism.
29. Blatherwick, N. R., Bell, M., and Hill, E., *J. Biol. Chem.*, 1924, lxi, 241. Some Effects of Insulin on the Carbohydrate and Phosphorus Metabolism of Normal Individuals.
30. Briggs, A. P., Koechig, I., Doisy, E. A., and Weber, C. J., *J. Biol. Chem.*, 1923-24, lviii, 721. Some Changes in the Composition of the Blood Due to the Injection of Insulin.
31. Brougher, J. C., *Am. J. Physiol.*, 1927, lxxx, 411. Blood Calcium as Affected by Insulin.
32. Davies, D. T., Dickens, F., and Dodds, E. C., *Biochem. J.*, 1926, xx, 695. Observations on the Preparation, Properties and Source of the Parathyroid Hormone.
33. Ellsworth, R., *J. Clin. Invest.*, 1930, viii, 139. Secondary Alterations in Total Serum Calcium After Administration of Glucose and Insulin.
34. Schulz, F., *Ann. Int. Med.*, 1930, iii, 667. Phosphorus in the Blood and Urine; A Study of the Excretion and Retention of Phosphorus in a Normal Subject and in Patients with Renal Disease.