

4. Kety, S. S., and Schmidt, C. F., The nitrous oxide method for the quantitative determination of cerebral blood flow in man: theory, procedure and normal values. *J. Clin. Invest.*, 1948, 27, 476.
5. Kety, S. S., Hafkenschiel, J. H., Jeffers, W. A., Leopold, I. H., and Shenkin, H. A., The blood flow, vascular resistance and oxygen consumption of the brain in essential hypertension. *J. Clin. Invest.*, 1948, 27, 511.
6. Scheinberg, P., and Stead, E. A., Jr., The cerebral blood flow in male subjects as measured by the nitrous oxide technique. Normal values for blood flow, oxygen utilization, glucose utilization and peripheral resistance, with observations on the effect of tilting and anxiety. *J. Clin. Invest.*, 1949, 28, 1163.
7. Kety, S. S., King, B. D., Horvath, S. M., Jeffers, W. A., and Hafkenschiel, J. H., The effects of an acute reduction in blood pressure by means of differential spinal sympathetic block on the cerebral circulation of hypertensive patients. *J. Clin. Invest.*, 1950, 29, 402.
8. Harmel, M. H., Hafkenschiel, J. H., Austin, G. M., Crumpton, C. W., and Kety, S. S., The effect of bilateral stellate ganglion block on the cerebral circulation in normotensive and hypertensive patients. *J. Clin. Invest.*, 1949, 28, 415.
9. Kety, S. S., Circulation and metabolism of the human brain in health and disease. *Am. J. Med.*, 1950, 8, 205.
10. Schmidt, C. F., *The Cerebral Circulation in Health and Disease*. Charles C. Thomas, Springfield, Ill., 1950, Ed. 1, p. 67.
11. Hafkenschiel, J. H., Crumpton, C. W., Moyer, J. H., and Jeffers, W. A., The effects of dihydroergocornine on the cerebral circulation of patients with essential hypertension. *J. Clin. Invest.*, 1950, 29, 408.

ERRATUM

In the article entitled "The Measurement of C-reactive Protein in Human Sera. Comparison of the Clinical Tests on the Basis of a Quantitative Method," by Harrison F. Wood and Maclyn McCarty, in the June 1951 issue (pp. 616-622), the data giving the concentration of C-reactive protein in human sera were calculated without allowance for the twofold concentration employed in the final step of the quantitative method. Consequently, the values for the concentration of the protein in human sera found in Table IV and in the text should be divided by two to obtain the correct figures.