correlation between erythrocyte sedimentation rate and the polysaccharide percentage of serum protein was found in rheumatoid arthritis patients.

Elevations of serum polysaccharides were found in two cases of active gout and in one case of lupus erythematosus, while essentially normal levels were noted in a limited number of patients with inactive gout, dermatomyositis, periarteritis nodosa, and Weber Christian Disease. The use of the determination of serum polysaccharide as a diagnostic aid to distinguish degenerative joint disease from rheumatoid arthritis is suggested.

REFERENCES

1. Ehrich, W. E., Nature of collagen diseases. Am. Heart J., 1952, **43**, 121.
2. Shetlar, M. R., Schmidt, H. L., Jr., Lincoln, R. B., DeVore, J. K., Bullock, J. A., and Hellbaum, A. A., Response of the serum polysaccharide fractions following cortisone treatment of patients with rheumatic fever. J. Lab. & Clin. Med., 1952, **39**, 372.
3. Shetlar, M. R., Shetlar, C. L., Richmond, V., and Everett, M. R., The polysaccharide content of serum fractions in carcinoma, arthritis, and infections. Cancer Research, 1950, **10**, 681.
4. Ludwig, A. W., Boas, N. F., and Soffer, L. J., Role of mucopolysaccharides in pathogenesis of experimental exophthalmos. Proc. Soc. Exper. Biol. & Med., 1950, **73**, 137.
5. Catchpole, H. R., Serum and tissue glycoproteins in mice bearing transplantable tumors. Proc. Soc. Exper. Biol. & Med., 1950, **75**, 221.
6. Milne, J., Serum protein fractionation: A comparison of sodium sulfate precipitation and electrophoresis. J. Biol. Chem., 1947, **169**, 595.
7. Kingsley, G. R., A rapid method for the separation of serum albumin and globulin. J. Biol. Chem., 1940, **133**, 731.
8. Weichselbaum, T. E., and Shapiro, I., A rapid and simple method for the determination of salicylic acid in small amounts of blood plasma. Am. J. Clin. Path. (Tech. Section), 1946, **10**, 42.
9. Winzler, R. J., Devor, A. W., Mehl, J. W., and Smyth, I. M., Studies on the mucoproteins of human plasma. I. Determination and isolation. J. Clin. Invest., 1948, **27**, 609.
10. Shetlar, M. R., Foster, J. V., and Everett, M. R., Determination of serum polysaccharides by the tryptophane reaction. Proc. Soc. Exper. Biol. & Med., 1948, **67**, 125.
11. Jager, B. V., and Nickerson, M., A simple quantitative chemical method for estimating γ-globulin in human serum. J. Biol. Chem., 1948, **173**, 683.
12. Gutman, A. B., The Plasma Proteins in Disease. 15. Rheumatoid Arthritis. Advances in Protein Chemistry, New York, Academic Press, Inc., 1948, **4**, 218.
13. Effersoe, P., Effect of adrenocorticotropic hormone (ACTH) and cortisone on serum proteins in various diseases. Scandinavian J. Clin. & Lab. Invest., 1950, **2**, 265.
14. Seibert, F. B., Seibert, M. V., Atno, A. J., and Campbell, H. W., Variation in protein and polysaccharide content of sera in the chronic diseases, tuberculosis, sarcoidosis, and carcinoma. J. Clin. Invest., 1947, **26**, 90.
15. Shetlar, M. R., The polysaccharides of serum proteins in cancer. Texas Rep. Biol. & Med., 1952, **10**, 228.

ERRATUM

Pulmonary Function in Boeck's Sarcoid, by M. Henry Williams, Jr., J. Clin. Invest., 32: 9, 909, 1953. Table II.

Ventilation
L./min./sq. meter B.S.A.
BTPS
$\dot{V}E$