THE FATE OF P^{32} LABELLED DIISOPROPYLFLUOROPHOSPHONATE IN THE HUMAN BODY AND ITS USE AS A LABELLING AGENT IN THE STUDY OF THE TURNOVER OF BLOOD PLASMA AND RED CELLS

By J. A. COHEN AND MIA G. P. J. WARRINGA

(From the Medical Biological Laboratory of the National Defence Research Council T. N. O., Rijswijk, Netherlands)

(Submitted for publication May 12, 1953; accepted October 28, 1953)

In recent years diisopropylfluorophosphonate (DFP) has established itself as a useful drug in such conditions as intestinal and vesical atonia, glaucoma and myasthenia gravis. With the consolidation of its place in therapy the desirability of more precise knowledge of the fate of this compound in the human organism became obvious. In the present study the absorption, the distribution in the blood and the excretion of DFP have been studied after intramuscular injection of the radioactive labelled substance (DFP^{32}) into three patients suffering from myasthenia gravis (two cases) and arteriosclerosis cerebri, respectively, and into two normal persons. There are no theoretical reasons why the metabolic fate of DFP should be different from normal in the patients studied. Indeed, no essential differences were observed between results obtained from the three patients and those from the normal persons. Therefore, all results to be described should be considered as applicable to the metabolism of DFP in the normal human organism.

In the course of this work it became obvious that the phosphorus of DFP^{32} was irreversibly bound to the proteins of blood plasma and red cells and that DFP was exclusively broken down to diisopropylphosphate, a substance which is not further metabolised nor taken up into any normal metabolic pool. Such properties suggested that DFP^{32} might prove to be a useful tool in the study of the turnover of blood plasma proteins and red cells. Preliminary results of the use of DFP^{32} in the determination of the life span of red cells and the half life of a plasma protein will be included in this study.

EXPERIMENTAL METHODS

Synthesis of DFP^{32} . DFP^{32} was prepared from $H_3P^{32}O_4$, purchased from the Atomic Energy Research Establish-

ment at Harwell by a method to be described elsewhere. It was injected intramuscularly as a solute in 1 to 2 ml. of peanut oil.

Radioactivity measurements. In the absorption experiments, samples of 0.1 ml. blood obtained from the fingertip were pipetted into flat counting dishes, which were provided with a snugly fitting piece of filter paper in the bottom. After drying, the samples were counted with a conventional Geiger Müller tube. All other radioactivity measurements were carried out with a G.M. liquid counter (M_6 of 20th Century Electronics). All measurements were related to standards consisting of diisopropylphosphate obtained by alkaline hydrolysis of a sample of DFP^{32} of known quantity and radioactivity.

Treatment of blood, urine, and feces for radioactivity measurements. Ten milliliters of blood obtained by venepuncture were collected in sodium citrate (final concentration 0.76 per cent). Red cells were separated from the plasma and washed thrice in normal saline. Counting was carried out on the hemolysate in distilled water. Samples, diluted if necessary, from 24-hour portions of urine were counted directly in the liquid counter. Feces were collected in aliquots over a period of 24 hours and mixed. Samples of 2 Gm. were suspended in a small volume of water and hydrolyzed during 15 minutes at $100^\circ C$. in molar NaOH.

Cholinesterase activity. Cholinesterase activity was measured manometrically at $37^\circ C$. according to Ammon (1) using acetylcholine as a substrate.

Paper chromatography. Paper chromatography was carried out on samples of 40 μ l. of urine using an ascending method with butanol-acetic acid-water mixtures as solvent. The development period lasted 16 hours at a temperature of $23^\circ C$.

Organic phosphates were made visible on the paper by a method which was essentially that of Hanes and Isherwood (2) with slight modifications.

Radioactivity on the paper was assessed by exposing x-ray films in direct contact with the chromatogram in the dark room for 72 hours.

RESULTS

The first series of experiments were performed on three young women. Two of these were treated in the Neurological Clinic of the Leyden Uni-

FIG. 1. RADIOACTIVITY IN WHOLE BLOOD AFTER INJECTION OF DFP^{32}
Activity expressed in counts per minute per 0.1 ml. sample of blood of patients A and B.

versity Hospital for myasthenia gravis (A and B), the third one (C) was a healthy volunteer. A, B, and C received dosages of 1.990, 0.240, and 0.433 mg. of DFP^{32} , respectively. The specific activity of the preparation used was approximately $40 \mu\text{c}$. per mg. DFP. One of the patients (A) had been

treated with OMPA (octamethylpyrophosphoramide) prior to the experiment. At the start of the experiment practically no cholinesterase activity was present in this case in blood plasma and red cells.

After completion of this series it became de-

FIG. 2. PLASMA BOUND RADIOACTIVITY AFTER INJECTION OF DFP^{32}
Activity expressed as the logarithm of the weight in μg . of DFP bound to 10 Gm. of plasma nitrogen of Subjects A, B, D, and E. Curves drawn by method of least squares.

FIG. 3. RED CELL BOUND RADIOACTIVITY AFTER INJECTION OF DFP³²

Activity expressed in $\mu\text{g. of DFP reacting per Gm. of erythrocyte nitrogen}$ in the blood of Subjects D and E. Curves drawn by method of least squares.

FIG. 4. PLASMA AND RED CELL CHOLINESTERASE

I, Red cell cholinesterase of Case B expressed in $\mu\text{l. CO}_2 \text{ per hour per 100 Gm. Hb.}$ II, Plasma cholinesterase of same person in $\mu\text{l. CO}_2 \text{ per hour per 100 Gm. N.}$ The arrow marks the moment of injection of 240 $\mu\text{g. DFP}^{32}$. P indicates the moment of neostigmine injection.

FIG. 5. EXCRETION OF P^{32} IN URINE

The excretion of radioactivity is expressed as percentage of the dose injected.

sirable to repeat in particular the experiments on plasma and red cell turnover using longer periods of observation. For this purpose a higher initial radioactivity of the labelled material was required. This could be achieved by injecting a preparation of a specific activity of approximately 200 μ c. per mg. DFP. Two patients of the City Hospital of the Hague were thus treated. One was an 80-year-old man suffering from arteriosclerosis cerebri (D); the other (E) was a 63-year-old woman who, at the time the experiment was started, had completely recovered from an attack of apoplexia and was considered normal. They received dosages of 0.44 and 0.51 mg. of DFP³², respectively.

Resorption of DFP

Blood samples from the fingertip from A and B were examined for radioactivity at 15-minute in-

tervals. Figure 1 shows the course of the radioactivity of the whole blood in counts per minute during an observation period of 130 minutes. There are two main differences between the curves obtained from A and B. Firstly, the maximal activity is reached after 30 minutes in Case A and only after 90 minutes in Case B. Secondly, the level of radioactivity is much higher in the latter case in spite of the fact that only one-eighth of the amount of radioactivity had been injected. These differences are readily explained as the result of the pretreatment with OMPA to which A had been subjected. This treatment has certainly resulted in blocking of the majority of the sites present on proteins of blood plasma and cells which are capable of reacting with DFP. These sites are readily accessible in Case B where they react with DFP to form the irreversible protein-DFP-reaction products well known from cholinesterase inhibition studies. The unbound DFP is assumed to be rapidly removed by the liver, the kidney, and other organs and the difference in level of curves A and B is thus explained on account of differences in bound DFP content of the blood. If the supposition is right that the protein-DFP linkages are irreversible, radioactivity will disappear at the rate of replacement of those proteins by newly formed molecules.

Binding of P^{32} to elements of the blood

For this experiment blood samples from A, B, D, and E were drawn at intervals. After separation of red cells and plasma the amount of DFP in micrograms that had reacted with plasma and cells, respectively, was calculated from the observed radioactivity.

Results are expressed in Figures 2 and 3. It is apparent that, in spite of the larger amount of DFP injected into A, the plasma and red cells harbour considerably less radioactivity than those of B, D, and E, as would be expected from the results presented in the previous passage.

Red cell and plasma cholinesterase activity was measured in the blood of B (Figure 4). A marked inhibition of plasma cholinesterase is observed down to approximately 10 per cent of the initial value. Low values are still observed on the eighth day. Unfortunately at that time it was necessary to start neostigmine treatment so that the further

FIG. 6. EXCRETION OF P^{32} IN FECES

The excretion of radioactivity is expressed as percentage of the dose injected.

course of the curve is not fit for detailed analysis. Slow restoration of activity seems however apparent after the twelfth day. The curve for the red cell cholinesterase, which is known to be less sensitive to DFP than the plasma enzyme, shows only a shallow and transitory dip. Moreover it occurred at least 24 hours after the injection. It seems highly unlikely that this transitory dip, occurring after a latent period on a dose of DFP of only 240 micrograms should be attributed to a true DFP inhibition of red cell cholinesterase.

Excretion of DFP^{32}

Radioactivity was measured in the urine of A, B, and C and also in the feces of A and B. In all cases, 60 to 70 per cent of the P^{32} injected as DFP appeared in the urine during the first 10 days (Figure 5). One-third was found in the urine in the first 24 hours. The amount of P^{32} excreted in the urine of A was slightly greater in the first few days and slightly less after the first three to four

days compared with B and C. This difference may be due to the OMPA treatment in case A, resulting in a modification of the ratio protein-bound/free P^{32} .

During the observation period less than 5 per cent of the injected radioactivity was recovered in the feces (Figure 6).

Paper chromatography of urine

The results so far reported provide only information on the distribution of the P^{32} incorporated in the injected DFP molecules. They convey nothing concerning the chemical alterations of the DFP molecule between its injection and its excretion. Therefore, the urine was further analysed by paper chromatography to elucidate the chemical character of the radioactive compounds it contained. Portions of urine of B and C collected during the first days after the injection were analysed. Portions of the same urine mixed with diisopropylphosphate (DIP) monoisopropylphos-

FIG. 7a. PAPER CHROMATOGRAM OF THE URINE OF C
Solvent: butanol-acetic acid-water.

- 1 urine
- 2 urine + DIP (diisopropylphosphate) + MIP (mono-isopropylphosphate) + AP (anorganic phosphate)
- 3 urine + DIP

FIG. 7b. RADIOGRAM OF THE CHROMATOGRAM OF
FIGURE 7a

- 4 urine + MIP
- 5 urine + AP
- 6 water + DIP + MIP + AP

phate (MIP) and inorganic phosphate were run simultaneously. These substances were considered to be the most likely phosphates to occur in the urine as a result of DFP break-down in the body.

In Figure 7 a chromatogram is presented of five samples of urine of C with additions as indicated. Number 6 is a control in water.

Unmixed urine (No. 1 in Figure 7) shows two large unidentified spots. Comparison with numbers 2, 5, and 6 suggests that the upper part of the lower spot contains probably inorganic phosphate. MIP is located immediately under the upper spot (2, 4, and 6) whereas DIP travels in the front (2, 3, and 6). The original urine (1) cannot be expected to give readily demonstrable spots resulting from DFP or its break-down products because the technique used is not sensitive enough to give a colour with the traces of phosphate³² excreted in 40 μ l. of urine (see Figure 5).

If, however, an x-ray film is exposed to the same chromatogram, an autoradiogram is obtained. A drawing of this radiogram is presented in Figure 7b. It proves conclusively that the injected DFP appears in the urine in only one form, DIP. It is concluded that at least the main part of the injected DFP is transformed into DIP and excreted

as such. It should be noted that these results were obtained on urine of the first days following injection of DFP³². Later, radioactivity becomes too slight to measure dependably. The possibility that in later periods other products appear can, therefore, not be entirely excluded. It could be argued that some of the injected DFP³² might be eliminated as such. This product would not appear on the chromatogram because of its volatility. A watery solution of DFP³² brought on filter paper rapidly loses its radioactivity due to this volatility. When a drop of fresh urine of C was treated in this way, no radioactivity disappeared on drying, showing that no free DFP was present nor any other volatile P³² containing compound.

DISCUSSION

The results obtained on the normal individuals C and E did not differ essentially from those observed in the patients A, B, and D. Such differences as were found between A on the one hand and B, C, D, and E on the other could be easily explained by the OMPA treatment of A prior to the administration of DFP.

The following picture of the fate of DFP may be tentatively suggested as a result of the present

experience with humans and by earlier experiments on animals by other authors (Figure 8). DFP is rapidly absorbed from the intramuscular depot. The presence of radioactive material in the blood can be demonstrated a few minutes after the injection. At the dosage used, maximal radioactivity in the blood is observed after 90 minutes, after which a slow decline occurs, indicating that removal now is predominant over resorption. Free DFP disappears rapidly from the circulation into such organs as kidney, liver, and lung which have been shown to contain radioactivity in similar experiments on rabbits described by Jandorf and McNamarra (3). A large fraction is probably hydrolyzed in the tissues to DIP and excreted into the urine and to a far less extent into the feces.

The hydrolysis may be spontaneous but is more likely to be essentially enzymatic. The enzyme system responsible is probably the fluorophosphatase, present in liver and kidney, as described by Mazur (4).

A certain amount of DFP³² reacts with the proteins of red cells, blood plasma and tissues. It is implied in the scheme of Figure 8 that this reaction is irreversible and that the reaction products are stable. This assumption is based on many studies reported in the literature, where it has been shown that a number of animal proteins, such as chymotrypsin, trypsin, cholinesterases, aliesterases etc. react with DFP. In all these instances the reaction was irreversible and the reaction product stable.

FIG. 8. TENTATIVE DIAGRAM OF THE METABOLIC FATE OF DFP IN THE HUMAN BODY

products. This possibility has not been explored but seems worth considering. This procedure might also be useful for the determination of red cell and plasma volume, particularly because of the irreversibility of the P^{32} label.

The possibility has not been ruled out as yet that the reaction-products of DFP and plasma or red cells do not behave entirely like the unlabelled structures. More evidence is required before the method can be confidently recommended as a reliable general technique. The values obtained seem reasonable compared with those described in the literature and the regular pattern of the curves provide some confidence that the methods might be useful.

With the dosages used no symptoms of toxicity have been observed.

SUMMARY

1. Five human subjects were injected intramuscularly with varying non-toxic doses of DFP³² (diisopropylfluorophosphonate labelled with P^{32}).

2. Under normal conditions radioactivity originating from DFP could be demonstrated in the peripheral blood within a few minutes; maximum radioactivity was reached in 90 minutes.

3. A small portion of the injected P^{32} was irreversibly bound to red cells and plasma. This fraction disappeared from the blood at a rate corresponding to the normal replacement of these elements.

4. This relationship could be exploited to calculate the rate of replacement of a plasma protein and the mean life span of red cells.

Values of 12 to 14 days were found for the half life of this plasma component in four persons.

A mean red cell life span of 116 and 129 days, respectively, was found in two subjects.

5. Injected DFP is excreted in the urine as diisopropylphosphate (60 to 70 per cent in the first ten days); less than 5 per cent of the P^{32} injected as DFP is recovered in the feces during that period.

6. In one person DFP receptors were blocked by treatment with OMPA (octamethylpyrophosphor-

amide) prior to the injection of DFP³². In this case less radioactivity was bound to plasma and red cells and maximal radioactivity in the blood was reached as soon as 30 minutes after the injection of DFP. Excretion was slightly accelerated. These data are understandable on account of the relative decrease of firmly bound and increase of free DFP³² as a result of the OMPA pretreatment. The free DFP is more readily removed from the blood than the bound fraction.

7. It is suggested that labelling by means of DFP³² may also be used for the evaluation of red cell and plasma volume.

REFERENCES

1. Ammon, R., Die fermentative Spaltung des Acetylcholin. *Pflüger's Arch. f. d. ges. Physiol.*, 1933, **233**, 486.
2. Hanes, C. S., and Isherwood, F. A., Separation of the phosphoric esters on the filter paper chromatogram. *Nature*, 1949, **164**, 1107.
3. Jandorf, B. J., and McNamarra, P. D., Distribution of radiophosphorus in rabbit tissues after injection of phosphorus-labelled diisopropyl fluorophosphate. *J. Pharmacol. Exper. Therap.*, 1950, **98**, 77.
4. Mazur, A., An enzyme in animal tissues capable of hydrolyzing the phosphorus-fluorine bond of alkyl fluorophosphates. *J. Biol. Chem.*, 1946, **164**, 271.
5. Jansen, E. F., Nutting, M.-D. F., Jang, R., and Balls, A. K., Mode of inhibition of chymotrypsin by diisopropyl fluorophosphate. II. Introduction of isopropyl and elimination of fluorine as hydrogen fluoride. *J. Biol. Chem.*, 1950, **185**, 209.
6. Jansen, E. F., and Balls, A. K., The inhibition of β - and γ -chymotrypsin and trypsin by diisopropyl fluorophosphate. *J. Biol. Chem.*, 1952, **194**, 721.
7. London, I. M., Shemin, D., West, R., and Rittenberg, D., Heme synthesis and red blood cell dynamics in normal humans and in subjects with polycythemia vera, sickle-cell anemia, and pernicious anemia. *J. Biol. Chem.*, 1949, **179**, 463.
8. Dixon, F. J., Talmage, D. W., Maurer, P. H., and Deichmiller, M., The half-life of homologous gamma globulin (antibody) in several species. *J. Exper. Med.*, 1952, **96**, 313.
9. London, I. M., in *Symposia on nutrition of the Robert Gould Research Foundation, Inc.*, J. B. Youmans, ed., Springfield, Thomas, 1950, Volume II. Plasma Proteins. Studies on the rates of formation of serum proteins in man, p. 72.