

ACKNOWLEDGMENTS

We would like to thank Dr. Cecil A. Krakower for isolating glomeruli from one patient by screening and centrifugation. We would also like to thank Drs. R. M. Kark, D. S. Jackson and H. R. Catchpole for critical reviews of the manuscript.

REFERENCES

1. Allen, A. C. *The Kidney, Medical and Surgical Diseases*. New York, Grune and Stratton, 1951.
2. Jones, D. B. Inflammatory and vascular disease of the glomerulus *in* *Analytical Pathology. Treatises in the Perspective of Biology, Chemistry and Physics*, A. C. Mellors, Ed. New York, Blakiston, 1957.
3. Rinehart, J. F., Farquhar, M. G., Jung, H. C., and Abul-Haj, S. K. The normal glomerulus and its basic reactions in disease. *Amer. J. Path.* 1953, **29**, 21.
4. Hall, B. V. Further studies of the normal structure of the renal glomerulus *in* *Proceedings of the Sixth Annual Conference on the Nephrotic Syndrome*, J. Metcoff, Ed. New York, National Nephrosis Foundation, 1954, p. 1.
5. Pease, D. C. Fine structures of the kidney seen by electron microscopy. *J. Histochem. Cytochem.* 1955, **3**, 295.
6. Mueller, C. B., Mason, A. D., Jr., and Stout, D. G. *Anatomy of the glomerulus*. *Amer. J. Med.* 1955, **18**, 267.
7. Gross, J. Studies on the formation of collagen. I. Properties and fractionation of neutral salt extracts of normal guinea pig connective tissue. *J. exp. Med.* 1958, **107**, 247.
8. Gustavson, K. H. *The Chemistry and Reactivity of Collagen*. New York, Academic Press, 1956.
9. Goodman, M., Greenspon, S. A., and Krakower, C. A. The antigenic composition of the various anatomic structures of the canine kidney. *J. Immunol.* 1955, **75**, 96.
10. Bonting, S. L., Pollak, V. E., Muehrcke, R. C., and Kark, R. M. Quantitative histochemistry of the nephron. *Science* 1958, **127**, 1342.
11. Bonting, S. L., Pollak, V. E., Muehrcke, R. C. and Kark, R. M. Quantitative histochemistry of the nephron. II. Alkaline phosphatase activity in man and other species. *J. clin. Invest.* 1960, **39**, 1372.
12. Grunbaum, B. W., and Glick, D. Studies in histochemistry. XLV. Determination of hydroxyproline in microgram amounts of tissue. *Arch. Biochem.* 1956, **65**, 260.
13. Neumann, R. E., and Logan, M. A. The determination of hydroxyproline. *J. biol. Chem.* 1950, **184**, 299.
14. Vernier, R. L. Personal communication.
15. Pollak, V. E., Bonting, S. L., Muehrcke, R. C., and Kark, R. M. Quantitative histochemistry of the nephron. V. Alkaline phosphatase and lactic dehydrogenase activity in lupus nephritis. *J. clin. Invest.* 1960, **39**, 1394.
16. Greenspon, S. A., Bollinger, F. W., and Krakower, C. A. Some chemical properties of nephrotoxic antigen. *Fed. Proc.* 1952, **11**, 416.
17. Jackson, D. S. The formation and breakdown of connective tissue *in* *Connective Tissue*, R. E. Tunbridge, Ed. Oxford, Blackwells, 1957.
18. Gross, J. Studies on the fibrogenesis of collagen. Some properties of neutral extracts of connective tissue *in* *Connective Tissue*, R. E. Tunbridge, Ed. Oxford, Blackwells, 1957.

CORRECTION

On page 1597 of the paper entitled "Serum factors in lupus erythematosus and other diseases reacting with cell nuclei and nucleoprotein extracts: Electrophoretic, ultracentrifugal and chromatographic studies" by Howard C. Goodman, John L. Fahey and Richard A. Malmgren (*J. clin. Invest.* 1960, **39**, 1595), lines 15-18 should read: "Anti-calf thymus nucleoprotein extract activity was present in 85 per cent of SLE sera, 10 per cent of the RA sera and 25 per cent of the scleroderma sera." An error in typesetting omitted a line which confused the meaning of this sentence. Also, on page 1600 of the same article, the first sentence at the top of column 2 should read: "The proteins eluted first are seen to sediment less rapidly ($s_{20,w}^0 \cong 6.6S$) than those eluted in the 60 to 80 per cent effluent region ($s_{20,w}^0 \cong 18S$)."