

## Role of the sympathetic nervous system in the renal response to hemorrhage

John R. Gill Jr., Alfred G. T. Casper

*J Clin Invest.* 1969;48(5):915-922. <https://doi.org/10.1172/JCI106050>.

### Research Article

In 12 studies, a femoral artery and vein of a donor dog treated with desoxycorticosterone were connected by tubing to a renal artery and vein of a recipient dog treated with desoxycorticosterone, and the kidney with its nerve supply intact was perfused at femoral arterial pressure. Infusion of normal saline, which contained albumin, from 2.7 to 3.1 g/100 ml, in the donor produced significant natriuresis in a kidney of the donor (from 112 to 532  $\mu\text{Eq}/\text{min}$ ) and in the perfused kidney (from 60 to 301  $\mu\text{Eq}/\text{min}$ ) of the recipient. Increased sodium excretion in the perfused kidney was associated with an increase in the clearances of inulin and *para*-aminohippurate ( $P < 0.01$ ) and a decrease in hematocrit of perfusing blood ( $P < 0.01$ ).

Infusion was continued in the donor while recipient was bled 23 ml/kg, with a decrease in mean arterial pressure from 152 to 130 mm Hg. Sodium excretion in perfused kidney decreased from 301 to 142  $\mu\text{Eq}/\text{min}$  ( $P < 0.01$ ), whereas sodium excretion in donor was unchanged (506 VS. 532  $\mu\text{Eq}/\text{min}$ ;  $P > 0.3$ ). Clearance of inulin by perfused kidney was not significantly affected by bleeding ( $26 \pm \text{SE } 2$  VS.  $25 \pm \text{SE } 2$ ;  $P > 0.2$ ), but the clearance of *para*-aminohippurate was decreased by bleeding ( $P < 0.01$ ), so that filtration fraction increased.

As the perfused kidney of the [...]

**Find the latest version:**

<https://jci.me/106050/pdf>



# Role of the Sympathetic Nervous System in the Renal Response to Hemorrhage

JOHN R. GILL, JR., and ALFRED G. T. CASPER

*From the Endocrinology Branch, National Heart Institute, National Institutes of Health, Bethesda, Maryland 20014*

**ABSTRACT** In 12 studies, a femoral artery and vein of a donor dog treated with desoxycorticosterone were connected by tubing to a renal artery and vein of a recipient dog treated with desoxycorticosterone, and the kidney with its nerve supply intact was perfused at femoral arterial pressure. Infusion of normal saline, which contained albumin, from 2.7 to 3.1 g/100 ml, in the donor produced significant natriuresis in a kidney of the donor (from 112 to 532  $\mu\text{Eq}/\text{min}$ ) and in the perfused kidney (from 60 to 301  $\mu\text{Eq}/\text{min}$ ) of the recipient. Increased sodium excretion in the perfused kidney was associated with an increase in the clearances of inulin and *para*-aminohippurate ( $P < 0.01$ ) and a decrease in hematocrit of perfusing blood ( $P < 0.01$ ).

Infusion was continued in the donor while recipient was bled 23 ml/kg, with a decrease in mean arterial pressure from 152 to 130 mm Hg. Sodium excretion in perfused kidney decreased from 301 to 142  $\mu\text{Eq}/\text{min}$  ( $P < 0.01$ ), whereas sodium excretion in donor was unchanged (506 vs. 532  $\mu\text{Eq}/\text{min}$ ;  $P > 0.3$ ). Clearance of inulin by perfused kidney was not significantly affected by bleeding ( $26 \pm \text{SE } 2$  vs.  $25 \pm \text{SE } 2$ ;  $P > 0.2$ ), but the clearance of *para*-aminohippurate was decreased by bleeding ( $P < 0.01$ ), so that filtration fraction increased.

As the perfused kidney of the recipient dog continued to receive blood from the natriuretic donor dog when the recipient dog was bled, the decrease in sodium excretion that bleeding produced in the perfused kidney was presumably mediated by renal nerves. Thus, an increase in nervous stimuli to the kidney that is not sufficient to decrease glomerular filtration rate can increase the tubular reabsorption of sodium and thereby significantly decrease its excretion. This property of the sympathetic nervous system to affect tubular reabsorption of sodium suggests

that an increase in sympathetic activity may constitute an important mechanism for the renal conservation of sodium when intravascular volume is contracted by hemorrhage or other cause.

## INTRODUCTION

Moderate hemorrhage increases sympathetic activity, most likely as a result of the effects of a decrease in effective circulating blood volume on baroreceptors (1). The consequent constriction of arterioles and veins and augmentation of cardiac contraction tend to minimize the decrease in arterial pressure which would occur were it not for an increase in sympathetic activity (2).

If the increase in sympathetic activity does not prevent a decrease in arterial pressure, then a decrease in glomerular filtration occurs and this decrease correlates well with the decrease in arterial pressure (3).

The increase in sympathetic activity also appears to involve the kidneys and to contribute to the decreases in glomerular filtration rate and renal blood flow. Thus, when Dibenzyline, an alpha adrenergic blocking agent, was injected into a renal artery before bleeding, it partially prevented a decrease in glomerular filtration rate and renal blood flow ipsilaterally (4).

A decrease in arterial pressure and an increase in sympathetic activity together provide a potent stimulus to increase the activity of the renin-angiotensin system (5), and, in turn, the production of aldosterone (6, 7). An interaction of these several factors is the probable cause of the decreased renal excretion of sodium that hemorrhage produces.

The present studies were undertaken to evaluate the effect of the sympathetic nervous system per se on renal function in hemorrhage, and to extend the previous observations which suggest that the sympathetic nervous system is part of the mechanism for the renal conservation of sodium (8). In the design of such an experiment, it is necessary to avoid the effects of a decrease in

This work was presented in part at the 1st Annual Meeting of the American Society of Nephrology, October 1967, Los Angeles, Calif.

Received for publication 25 October 1968 and in revised form 27 December 1968.

arterial pressure, and, in so far as possible, to minimize the effects of changes in circulating angiotensin and aldosterone. To accomplish this objective, a donor dog treated with desoxycorticosterone was used to perfuse a kidney of a desoxycorticosterone-treated recipient dog without destruction of the nerve supply to the perfused kidney. Natriuresis was established in the donor dog and in the perfused kidney to permit better characterization of renal nerve discharge as a stimulus to decrease the renal excretion of sodium. The recipient dog was then bled to increase the sympathetic nerve discharge to the perfused kidney that was protected from the other changes that accompany bleeding.

## METHODS

For each study, two mongrel dogs weighing 17–28 kg were given a sodium intake of 50 mEq/day for 4 days and desoxycorticosterone, 5 mg i.m., the day before and morning of the study. After both dogs were anesthetized with Pentothal, hydrated with 400–800 ml of normal saline, and anticoagulated with heparin, a femoral artery and vein of the donor dog were connected by polyethylene tubing (PE 280 graduated to PE 400) to the left renal artery and vein of the recipient dog, whose kidney was thus perfused at femoral arterial pressure. Care was taken to preserve the nerve supply to the perfused kidney, except for four studies in which the nerve supply to the perfused kidney was interrupted at the time of cannulation of the renal vessels (experiments 16, 21, 22, and 39). Mean arterial pressure (MAP) was measured in both dogs from catheters in the aorta (9) and urine was collected through ureteral catheters from both kidneys of the recipient dog and one kidney of the donor.

Normal saline containing inulin, *para*-aminohippurate and pitressin, 0.02 U/ml, was infused into both dogs at 1 ml/min by a constant infusion pump. The experimental design is shown schematically in Fig. 1. Clearance measurements were started approximately 1 hr after the operative procedure; three control periods were obtained, then normal saline, 200–1000 ml, was rapidly infused into the donor dog, fol-




FIGURE 1 Schematic presentation of the experimental procedure. MAP = mean arterial pressure.

lowed by a continuous infusion of normal saline that contained human albumin, from 2.7 to 3.1 g/100 ml at 15 ml per min, for the remainder of the experiment. The total volume given to the donor dog during an experiment ranged from 2700 to 4000 ml. Blood loss in the recipient dog was replaced by normal saline to prevent an increase in sympathetic activity before experimental hemorrhage. After urine flow from the donor dog and from the perfused kidney of the recipient dog had reached maximal rates, the recipient dog was bled until there was an obvious decrease in urine flow from the perfused innervated kidney. The mean volume of blood shed approximated 2.3% of body weight (mean, 22.7 kg) except in the four experiments in which the perfused kidney was denervated and in these studies it was greater and approximated 3.2% of body weight (mean, 22.8 kg). Each study consisted of 11–16 clearance periods; all clearance periods were 20 min in duration. The clearances of inulin ( $C_{In}$ ) and *para*-aminohippurate ( $C_{PAH}$ ) were determined as previously described (9).

The values presented are the means of from two to four periods: *a*) immediately before infusion (control), *b*) during infusion when sodium excretion by the perfused kidney was maximal, *c*) during and immediately after hemorrhage (seven experiments plus the four experiments in which the perfused kidney was denervated) and 20–35 min after completion of hemorrhage (experiments 55, 56, 57, 62, and 64). Standard errors of the mean (SE) are presented for  $C_{In}$ . The significance of the data was determined by paired analysis.

## RESULTS

Fig. 2 shows the results of a study in which natriuresis in a donor dog was maintained by continued infusion while the recipient dog was bled. In the donor, during infusion, sodium excretion increased from 248 to 557  $\mu$ Eq/min with an increase in  $C_{PAH}$  and  $C_{In}$  but not in MAP. In the recipient dog, sodium excretion in the perfused kidney increased from 73 to 298  $\mu$ Eq/min with an increase in  $C_{PAH}$  but not in  $C_{In}$ . Sodium excretion in the contralateral kidney was not affected. When the recipient dog was bled, MAP decreased slightly from 145 to 140 mm Hg, and sodium excretion decreased from 143  $\mu$ Eq/min to 87  $\mu$ Eq/min, but  $C_{PAH}$  and  $C_{In}$  were unchanged. Sodium excretion in the perfused kidney decreased from 298 to 174  $\mu$ Eq/min without a change in  $C_{PAH}$  and  $C_{In}$ . In the donor dog sodium excretion, MAP, and  $C_{In}$  were essentially unchanged, whereas  $C_{PAH}$  increased from 141 to 153 ml/min.

The results of 12 such studies are presented in Table I. Infusion in the donor dog increased sodium excretion from a mean value of 112  $\mu$ Eq/min to 532  $\mu$ Eq/min with an increase in mean  $C_{PAH}$  from 91 ml/min to 125 ml/min ( $P < 0.01$ ), an increase in mean  $C_{In}$  from  $33 \pm SE 3$  to  $37 \pm SE 3$  ( $P < 0.05$ ), and an increase in mean MAP from 154 mm Hg to 169 mm Hg ( $P < 0.02$ ). Mean hematocrit in the donor dog decreased from 46% to 29% ( $P < 0.01$ ).

In the recipient dog, sodium excretion in the perfused kidney increased in response to infusion in the donor dog from a mean value of 60  $\mu$ Eq/min to 301  $\mu$ Eq/min




FIGURE 2 Effect of infusion in donor dog (*D*) and of infusion in donor dog together with bleeding of recipient dog (*R*) on *MAP* (mean arterial pressure), *C*<sub>PAH</sub> (clearance of *para*-aminohippurate), *C*<sub>IN</sub> (clearance of inulin) and *U*<sub>NaV</sub> (sodium excretion) in the donor dog and in the recipient dog. The solid dots under recipient dog denote the perfused kidney. Note that bleeding the recipient dog decreased sodium excretion but did not affect *C*<sub>IN</sub> in the perfused kidney, and did not change *U*<sub>NaV</sub> in donor dog.

( $P < 0.01$ ), with an increase in mean *C*<sub>PAH</sub> from 66 ml/min to 91 ml/min ( $P < 0.01$ ) and in mean *C*<sub>IN</sub> from  $23 \pm SE 2$  to  $26 \pm SE 2$  ( $P < 0.01$ ).

The recipient dog was then bled 23 ml/kg. The results of observations made during and immediately after hemorrhage were similar to the results obtained after a wait of 20–35 min from the end of hemorrhage (experiments 55–57, 62, 64). Mean *MAP* in the recipient dog decreased with bleeding from 152 to 130 mm Hg ( $P < 0.01$ ). Mean *C*<sub>PAH</sub> decreased from 80 ml/min to 69 ml/min ( $P < 0.05$ ), mean *C*<sub>IN</sub> decreased from  $35 \pm SE 3$  ml/min to  $28 \pm SE 4$  ml/min ( $P < 0.01$ ) and mean sodium excretion decreased from 133  $\mu$ Eq/min to 46  $\mu$ Eq/min ( $P < 0.01$ ). In the perfused kidney, mean sodium excretion decreased from 301  $\mu$ Eq/min to 142  $\mu$ Eq/min ( $P < 0.01$ ) with a decrease in mean *C*<sub>PAH</sub> from 91 ml/min to 81 ml/min ( $P < 0.01$ ), but without a change in mean *C*<sub>IN</sub> ( $26 \pm SE 2$  vs.  $25 \pm SE 2$  ml/min) ( $P > 0.2$ ). In the

donor dog, mean sodium excretion was sustained (532 vs. 506  $\mu$ Eq/min) as mean *C*<sub>PAH</sub>, mean *MAP*, and mean hematocrit decreased (the latter two variables, significantly,  $P < 0.01$ ), and *C*<sub>IN</sub> increased ( $P < 0.01$ ).

The results of four studies in which the perfused kidney lacked a normal nerve supply are presented in Table II. Infusion in the donor dog increased sodium excretion in the donor dog from a mean value of 22  $\mu$ Eq/min to 461  $\mu$ Eq/min and in the perfused kidney of the recipient dog from a mean value of 9  $\mu$ Eq/min to 258  $\mu$ Eq/min.

The recipient dog was then bled 32 ml/kg. Mean *MAP* in the recipient dog decreased with bleeding from 156 to 91 mm Hg. In the two studies in which they were determined, mean *C*<sub>PAH</sub> decreased from 70 ml/min to 28 ml/min and mean *C*<sub>IN</sub> decreased from 34 ml/min to 13 ml/min. Mean sodium excretion decreased from 61  $\mu$ Eq/min to 16  $\mu$ Eq/min. In the perfused kidney, mean sodium

TABLE I  
Effects of Infusion in a Donor Dog and of Hemorrhage in the Recipient Dog on Renal Function in the Two Dogs

Experiment	Regimen	Donor dog					Recipient dog									
		Right kidney					Perfused kidney					Contralateral kidney				
		UNaV	C <sub>IN</sub>	C <sub>PAH</sub>	MAP	Hct	UNaV	C <sub>IN</sub>	C <sub>PAH</sub>	MAP	Hct	UNaV	C <sub>IN</sub>	C <sub>PAH</sub>	MAP	
		$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{mm Hg}$	%	$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{mm Hg}$			$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{mm Hg}$			
55	C*	109	22 $\pm$ 0.5	59	33	24	—	—	—	—	7	—	—	—	—	
	I-D†	724	32 $\pm$ 0	56	21	268	13 $\pm$ 0	48	145	16 $\pm$ 0.5	145	16 $\pm$ 0.5	44	150	—	
	H-R§	756	34 $\pm$ 1.5	53	18	80	12 $\pm$ 0.5	48	52	16 $\pm$ 0.5	52	16 $\pm$ 0.5	48	152	—	
56	C	44	33 $\pm$ 2	62	39	56	24 $\pm$ 0.5	60	245	49 $\pm$ 1	245	49 $\pm$ 1	116	157	—	
	I-D	625	44 $\pm$ 3	111	25	343	29 $\pm$ 0.6	96	154	46 $\pm$ 1	154	46 $\pm$ 1	111	152	—	
	H-R	670	51 $\pm$ 2	102	20	130	28 $\pm$ 0.5	76	8	32 $\pm$ 2	8	32 $\pm$ 2	84	140	—	
57	C	7	45 $\pm$ 11	176	49	66	26 $\pm$ 4	80	224	33 $\pm$ 1	224	33 $\pm$ 1	80	170	—	
	I-D	140	39 $\pm$ 2	174	21	392	29 $\pm$ 1	120	356	39 $\pm$ 2	356	39 $\pm$ 2	80	158	—	
	H-R	259	42 $\pm$ 0.5	138	15	193	30 $\pm$ 4	119	86	35 $\pm$ 2	86	35 $\pm$ 2	75	125	—	
60	C	66	30 $\pm$ 2	73	45	5	26 $\pm$ 1	93	16	38 $\pm$ 1	16	38 $\pm$ 1	88	142	—	
	I-D	492	38 $\pm$ 4.5	129	25	180	30 $\pm$ 2.5	134	98	34 $\pm$ 1.5	98	34 $\pm$ 1.5	82	136	—	
	H-R	496	45 $\pm$ 1	132	20	73	28 $\pm$ 1	114	26	26 $\pm$ 0	26	26 $\pm$ 0	68	112	—	
62	C	124	50	104	44	6	—	—	56	—	56	—	—	138	—	
	I-D	678	50 $\pm$ 4	96	19	156	18 $\pm$ 3	50	137	42 $\pm$ 1.5	137	42 $\pm$ 1.5	76	138	—	
	H-R	763	50 $\pm$ 1.5	100	16	88	18 $\pm$ 0.5	56	16	26 $\pm$ 0.5	16	26 $\pm$ 0.5	68	120	—	
64	C	9	16 $\pm$ 2	48	43	143	18 $\pm$ 1.5	40	24	14 $\pm$ 2.5	24	14 $\pm$ 2.5	43	158	—	
	I-D	393	27 $\pm$ 2.7	111	20	700	22 $\pm$ 2.7	90	53	32 $\pm$ 7	53	32 $\pm$ 7	66	158	—	
	H-R	352	28 $\pm$ 2.5	70	13	304	22 $\pm$ 0.5	62	1	—	1	—	—	92	—	

C<sub>IN</sub> = mean clearance of inulin  $\pm$ SE; C<sub>PAH</sub> = mean clearance of p-aminohippurate; MAP = mean arterial pressure; Hct = hematocrit; U<sub>Na</sub>V = sodium excretion  
\* C = control

† I-D = infusion of normal saline which contained albumin, from 2.7 to 3.1 g/100 ml in the donor dog.

§ H-R = bleeding of recipient dog plus infusion in donor dog.

TABLE I—(Continued)

Experiment	Regimen	Donor dog				Recipient dog							
		Right kidney				Perfused kidney				Contralateral kidney			
		UNaV	C <sub>in</sub>	CPAH	MAP	Hct	UNaV	C <sub>in</sub>	CPAH	UNaV	C <sub>in</sub>	CPAH	MAP
		$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{mm Hg}$	%	$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{ml}/\text{min}$	$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{ml}/\text{min}$	$\text{mm Hg}$	
65	C	94	29 ± 1	93	48	72	17 ± 0.3	60	61	27 ± 1	78	173	
	I-D	352	31 ± 1.3	126	25	148	20 ± 0.3	107	109	25 ± 0.7	65	175	
	H-R	398	30 ± 0.5	111	23	82	19 ± 4	92	11	17 ± 2	38	168	
66	C	162	32 ± 2.5	72	52	94	24 ± 1	81	35	40 ± 3	124	178	
	I-D	650	24 ± 0.5	112	44	228	26 ± 2	100	30	28 ± 8.5	84	150	
	H-R	566	30 ± 1.5	113	28	106	24 ± 4	87	35	12 ± 9.5	48	85	
68	C	130	31 ± 3	71	50	94	25 ± 1	64	158	43 ± 1	117	145	
	I-D	656	41 ± 2	158	34	367	33 ± 4	107	93	45 ± 4	104	145	
	H-R	604	45 ± 1	164	28	198	33 ± 1	94	58	37 ± 0	90	120	
69	C	109	24	67	53	59	17	48	228	30	79	150	
	I-D	614	29 ± 2	143	34	238	30 ± 0.5	94	91	34 ± 0.5	78	155	
	H-R	462	34 ± 2.5	136	27	122	28 ± 0.5	88	34	23 ± 1	46	148	
71	C	248	48 ± 3	100	48	73	33 ± 3	71	140	30 ± 1.3	68	143	
	I-D	557	55 ± 4	141	36	298	35 ± 1.3	85	143	29 ± 2	59	145	
	H-R	531	54 ± 0.7	153	30	174	34 ± 0.7	92	87	32 ± 2	59	140	
74	C	236	35 ± 0.7	83	43	32	—	—	125	—	—	153	
	I-D	497	36 ± 1.5	142	40	294	22 ± 2	60	182	53 ± 7	112	156	
	H-R	318	40 ± 3.5	122	29	148	23 ± 4	44	140	52 ± 0.5	133	162	
Mean ± SEM	C	112	33 ± 3	91	46	60	23 ± 2	66	110	34 ± 4	88	154	
	I-D	532	37 ± 3	125	29	301	26 ± 2	91	133	35 ± 3	80	152	
	H-R	506	40 ± 2	116	22	142	25 ± 2	81	46	28 ± 4	69	130	
P	C vs. I-D	<0.01	<0.05	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	
	I-D vs. H-R	>0.3	<0.01	>0.05	<0.01	<0.01	>0.2	<0.01	<0.01	<0.01	<0.05	<0.01	

TABLE II  
Effect of Denervation of Perfused Kidney

Experiment	Regimen	Donor dog				Recipient dog							
		Right kidney				Perfused kidney			Contralateral kidney				
		U <sub>NAV</sub>	C <sub>IN</sub>	C <sub>PAH</sub>	MAP	Hct	U <sub>NAV</sub>	C <sub>IN</sub>	C <sub>PAH</sub>	U <sub>NAV</sub>	C <sub>IN</sub>	C <sub>PAH</sub>	MAP
	$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{mm Hg}$	%	$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$		$\mu\text{Eq}/\text{min}$	$\text{ml}/\text{min}$	$\text{mm Hg}$			
16	C*	0			133				14			133	
	I-D‡	344	20	52	145	376	36	49	109	39	81	132	
	H-R§	267	19	46	145	408	33	44	42	22	50	100	
21	C	27			162	4			20			183	
	I-D	370			185	156			55			180	
	H-R	466			178	157			6			82	
22	C	16			153	16			50			140	
	I-D	482			148	218			55			158	
	H-R	450			150	230			13			102	
39	C	46	24	46	165	50	5		92	34	54	150	
	I-D	648	32	67	156	34	282	40	79	26	29	60	152
	H-R	557	36	69	150	32	277	36	84	4	4	7	80
Mean	C	22			153		9		44			152	
	I-D	461	26	60	158		258	38	64	61	34	70	156
	H-R	435	28	58	156		268	34	64	16	13	28	91

Specimens were lost before determination of C<sub>IN</sub> and C<sub>PAH</sub> for experiments 21 and 22.

\* C = control

‡ I-D = infusion of normal saline which contained albumin, from 2.7 to 3.1 g/100 ml in the donor dog.

§ H-R = bleeding of recipient dog plus infusion in donor dog.

C<sub>IN</sub> = mean clearance of inulin; C<sub>PAH</sub> = mean clearance of *p*-aminohippurate; MAP = mean arterial pressure; Hct = hematocrit; U<sub>NAV</sub> = sodium excretion.

excretion, mean C<sub>IN</sub>, and mean C<sub>PAH</sub> showed little change (258 vs. 268  $\mu\text{Eq}/\text{min}$ , 38 vs. 34 ml/min, and 64 vs. 64 ml/min, respectively). In the donor dog mean sodium excretion, mean C<sub>IN</sub>, mean C<sub>PAH</sub>, and mean MAP were sustained (461 vs. 435  $\mu\text{Eq}/\text{min}$ , 26 vs. 28 ml/min, 60 vs. 58 ml/min, and 158 vs. 156 mm Hg, respectively).

## DISCUSSION

In 12 studies in which a kidney of a recipient dog with its nerve supply intact was perfused by blood from a donor dog at femoral arterial pressure, the recipient dog showed a significantly lower C<sub>PAH</sub> ( $P < 0.02$ ) and C<sub>IN</sub> ( $P < 0.02$ ) in the perfused kidney than in the contralateral kidney during the control observations (Table I). Operative manipulation of a kidney can produce as great a decrease in C<sub>PAH</sub> and C<sub>IN</sub> as that observed in the present studies, and is very likely the reason for the diminished function of the perfused kidney (4). The effect of manipulation on C<sub>PAH</sub> and C<sub>IN</sub> can be abolished by an injection of Dibenzylamine, an alpha adrenergic blocking agent, into the renal artery (4). When the nerve supply to the perfused kidney was interrupted, mean C<sub>PAH</sub> and C<sub>IN</sub> in the perfused kidney were com-

parable to those in the contralateral kidney (Table II). Renal denervation at the time of manipulation also appears to prevent alterations in the pattern of renal blood flow that accompany manipulation of the kidney (10). The effects of manipulation therefore probably result from increased renal nerve discharge with vasoconstriction.

Infusion of normal saline, which contained albumin, from 2.7 to 3.1 g/100 ml, in the donor dog produced a natriuresis in the donor dog (532  $\mu\text{Eq}/\text{min}$ ) and in the perfused (301  $\mu\text{Eq}/\text{min}$ ) but not in the contralateral kidney of the recipient dog (Table I). The natriuresis was accompanied by significant increases in MAP, C<sub>PAH</sub>, and C<sub>IN</sub>, and a significant decrease in hematocrit in the donor dog, and by significant increases in both C<sub>PAH</sub> and C<sub>IN</sub> in the perfused kidney.

Recent studies suggest that physical factors (increase in mean arterial pressure, decrease in hematocrit, and increase in renal blood flow) (11) and release of a "natriuretic substance" (12) are the principal determinants of the decrease in tubular reabsorption of sodium (13) and of the increased excretion of sodium that occurs in response to expansion of the intravascular vol-

ume with saline or plasma. All of these factors, as well as an increase in glomerular filtration rate, could have mediated the natriuretic response of the donor dog and of the perfused kidney of the recipient dog.

While the infusion was continued in the donor dog, the recipient dog was bled until there was an appreciable decrease in urine flow from the perfused kidney. The result was a 50% decrease in sodium excretion by the perfused kidney from 301 to 142  $\mu\text{Eq}/\text{min}$  ( $P < 0.01$ ), without a significant change in sodium excretion by the donor dog (532 vs. 506  $\mu\text{Eq}/\text{min}$  ( $P > 0.3$ ) (Fig. 2, Table I). The decrease in sodium excretion thus occurred despite continued exposure of the perfused kidney to natriuretic stimuli such as hemodilution (14) and, possibly, a circulating "natriuretic substance" (12), which would tend to oppose such a decrease.

Presumably, the sympathetic nerves of the recipient dog which supplied the perfused kidney mediated its response to a blood loss of 23 ml/kg with a decrease in MAP of 14% in the recipient dog. A blood loss of from 10 to 12 ml/kg is sufficient to increase renal resistance and, thus, is probably a sufficient stimulus to increase the rate of discharge of the renal nerves (15). When the sympathetic nerve supply from the recipient dog to the perfused kidney had been interrupted, however, the perfused kidney was unaffected by a blood loss of 32 ml/kg with a decrease in MAP of 42% in the recipient dog.

Whereas the effect of hemorrhage on blood flow was comparable in the innervated perfused and contralateral kidneys (a decrease of 10 ml/min, or 11%, and 11 ml/min, or 14%, respectively), the effect on glomerular filtration rate was considerably greater in the contralateral kidney (a decrease of 7 ml/min, or 20%, as opposed to 1 ml per min) (Table I). Previous studies on the renal response to hemorrhage indicate that the degree of decrease in glomerular filtration rate correlates well with the degree of decrease in arterial pressure (3). Therefore, one would expect the decrease in mean arterial pressure of 22 to 130 mm Hg in the recipient dog to have a greater effect on glomerular filtration rate than the decrease of 10 to 159 mm Hg in the donor dog (Table I). Normally, changes in mean arterial pressure of this magnitude have very little effect on renal blood flow or glomerular filtration rate when vascular tone is normal (16). Thus, the increased renal vascular tone associated with increased sympathetic activity probably accounts for the loss of the normal stability of glomerular filtration rate and its greater dependence upon arterial pressure following hemorrhage.

The decrease in sodium excretion in the perfused innervated kidney when the recipient dog was bled was associated with a significant decrease in mean  $C_{\text{PAH}}$  ( $P < 0.01$ ), but not in mean  $C_{\text{Tb}}$  ( $P > 0.2$ ) in the perfused kidney. These findings suggest that the change in sodium

excretion in the perfused kidney which accompanied hemorrhage in the recipient dog resulted from a change in the tubular reabsorption of sodium.

As suggested above, an increase in nervous stimuli to the perfused kidney best explains an increase in sodium reabsorption when it occurs in response to hemorrhage in a dog connected by renal nerves but not by blood supply to that kidney. In this case, the blood was supplied by a donor dog with stable sodium excretion.

An increase in sympathetic nerve discharge could increase the tubular reabsorption of sodium in several ways: first, a decrease in the volume of interstitial fluid around the proximal tubules as a result of an increase in oncotic pressure of the blood in the peritubular capillaries could indirectly lead to an increase in the tubular reabsorption of sodium (17). When the recipient dog was bled, filtration fraction increased from 0.29 to 0.31 in the perfused kidney and this change is compatible with an increase in peritubular capillary oncotic pressure. If, in addition, hemorrhage decreased peritubular capillary hydrostatic pressure, then this change would also tend to decrease the volume of interstitial fluid and to augment the effect of increased capillary oncotic pressure.

Second, an increase in the proportion of total renal glomerular filtrate that traverses juxtamedullary nephrons with long loops and, presumably, with a greater capacity for sodium reabsorption could lead to an increase in the tubular reabsorption of sodium (18). This hypothesis is based on the reports that hemorrhagic hypotension decreased blood flow through the superficial cortex but had little effect on blood flow through the juxtamedullary area (19, 20). More recent studies, however, indicate that hemorrhagic hypotension decreases cortical and medullary blood flow to the same extent and thus do not support the concept of a selective resistance of the juxtamedullary circulation to hemorrhage (21). Conceivably, glomerular filtration rate could increase in the juxtamedullary area relative to the superficial cortical area in response to hemorrhage despite a comparable decrease in blood flow in the two areas. It has recently been suggested that a decrease in the intake of sodium leads to a decrease in glomerular filtration rate in superficial cortical nephrons but an increase in glomerular filtration rate in the juxtamedullary nephrons (22).

Third, an increase in sodium reabsorption could result from a direct effect of the renal nerves or catechols on tubular transport processes. The extent to which any or all of these three mechanisms are operative remains to be established.

Thus, in the present study, a decrease in the volume of extracellular fluid produced by hemorrhage produced a sizable decrease in the renal excretion of sodium with-



out a decrease in glomerular filtration rate, presumably as a result of an increase in the tubular reabsorption of sodium mediated by an increase in sympathetic nervous system activity. In a previous study, a decrease in the volume of extracellular fluid produced by restriction of dietary sodium did not decrease the renal excretion of sodium in a normal fashion despite a decrease in glomerular filtration rate and mean arterial pressure, when sympathetic nervous system activity was blocked, presumably as a result of the inability of the sympathetic nervous system to increase the tubular reabsorption of sodium (8). Together, these two studies provide evidence that the sympathetic nervous system may play an important role in the normal renal conservation of sodium through its ability to increase the tubular reabsorption of sodium.

#### ACKNOWLEDGMENT

We are grateful for the invaluable technical assistance of John Tate, James Cox, Belle Ruskin, and Harvey J. McDonald.

#### REFERENCES

- Chien, S. 1967. Role of the sympathetic nervous system in hemorrhage. *Physiol. Rev.* **47**: 214.
- Wagner, H. N. 1957. The influence of autonomic vasoregulatory reflexes on the rate of sodium and water excretion in man. *J. Clin. Invest.* **36**: 1319.
- Coelho, J. B., and S. E. Bradley. 1964. Function of the nephron population during hemorrhagic hypotension in the dog, with special reference to the effects of osmotic diuresis. *J. Clin. Invest.* **43**: 386.
- Handley, C. A., and J. H. Moyer. 1954. Unilateral renal adrenergic blockade and the renal response to vasopressor agents and to hemorrhage. *J. Pharmacol. Exp. Ther.* **112**: 1.
- Vander, A. J. 1967. Control of renin release. *Physiol. Rev.* **47**: 359.
- Carpenter, C. C. J., J. O. Davis, and C. R. Ayers. 1961. Relation of renin, angiotensin II, and experimental renal hypertension to aldosterone secretion. *J. Clin. Invest.* **40**: 2026.
- Slater, J. D. H., B. H. Barbour, H. H. Henderson, A. G. T. Casper, and F. C. Bartter. 1963. Influence of the pituitary and the renin-angiotensin system on the secretion of aldosterone, cortisol and corticosterone. *J. Clin. Invest.* **42**: 1504.
- Gill, J. R. Jr., and F. C. Bartter. 1966. Adrenergic nervous system in sodium metabolism. II. Effects of guanethidine on the renal response to sodium deprivation in normal man. *N. Engl. J. Med.* **275**: 1466.
- Gill, J. R. Jr., A. A. Carr, L. E. Fleischman, A. G. T. Casper, and F. C. Bartter. 1967. Effects of pentolinium on sodium excretion in dogs with constriction of the vena cava. *Amer. J. Physiol.* **212**: 191.
- Rosen, S. M., B. P. Truniger, H. R. Kriek, J. E. Murray, and J. P. Merrill. 1967. Intrarenal distribution of blood flow in the transplanted dog kidney: effect of denervation and rejection. *J. Clin. Invest.* **46**: 1239.
- Martino, J. A., and L. E. Earley. 1967. Demonstration of a role of physical factors as determinants of the natriuretic response to volume expansion. *J. Clin. Invest.* **46**: 1963.
- Rector, F. C. Jr., M. Martinez-Maldonado, N. A. Kurtzman, J. C. Sellman, F. Oerther, and D. W. Seldin. 1968. Demonstration of a hormonal inhibitor of proximal tubular reabsorption during expansion of extracellular volume with isotonic saline. *J. Clin. Invest.* **47**: 761.
- Dirks, J. H., W. J. Cirksena, and R. W. Berliner. 1965. The effect of saline infusion on sodium reabsorption by the proximal tubule of the dog. *J. Clin. Invest.* **44**: 1160.
- Craig, G. M., I. H. Mills, G. W. Osbeldiston, and B. L. Wise. 1966. The effect of change in perfusion pressure and haematocrit in the perfused isolated kidney of the dog. *J. Physiol. (London)*. **186**: 113 P.
- Abel, F. L., and Q. R. Murphy. 1962. Mesenteric, renal and iliac vascular resistance in dogs after hemorrhage. *Amer. J. Physiol.* **202**: 978.
- Thurau, K., and K. Kramer. 1959. Weitere untersuchungen zur myogen Natur der Autoregulation des Nierenkreislaufes, Aufhebung der Autoregulation durch muskulotrope Substanzen und druckpassives Verhalten des Glomerulusfiltrates. *Pflueger Arch. Ges. Physiol.* **269**: 77.
- Lewy, J. E., and E. E. Windhager. 1968. Peritubular control of proximal tubular fluid reabsorption in the rat kidney. *Amer. J. Physiol.* **214**: 943.
- Barger, A. C. 1966. Renal hemodynamic factors in congestive heart failure. *Ann. N. Y. Acad. Sci.* **139**: 276.
- Selkurt, E. E., and M. J. Elpers. 1963. Influence of hemorrhagic shock on renal hemodynamics and osmolar clearance in the dog. *Amer. J. Physiol.* **205**: 147.
- Carriere, S., G. D. Thorburn, C. C. C. O'Morcloe, and A. C. Barger. 1966. Intrarenal distribution of blood flow in dogs during hemorrhagic hypotension. *Circ. Res.* **19**: 167.
- Aukland, K., and M. Wolgast. 1968. Effect of hemorrhage and retransfusion on intrarenal distribution of blood flow in dogs. *J. Clin. Invest.* **47**: 488.
- Horster, M., and K. Thurau. 1968. Micropuncture studies on the filtration rate of single superficial and juxtamedullary glomeruli in the rat kidney. *Pflueger Arch. Ges. Physiol.* **301**: 162.