

ERRATUM

In Volume 48, No. 8, August 1969
p. 1360, Table V, top column heads corrected
and middle column heads added
as shown below

TABLE V
Test of Significance for Mean Difference ($\bar{D}$) between Infusion Periods for Paired Comparisons in Individual Animals

Sequence	C*	IV	IC	R	Differences	C-IV	IV-IC	IC-R	C-IC
FFA	1.16	1.07	0.95	1.05	$\bar{D} \pm SD$	-0.089 ± 0.10	-0.12 ± 0.06	$+0.11 \pm 0.13$	-0.21 ± 0.10
Gly	0.33	0.30	0.25	0.27	t	3.37	8.51	3.04	7.95
Glu	76	76	74	70	P	<0.01	<0.001	0.01	<0.001
IRI	39	38	31	26					
Dose range, 0.50-0.663 mg/kg per min					Gly $\bar{D} \pm SD$	-0.028 ± 0.073	-0.050 ± 0.030	$+0.020 \pm 0.035$	-0.070 ± 0.060
Number of animals, 13					t	1.38	6.01	2.06	4.96
					P	>0.1	<0.001	>0.05	<0.001
					Glu $\bar{D} \pm SD$	-0.90 ± 4.9	-1.8 ± 3.9	-2.8 ± 5.0	-2.0 ± 5.7
					t	0.66	1.65	2.03	1.27
					P	>0.5	>0.1	>0.05	>0.2
					IRI $\bar{D} \pm SD$	-0.25 ± 6.6	-7.7 ± 11	-4.7 ± 9.9	-8.1 ± 12
					t	0.13	2.42	1.65	2.32
					P	>0.5	<0.05	>0.1	<0.05
Sequence	C*	IC	IV	R	Differences	C-IC	IC-IV	IV-R	C-IV
FFA	1.20	0.90	1.01	0.99	$\bar{D} \pm SD$	-0.30 ± 0.14	$+0.11 \pm 0.12$	$+0.032 \pm 0.10$	-0.19 ± 0.26
Gly	0.33	0.24	0.27	0.26	t	4.81	2.08	0.72	1.64
Glu	85	84	79	77	P	<0.01	>0.1	>0.5	>0.2
IRI	52	41	29	19					
Dose range, 0.50-0.64 mg/kg min					Gly $\bar{D} \pm SD$	-0.09 ± 0.06	$+0.03 \pm 0.04$	$+0.006 \pm 0.073$	-0.06 ± 0.09
Number of animals, 5					t	3.41	1.73	0.19	1.50
					P	0.025	>0.1	>0.5	>0.2
					Glu $\bar{D} \pm SD$	-0.80 ± 2.7	-5.0 ± 2.7	-2.6 ± 3.7	-5.8 ± 4.7
					t	0.66	4.17	1.57	2.76
					P	>0.5	<0.025	>0.2	>0.05
					IRI $\bar{D} \pm SD$	-10 ± 11	-12 ± 13	-10 ± 6	-22 ± 22
					t	2.19	2.06	3.78	2.32
					P	>0.05	>0.1	<0.025	>0.05

* C = saline control, three samples, 30 min; IV = intravenous infusion, four samples, 120 min; IC = intracarotid infusion four samples, 120 min; R = recovery period, two samples, 60 min.