Salicylate-Induced Increases in Free Triiodothyronine in Human Serum

EVIDENCE OF INHIBITION OF TRIIODOTHYRONINE BINDING TO THYROXINE-BINDING GLOBULIN AND THYROXINE-BINDING PREALBUMIN

P. R. LARSEN

From the Division of Endocrinology and Metabolism, Department of Medicine, School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania 15213

ABSTRACT Addition of sodium salicylate to human serum at concentrations often obtained during aspirin therapy causes 100–200% increases in free triiodothyronine (T_3) and free thyroxine (T_4) as estimated by ultrafiltration. The increase in free T_3 was unexpected since previous data had suggested that salicylate inhibits binding of T_4 only to thyroxine-binding prealbumin (TBPA) and that T_3 is not bound to this protein. Using ultrafiltration techniques, we demonstrated binding of T_3 to TBPA. The affinity constant for T_3 -TBPA binding appears to be slightly greater than that for albumin- T_3 binding. While salicylate inhibits the binding of T_3 (and T_4) to TBPA, it can be predicted that little change will be observed in the free T_3 (or free T_4) without inhibition of thyroid hormone binding to thyroxine-binding globulin (TBG). Using a competitive-binding protein displacement technique, it has been shown that sodium salicylate, like diphenylhydantoin (DPH), inhibits the binding of T_3 and T_4 to TBG. The magnitude of the increase in free T_3 and free T_4 induced by salicylates suggests that interference with TBG binding is its major effect.

Aspirin was administered orally to two normal subjects in quantities sufficient to obtain serum salicylate levels of 20–25 mg/100 ml. This resulted in a decrease of 20–30% in total serum T_3 and T_4 levels. This decrease in T_4 levels is similar in magnitude to that previously observed in subjects receiving DPH. Unlike what has

been observed with DPH treatment, therapeutic salicylate levels are associated with increases of 50–75% in the unbound fraction of both T_3 and T_4 which persist throughout an 8–10 day treatment period.

INTRODUCTION

The recent availability of techniques for measuring triiodothyronine (T_3)¹ in human serum has provoked new interest in this thyroid hormone (1–3). It is speculated that T_3 may provide as much as 50–60% of the body's thyroid hormone requirements (2). Because of the apparent importance of this hormone, we have been interested in the effects of various drugs which might interfere with binding of T_3 to the proteins in human serum leading to subsequent increases in the free form of this hormone.

Extensive prior investigations with thyroxine (T_4) have shown that the small, unbound fraction, free T_4 , is the most satisfactory peripheral index of the thyroid status (4). It is postulated that free T_4 is the only, or at least the most readily available, fraction of T_4 present in extracellular fluid that can be utilized by the cells. Similar theoretical arguments are applicable to free T_3 . Therefore, agents which interfere with the binding of T_3 to its binding proteins would lead to at least a temporary increase in free T_3 . This might be expected to increase the amount of T_3 available to peripheral tissues

This material was presented in part at the 53d Annual Meeting of the Endocrine Society, 24–26 June 1971 in San Francisco, Calif.

Received for publication 21 September 1971 and in revised form 3 December 1971.

¹Abbreviations used in this paper: DPH, diphenylhydantoin; T_3 , triiodothyronine; T_4 , thyroxine; TBG, thyroxine-binding globulin; TBPA, thyroxine-binding prealbumin; UFT₃ and UFT₄, ultrafiltrable fraction of tracer hormones in serum after labeling with T_3 and T_4 , respectively.

and perhaps, because of the rapid onset and high potency of T_3 , to cause hypermetabolism.

Previous studies have shown that salicylate and its congeners decrease the binding of T_4 to the binding proteins in human serum (5-7). These drugs were found to interfere specifically with the binding of T_4 to thyroxine-binding prealbumin (TBPA). This conclusion was largely based on the alterations in the distribution of labeled T_4 after paper electrophoresis of human serum. Since T_3 is not thought to be bound to TBPA, we anticipated that addition of salicylate to human serum would cause an increase in free T_4 with no change in free T_3 . Administration of this drug might, therefore, offer a way to compare the effects of changes in the free form of only one of the two thyroid hormones. However, preliminary studies showed that addition of sodium salicylate to human serum caused an increase in free T_3 that was nearly as great as the salicylate-induced increase in free T_4 .

Simultaneous experiments with serum enriched with T_4 showed that the increased free T_3 induced by salicylates could not be explained by displacement of T_4 from TBPA to TBG. The salicylate effect on free T_3 suggested either that this drug could interfere with TBG- T_3 interaction or that T_3 was, in fact, bound to TBPA, or, perhaps, both. This seemed inconsistent with the previously mentioned conclusions. The following studies were performed to clarify the mechanism of action of salicylate on the protein binding of the two thyroid hormones.

METHODS

Free triiodothyronine and free thyroxine. These values were estimated by determining the ultrafiltrable fraction (UFT₃ and UFT₄) of tracer hormones present after enrichment of human serum with labeled T_3 and T_4 by methods previously described (8). In order to obtain simultaneous measurements, T_3 -¹²⁵I, 0.05-0.20 μ g/100 ml, and T_4 -¹²⁵I, 1-3 μ g/100 ml serum were added. Both isotopes were obtained from Abbott Laboratories, North Chicago, Ill. The T_4 -¹²⁵I preparations used contained less than 0.3% T_3 -¹²⁵I as a contaminant as determined by paper chromatography in tertiary amyl alcohol-hexane-NH₄OH by methods described previously (9). Overnight dialysis of the 1:1 dilution of T_4 -¹²⁵I with serum removes about 20% of this contaminant as determined by appropriate studies with tracer T_3 . Therefore, the over-all contamination of T_4 -¹²⁵I with T_3 -¹²⁵I was less than 0.25%. Since the ratio UFT₃/UFT₄ in pooled serum is about 20:1 the T_3 contaminant in the tracer leads to an artifactual increment in UFT₄ of about 5% which is negligible for the purposes of these studies. Since the salicylate-induced increase in UFT₄ is greater than the increment in UFT₃, the percentage artifactual elevation in the UFT₄ due to contaminating T_3 in the presence of salicylates is less than under base line conditions.

Ultrafiltrate was obtained by centrifugation at 37°C with the pH in whole serum maintained at 7.4-7.6 by an atmosphere of 10% CO₂. The UFT₃ and UFT₄ were corrected for the yield of 89 and 93%, respectively, during the magnesium chloride precipitation step and are expressed as fractions.

Serum T_3 levels were measured in duplicate by a modification of a recently described T_3 immunoassay using unextracted serum and a specific anti- T_3 antibody (10).^{*} Normal values for this method in our laboratory are 1.1 ± 0.25 ng/ml (mean $\pm$ SD).

Thyroxine determinations were performed by the method of Murphy with minor modifications (11). The normal range in our laboratory is 4.5 - 12 μ g T_4 /100 ml serum. The values are corrected for the 80% yield in the ethanol extraction step. The values for absolute free T_3 and free T_4 were determined by multiplication of the ultrafiltrable fraction by the concentration of total hormone. In the sequential studies of the effects of aspirin administration, all determinations of each parameter were carried out simultaneously to minimize technical variations.

In the studies performed to demonstrate displacement of T_3 and T_4 from TBG, methods identical with those previously described for quantitation of T_3 using TBG were used (9). In this system, 4% human serum in glycine-acetate or barbitol buffer, pH 8.6, is enriched with tracer T_3 and T_4 . 1-ml aliquots are added to tubes containing various quantities of T_3 , T_4 , and Na salicylate. The tubes are then allowed to incubate overnight and the bound and free hormones are separated by adsorption of the free tracer hormones to dextran-coated charcoal. The per cent of the tracer remaining in the supernate is expressed as a fraction of the total counts for each isotope.

TBG-binding capacities were estimated by the reverse flow electrophoresis method of Elzinga, Carr, and Beierwaltes and TBPA by the method of Oppenheimer, Martinez, and Bernstein with minor modifications (12, 13).

All solutions were made using distilled, deionized water. Sodium salicylate was obtained from Fisher Scientific Co., Pittsburgh, Pa. Sodium thyroxine pentahydrate, sodium triiodothyronine, crystalline human serum albumin and 5,5-diphenylhydantoin were obtained from Mann Research Labs, Inc., Orangeburg, N. Y. Purified TBPA was obtained from Behring Diagnostics, Inc., Woodbury, N. Y. Statistical analyses were performed using standard methods (14). Tests involving humans were performed with the informed, written consent of the subjects, some of whom were hospitalized on the General Clinical Research Unit. Pooled human serum was obtained through the kindness of the Clinical Chemistry laboratory of the Presbyterian-University Hospital in Pittsburgh.

RESULTS

Effect of in vitro addition of sodium salicylate on UFT₃ and UFT₄. In vitro addition of increasing amounts of sodium salicylate to human serum results in progressive increases in both UFT₃ and UFT₄ (Fig. 1). At a concentration of 60 mg/100 ml there is an increase of 155 and 241% over control values respectively. In this system, addition of 60-80 μ g/100 ml of T_4 are required to obtain similar increases in UFT₃ and UFT₄. Therefore, the increase in UFT₃ cannot be attributed to displacement of a small amount of T_4 from TBPA to TBG with a subsequent displacement of T_3 . The concentrations of salicylate used are within the therapeutic (20-30

^{*} Larsen, P. R. 1971. The direct immunoassay of triiodothyronine in human serum. Submitted for publication.

FIGURE 1 Effect of in vitro addition of sodium salicylate or diphenylhydantoin to pooled human serum on the ultrafiltrable fraction of T_3 (UFT₃) and T_4 (UFT₄). The mean of quadruplicate determinations is given. The SEM is too small to be depicted on this scale being $< 3\%$ of the corresponding value in all cases.

mg/100 ml) or toxic (60, or greater, mg/100 ml) ranges observed in man.

For further comparison the effects of the addition of diphenylhydantoin (DPH) are also shown. This agent causes both displacement of T_4 and presumably T_3 from TBG (6). At therapeutic levels (10–20 $\mu\text{g/ml}$), DPH addition results in considerably less dramatic increases in UFT₃ and UFT₄ though the changes are statistically significant ($P < 0.05$ at 10 $\mu\text{g/ml}$, $P < 0.01$ at 20 and 40 $\mu\text{g/ml}$).

Effect of dilution of human serum containing salicylate on UFT₃ and UFT₄. It would be anticipated from the mass law equations that dilution of a weakly bound ligand would decrease its competition with one more firmly bound. For example, dilution of serum containing DPH results in a marked decrease in the apparent effect of this agent on T_4 binding in human serum (15). A similar phenomenon was observed with sodium salicylate (Fig. 2). In these studies, control and salicylate-enriched human sera were progressively diluted as indicated and UFT₃ and UFT₄ determinations made simultaneously. There is a progressive decrease in the apparent effect of salicylate with dilution which is significant for UFT₃ even at a 1:2 dilution. The increase in both UFT₃ and UFT₄, over twofold in the undiluted control specimen, is nearly eliminated at a 1:50 dilution of human serum. These studies underline the necessity for performing studies of weak binding inhibitors in whole serum. In addition, these data demonstrate that the effect of salicylates on the UFT₃ and UFT₄ is reversible.

Comparison of the effects of sodium salicylate and barbital on the binding of T_3 and T_4 in human serum and

to human serum albumin. Since the results of previous studies argued against an effect of salicylate either on interference with T_3 -TBG binding or T_3 -TBPA binding, we initially examined the effect of this agent on albumin- T_3 binding. Both salicylate and the thyroid hormones are known to be bound to this protein. In addition, the effects of barbital, another agent previously demonstrated to interfere with the binding of T_4 to TBPA, were compared with those of salicylate (7). For technical reasons, diluted serum or human serum albumin was used in these studies and sodium salicylate was added to a final concentration of 15 mg/100 ml. In the first entry of Table I, it is seen that both salicylate and barbital cause an increase in the UFT₄. The increase in UFT₄ after dilution of serum in barbital buffer is somewhat greater than that caused by salicylate; however, this may be only a matter of concentration differences. The inhibitory effect of salicylate and barbital on albumin- T_4 binding is considerably greater than that in serum as would be anticipated. The effect of salicylate and barbital on the binding of T_3 in human serum is similar to the effects of these agents on T_4 binding. Thus, both salicylate and barbital displace T_3 from its binding proteins. However, it is clear from the last entry in Table I that the effect of these agents on T_3 binding in whole serum cannot be explained by inhibition of T_3 binding to human serum albumin. The per cent increase in the UFT₃ caused by salicylate and barbital is less with purified human albumin than it is with whole serum. Thus, it

FIGURE 2 Effect of dilution of control and sodium salicylate-enriched samples of human serum in 0.1 M PO_4 buffer, pH 7.4, on the UFT₃ and UFT₄. The salicylate was added and then successive dilutions of this and the control serum were made. Ultrafiltrate was then collected simultaneously from the eight samples. The results are the mean of three experiments.

TABLE I
Effects of Sodium Salicylate and Barbitol on the Binding of
 T_4 and T_3 to Whole Human Serum and to Purified
Human Serum Albumin

	PO ₄ *	Salicylate	Barbitol†
T_4			
Serum			
UFT ₄ ($\times 10^{-4}$)§	2.3 $\pm$ 0.2	6.4 $\pm$ 0.2	8.7 $\pm$ 0.3
Increase, %	—	178	278
Albumin			
UFT ₄ ($\times 10^{-3}$)	0.9 $\pm$ 0.2	4.9 $\pm$ 0.2	7.0 $\pm$ 0.01
Increase, %	—	445	678
T_3			
Serum			
UFT ₃ ($\times 10^{-3}$)	3.0 $\pm$ 0.1	6.3 $\pm$ 0.3	11.2 $\pm$ 0.3
Increase, %	—	110	274
Albumin			
UFT ₃ ($\times 10^{-3}$)	8.7 $\pm$ 0.2	15.3 $\pm$ 0.3	19.4 $\pm$ 0.1
Increase, %	—	76	123

* Serum or human serum albumin (4% in 0.15 M NaCl) was diluted 1:5 in 0.1 M phosphate buffer, pH 7.8. When indicated, sodium salicylate was added to a final concentration of 15 mg/100 ml.

† Serum or human serum albumin (4% in 0.15 M NaCl) was diluted 1:5 in 0.08 M barbitol buffer, pH 7.8.

§ Ultrafiltrable tracer T_4 and T_3 are expressed as fractions of the total.

|| Mean $\pm$ SEM of quadruplicate samples.

was necessary to examine both the question of salicylate interference with T_3 -TBG binding as well as to determine whether or not T_3 was bound to TBPA.

Effects of salicylate on T_3 -TBPA binding. Using the ultrafiltration system and simultaneous labeling with T_3 -¹²⁵I and T_4 -¹²⁵I, we found the UFT₃ to be consistently in the range of 0.71 while the UFT₄ was approximately 0.04 in the presence of 1.3×10^{-6} M TBPA (Table II). While T_3 is bound to human TBPA as indicated by the UFT₃ of < 1.0 , the binding was considerably less strong than that of T_4 . Paper electrophoretic studies of the commercial preparation of TBPA after addition of tracer quantities of T_4 -¹²⁵I did not show any evidence of TBG contamination.

As a control for these studies, similar concentrations of human serum albumin were tested. The binding of T_3 to TBPA seems to be slightly greater than the binding of T_3 to human serum albumin as indicated by the higher UFT₃ in the presence of albumin. It is seen that both the UFT₃ and UFT₄ are increased by salicylate addition to either protein solution indicating this compound interferes with T_3 and T_4 binding to both proteins. The studies in this table were performed in gly-

cine-acetate buffer at pH 8.6. Similar results were obtained in PO₄ buffer at physiological pH, although the binding of T_3 to TBPA appeared somewhat lower in magnitude. The initial UFT₃ was 0.80 in this system.

While these data could explain the salicylate effect, further considerations suggested that this could only be a partial cause. First, while the affinity of T_3 for TBPA is only slightly greater than that of albumin, the concentration of albumin in human serum is at least 200-fold greater than that of TBPA (17). One would anticipate, therefore, at physiological protein concentrations, more T_3 would be bound to albumin than to TBPA. Yet, as previously demonstrated, the effects of salicylate on albumin- T_3 binding were considerably less than they were on the binding of T_3 in whole human serum. Secondly, the affinity of T_3 for TBG is thought to be on the order of 10^9 , much greater than the estimated affinity of T_3 for either albumin or TBPA (18). Only about $\frac{1}{3}$ of the binding sites of TBG are occupied by T_4 and T_3 at the concentrations present in normal human serum. Therefore, the small quantities of T_3 (and T_4) displaced from albumin and TBPA by salicylates could be accommodated by TBG with minimal changes in the free hormone fraction. It is, then, difficult to explain how significant elevations in the free T_3 can occur in the presence of salicylates without interference with TBG- T_3 interaction. For these reasons the effect of salicylate on the binding of T_3 and T_4 to TBG was examined.

Effects of sodium salicylate on T_3 and T_4 binding to TBG. The effects of salicylate on TBG binding were examined in an assay system similar to that used in the competitive-binding displacement assay used for T_3 determinations (9). This was less cumbersome than the ultrafiltration system, and provided similar qualitative information. Initially, whole human serum was diluted 1:25 in barbitol buffer to minimize TBPA and albumin binding of tracer T_3 and T_4 but similar results have been obtained in both glycine-acetate buffer, pH 8.6, and phosphate buffer, pH 7.4. After addition of various

TABLE II
Inhibition of T_3 and T_4 Binding to TBPA and Human Serum
Albumin by Sodium Salicylate

	Control		Salicylate (15 mg/100 ml)	
	UFT ₃ *	UFT ₄ *	UFT ₃	UFT ₄
TBPA (1.3×10^{-6} M)§	0.71 $\pm$ 0.01‡	0.040 $\pm$ 0.006	0.90 $\pm$ 0.02	0.14 $\pm$ 0.01
Albumin (1.2×10^{-6} M)	0.80 $\pm$ 0.03	0.36 $\pm$ 0.01	0.90 $\pm$ 0.03	0.57 $\pm$ 0.01

* Ultrafiltrable tracer T_3 and T_4 are expressed as fractions of the total.

‡ Mean $\pm$ SEM of quadruplicate determinations.

§ Calculated using molecular weight of 50,000 (16).

FIGURE 3 Displacement of T_4 - ^{125}I and T_3 - ^{131}I from the binding proteins in human serum diluted 1:25 in barbital buffer 0.08 M, pH 8.6. Bound and free tracer hormone was separated by adsorption of unbound hormone to dextran-coated charcoal. The mean of triplicate determinations is given, the SEM is <1.5% for all points. The concentration of added binding inhibitor is given as moles/liter.

amounts of T_3 , T_4 or sodium salicylate, 1-ml samples of the mixture were equilibrated overnight and then bound and free hormones were separated by addition of dextran-coated charcoal. As is seen in Fig. 3, both T_3 and T_4 displace the labeled iodothyronines from the binding proteins present in human serum. In addition, sodium salicylate readily displaces T_3 and, at higher concentrations, T_4 from these binding proteins.

While physicochemical considerations suggest that TBG is the only binding protein present in human serum with a sufficiently high affinity for T_3 and T_4 to act as a competitive binding protein, it was necessary to prove this conclusively. It has been demonstrated by Pensky and Marshall that TBG can be isolated from human serum using affinity chromatography with T_4 coupled to Sepharose (Pharmacia Fine Chemicals, Inc., Uppsala, Sweden) (19). Applying the same principle, T_4 -Sepharose, prepared as described by these authors, was added to human serum and incubated at 4°C overnight with stirring. Comparison of paper electrophoretic patterns

after addition of T_4 - ^{125}I in tracer amounts showed that TBG binding was almost eliminated in serum treated in this way (Fig. 4). If the small deflections in the TBG

FIGURE 4 Distribution of tracer T_4 in human serum before and after TBG adsorption with T_4 -Sepharose. Reverse flow electrophoresis of serum was performed in glycine-acetate buffer pH 8.6 after addition of tracer quantities of T_4 - ^{125}I (<2 μ g/100 ml).

area of the T_4 -Sephadex adsorbed sera are due to TBG-bound T_4 , by planimetry, it represents less than 8% of the original TBG-binding capacity.

TBPA-binding capacity in serum was unchanged by T_4 -Sephadex adsorption and the T_4 concentration increased only 2 $\mu\text{g}/100\text{ ml}$. The latter occurred due to the fact that even with repeated attempts to purify the T_4 -Sephadex, small quantities of presumably noncovalently bound T_4 adhere to the material and are subsequently bound to the serum proteins. It is obvious that enrichment of serum with large quantities of T_4 , such as might occur with unwashed T_4 -Sephadex preparations, could lead to a similar decrease in the percentage of tracer bound to TBG that is shown in Fig. 4 (though one would expect to see much greater binding in the albumin area under these conditions).

Since the same serum could be compared before and after TBG elimination, its role in this assay system could be clarified. Table III shows comparison of T_3 and T_4 binding of these sera in both glycine-acetate and barbital buffer in the same system described for Fig. 3. It is apparent that the binding of T_3 is inhibited by barbital. There is also slight inhibition of T_4 binding in barbital buffer. After removal of TBG, the per cent T_3 bound falls to a level of 13.3 in glycine acetate and 6.5 in barbital buffer demonstrating that TBG is the critical T_3 -binding protein in this system. While there is a significant decrease in the per cent T_4 bound after removal of TBG, it is only after inhibition of TBPA and albumin binding in barbital buffer that the importance of TBG- T_4 binding in this system is observed. The control data demonstrate that there is no significant difference in

TABLE III
Binding of T_3 - ^{125}I and T_4 - ^{125}I in Dilute Human Serum before and after Removal of Thyroxine Binding Globulin (TBG) as Assessed by Dextran-Coated Charcoal Separation of Bound and Free Tracer

Buffer*		Amount bound to serum proteins	
		T_3 - ^{125}I	T_4 - ^{125}I
Control	Glycine acetate	70.2 $\pm$ 0.7§	96.4 $\pm$ 0.6
	Barbital	43.3 $\pm$ 0.7	94.6 $\pm$ 0.3
After removal of TBG	Glycine acetate	13.3 $\pm$ 0.3	82.6 $\pm$ 0.6
	Barbital	6.5 $\pm$ 0.4	22.6 $\pm$ 0.7
Buffer control (No serum)	Glycine acetate	2.7 $\pm$ 0.3	3.1 $\pm$ 0.7
	Barbital	3.4 $\pm$ 0.3	3.7 $\pm$ 0.1

* pH 8.6.

† Mean $\pm$ SEM of triplicate determinations in two experiments.

FIGURE 5 UFT $_3$ and UFT $_4$ in a subject receiving chronic aspirin therapy. Values given are single determinations which were estimated simultaneously using the same tracer.

the two systems with respect to the binding of free tracer to the dextran-charcoal in the absence of serum. The data in Table III show that in this system (the same as that of Fig. 3), a maximum of about 6.5% of the tracer T_3 - ^{125}I is bound to TBPA and albumin and likewise about 22.6% of the tracer T_4 is bound to these two proteins under base line conditions. Therefore, any decrease in the percentage of tracer T_3 bound greater than 6.5% and of T_4 greater than 22.6% implies interference with TBG binding of these two hormones. Since sodium salicylate at $6 \times 10^{-3}\text{ M}$ causes a net displacement of about 41% of the T_3 - ^{125}I and at $6 \times 10^{-3}\text{ M}$ displaces about 52% of the T_4 - ^{125}I (Fig. 3), one may conclude that, in this system, salicylate inhibits the binding of T_3 and T_4 to TBG.

Effect of salicylate on T_3 and T_4 binding in vivo. While previous studies have shown decreases in the PBI in patients treated with salicylates, no reports of ultrafiltrable T_3 or T_4 have appeared (20). If salicylates interfere with TBG- T_4 and T_3 binding one would anticipate a decrease in the total T_3 and T_4 and increases in the UFT $_3$ and UFT $_4$. We have studied the effects of aspirin administration in two subjects under controlled conditions. Three base line determinations were obtained during a 6 day control period. Specimens during aspirin administration were obtained every other day over a subsequent 8 to 10 day period. There was an immediate and persistent increase in the UFT $_3$ and UFT $_4$ in both subjects. Fig. 5 shows the relationship of these changes to the salicylate levels in one of these. Maintenance of a salicylate concentration on the order of 20–25 mg/100 ml was associated with increases in the free fraction comparable to those obtained in vitro at this concentration. Unlike what has been observed during DPH therapy, this increase in the free fraction of both T_3 and T_4 appears to be maintained (8, 15).

TABLE IV
Effect of Aspirin Administration on Free Triiodothyronine (FT₃) and Free Thyroxine (FT₄)

	Subject T. L.						Subject R. R.					
	T ₃	UFT ₃	FT ₃	T ₄	UFT ₄	FT ₄	T ₃	UFT ₃	FT ₃	T ₄	UFT ₄	FT ₄
	ng/ml	×10 ⁻³	ng/100 ml	μg/100 ml	×10 ⁻⁴	ng/100 ml	ng/ml	×10 ⁻³	ng/100 ml	μg/100 ml	×10 ⁻⁴	ng/100 ml
Control period												
1	1.05	3.7	0.39	7.7	2.2	1.7	1.13	4.6	0.52	9.0	3.0	2.7
2	0.80	3.7	0.30	7.7	2.5	1.9	0.98	3.7	0.36	8.9	2.6	2.3
3	0.80	3.4	0.27	7.3	2.2	1.6	0.93	4.8	0.45	8.5	2.9	2.4
Mean ± SEM	0.88 ± 0.08	3.6 ± 0.1	0.32 ± 0.04	7.6 ± 0.1	2.3 ± 0.1	1.7 ± 0.1	1.01 ± 0.06	4.4 ± 0.3	0.44 ± 0.05	8.8 ± 0.2	2.8 ± 0.1	2.5 ± 0.1
Aspirin period												
4	0.70	6.0	0.42	5.9	3.9	2.2	0.85	9.8	0.83	6.3	5.3	3.3
5	0.70	4.7	0.33	5.6	3.4	1.9	0.70	5.9	0.41	6.5	4.4	2.9
6	0.62	5.7	0.35	4.4	3.5	1.5	0.76	7.8	0.59	7.0	5.3	3.7
7	0.78	5.3	0.41	5.8	3.2	1.9	0.58	6.4	0.37	—	4.4	—
Mean ± SEM	0.70 ± 0.03	5.4 ± 0.3	0.38 ± 0.02	5.4 ± 0.4	3.5 ± 0.1	1.9 ± 0.1	0.72 ± 0.06	7.5 ± 0.9	0.55 ± 0.10	6.6 ± 0.2	4.9 ± 0.3	3.3 ± 0.2
Treatment/Control, %	80	150	119	71	152	112	71	170	125	75	175	132
P*	<0.05	<0.01	NS	<0.01	<0.01	NS	<0.02	<0.05	NS	<0.01	<0.01	<0.05

* P determined by *t* test for unpaired samples.

In Table IV are shown the effects of aspirin administration on the absolute free T₃ and free T₄ in the two subjects. While it is difficult to know whether it is accurate to average the samples during the entire treatment period due to disequilibrium conditions, this has been done for the purposes of these initial studies. With regard to T₃ levels, there is a suggestion of a decrease in the total amount of T₃ to about 76% of control in both subjects. This compensates to some extent for the increased UFT₃ resulting in an average increase in free T₃ of only about 23% in these subjects. The decrease in T₄ is substantial, about 25–30%, similar to the effect of DPH therapy (8, 15, 21). Again this compensation results in only a slight increase in the absolute free T₄. These two studies were carried out for a period of about 8 to 10 days of salicylate administration. Longer periods of aspirin treatment will be necessary to determine whether this slightly elevated free hormone concentration persists under steady-state conditions.

DISCUSSION

The above data indicate that sodium salicylate interferes with T₃ and T₄ binding to TBG. This interference was first demonstrated in human serum by Osorio in 1962 (22). In these studies, paper electrophoresis was used but subsequent studies using this technique have emphasized only the salicylate effect on TBPA-T₄ binding (7).

Apparently, the effect of salicylates on the distribution of labeled hormones during paper electrophoresis is markedly greater on TBPA than it is on TBG. Thus, a different system is desirable to demonstrate the magnitude of the inhibition of TBG binding of T₃ and T₄. This could conceivably be done in paper electrophoretic studies with serum completely devoid of TBPA.

While surprising in light of current concepts, one could predict the interference of salicylates with T₃ and T₄ binding to TBG from theoretical considerations based on recently available data. Woeber and Ingbar have shown that complete elimination of TBPA by immuno-adsorption increases the free T₄ only an average of 22% (23). Yet at a concentration of 60 mg salicylate/100 ml, the UFT₄ in our system is increased 241%. Therefore, salicylate inhibition of TBPA binding is not playing a major role at this salicylate level. In addition, using assumptions based on the data of these and other investigators regarding T₄ distribution and current estimates of affinity constants for the various binding proteins, one can calculate the theoretical effect of complete elimination of TBPA and albumin binding using the mass action equation developed by Robbins and Rall (reference 4, see Appendix). Using this formula, one estimates that after complete inhibition of T₄ binding to both albumin and TBPA, there will be only a 30% increase in the free T₄. This emphasizes the importance of TBG-binding in the determination of the amount of T₄ that is free. Thus, at higher salicylate levels, interference with TBG-T₄ binding must be a factor and it is presumably present at lower concentrations to some extent as well. Since the affinity of T₃ for all the binding proteins is considerably less than that of T₄, T₃-TBG interaction will be inhibited at lower salicylate levels.

The per cent increase in free T₄ due to salicylate addition is moderately greater than that in free T₃ (241% vs. 155% at 60 mg/100 ml salicylate). However, the fraction of bound T₃ displaced by salicylate is greater than that of T₄. This would be anticipated from the data in Fig. 3 and is a result of the lower binding affinity of T₃ vs. T₄ for TBG. This can be quantitated by comparing the net increase in the UFT₃ in the presence of 60

mg/100 ml salicylate (approximately 6.7×10^{-8}) with the net increase in the UFT_4 (approximately 5.3×10^{-4}). The ratio of the two fractional increases is about 13:1, $UFT_3:UFT_4$. Nevertheless, the original fraction of T_3 free is about 20-fold greater than the fraction of T_4 free (4.3×10^{-8} vs. 2.2×10^{-4} , Fig. 1). Therefore, sodium salicylate does cause a disproportionate increase in UFT_4 compared with the base line relationships. This could be a result of salicylate interference with T_4 -TBPA binding which is probably more quantitatively significant than T_3 -TBPA binding. It is postulated that T_3 and T_4 are bound to the same site on TBG (18). In a simple system containing a single protein with one binding site, the free $T_4:T_3$ ratio should remain constant in the presence of a competitive binding inhibitor.

The at least temporary increases in free T_3 and free T_4 in human subjects during 1 wk of aspirin therapy is of considerable interest both clinically and theoretically. For many years the similarity of the effects of salicylate and thyroid hormones on oxygen consumption have been recognized (24). This similarity of action extends even to the demonstration in rats that salicylate will, at least temporarily, suppress TSH (25). If free hormone is an index of the quantity of thyroid hormone available to the cells, then elevation of free T_4 , and particularly free T_3 with its more rapid action and greater potency, may lead to symptoms of hyperthyroidism. While it is recognized that salicylates have T_4 -like effects on subcellular organelles, e.g. to uncouple oxidative phosphorylation in mitochondria, the possibility that increases in the utilization of T_3 and T_4 due to unbinding of these hormones from their binding proteins also occurs during salicylate therapy cannot be excluded (26). The most likely situation in which this could occur would be during salicylate intoxication in children where levels of 50-100 mg/100 ml salicylate are commonly seen. We are currently evaluating the possible significance of drug-induced hyperthyroidism in these patients. It is to be emphasized that at new steady-state concentrations of salicylate, T_3 , and T_4 , a return to pretreatment levels of absolute free T_3 and free T_4 concentrations would be anticipated. As pointed out previously under Results, the small increase in absolute free hormone over the 1st wk of therapy may only be temporary. Only if salicylates affect some other, perhaps cellular, binding site, could a persistent increase in free hormone occur. If only extracellular binding proteins are affected, one would anticipate that the increases in UFT_3 and UFT_4 would be completely balanced by decreases in total T_3 and T_4 . There is evidence that this is occurring in these two subjects and that only more time would be required for complete equilibration.

The inability of previous investigators to demonstrate T_3 binding to TBPA is not surprising in that studies

of this hormone using various electrophoretic techniques are complicated by the relatively low affinity of T_3 for the binding protein compared with its affinity for the supporting media. Purified TBPA has not been studied for T_3 binding in ultrafiltration or equilibrium dialysis systems. It would appear from these preliminary data that the binding of T_3 to TBPA is slightly stronger than that of T_3 to albumin. The affinity constant for T_3 -albumin interactions is on the order of 3×10^5 (18). However, the approximately 200-fold concentration excess of albumin relative to TBPA would suggest that more T_3 would be transported bound to this protein than to TBPA. Further modifications in electrophoretic techniques will be required to evaluate this question. However, in the patient with TBG deficiency, TBPA should be an important T_3 carrier.

While there are various other theoretical aspects of these data, we have recently been able to apply these observations to a practical problem. In the direct immunoassay of T_3 in human serum, there is considerable competition for T_3 between the antibody and the TBG present in human serum. While this is more important in systems in which bound and free tracer T_3 are separated by adsorption of the free T_3 to various agents, the presence of TBG is also theoretically disadvantageous in the double-antibody system. We have found that salicylate will block the binding of 90 to 95% of T_3 to TBG and this agent currently is being used in our direct immunoassay for T_3 (10). It also has the advantage of very low cross-reactivity with the T_3 antibody, is readily available and water soluble.

In summary, it would appear that, as might have been predicted from theoretical considerations, salicylate interferes with the binding of T_3 and T_4 to both TBPA and TBG. The increase in available free T_3 and free T_4 during the initial phases of aspirin administration may contribute to hypermetabolism associated with the administration of this agent. High concentrations of sodium salicylate can also be used to block the binding of T_3 to TBG during the direct immunoassay of T_3 in human serum.

APPENDIX

The expression describing the concentration of free T_4 originally formulated by Robbins and Rall is as follows (4):

$$(T_4) = \frac{(T_4 \cdot TBG) + (T_4 \cdot TBPA) + (T_4 \cdot ALB)}{[K_{TBG}(TBG)] + [K_{TBPA}(TBPA)] + [K_{ALB}(ALB)]}$$

where (T_4) = free T_4 , K_{TBG} , K_{TBPA} , K_{ALB} are the binding affinity constants for the respective proteins, (TBG), (TBPA), and (ALB) are the concentrations of unoccupied binding sites on the three binding proteins and $T_4 \cdot TBG$, $T_4 \cdot TBPA$ and $T_4 \cdot ALB$ are the concentrations of T_4 bound to the respective binding proteins. All concentrations are moles/liter. The following affinity constants, binding capacities, and distribution of T_4 are assumptions based on previously published data

using a T_4 concentration of $7.7 \mu\text{g}/100 \text{ ml}$ (16, 18, 23). In addition, it is assumed that there is one primary binding site/mole of binding protein.

	K	T_4 binding capacity	T_4 bound
		moles/liter	moles/liter
TBG	1.6×10^{10}	26×10^{-8}	7.5×10^{-8}
TBPA	1.3×10^7	320×10^{-8}	1.5×10^{-8}
ALB	6.2×10^5	5.7×10^{-4}	1.0×10^{-8}
Total $T_4 = 10 \times 10^{-8}$			

Substitution in the above equation gives the following:

$$(T_4) = \frac{10 \times 10^{-8}}{[(1.6 \times 10^{10})(18.5 \times 10^{-8})] + [(1.3 \times 10^7)(318 \times 10^{-8})] + [(6.2 \times 10^5)(5.7 \times 10^{-4})]}$$

and

$$(T_4) = 3.0 \times 10^{-11}$$

and

$$\text{UFT}_4 = \frac{(T_4)}{\text{Total } T_4} = \frac{3.0 \times 10^{-11}}{10 \times 10^{-8}} = 3.0 \times 10^{-4}.$$

After complete inhibition of T_4 binding to TBPA and albumin, the following conditions obtain in vitro:

$$(T_4) = \frac{10 \times 10^{-8} \text{ M}}{(1.6 \times 10^{10})(16 \times 10^{-8})}$$

and

$$(T_4) = 3.9 \times 10^{-11}$$

and

$$\text{UFT}_4 = \frac{3.9 \times 10^{-11}}{10 \times 10^{-8}} = 3.9 \times 10^{-4}.$$

Therefore, the relative increase in UFT_4 predicted based on current estimates of the binding affinities and protein concentrations can be calculated:

$$\frac{3.9 \times 10^{-4}}{3.0 \times 10^{-4}} \times 100 = 130\% \text{ of original value.}$$

ACKNOWLEDGMENTS

The author would like to express his gratitude to Mrs. Darina Sipula, Miss Jitka Dockalova, and Miss Regina Oswalt for their expert technical assistance.

This work was supported by NIH Grant No. AM14283 from the National Institute of Arthritis and Metabolic Diseases, Grant M10 from the Health Research Services Foundation of Pittsburgh, The Dreyfus Charitable Fund, and Grant FR56, General Clinical Research Center Grant from the National Institutes of Health.

REFERENCES

- Nauman, J. A., A. Nauman, and S. C. Werner. 1967. Total and free triiodothyronine in human serum. *J. Clin. Invest.* **46**: 1346.
- Sterling, K., D. Bellabarba, E. S. Newman, and M. A. Brenner. 1969. Determination of triiodothyronine concentration in human serum. *J. Clin. Invest.* **48**: 1150.
- Mitsuma, T., M. Gershengorn, J. Colucci, and C. S. Hollander. 1971. Radioimmunoassay of triiodothyronine in unextracted human serum. *J. Clin. Endocrinol. Metab.* **33**: 364.
- Robbins, J., and J. E. Rall. 1960. Proteins associated with the thyroid hormones. *Physiol. Rev.* **40**: 415.
- Christensen, L. K. 1959. Thyroxine-releasing effect of salicylate and of 2,4 dinitrophenol. *Nature*. **183**: 1189.
- Wolff, J., M. E. Standaert, and J. E. Rall. 1961. Thyroxine displacement from serum proteins and depression of serum protein-bound iodine by certain drugs. *J. Clin. Invest.* **40**: 1373.
- Ingbar, S. H. 1963. Observations concerning the binding of thyroid hormones by human serum prealbumin. *J. Clin. Invest.* **42**: 143.
- Larsen, P. R., A. J. Atkinson, Jr., H. N. Wellman, and R. E. Goldsmith. 1970. The effect of diphenylhydantoin on thyroxine metabolism in man. *J. Clin. Invest.* **49**: 1266.
- Larsen, P. R. 1971. Technical aspects of the estimation of triiodothyronine in human serum: evidence of conversion of thyroxine to triiodothyronine during assay. *Metabolism*. **20**: 609.
- Larsen, P. R. 1971. Inhibition of triiodothyronine (T_3) binding to thyroxine-binding globulin by sodium salicylate and its application to the immunoassay of T_3 in human serum. *Metabolism*. **20**: 976.
- Murphy, B. P. 1965. The determination of thyroxine by competitive protein-binding analysis employing an anion exchange resin and radiothyroxine. *J. Lab. Clin. Med.* **66**: 161.
- Elzinga, K. E., and E. A. Carr, Jr., and W. H. Beierwaltes. 1961. Adaptation of the standard durrum type cell for reverse-flow electrophoresis. *Amer. J. Clin. Pathol.* **36**: 125.
- Oppenheimer, J. H., M. Martinez, and G. Bernstein. 1966. Determination of maximal binding capacity in protein concentration of thyroxine-binding prealbumin in human serum. *J. Lab. Clin. Med.* **67**: 500.
- Snedecor, G., and W. G. Cochran. 1967. Statistical Methods. Iowa State University Press, Ames, Iowa. 6th edition.
- Chin, W., and G. C. Schussler. 1968. Decreased serum free thyroxine concentration in patients treated with diphenylhydantoin. *J. Clin. Endocrinol. Metab.* **28**: 181.
- Raz, A., and D. S. Goodman. 1969. The interaction of thyroxine with human plasma prealbumin and with the prealbumin-retinol-binding protein complex. *J. Biol. Chem.* **244**: 3230.
- Oppenheimer, J. H., and M. I. Surks, J. C. Smith, and R. Squef. 1965. Isolation and characterization of human thyroxine-binding prealbumin. *J. Biol. Chem.* **240**: 173.
- Robbins, J., and J. E. Rall. 1967. The iodine-containing hormones. In *Hormones in Blood*. C. H. Gray and A. L. Bacharach, editors. Academic Press Inc., Ltd., London. 2nd edition. 1: 383.
- Pensky, J., and J. S. Marshall. 1969. Studies on thyroxine-binding globulin (TBG) II. Separation from human serum by affinity chromatography. *Arch. Biochem. Biophys.* **135**: 304.
- Austen, F. K., M. E. Rubini, W. H. Meroney, and J. Wolff. 1958. Salicylates and thyroid function. I. Depression of thyroid function. *J. Clin. Invest.* **37**: 131.
- Oppenheimer, J. H., L. V. Fisher, K. M. Nelson, and J. W. Jailer. 1961. Depression of the serum pro-

- tein-bound iodine level by diphenylhydantoin. *J. Clin. Endocrinol. Metab.* 21: 252.
22. Osorio, C. 1962. Effect of salicylate and dinitrophenol on the binding of thyroid hormones by human and rat serum proteins at pH 7.4. *J. Physiol. (London)*. 163: 151.
 23. Woeber, A., and S. H. Ingbar. 1968. A contribution of thyroxine binding prealbumin to the binding of thyroxine in human serum as assessed by immunoadsorption. *J. Clin. Invest.* 47: 1710.
 24. Denis, W., and J. H. Means. 1916. The influence of salicylate on metabolism in man. *J. Pharmacol. Exper. Ther.* 8: 273.
 25. Good, B. F., and B. S. Hetzel, and B. M. Hogg. 1965. Studies of the control of thyroid function in rats: effects of salicylate and related drugs. *Endocrinology*. 77: 674.
 26. Brody, T. M. 1956. Action of sodium salicylate and related compounds on tissue metabolism in vitro. *J. Pharmacol. Exp. Ther.* 117: 39.