Relationship between Phosphaturia and Acute Hypercapnia in the Rat

R. KENT WEBB, PHILIP B. WOODHALL, C. CRAIG TISHER, GEORGE GLAUBIGER, FRANK A. NEELON, and ROSCOE R. ROBINSON, *The Divisions of Nephrology and Endocrinology, Department of Medicine, Duke University Medical Center, Durham, North Carolina 27710*

ABSTRACT Standard clearance studies were performed in mechanically ventilated intact and acutely thyroparathyroidectomized (TPTX) rats to document and characterize the effect of hypercapnia (HC) on urinary phosphorus excretion (U_PV). HC as compared to normocapnia (NC) was associated with an increase in U_PV in intact (62.5 vs. 7.93 $\mu\text{g}/\text{min}$) and TPTX (30.5 vs. 0.59 $\mu\text{g}/\text{min}$) rats, an increase in filtered load of phosphorus in intact (218 vs. 191 $\mu\text{g}/\text{min}$) and TPTX (243 vs. 146 $\mu\text{g}/\text{min}$) rats, an increase in blood bicarbonate concentration in intact (27.8 vs. 26.0 meq/liter) and TPTX (24.5 vs. 22.3 meq/liter) animals, and a decrease in blood pH in intact (7.15 vs. 7.42) and TPTX (7.07 vs. 7.39) rats. Additional TPTX rats with NC and HC were studied during phosphorus infusion at a comparable filtered load of phosphorus (NC = 307 $\mu\text{g}/\text{min}$ and HC = 328 $\mu\text{g}/\text{min}$). U_PV was 18.5 $\mu\text{g}/\text{min}$ in NC and 85.2 $\mu\text{g}/\text{min}$ in HC animals. Intact NC animals infused with NaHCO_3 achieved a blood bicarbonate of 45.9 meq/liter compared to 26.0 meq/liter in intact NC NaCl-infused rats. U_PV was 10.0 $\mu\text{g}/\text{min}$ in the NaHCO_3 and 7.93 $\mu\text{g}/\text{min}$ in NaCl-infused animals. In intact HC animals infused with NaHCO_3 , blood pH was 7.36 compared to 7.42 in NC intact NaCl-infused animals. U_PV was 83.2 $\mu\text{g}/\text{min}$ in the HC bicarbonate-infused and 7.93 $\mu\text{g}/\text{min}$ in the NC NaCl-infused rats. These experiments demonstrate that elevated blood carbon dioxide tension per se increases U_PV . Increases in filtered load of phosphorus and blood bicarbonate which are associated with HC contribute to the phosphaturia as does parathyroid hormone. The phosphaturia is not dependent upon reduction of extracellular pH.

Presented in part before the Southern Society for Clinical Investigation, New Orleans, Louisiana, January 1976.

Reprint requests should be addressed to, C. Craig Tisher, M.D., Duke University Medical Center, Durham, North Carolina 27710.

Received for publication 21 February 1977 and in revised form 15 June 1977.

INTRODUCTION

Previous studies have demonstrated that acute hypercapnia in animals with intact parathyroid glands is associated with an increase in the urinary excretion of phosphorus (1-9). The exact mechanism responsible for this phosphaturia remains unknown. A rise in the plasma phosphorus has been noted by some investigators (1, 4, 10-12) and consequently the phosphaturia has been attributed to an increase in the filtered load of phosphorus (1, 4, 9). However, in those studies a possible contributory role by other major determinants of phosphorus excretion either was not evaluated or the studies were not designed in such a way as to permit an adequate evaluation of their potential contribution. Factors known to increase urine phosphorus excretion which might become operative during acute hypercapnia include an increase in the release of parathyroid hormone (PTH)¹ (13-15) an enhancement of PTH effect on the kidney (16-18), and extracellular fluid (ECF) volume expansion (19-21). Furthermore, changes in either the blood or urine pH (22-26) or an increase in the blood bicarbonate concentration or renal reabsorption of bicarbonate (23, 24, 27, 28) might also influence renal phosphate reabsorption.

The present experiments were designed to examine the nature of the phosphaturia that occurs in association with acute hypercapnia. The results indicate that

¹Abbreviations used in this paper: B, bicarbonate infused; B_{HCO_3} , blood bicarbonate concentration; B_{pCO_2} , blood carbon dioxide tension; P_{pH} , blood pH; ECF, extracellular fluid; EF_{Na} , excreted fraction of filtered sodium; EF_{P} , excreted fraction of filtered phosphorus; F_{P} , filtered load of phosphorus; GFR, glomerular filtration rate; HC, hypercapnia; NC, normocapnia; P, phosphorus infused; P_{Ca} , ionized plasma calcium; P_{CaT} , total plasma calcium; P_{P} , plasma phosphorus; PTH, parathyroid hormone; RBF, renal blood flow; T_{HCO_3} , tubular bicarbonate reabsorption; TPTX, thyroparathyroidectomized; U_{cAMPV} , urinary cyclic AMP excretion; U_{CaV} , urinary calcium excretion; U_{ClV} , urinary chloride excretion; U_{NaV} , urinary sodium excretion; U_{PV} , urinary phosphorus excretion; V, urine flow.

an increase in the filtered load of phosphorus (F_p), stimulation of endogenous PTH release or enhancement of its effect on the kidney, ECF volume expansion, an increase in blood or renal reabsorption of bicarbonate, and extracellular acidosis are not the major mechanisms responsible for the phosphaturia. Instead, a direct or indirect renal effect of the increase in blood carbon dioxide tension appears to be a likely determinant of the phosphaturia observed during acute hypercapnia.

METHODS

The studies were performed on 40 female Holtzman Sprague-Dawley rats weighing 200–260 g. Animals were fed a diet (Purina Lab Chow, Ralston Purina Co., St. Louis, Mo.) containing 1.20 g of calcium, 0.86 g of phosphorus, and 530 IU of vitamin D per 100 g, and were maintained on a time cycle of 12 h of light and 12 h of darkness. Experiments were always initiated on nonfasted rats between the hours of 8:00 and 11:00 a.m. Anesthesia was induced with intraperitoneal sodium pentobarbital (45 mg/kg body wt) as a solution of 15 mg in 1 ml of a NaCl solution (osmolality: 300 mosmol/kg H_2O) with administration of small additional doses as needed. Animals were placed on a thermostatically controlled warming table to maintain their body temperature between 37° and 38°C. Intubation was accomplished via a tracheostomy with a polyethylene (PE 240) endotracheal tube, and animals breathed room air supplemented with 100% O_2 during the subsequent surgery. Thyroparathyroidectomy, if required, was then performed utilizing blunt dissection. The right femoral artery and vein were then cannulated with polyethylene tubing (PE 50) through an inguinal incision. Mean arterial blood pressure was monitored continuously with a Statham pressure transducer (model no. P23Db, Statham Instruments, Div. Gould Inc., Oxnard, Calif.). All blood samples for chemical determinations were obtained from the arterial line. The bladder was cannulated with PE 100 tubing. A hypertonic NaCl solution (osmolality: 400 mosmol/kg H_2O) was given i.v. in small intermittent doses (total dose: 1 ml/100 g body wt) to replace estimated fluid losses during surgical preparation. Upon the completion of surgery, 0.5 ml of the hypertonic NaCl solution containing 1 μ Ci [carboxyl- ^{14}C]inulin (New England Nuclear, Boston, Mass.) and 2.5 μ Ci 3H -*p*-aminohippuric acid (New England Nuclear) was injected i.v. This was followed by a continuous

sustaining infusion of the same hypertonic NaCl solution administered at a rate calculated to deliver a volume equivalent to 4.5% of the body weight/h. The latter solution contained 0.032 μ Ci [carboxyl- ^{14}C]inulin/ml and 2.8 μ Ci 3H -*p*-aminohippuric acid/ml. In some experiments a sustaining infusion of different composition was utilized (see below). At this point muscle relaxation was induced with i.v. tubocurarine chloride (E. R. Squibb & Sons, Princeton, N. J.) at an initial dose of 0.12 mg, and the animal was placed on a mechanical respirator (small animal respirator, model no. 680, Harvard Apparatus Co., Inc., Millis, Mass.). Additional doses of tubocurarine chloride were given as needed.

After completion of all surgery and placement of the animal on assisted ventilation, the following experimental protocol was initiated. Each experiment included a 40-min period of equilibration followed by four 20-min clearance periods. In those animals subjected to thyroparathyroidectomy at least 2 h elapsed from the completion of the surgical procedure to initiation of the first clearance period. At the beginning of the equilibration period, the sustaining infusion administered during surgery was altered as described below, depending on the experimental condition under study. Blood samples were drawn at the completion of surgery, at the midpoint of the equilibration and clearance periods, and at the end of the fourth clearance period. Urine was always collected under oil during the equilibration period and each of the four clearance periods. Only animals with a blood oxygen saturation of at least 90 mm Hg were studied as described below. Eight groups of animals were studied.

Healthy control animals (group NC [normocapnia]). Five animals were studied during the constant infusion of hypertonic NaCl alone (osmolality: 400 mosmol/kg H_2O). The inspired gas mixture was 30% O_2 and 70% N_2 , and the respirator was adjusted to maintain the blood carbon dioxide tension ($BpCO_2$) at approximately 40 mm Hg.

Acute hypercapnia (group HC). Five animals were studied in the same manner as NC animals except the inspired gas mixture was adjusted to contain 10% CO_2 , 30% O_2 , and 60% N_2 with the respirator adjusted to maintain the $BpCO_2$ at approximately 80 mm Hg.

Thyroparathyroidectomized animals (group TPTX). Five animals were studied in a manner identical to the NC animals except they underwent surgical thyroparathyroidectomy at least 2 h before the start of the clearance periods.

Thyroparathyroidectomy with acute hypercapnia (group HC-TPTX). Five animals were studied as in group HC except an acute thyroparathyroidectomy was performed as in the TPTX animals.

Bicarbonate-loaded animals (group B). Five rats were

TABLE I
Effects of HC

	B_{pCO_2}	B_{pH}	B_{HCO_3}	T_{HCO_3}	GFR	P_p	F_p	U_{pV}
	mm Hg		meq/ liter	μ eq/ min	ml/min	mg/ 100 ml	μ g/min	μ g/min
NC	41.2	7.42	26.0	80.7	3.08	6.19	191	7.93
$\pm$ SD (n = 5)	3.0		1.3	21.7	0.74	1.01	51	6.01
HC	81.5	7.15	27.8	89.2	3.19	6.89	218	62.5
$\pm$ SD (n = 5)	3.5		2.6	22.3	0.58	0.97	41	9.7
P*	<0.001	<0.001	<0.02	>0.2	>0.6	<0.05	>0.05	<0.001

* NC vs. HC animals.

studied in a manner identical to NC animals except the rats received a constant infusion of NaHCO_3 (osmolality: 400 mosmol/kg H_2O) in place of the hypertonic NaCl solution.

Compensated metabolic alkalosis (group HC-B). Five animals were studied in a manner similar to those in group B (sodium bicarbonate infusion) except the inspired gas mixture was 10% CO_2 , 30% O_2 , and 60% N_2 .

Phosphate-loaded thyroparathyroidectomized animals (group P-TPTX). Five animals were studied in a manner identical to the TPTX animals except the infusate contained sodium phosphate (40 mg/100 ml) and the osmolality was adjusted to 400 mosmol/kg H_2O by the addition of NaCl . The pH of the solution was 7.4 with the molar ratio of $\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$ being 4:1.

Phosphate-loaded thyroparathyroidectomized animals with acute hypercapnia (group HC-P-TPTX). Five animals were studied in a manner identical to the P-TPTX animals (sodium phosphate infusion) except the inspired gas mixture was 10% CO_2 , 30% O_2 , and 60% N_2 .

Glomerular filtration rate (GFR) was determined by inulin clearance which was calculated by dividing urinary excretion rate of ^{14}C -radioactivity by plasma ^{14}C -radioactivity. Renal plasma flow (RPF) was determined by *p*-aminohippuric acid clearance which was calculated by dividing the urinary excretion rate of ^3H -radioactivity by plasma ^3H -radioactivity. Renal blood flow (RBF) was calculated from the relationship: $\text{RBF} = \text{RPF}/1\text{-hematocrit}$. The radioactivity in the blood and urine specimens was counted by liquid scintillation spectrometry (Isocap/300, Searle Analytic Inc., Des Plaines, Ill.).

Concentration of sodium and potassium in plasma and urine were determined by flame photometry (model 143, Instrumentation Laboratory, Inc., Lexington, Mass.). Plasma and urine osmolalities were determined with a vapor pressure osmometer (model 5130, Wescor Inc., Logan, Utah). Total calcium concentration in urine and plasma was determined with a fluorometric titrator (Calchette, Precision Systems, Inc., Sudbury, Mass.). Ionized calcium in plasma was measured on blood specimens drawn and centrifuged anaerobically with a flow-through calcium electrode (model 99-20, Orion Research Inc., Cambridge, Mass.). Plasma and urine phosphorus concentrations were measured with a Pierce Phosphorus Auto/Stat TM Kit (Pierce Chemical Co., Rockford, Ill.).

Urine pH and pCO_2 and blood pH, pCO_2 , and pO_2 determinations were performed with a pH blood gas analyzer (model 213, Instrumentation Laboratory). Plasma and urine bicarbonate concentrations were calculated from the pH and pCO_2 data by the Henderson-Hasselbalch equation. For calculation of blood bicarbonate, a pKa of 6.1 and a CO_2

solubility coefficient of 0.0301 were used. A CO_2 solubility coefficient of 0.0309 was used for the calculation of urine bicarbonate. The pKa used for urinary bicarbonate was derived from the relationship: $\text{pKa} = 6.33 - 0.5\sqrt{[\text{Na}^+ + \text{K}^+]}$ with Na^+ and K^+ representing the urinary concentration of these ions in Eq/l. Urinary cyclic AMP concentrations were measured by the Gilman competitive binding assay (29).

Statistical methods. Statistical comparisons were made utilizing Student's unpaired *t* test. All data listed in the tables represent the mean $\pm$ SD of 20 observations derived from five animals in each group studied.

RESULTS

The results of the studies in intact NC and HC animals are summarized in Table I. The presence of respiratory acidosis is obvious. Although the blood bicarbonate concentration (B_{HCO_3}) was slightly greater in HC animals, the mean values for the tubular reabsorption of bicarbonate (T_{HCO_3}) did not differ significantly between the two groups. Urine pH averaged 5.63 in NC and 5.78 in HC animals ($P > 0.05$). Plasma phosphorus concentration (P_p) was significantly greater in HC animals, but the filtered load of phosphorus, although greater in HC animals, was not statistically different from that measured in NC animals. A marked increase in absolute urinary phosphorus excretion (U_pV) and the excreted fraction of filtered phosphorus (EF_p) was observed in HC animals. GFR, total plasma calcium (P_{CaT}), ionized plasma calcium (P_{CaI}), urinary calcium excretion (U_{CaV}), and urinary cyclic AMP excretion (U_{cAMPV}) were not significantly different in the two groups of animals. Mean values for RBF, urine flow (*V*), urinary sodium excretion (U_{NaV}), the excreted fraction of filtered sodium (EF_{Na}), and urinary chloride excretion (U_{ClV}) were not significantly different in the two groups, thus indicating that the degree of ECF volume expansion was similar in HC and NC animals. However, urinary potassium excretion (U_{KV}) was greater in HC animals ($P < 0.02$) and there was a small but significant increase in the blood bicarbonate concentration. Thus, acute HC in intact animals produced

in Intact Rats

EF_p	P_{CaT}	P_{CaI}	U_{CaV}	U_{cAMPV}	RBF	<i>V</i>	U_{NaV}	EF_{Na}	U_{ClV}	U_{KV}
%	mg/ 100 ml	mg/ 100 ml	$\mu\text{g}/\text{min}$	pmol/min	ml/min	$\mu\text{l}/\text{min}$	$\mu\text{eq}/\text{min}$	%	$\mu\text{eq}/\text{min}$	$\mu\text{eq}/\text{min}$
4.1	8.12	4.29	7.40	259	12.3	96.2	20.7	4.62	25.4	3.15
2.7	0.44	0.33	3.92	133	2.1	42.1	8.4	1.70	9.6	0.43
29.7	8.52	4.30	7.64	239	12.7	82.0	20.2	4.37	24.2	3.54
7.3	0.78	0.50	2.34	110	2.3	27.1	5.8	0.87	7.3	0.56
<0.001	>0.05	>0.9	>0.8	>0.6	>0.5	>0.2	>0.8	>0.5	>0.6	<0.02

TABLE II
Effects of HC

	B _{PCO₂}	B _{pH}	B _{HCO₃}	T _{HCO₃}	GFR	P _p	F _p	U _{pV}
	mm Hg		meq/ liter	μeq/ min	ml/ min	mg/ 100 ml	μg/ min	μg/ min
NC-TPTX	37.9	7.39	22.3	60.0	2.68	5.45	146	0.59
±SD	2.7		1.4	10.2	0.38	0.67	29	0.45
(n = 5)								
HC-TPTX	86.2	7.07	24.5	68.8	2.80	8.73	243	30.51
±SD	5.4		1.9	11.2	0.34	0.98	29	6.43
(n = 5)								
P*	<0.001	<0.001	<0.001	<0.02	>0.2	<0.001	<0.001	<0.001

* NC-TPTX vs. HC-TPTX animals.

hyperphosphatemia and a marked increase in U_{pV}, largely in the absence of other significant alterations in renal function.

To assess the contribution of endogenous PTH to the phosphaturia associated with HC, NC-TPTX and HC-TPTX animals were studied (Table II). As was true of the intact animals, B_{HCO₃} was slightly but significantly greater in HC-TPTX animals. T_{HCO₃} was also slightly greater in HC-TPTX animals, undoubtedly reflecting the increase in filtered bicarbonate secondary to an increase in B_{HCO₃} and GFR. Mean urine pH was 5.52 in NC-TPTX and 5.65 in HC-TPTX animals, the difference just achieving statistical significance ($P < 0.05$). P_p, F_p, U_{pV}, and EF_p were all significantly greater in HC-TPTX animals, but there were no significant differences in mean P_{CaT}, P_{CaI}, and U_{CaV}. U_{CAMPV} was significantly greater in HC-TPTX rats. Mean values for RBF, V, U_{NaV}, EF_{Na}, and U_{ClV} were not significantly different in the two groups of animals. Mean U_{KV} was similar in the two groups ($P < 0.02$). These experiments documented the occurrence of hyperphosphatemia and phosphaturia during acute HC, even when the release of PTH had been prevented.

To assess the possible contribution of the increase

in F_p to the phosphaturia of HC, studies were performed in normocapnic phosphate-infused TPTX (NC-P-TPTX) and similarly prepared hypercapnic (HC-P-TPTX) animals (Table III). B_{HCO₃} and T_{HCO₃} were slightly, although not significantly greater in the HC-P-TPTX animals. Average urine pH was 5.73 in NC-P-TPTX and 5.85 in HC-P-TPTX animals ($P > 0.01$). P_p and F_p were not significantly different in the two groups, yet U_{pV} and EF_p were again significantly greater in the HC animals. P_{CaT} and U_{CaV} were not different in the two groups; however, P_{CaI} was slightly greater in the HC-P-TPTX animals, the values just achieving statistical significance. U_{CAMPV} was greater in HC-P-TPTX animals. Mean values for RBF, V, U_{NaV}, EF_{Na}, U_{ClV}, and U_{KV} were not significantly different between the two groups of animals. These studies demonstrated that the phosphaturia of acute HC does not occur merely as a consequence of an increase in F_p.

Next, studies were performed in intact bicarbonate-infused normocapnic (NC-B) and bicarbonate-infused hypercapnic (HC-B) animals to assess the possible contribution of the increased concentration of B_{HCO₃} to the phosphaturia (Table IV). The results are compared with those of experiments performed in intact NC animals. As expected, a marked metabolic alkalosis

TABLE III
Effects of HC

	B _{PCO₂}	B _{pH}	B _{HCO₃}	T _{HCO₃}	GFR	P _p	F _p	U _{pV}
	mm Hg		meq/ liter	μeq/ min	ml/ min	mg/ 100 ml	μg/ min	μg/ min
NC-P-TPTX	38.7	7.41	23.6	71.3	2.99	10.1	307	18.5
±SD	3.0		3.1	16.3	0.35	1.6	70	6.4
(n = 5)								
HC-P-TPTX	81.6	7.09	24.3	75.5	3.07	10.6	328	85.2
±SD	7.1		3.5	18.8	0.39	1.2	68	8.8
(n = 5)								
P*	<0.001	<0.001	>0.5	>0.4	>0.5	>0.2	>0.3	<0.001

* NC-P-TPTX vs. HC-P-TPTX animals.

in TPTX Rats

EF _P	P _{Ca_T}	P _{Ca_I}	U _{Ca} V	U _{CAMP} V	RBF	V	U _{Na} V	EF _{Na}	U _{Cl} V	U _K V
%	mg/ 100 ml	mg/ 100 ml	μg/ min	pmol/ min	ml/ min	μl/ min	μeq/ min	%	μeq/ min	μeq/ min
0.4	8.91	4.24	8.05	106	10.2	72.4	16.7	4.29	22.4	3.53
0.3	0.87	0.46	2.54	38	1.4	30.4	6.5	1.71	7.9	0.77
12.6	8.66	4.34	8.80	157	10.6	80.9	18.1	4.38	23.4	2.95
2.4	0.93	0.79	1.40	60	1.5	26.7	6.1	1.48	5.6	0.71
<0.001	>0.3	>0.8	>0.2	<0.01	>0.3	>0.3	>0.4	>0.8	>0.6	<0.02

was induced in the NC-B animals with bicarbonate infusion. Urine pH was 7.98 in NC-B and 5.63 in NC animals. The B_{pCO₂}, however, was not significantly different from that observed in NC animals. As a consequence of the increased B_{HCO₃}, the T_{HCO₃} was significantly greater in NC-B rats. P_P was significantly less in the alkalotic animals and as a result of this change and the slight decrease in GFR, F_P was also significantly less in NC-B as compared to NC animals. U_PV was not different between the two groups (NC vs. NC-B), but EF_P was slightly greater in NC-B rats. P_{Ca_T} was not different, but P_{Ca_I} and U_{Ca}V were decreased in the NC-B animals. U_{CAMP}V was markedly decreased in the animals with metabolic alkalosis. Mean values for RBF, V, and U_{Cl}V were significantly less in NC-B animals, whereas U_{Na}V, EF_{Na}, and U_KV were significantly greater ($P < 0.01$). Because EF_P was only slightly greater in the animals with marked increases in B_{HCO₃}, it seems unlikely that the increase in B_{HCO₃} contributed in a major way to the phosphaturia of HC.

To evaluate the possible relationship of extracellular pH or H⁺ concentration to the phosphaturia, the extracellular acidosis of HC was minimized by infusion of sodium bicarbonate (Table IV). The blood pH (B_{pH}) was

almost restored to levels comparable to those observed in NC animals, although the difference between the two groups was still statistically significant. B_{pCO₂}, B_{HCO₃}, and T_{HCO₃} were all significantly greater in HC-B animals. Urine pH was 7.84 in HC-B and 5.63 in NC animals. P_P, U_PV, and EF_P were increased in the HC-B animals; however, F_P was not significantly greater in the HC-B rats. P_{Ca_I} and U_{Ca}V were decreased but P_{Ca_T} was not different in the two groups. U_{CAMP}V was not significantly different between the two groups (HC-B vs. NC). The mean values for RBF and U_{Cl}V were significantly less in HC-B animals when compared to NC animals, but U_{Na}V, EF_{Na}, and U_KV were significantly greater in HC-B animals. V was not different in the two groups. These experiments demonstrate that the phosphaturia of HC occurs even when the associated extracellular acidosis has been minimized.

DISCUSSION

The present studies demonstrate that acute HC is associated with an increased U_PV in rats with intact parathyroid glands. Several known factors could contribute to such a phosphaturia. The possible role of

in P-TPTX rats

EF _P	P _{Ca_T}	P _{Ca_I}	U _{Ca} V	U _{CAMP} V	RBF	V	U _{Na} V	EF _{Na}	U _{Cl} V	U _K V
%	mg/ 100 ml	mg/ 100 ml	μg/ min	pmol/ min	ml/ min	μl/ min	μeq/ min	%	μeq/ min	μeq/ min
6.3	7.87	3.09	4.59	183	10.9	77.8	18.0	3.97	21.7	2.74
2.6	0.58	0.18	1.15	49	1.4	44.4	3.4	1.21	4.1	0.60
26.7	7.99	3.52	4.27	225	10.4	58.9	18.6	4.03	19.4	2.77
4.5	0.48	0.38	1.29	66	1.0	17.5	4.9	0.92	5.4	0.44
<0.001	>0.5	<0.05	>0.4	<0.05	>0.1	>0.05	>0.6	>0.8	>0.1	>0.8

TABLE IV
Effects of Bicarbonate Infusion

	B _{PCO₂}	B _{pH}	B _{HCO₃}	T _{HCO₃}	GFR	P _p	F _p	U _{pV}
	mm Hg		meq/ liter	μeq/ min	ml/ min	mg/ 100 ml	μg/ min	μg/ min
NC	41.2	7.42	26.0	80.7	3.08	6.19	191	7.93
±SD (n = 5)	3.0		1.3	21.7	0.74	1.01	51	6.01
NC-B	41.3	7.66	45.9	105	2.70	5.05	136	10.0
±SD (n = 5)	3.4		7.8	17	0.30	0.58	20	4.6
P*	>0.9	<0.001	<0.001	<0.001	<0.05	<0.001	<0.001	>0.2
HC-B	84.7	7.36	47.0	118	2.99	7.14	213	83.2
±SD (n = 5)	4.2		4.8	14	0.35	1.03	35	11.4
P*	<0.001	<0.001	<0.001	<0.001	>0.6	<0.01	>0.1	<0.001

* Compared to NC animals.

those factors during HC was explored in some detail. It seemed most likely that the phosphaturia might be related to release of endogenous PTH with subsequent reduction of renal phosphorus reabsorption (13–15). However, several findings offered no support for such a mechanism. First, a major stimulus for PTH release, lowering of the plasma calcium concentration (30), was not observed in intact HC animals. Second, an appropriate end-organ response to elevated levels of circulating PTH did not occur, inasmuch as U_{CAMPV} remained unchanged during HC. The most compelling evidence against a major role for PTH was the demonstration that HC still provided a potent phosphaturic stimulus when PTH release was prevented by acute thyroparathyroidectomy. The effectiveness of acute thyroparathyroidectomy in reducing plasma PTH activity is supported by the finding that EF_p and U_{CAMPV} were lower in NC and HC animals subjected to thyroparathyroidectomy than in their counterparts. These observations suggest that PTH does not play a major role in mediating the phosphaturia of acute HC.

Extracellular volume expansion is also known to increase U_{pV} (19–21). This mechanism appears to be an unlikely possibility inasmuch as NC and HC animals were subjected to the same degree of volume expansion using identical rates of infusion with infusates of the same composition. That success was achieved in the maintenance of comparable extracellular volumes in all groups of animals is supported by the similarity of values for V, U_{NaV}, EF_{Na}, GFR, and RBF in all experimental groups.

F_p was found to be increased in HC animals, a factor that could possibly contribute to the phosphaturia. Others have reported a similar finding and have suggested that the increased F_p was responsible for the phosphaturia (1, 4, 9). Indeed, an elevation of the P_p concentration was observed by us in intact HC

animals when compared with their NC counterparts, although the rise in F_p did not achieve statistical significance. However, a statistically significant increase in both the P_p concentration and F_p was found in HC animals subjected to acute thyroparathyroidectomy when compared to their NC counterparts. Thus, it was still felt necessary to examine the potential role of an increased P_p concentration and F_p on the phosphaturia of HC. NC- and HC-TPTX animals were studied at comparable levels of plasma and filtered phosphorus which were achieved by sodium phosphate infusion. The results indicated that HC produces phosphaturia independent of the increase of the P_p concentration or F_p, although it must be admitted that elevations of either may contribute to phosphaturia in a minimal way.

The possibility that the phosphaturia could be the result of an increase in the ultrafilterability of phosphate occurring as a direct consequence of HC was also entertained, but considered highly unlikely. Previous studies in rats have demonstrated that the ultrafilterable fraction is at least 0.93 (31). Parathyroidectomy (32, 33) and phosphate loading (31–33) appear to have no discernible effect on the ultrafilterable fraction of phosphate as determined in *in vitro* systems. Moreover, LeGrimellec et al. (34) have shown that the filterability of phosphate across the glomerular filtration barrier is essentially identical to the value measured across cuprophane. Thus, it appears that the maximum increment in filtered load which could be attributed to a change in phosphate ultrafilterability must be less than 10%. A change of this magnitude could not produce the phosphaturia observed in HC animals.

The relationship between a decrease in ECF pH and U_{pV} is unclear. This and earlier studies have demonstrated that respiratory and metabolic acidosis are

EF _P	P _{CaT}	P _{CaI}	U _{CaV}	U _{CAMPV}	RBF	V	U _{NaV}	EF _{Na}	U _{ClV}	U _{KV}
%	mg/ 100 ml	mg/ 100 ml	μg/ min	pmol/ min	ml/ min	μl/ min	μeq/ min	%	μeq/ min	μeq/ min
4.07	8.12	4.29	7.40	259	12.3	96.2	20.7	4.62	25.4	3.15
2.70	0.44	0.33	3.92	133	2.1	42.1	8.4	1.70	9.6	0.43
7.41	8.49	1.99	4.49	80	9.49	115.7	26.8	6.73	7.38	3.76
3.32	0.82	0.32	1.42	62	1.7	24.7	4.3	0.11	5.99	0.82
<0.001	>0.05	<0.001	<0.01	<0.001	<0.001	<0.05	<0.01	<0.001	<0.001	<0.01
39.6	8.12	3.40	4.68	195	11.0	111.6	30.4	6.83	4.47	3.55
5.8	1.20	0.53	2.80	62	1.2	35.4	7.2	1.41	2.24	0.63
<0.001	>0.9	<0.01	<0.02	>0.05	<0.05	>0.2	<0.001	<0.001	<0.001	<0.05

accompanied by phosphaturia (1–9, 35–38). However, the earlier studies were not designed to assess the influence of major known determinants of U_{PV}. In particular, studies of metabolic acidosis (35–38) did not attempt to define the role of compensatory changes in BpCO₂. Some investigators (23–25) have suggested that in the presence of a more acid renal tubular fluid at the site or sites of phosphate reabsorption, H₂PO₄⁻ relative to HPO₄⁼, would be increased and that, assuming H₂PO₄⁻ to be the more readily reabsorbed ionic species, tubular reabsorption of phosphate would increase. Therefore, if this mechanism is operative and important, a decrease in U_{PV} would be expected to occur during respiratory or metabolic acidosis if one assumes that the luminal fluid at the site or sites of phosphate reabsorption is more acid than in the nonacidotic state. The early proximal tubule represents one site for phosphate reabsorption (39–43) and, at this site, the luminal fluid pH should approach systemic pH. The distal nephron has also been shown by some investigators (39, 40, 44, 45) to reabsorb phosphate, and, for its more distal portion, urine pH should reflect tubular fluid pH. Thus, experimental conditions in which the pH of the blood and urine are lower should be associated with a decreased U_{PV}. Direct attempts to evaluate the importance of this mechanism in the proximal tubule have yielded conflicting results. Bank et al. (25), utilizing the micropfusion technique in the rat, demonstrated enhanced reabsorption of the H₂PO₄⁻ ion species, whereas Baumann et al. (26) demonstrated greater reabsorption of the HPO₄⁼ ion species.

To assess the possible effects of acid-base alterations on U_{PV}, HC-B and NC-B animals were studied (Tables I and IV). Phosphaturia was not prevented by normalization of the ECF pH with bicarbonate infusion (HC-B animals). These results suggest that changes in

the ionic species (H₂PO₄⁻ vs. HPO₄⁼) in the early proximal tubule or ECF acidosis per se provide an unlikely explanation for the phosphaturia of HC. In addition, inasmuch as urine pH was not significantly different ($P > 0.05$) in intact NC and HC animals, it appears unlikely that changes in tubular fluid pH in the terminal nephron can account for the phosphaturia. Interestingly, U_{PV} was actually greater in HC-B than in HC animals ($P < 0.001$), suggesting that some function of bicarbonate infusion served to enhance the phosphaturia associated with HC. This could have been the result of a change in ECF pH, B_{HCO₃}, or urine pH because all were significantly greater ($P < 0.001$) in the HC-B animals. However, NC-B animals also had significantly different B_{pH}, B_{HCO₃}, and urine pH than HC animals, but did not demonstrate phosphaturia. Therefore, the only acid-base disturbance that was consistently associated with phosphaturia was an elevation of the B_{pCO₂}. It would thus appear that alterations of B_{HCO₃}, B_{pH}, urine pH, and presumably of tubular fluid pH cannot account for the phosphaturia of HC. On the other hand, although ECF pH was normalized in HC-B animals, intracellular fluid pH may still have been less than normal because in certain *in vitro* studies where intracellular pH has been measured, poor cellular penetration of bicarbonate has been demonstrated in some cell types (46). It is possible that the phosphaturic effect of the increase in B_{pCO₂} is mediated via a reduction of intracellular pH in those tubular cells responsible for phosphate reabsorption.

The present studies demonstrated that an elevation in P_p concentration was associated with an increase in U_{CAMPV} in the presumed absence of endogenous PTH. NC-P-TPTX animals had a significantly greater U_{CAMPV} than NC-TPTX animals not receiving a phosphate infusion (183 vs. 106 pmol/min; $P < 0.01$). In addition,

HC-P-TPTX animals had significantly greater $U_{\text{cAMP}}V$ than HC-TPTX animals not receiving phosphate (225 vs. 157 pmol/min; $P < 0.01$). This study was not designed to determine whether increases in U_{cAMP} were of renal origin; however, it is possible that phosphate infusion could raise intracellular phosphate levels thereby stimulating increased production of cAMP with a resultant increase in $U_{\text{cAMP}}V$. One might speculate that by activating the cellular cAMP system, phosphate infusion could ultimately lead to enhanced excretion of phosphate providing a convenient mechanism for preventing phosphate retention during phosphate loading.

An interesting association was also observed between $U_{\text{cAMP}}V$ and bicarbonate infusion. Intact NC-B animals had a markedly reduced $U_{\text{cAMP}}V$ when compared to intact NC animals not receiving bicarbonate ($P < 0.001$). Yet, $U_{\text{p}}V$ did not decrease in the NC-B animals. Similar findings have been reported by Rodriguez et al. (47). When compared to NC animals, P_{Ca} fell significantly in the NC-B animals ($P < 0.001$). The expected finding would have been an increase both in the $U_{\text{cAMP}}V$ and $U_{\text{p}}V$ as PTH was released in response to the decrease in ionized calcium. Although the mechanism is obscure, bicarbonate infusion in animals with intact parathyroid glands appears to dissociate $U_{\text{cAMP}}V$ from $U_{\text{p}}V$.

The present results failed to confirm earlier in vitro observations which have demonstrated an inverse relationship between P_{Ca} concentration and B_{pH} (48). There was no significant difference noted in the P_{Ca} concentrations between intact HC ($B_{\text{pH}} - 7.15$) and intact NC animals ($B_{\text{pH}} - 7.42$). Our findings are consistent with those of Höffken et al. (49) who failed to observe a significant rise in P_{Ca} levels in intact rats in acute respiratory acidosis. The failure to observe such an inverse relationship may have been due to the effect of plasma phosphate on plasma concentrations of P_{Ca} . Elevation of the P_{p} concentration was associated with depression of P_{Ca} concentration in both NC- and HC-TPTX animals (NC-P-TPTX vs. NC-TPTX, $P < 0.01$; HC-P-TPTX vs. HC-TPTX, $P < 0.01$). Inasmuch as HC animals had a significant rise in their P_{p} concentrations in comparison with NC animals, the anticipated rise in P_{Ca} in response to respiratory acidosis may have been prevented. Therefore, acidosis, at least of acute respiratory etiology, did not produce an increase in P_{Ca} levels under the conditions of this study.

The present experiments were not designed to identify the source of the phosphate excreted in the urine of HC animals. In animals not receiving a phosphate infusion, the source of the increased phosphate must have derived from an intracellular pool, but it is not known which cell types contribute to such a pool that responds so readily to HC. However, during metabolic acidosis it has been demonstrated that the intracellular phosphate content of erythrocytes is de-

creased (50) and such a phenomenon could conceivably occur during respiratory acidosis.

In summary, acute HC was found to produce hyperphosphatemia and phosphaturia. The phosphaturia was not dependent upon endogenous PTH, an increase in F_{p} , an associated extracellular acidosis, or an increase in B_{HCO_3} concentration. Instead the findings suggest that the increase in B_{pCO_2} has a direct effect on the kidney which is mainly responsible for the phosphaturia observed during acute HC. Because HC is an extremely potent stimulus of phosphaturia, future studies directed toward examination of renal phosphorus reabsorption should consider this variable in the design of the experimental protocol.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the technical assistance of Mrs. Helen Parks and the secretarial assistance of Mrs. Greta Owens.

This work was supported, in part, by U. S. Public Health Service grants AM 13845 and HL 05848.

REFERENCES

1. Haldane, J. B. S., V. B. Wigglesworth, and C. E. Woodrow. 1924. The effect of reaction changes on human inorganic metabolism. *Proc. R. Soc. Med.* **96**: 1-14.
2. Brassfield, C. R., and V. G. Behrmann. 1941. A correlation of the pH of arterial blood and urine as affected by changes in pulmonary ventilation. *Am. J. Physiol.* **132**: 272-280.
3. Denton, D. A., M. Maxwell, I. R. McDonald, J. Munro, and W. Williams. 1952. Renal regulation of the extracellular fluid in acute respiratory acidemia. *Aust. J. Exp. Biol. Med. Sci.* **80**: 489-510.
4. Freeman, S., A. B. Jacobsen, and B. J. Williamson. 1957. Acid-base balance and removal of injected calcium from the circulation. *Am. J. Physiol.* **191**: 377-383.
5. Barker, E. S., R. B. Singer, J. R. Elkinton, and J. K. Clark. 1957. The renal response in man to acute experimental respiratory alkalosis and acidosis. *J. Clin. Invest.* **36**: 515-529.
6. Levitin, H., W. Branscome, and F. H. Epstein. 1958. The pathogenesis of hypochloremia in respiratory acidosis. *J. Clin. Invest.* **37**: 1667-1675.
7. Polak, A., G. D. Haynie, R. M. Hays, and W. B. Schwartz. 1961. Effects of chronic hypercapnia on electrolyte and acid-base equilibrium. I. Adaptation. *J. Clin. Invest.* **40**: 1223-1237.
8. Gray, S. P., J. E. W. Morris, and C. J. Brooks. 1973. Renal handling of calcium, magnesium, inorganic phosphate and hydrogen ions during prolonged exposure to elevated carbon dioxide concentrations. *Clin. Sci. Mol. Med.* **45**: 751-764.
9. Schwartz, W. B., N. C. Brackett, Jr., and J. J. Cohen. 1965. The response of extracellular hydrogen ion concentration to graded degrees of chronic hypercapnia: the physiologic limits of the defense of pH. *J. Clin. Invest.* **44**: 291-301.
10. Giebisch, G., L. Berger, and R. F. Pitts. 1955. The extrarenal response to acute acid-base disturbances of respiratory origin. *J. Clin. Invest.* **34**: 231-245.
11. Cohen, J. J., N. C. Brackett, Jr., and W. B. Schwartz. 1964. The nature of the carbon dioxide titration curve in the normal dog. *J. Clin. Invest.* **43**: 777-786.
12. Waite, L. C., W. A. Volkert, and A. D. Kenny. 1970. Inhibi-

- tion of bone resorption by acetazolamide in the rat. *Endocrinology*. 87: 1129-1139.
13. Greenwald, I., and J. Gross. 1925. The effect of the administration of a potent parathyroid extract upon the excretion of nitrogen, phosphorus, calcium, and magnesium, with some remarks on the solubility of calcium phosphate in serum and on the pathogenesis of tetany. *J. Biol. Chem.* 66: 217-227.
 14. Bentner, E. H., and P. L. Munson. 1960. Time course of urinary excretion of inorganic phosphate by rats after parathyroidectomy and after injection of parathyroid extract. *Endocrinology*. 66: 610-616.
 15. Pullman, T. N., A. R. Lavender, I. Aho, and H. Rasmussen. 1960. Direct renal action of a purified parathyroid extract. *Endocrinology*. 67: 570-582.
 16. Chase, L. R., and G. D. Aurbach. 1967. Parathyroid function and the renal excretion of 3',5'-adenylic acid. *Proc. Natl. Acad. Sci. U. S. A.* 58: 518-525.
 17. Kaminsky, N. I., A. E. Broadus, J. G. Hardman, D. J. Jones, Jr., J. H. Ball, E. W. Sutherland, and G. W. Liddle. 1970. Effects of parathyroid hormone on plasma and urinary adenosine 3',5'-monophosphate in man. *J. Clin. Invest.* 49: 2387-2395.
 18. Butlen, D., and S. Jard. 1972. Renal handling of 3'-5'-cyclic AMP in the rat: the possible role of luminal 3'-5'-cyclic AMP in the tubular reabsorption of phosphate. *Pflügers Arch. Eur. J. Physiol.* 331: 172-190.
 19. Frick, A. 1968. Reabsorption of inorganic phosphate in the rat kidney. I. Saturation of transport mechanism. II. Suppression of fractional phosphate reabsorption due to expansion of extracellular fluid volume. *Pflügers Arch. Eur. J. Physiol.* 304: 351-364.
 20. Maesaka, J. K., M. F. Levitt, and R. G. Abramson. 1973. Effect of saline infusion on phosphate transport in intact and thyroparathyroidectomized rats. *Am. J. Physiol.* 225: 1421-1429.
 21. Schneider, E. G., R. S. Goldsmith, C. D. Arnaud, and F. G. Knox. 1975. Role of parathyroid hormone in the phosphaturia of extracellular fluid volume expansion. *Kidney Int.* 7: 317-324.
 22. Carrasquer, G., and W. A. Brodsky. 1961. Elimination of transient secretion of phosphate by alkalization of plasma in dogs. *Am. J. Physiol.* 201: 499-504.
 23. Fulop, M., and P. Brazeau. 1968. The phosphaturic effect of sodium bicarbonate and acetazolamide in dogs. 1968. *J. Clin. Invest.* 47: 983-991.
 24. Puschett, J. B., and M. Goldberg. 1969. The relationship between the renal handling of phosphate and bicarbonate in man. *J. Lab. Clin. Med.* 73: 956-969.
 25. Bank, N., H. S. Aynedjian, and S. W. Weinstein. 1974. A micropuncture study of phosphate reabsorption by the rat proximal renal tubule. *J. Clin. Invest.* 54: 1040-1048.
 26. Baumann, K., G. Rumrich, F. Papavassiliou, and S. Klöss. 1975. Letters and notes: pH dependence on phosphate reabsorption in the proximal tubule of rat kidney. *Pflügers Arch. Eur. J. Physiol.* 360: 183-187.
 27. Malvin, R. L., and W. D. Lotspeich. 1956. Relation between tubular transport of inorganic phosphate and bicarbonate in the dog. *Am. J. Physiol.* 187: 51-56.
 28. Mercado, A., E. Slatopolsky, and S. Klahr. 1975. On the mechanisms responsible for the phosphaturia of bicarbonate administration. *J. Clin. Invest.* 56: 1386-1395.
 29. Gilman, A. G. 1970. A protein binding assay for adenosine 3':5'-cyclic monophosphate. *Proc. Natl. Acad. Sci. U. S. A.* 67: 305-312.
 30. Sherwood, L. M., G. P. Mayer, C. F. Ramberg, Jr., D. S. Kronfeld, G. D. Aurbach, and J. T. Potts, Jr. 1968. Regulation of parathyroid hormone secretion: proportional control by calcium, lack of effect of phosphate. *Endocrinology*. 83: 1043-1051.
 31. Boudry, J-F., U. Troehler, M. Touabi, H. Fleisch, and J-P. Bonjour. 1975. Secretion of inorganic phosphate in the rat nephron. *Clin. Sci. Mol. Med.* 48: 475-489.
 32. Steele, T. H., and H. F. DeLuca. 1976. Influence of dietary phosphorus on renal phosphate reabsorption in the parathyroidectomized rat. *J. Clin. Invest.* 57: 867-874.
 33. Engle, J. E., and T. H. Steele. 1975. Renal phosphate reabsorption in the rat: effect of inhibitors. *Kidney Int.* 8: 98-104.
 34. Le Grimellec, C., P. Poujeol, and C. de Rouffignac. 1975. ³H-Inulin and electrolyte concentrations in Bowman's capsule in rat kidney: comparison with artificial ultrafiltration. *Pflügers Arch. Eur. J. Physiol.* 354: 117-131.
 35. Haldane, J. B. S., R. Hill, and J. M. Luck. 1923. Calcium chloride acidosis. *J. Physiol. (Lond.)* 57: 301-306.
 36. Guest, G. M. 1942. Organic phosphates of the blood and mineral metabolism in diabetic acidosis. *Am. J. Dis. Child.* 64: 401-412.
 37. Martin, H. E., and R. Jones. 1961. The effect of ammonium chloride and sodium bicarbonate on the urinary excretion of magnesium, calcium, and phosphate. *Am. Heart J.* 62: 206-210.
 38. Scott, D. 1972. Excretion of phosphorus and acid in the urine of sheep and calves fed either roughage or concentrate diets. *Q. J. Exp. Physiol. Cogn. Med. Sci.* 57: 379-392.
 39. Amiel, C., H. Kuntziger, and G. Richet. 1970. Micropuncture study of handling of phosphate by proximal and distal nephron in normal and parathyroidectomized rat: evidence for distal reabsorption. *Pflügers Arch. Eur. J. Physiol.* 317: 93-109.
 40. Amiel, C., and H. Kuntziger. 1972. Use of radiophosphate as a tracer in micropuncture study of handling of inorganic phosphate by the nephron of the rat. In *Radio-nuclides in Nephrology*. M. D. Blafox and J-L Funck-Brentano, editors. Grune & Stratton Inc., New York. 159-164.
 41. Staum, B. B., R. J. Hamburger, and M. Goldberg. 1972. Tracer microinjection study of renal tubular phosphate reabsorption in the rat. *J. Clin. Invest.* 51: 2271-2275.
 42. Brunette, M. G., L. Taleb, and S. Carriere. 1973. Effect of parathyroid hormone on phosphate reabsorption along the nephron of the rat. *Am. J. Physiol.* 225: 1076-1081.
 43. Le Grimellec, C. 1975. Micropuncture study along the proximal convoluted tubule: electrolyte reabsorption in first convolutions. *Pflügers Arch. Eur. J. Physiol.* 354: 133-150.
 44. Wen, S. F. 1974. Micropuncture studies of phosphate transport in the proximal tubule of the dog. *J. Clin. Invest.* 53: 143-153.
 45. Harris, C. A., P. G. Baer, E. Chirito, and J. H. Dirks. 1974. Composition of mammalian glomerular filtrate. *Am. J. Physiol.* 227: 972-976.
 46. Waddell, W. J., and R. G. Bates. 1969. Intracellular pH. *Physiol. Rev.* 49: 285-329.
 47. Rodriguez, H. J., J. Walls, J. Yates, and S. Klahr. 1974. Effects of acetazolamide on the urinary excretion of cyclic AMP and on the activity of renal adenylyl cyclase. *J. Clin. Invest.* 53: 122-130.
 48. Moore, E. W. 1970. Ionized calcium in normal serum, ultrafiltrates, and whole blood determined by ion-exchanged electrodes. *J. Clin. Invest.* 49: 318-334.
 49. Höffken, B., D. K. Parkinson, P. Storms, and I. C. Radde. 1971. Effects of alterations of blood pH on calcium ion activity in rat plasma. *Clin. Orthop. Relat. Res.* 78: 30-39.
 50. Guest, G. M., and S. Rapoport. 1940. Clinical studies of the organic acid-soluble phosphorus of red blood cells in different acidotic states. *J. Lab. Clin. Med.* 26: 190-198.