Chronic Hypercapnia Stimulates Proximal Bicarbonate Reabsorption in the Rat

Martin G. Cogan

Cardiovascular Research Institute and Department of Medicine,
University of California, San Francisco, California 94143

Abstract. The hyperbicarbonatemia of chronic respiratory acidosis might be maintained by a reduction in filtration rate or an enhancement of tubular bicarbonate reabsorption. To investigate this question, 12 Munich-Wistar rats were exposed to a 10% CO₂ atmosphere for 6–8 d. Chronic respiratory acidosis developed, with arterial pH 7.30 ± 0.01 , partial pressure of CO₂ (pCO₂) 80 ± 2 mmHg, and total CO₂ concentration 45 ± 1 mM. Single nephron glomerular filtration rate was normal (42 ± 1 nl/min). Chronic hypercapnia caused absolute proximal reabsorption to be significantly stimulated ($1,449 \pm 26$ pmol/min) as compared with reabsorption previously observed in normal animals ($1,075 \pm 74$ pmol/min) or in animals subjected to acute hypercapnia ($1,200 \pm 59$ pmol/min). This is the first demonstration that proximal bicarbonate reabsorption can be stimulated above normal euvoletic values. When eight animals were subsequently allowed to return toward a normocapnic state (arterial pCO₂ 46 ± 1 mmHg) over the course of 1–1.5 h, bicarbonate reabsorption was still significantly higher ($1,211 \pm 34$ pmol/min) than in similarly alkalotic, normocapnic control groups (994 ± 45 pmol/min). In conclusion, chronic, but not acute, hypercapnia stimulates absolute proximal bicarbonate reabsorption to exceed the level found in normal euvoletic rats.

Introduction

The high plasma bicarbonate concentration found in chronic respiratory acidosis might be maintained in two ways. First,

This work has been presented in part at the Sixteenth Annual Meeting of the American Society of Nephrology, Washington DC, 4–7 December 1983, and published in abstract (1984. *Kidney Int.* 25:273).

Address reprint requests to Dr. Cogan, 1065 HSE, University of California, San Francisco, CA 94143.

Received for publication 3 April 1984 and in revised form 25 June 1984.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/84/12/1942/06 \$1.00

Volume 74, December 1984, 1942–1947

glomerular filtration might fall as the plasma bicarbonate concentration rises, thus attenuating or preventing the increase in filtered bicarbonate load. Relatively normal rates of tubular hydrogen ion secretion would then suffice to prevent bicarbonate from appearing in the urine. This mechanism has been proposed to be the one that maintains the other major hyperbicarbonatemic acid-base disorder, chronic metabolic alkalosis (1). On the other hand, if glomerular filtration remained normal, enhanced tubular bicarbonate reabsorption, predominantly in the proximal nephron, would be needed to prevent bicarbonaturia.

Measurement of absolute rates of bicarbonate filtration and proximal reabsorption needed to differentiate these two pathophysiological mechanisms have not been reported to date. In previous free-flow micropuncture studies fractional proximal bicarbonate reabsorption in chronic hypercapnia appeared normal, a finding that does not allow discrimination between the two alternatives (2). Split-droplet studies during chronic hypercapnia have found a relatively normal rate of proximal acidification (3), which would support the first process discussed above (reduced filtration rate with normal bicarbonate reabsorption). Acute hypercapnia has also been found to have only a small effect on free-flow proximal bicarbonate reabsorption (4). In fact, no factor has yet been reported to reliably increase proximal bicarbonate reabsorption above the mean level found in the normal euvoletic state (1, 5).

Thus, the purpose of the present studies was to measure absolute rates of bicarbonate filtration and reabsorption in rats rendered chronically hypercapnic.

Methods

Protocols. To induce chronic respiratory acidosis, 12 Munich-Wistar rats were exposed to 10% CO₂ (air balance) in an environmental chamber for 6–8 d. Gas samples from the chamber were periodically measured to ensure that the 10% CO₂ atmosphere was constant. Rats were allowed chow diet and distilled water ad lib. while in the chamber. On the day of micropuncture study, the animal was rapidly weighed (mean weight was 184 ± 4 g), injected intraperitoneally with Inactin (100 mg/kg), and immediately placed into a holding chamber flooded with the same 10% CO₂ gas mixture. When anesthetized, the rat was removed from the holding chamber, a tracheostomy and intubation were performed as rapidly as possible (≤ 2 min), and 10% CO₂ was again administered via a T-tube apparatus, as previously described (4).

The rest of the surgical preparation and free-flow micropuncture techniques were performed as previously outlined (5). Surgically induced plasma volume losses, assumed to be similar to previously established values (1.3% body weight), and whole blood losses from sampling were quantitatively replaced (5). Donors for plasma and whole blood were also maintained in the environmental chamber with 10% CO₂ for 6–8 d.

After the first period of micropuncture, eight animals were extubated and allowed to breathe room air to allow arterial partial pressure of CO₂ (pCO₂)¹ to decrease toward normal. After a 30–60-min equilibration, the animals were again studied by micropuncture during this post-hypercapnic period. The time between extubation and the end of the second micropuncture period was ~1–1.5 h.

A separate group of 15 animals was rendered acutely alkalotic to serve as normocapnic controls for the post-hypercapnic animals. A 10% body weight bicarbonate solution (NaHCO₃ 200 mM plus KCl 25 mM) was infused over 1 h, and followed by a maintenance infusion of 0.1 ml/min. Micropuncture began after an equilibration of 30 min. Tubular chloride concentrations were not measured in this group.

Analysis and calculations. Plasma, urine, and tubular fluid measurements were made as previously described (1, 4, 5). [³H]inulin was measured by scintillation counting, total CO₂ by microcalorimetry (6), and chloride by the method of Ramsay (7). Calculations of filtration and reabsorptive rates were also as previous outlined (1, 4, 5). Results are presented as mean±SEM, and significance was assessed by the paired or unpaired *t* test as appropriate.

Results

Exposure to 10% CO₂ for 6–8 d resulted in chronic respiratory acidosis. Arterial measurements were: pH 7.30±0.01; pCO₂, 80±2 mmHg; plasma total CO₂ concentration, 45±1 mM; and plasma chloride concentration, 93±1 meq/liter. Other blood measurements and urinary excretion data, shown in Table I, were generally similar to those previously reported for normal euvoletic rats (5, 8). Kidney weight was 0.90±0.04 g.

Micropuncture results are shown in Table II. Single nephron glomerular filtration rate (SNGFR) (42.4±0.7 nl/min) was normal for rats of this size (5, 8). The elevated total CO₂ concentration in Bowman's space (47.9±0.9 mM) was therefore associated with a high filtered total CO₂ load (2,025±47 pmol/min). Chronic hypercapnia caused absolute proximal total CO₂ reabsorption to be stimulated to 1,449±26 pmol/min, a value 35% higher than that of a normal euvoletic animal (1,075±74 pmol/min, *P* < 0.001) (5).

Bicarbonate reabsorption during chronic hypercapnia was also higher than values previously reported after acute exposure to 10% CO₂ (4). The comparison of micropuncture data during chronic and acute hypercapnia is shown in Table III. In chronic hypercapnia, total CO₂ reabsorption was 23% higher than when acute hypercapnia was superimposed on chronic metabolic alkalosis, despite there being only a slight difference

Table I. Blood Composition and Urinary Excretion Rates in Chronic Hypercapnia and Post-Hypercapnia

	Chronic hypercapnia	Post-hypercapnia	<i>P</i>
Blood			
Arterial			
pH	7.30±0.01	7.50±0.01	<0.001
pCO ₂ (mmHg)	80±2	46±1	<0.001
Blood pressure (mmHg)	130±4	114±3	<0.005
Hematocrit (vol %)	47.5±0.7	46.5±0.5	<0.025
Plasma			
[Na ⁺] (meq/liter)	149±1	146±1	<0.01
[K ⁺] (meq/liter)	3.9±0.1	3.9±0.1	NS
[Total CO ₂] (mM)	45±1	37±1	<0.001
[Cl ⁻] (meq/liter)	93±1	96±1	<0.05
[Protein] (g/dl)	5.2±0.1	5.0±0.1	<0.25
Urine			
Glomerular filtration rate			
(ml/min · g kwt)	1.0±0.1	1.0±0.1	NS
Urinary excretion			
Na ⁺ (μeq/min)	0.2±0.1	0.2±0.1	NS
K ⁺ (μeq/min)	0.5±0.1	1.0±0.1	<0.01
Total CO ₂ (μmol/min)	0.05±0.02	0.4±0.1	<0.05
Cl ⁻ (μeq/min)	0.5±0.1	1.0±0.4	NS
Volume (μl/min)	2.9±0.4	2.4±0.2	NS

kwt, kidney weight; NS, not significant.

in the logarithmic mean luminal total CO₂ concentration, an important determinant of proximal acidification (5, 9, 10). Total CO₂ reabsorption was also 21% higher during chronic hypercapnia than when acute hypercapnia was superimposed on acute metabolic alkalosis. In this case, the acutely hypercapnic group had a higher SNGFR and mean luminal total CO₂ concentration, factors which, if anything, should accelerate acidification (9, 10). The stimulation of bicarbonate reabsorption over mean normal levels in the chronically hypercapnic group but not in the combined acutely hypercapnic groups is illustrated in Fig. 1.

Absolute chloride reabsorption during chronic respiratory acidosis (Table II) was slightly reduced from normal values, possibly as a result of the hypochloremic state (1), whereas absolute water reabsorption was approximately the same as in normal euvoletic animals (5, 8).

When eight of the animals were extubated and allowed to breathe room air, a stable post-hypercapnic metabolic alkalosis developed, with arterial pH of 7.50±0.01 and pCO₂ of 46±1 mmHg (Table I). Plasma total CO₂ concentration fell to 37±1 mM, associated with bicarbonaturia. SNGFR was stable (Table II). The resulting decline in filtered total CO₂ load (1,665±68 pmol/min) was associated with a fall in absolute proximal total CO₂ reabsorption to 1,211±34 pmol/min. However, this

1. Abbreviations used in this paper: pCO₂, partial pressure of CO₂; SNGFR, single nephron glomerular filtration rate.

Table II. Proximal Filtration and Reabsorption during Chronic Hypercapnia and Post-Hypercapnia

Group	Bowman's space			End-proximal			Absolute proximal reabsorption			Fractional proximal reabsorption		
	SNGFR*	[tCO ₂]	[Cl]	[tCO ₂]	[Cl]	V	tCO ₂	Cl	H ₂ O	tCO ₂	Cl	H ₂ O
	nl/min	mM	meq/liter	mM	meq/liter	nl/min	pmol/min	peq/min	nl/min			
Chronic hypercapnia												
(n = 12)	42.4±0.7	47.9±0.9	102.9±2.4	23.5±1.2	126.0±2.4	24.2±1.0	1,449±26	1,322±131	18.1±0.6	0.72±0.02	0.30±0.02	0.43±0.02
Post-hypercapnia												
(n = 8)	40.4±1.3	41.2±0.9	106.4±2.3	18.5±1.4	130.3±2.4	24.2±1.2	1,211±34	1,140±141	16.2±0.3	0.73±0.02	0.27±0.03	0.40±0.01
P‡	NS	<0.001	NS	<0.05	NS	NS	<0.01	NS	NS	NS	NS	NS
Acute metabolic alkalosis (n = 15)												
	46.3±2.2	45.1±1.0	—	35.5±1.3	—	30.1±1.7	1,015±49	—	16.2±0.8	0.49±0.01	—	0.35±0.01
P§	NS	<0.05	—	<0.001	—	<0.05	<0.01	—	NS	<0.001	—	<0.025

Means±SEM. * tCO₂, total CO₂; V, end-proximal flow rate; NS, not significant. ‡ P value comparing chronic hypercapnia with post-hypercapnia. § P value comparing post-hypercapnia with acute metabolic alkalosis.

Table III. Proximal Bicarbonate Reabsorption during Hypercapnic States

Group	Bowman's space [total CO ₂]	SNGFR	Filtered total CO ₂ load	End-proximal [total CO ₂]	Log mean [total CO ₂]	Absolute proximal total CO ₂ reabsorption
	mM	nl/min	pmol/min	mM	mM	pmol/min
Chronic hypercapnia (n = 12)	47.9±0.9	42.4±0.7	2,025±47	23.5±1.2	34±1	1,449±26
Acute hypercapnia*						
+ chronic metabolic alkalosis (n = 10)	50.4±1.7	27.9±1.7	1,385±98	15.5±1.1	30±1	1,176±77
P‡	NS	<0.001	<0.001	<0.001	<0.05	<0.005
+ Acute metabolic alkalosis (n = 7)	58.6±1.5	50.1±1.5	2,951±249	49.0±1.5	54±1	1,234±97
P‡	<0.001	<0.025	<0.001	<0.001	<0.001	<0.025

* From reference 4. ‡ As compared with chronic hypercapnia.

value for total CO₂ reabsorption in the post-hypercapnic period was still significantly higher than in a previous report on a group of 22 animals with chronic metabolic alkalosis (981±49 pmol/min, $P < 0.025$) that had both a similar arterial pCO₂ (43±1 mmHg) and a mean luminal total CO₂ concentration (32 mM) (1). The post-hypercapnic value was also higher than in the 15 animals in this study with acute metabolic alkalosis (1,015±49 pmol/min, $P < 0.01$), as shown in Table II. This

latter group had a slightly lower arterial pCO₂ (39±1 mmHg) but a higher SNGFR (46±2 nl/min) and mean luminal total CO₂ concentration (40 mM). Thus, as illustrated in Fig. 2, stimulation of proximal bicarbonate reabsorption by chronic hypercapnia appeared to persist, at least to some degree, after inhalation of CO₂ was terminated.

Discussion

It has long been known that several days are required for the full increase in renal bicarbonate reabsorption due to a sustained

Figure 1. Comparison of absolute proximal bicarbonate reabsorption in rats with chronic hypercapnia with reabsorption in normal euvo-lemic rats (from reference 5) and with reabsorption in combined groups of rats with acute hypercapnia superimposed on chronic or acute metabolic alkalosis (from reference 4).

Figure 2. Comparison of absolute proximal bicarbonate reabsorption in rats with post-hypercapnic alkalosis with reabsorption in rats with acute metabolic alkalosis (no pre-existing hypercapnic stimulus).

increment in arterial $p\text{CO}_2$ (11), irrespective of sodium intake (12), bicarbonate availability and rate of net acid excretion (13), potassium stores (14), or mineralocorticoid activity (15). The present studies demonstrate that this renal adaptive response is mediated not by altered hemodynamics (i.e., by a reduction in glomerular filtration rate) but rather by increased bicarbonate transport in the superficial proximal convoluted tubule. A simultaneous adaptation in bicarbonate reabsorption by sites distal to the superficial proximal convoluted tubule and/or by juxtamedullary nephrons is also possible. That such nephron segments contribute to the adaptive response is supported by the finding that when the arterial $p\text{CO}_2$ was allowed to return toward normal in the second period, bicarbonaturia appeared despite the fact that bicarbonate delivery out of the superficial proximal tubule was lower (454 ± 46 post-hypercapnia as compared with 577 ± 45 pmol/min during chronic hypercapnia).

The mechanism underlying this stimulation of proximal bicarbonate reabsorption during chronic hypercapnia might involve either a primary increase in Na^+/H^+ exchanger activity and/or in bicarbonate exit. Rector has argued for the former based on the time course of changes in ammonium excretion induced by hypercapnia (16). An increase in antiporter activity caused by chronic metabolic acidosis has been recently demonstrated (17–19). Studies using vesicle preparations or intracellular microelectrodes will be necessary to clarify whether a similar mechanism is induced by chronic respiratory acidosis.

The augmentation of proximal bicarbonate reabsorption persisted, to some degree, for as long as 1.5 h after the discontinuation of CO_2 inhalation. The possibility that cellular transport processes might have a “memory” of pre-existing conditions has been previously proposed for acidification in the distal tubule after hypocapnia (20), metabolic acid-base disturbances (21), or dietary manipulations (22). Changes in glycodiazine transport after acid-base disorders (3) and in volume adsorption measured in vitro after partial renal ablation in vivo (23) have also been documented in the proximal convoluted tubule. Such a memory effect in the proximal tubule after normalization of a high CO_2 tension might be at least partially responsible for maintaining post-hypercapnic metabolic alkalosis, previously attributed solely to extracellular volume depletion (13).

In conclusion, chronic but not acute hypercapnia increased absolute proximal bicarbonate reabsorption. This is the first demonstration that proximal bicarbonate reabsorption can be stimulated above the mean level found in a normal, euvoletic animal.

Acknowledgments

I wish to thank Dr. Edmond I. Eger and Mr. Brynte H. Johnson for the generous loan of the environmental chamber, Mr. Kenneth Wong

for excellent technical assistance, and Dr. Floyd C. Rector, Jr. for ongoing support and advice.

These studies were supported in part by a Clinical Investigator Award (1 KO8 AM-01015) and a grant (AM-27045) from the National Institute of Arthritis, Diabetes, and Digestive and Kidney Diseases.

References

1. Cogan, M. G., and F.-Y. Liu. 1983. Metabolic alkalosis in the rat. Evidence that reduced glomerular filtration rather than enhanced tubular bicarbonate reabsorption is responsible for maintaining the alkalotic state. *J. Clin. Invest.* 71:1141–1160.
2. Warren, Y., R. G. Luke, M. Kashgarian, and H. Levitin. 1970. Micropuncture studies of chloride and bicarbonate absorption in the proximal renal tubule of the rat in respiratory acidosis and in chloride depletion. *Clin. Sci.* 38:375–383.
3. Ullrich, K. J., G. Rumrich, and K. Baumann. 1975. Renal proximal tubular buffer-(glycodiazine) transport. Inhomogeneity of local transport rate, dependence on sodium, effect of inhibitors and chronic adaptation. *Pfluegers Arch.* 357:149–163.
4. Cogan, M. G. 1984. Effects of acute alterations in $p\text{CO}_2$ on proximal HCO_3^- , Cl^- and H_2O reabsorption. *Am. J. Physiol. (Renal Fluid and Electrolyte Physiol.)* 15: 246:F21–F26.
5. Cogan, M. G., D. A. Maddox, M. S. Lucci, and F. C. Rector, Jr. 1979. Control of proximal bicarbonate reabsorption in normal and acidotic rats. *J. Clin. Invest.* 64:1168–1180.
6. Vurek, G. G., D. G. Warnock, and R. Corsey. 1975. Measurement of picomole amounts of carbon dioxide by calorimetry. *Anal. Chem.* 47:765–767.
7. Ramsay, J. A., R. H. J. Brown, and P. C. Croghan. 1955. Electrometric titration of chloride in small volumes. *J. Exp. Biol.* 32:822–829.
8. Cogan, M. G. 1983. Volume expansion predominantly inhibits proximal reabsorption of NaCl rather than NaHCO_3 . *Am. J. Physiol. (Renal Fluid Electrolyte Physiol.)* 245:F272–F275.
9. Cogan, M. G., and R. J. Alpern. 1984. Regulation of proximal bicarbonate reabsorption. *Am. J. Physiol. (Renal Fluid and Electrolyte Physiol.)* 247:F387–F395.
10. Alpern, R. J., M. G. Cogan, and F. C. Rector, Jr. 1982. Effect of luminal bicarbonate concentration on proximal acidification in the rat. *Am. J. Physiol. (Renal Fluid Electrolyte Physiol.)* 243:F53–F59.
11. Sullivan, W. J., and P. J. Dorman. 1955. The renal response to chronic respiratory acidosis. *J. Clin. Invest.* 34:268–276.
12. Polak, A., G. D. Haynie, R. M. Hays, and W. B. Schwartz. 1961. Effects of chronic hypercapnia on electrolyte and acid-base equilibrium. I. Adaptation. *J. Clin. Invest.* 40:1223–1237.
13. Van Ypersele de Strihou, C., P. F. Gulyassy, and W. B. Schwartz. 1962. Effects of chronic hypercapnia on electrolyte and acid-base equilibrium. III. Characteristics of the adaptive and recovery process as evaluated by provision of alkali. *J. Clin. Invest.* 41:2246–2253.
14. Makoff, D., and B. J. Rosenbaum. 1971. Adaptation to chronic hypercapnia in the potassium-depleted dog. *Am. J. Physiol.* 220:1724–1727.
15. Luke, R. G., and H. Levitin. 1966. The renal and electrolyte

response to respiratory acidosis in the adrenalectomized rat. *Yale J. Biol. Med.* 39:27-37.

16. Rector, F. C., Jr. 1965. Role of the kidney in the homeostatic control of hydrogen ion concentration in body fluids. *Excerpta Medical Int. Congr. Ser.* 87:176-188.

17. Cohn, D. W., S. Klahr, and M. R. Hammerman. 1983. Metabolic acidosis and parathyroidectomy increase Na^+/H^+ exchange in brush border vesicles. *Am. J. Physiol. (Renal Fluid Electrolyte Physiol.)*. 245:F217-F222.

18. Kinsella, J. L., T. Cujkid, and B. Sacktor. 1984. Na^+/H^+ exchange activity in renal brush border membrane vesicles in response to metabolic acidosis: the role of glucocorticoids. *Proc. Natl. Acad. Sci. USA*. 81:630-634.

19. Tsai, C.-J., H. E. Ives, R. J. Alpern, V. J. Yee, D. G. Warnock, and F. C. Rector, Jr. 1984. Increased $V_{\max}$ for Na^+/H^+ antiporter activity in proximal tubule brush border vesicles from rabbits with

metabolic acidosis. *Am. J. Physiol. (Renal Fluid Electrolyte Physiol.)*. 247:F339-F343.

20. Battle, D. C., K. Itsarayoungyuen, M. Downer, R. Foley, J. A. L. Arruda, and N. A. Kurtzman. 1983. Suppression of distal urinary acidification after recovery from chronic hypocapnia. *Am. J. Physiol. (Renal Fluid Electrolyte Physiol.)*. 245:F433-F442.

21. Ullrich, K. J., and F. Papavassiliou. 1981. Bicarbonate reabsorption in the papillary collecting duct of rats. *Pfluegers Arch.* 389:271-275.

22. McKinney, T. D., and M. B. Burg. 1977. Bicarbonate transport by rabbit cortical collecting tubules. Effect of acid and alkali loads in vivo on transport in vitro. *J. Clin. Invest.* 60:766-768.

23. Trizna, W., N. Yanagawa, Y. Bar-Khayim, B. Houston, and L. G. Fine. 1981. Functional profile of the uremic nephron. Evidence of proximal tubular "memory" in experimental renal disease. *J. Clin. Invest.* 68:760-767.

/