

Relationship of Oxygen and Glutathione in Protection against Carbon Tetrachloride-induced Hepatic Microsomal Lipid Peroxidation and Covalent Binding in the Rat

Rationale for the Use of Hyperbaric Oxygen to Treat Carbon Tetrachloride Ingestion

Raymond F. Burk, James M. Lane, and Kuldeep Patel
Department of Medicine, Division of Gastroenterology & Nutrition, University of Texas Health Science Center, San Antonio, Texas 78284

Abstract. CCl_4 exerts its toxicity through its metabolites, including the free radicals $\text{CCl}_3\cdot$ and $\text{CCl}_3\text{OO}\cdot$. Oxygen strongly inhibits the hepatic cytochrome P-450-mediated formation of $\text{CCl}_3\cdot$ from CCl_4 and promotes the conversion of $\text{CCl}_3\cdot$ to $\text{CCl}_3\text{OO}\cdot$. Both these free radicals injure the hepatocyte by causing lipid peroxidation and binding covalently to cell structures. A reduced glutathione (GSH)-dependent mechanism can protect the liver microsomal membrane against CCl_4 -induced damage under aerobic conditions but not under anaerobic conditions (Burk, R. F., K. Patel, and J. M. Lane, 1983, *Biochem. J.*, 215:441-445). Experiments were carried out using rat liver microsomes to examine the effect of O_2 tensions found in the liver and of GSH on CCl_4 -induced covalent binding and lipid peroxidation. An NADPH-supplemented microsomal system was used. CCl_4 or $^{14}\text{CCl}_4$ was added to the sealed flask that contained the system, and after 20 min CHCl_3 production, thiobarbituric acid-reactive substances (an index of lipid peroxidation), and covalent binding of ^{14}C were measured. O_2 tensions of 0, 1, 3, 5, and 21% were studied. Increases in O_2 tension caused a fall in CHCl_3 production, which indicated that it decreased $\text{CCl}_3\cdot$. GSH had no significant effect on CHCl_3 production at any O_2 tension. Lipid peroxidation and covalent binding of ^{14}C fell progressively as O_2 tension was increased from 1 to 21%. The addition of GSH decreased both lipid peroxidation and covalent binding, but did so better at the higher O_2 tensions than at the lower ones.

Received for publication 21 February 1984 and in revised form 22 May 1984.


J. Clin. Invest.
© The American Society for Clinical Investigation, Inc.
0021-9738/84/12/1996/06 \$1.00
Volume 74, December 1984, 1996-2001

These results indicate that low O_2 tensions such as are found in the centrilobular areas of the liver favor conversion of CCl_4 to free radical products which cannot be detoxified by the GSH-dependent mechanism. They suggest that hyperbaric O_2 might decrease free radical formation in the liver in vivo and promote formation of $\text{CCl}_3\text{OO}\cdot$ from $\text{CCl}_3\cdot$. This should result in diminished CCl_4 -induced lipid peroxidation and liver damage. Rats given CCl_4 (2.5 ml/kg) were studied in metabolic chambers. Production of CHCl_3 and ethane, the latter an index of lipid peroxidation, were measured. Rats in two atmospheres of 100% O_2 produced much less CHCl_3 and ethane than rats in air. This strongly suggests that hyperbaric O_2 is decreasing free radical formation from CCl_4 and/or promoting the formation of $\text{CCl}_3\text{OO}\cdot$ from $\text{CCl}_3\cdot$. These results provide the rationale for the use of hyperbaric O_2 in the treatment of CCl_4 ingestion.

Introduction

The hepatotoxicity of CCl_4 results from its metabolic conversion to free radical products (1). Evidence of $\text{CCl}_3\cdot$ formation from CCl_4 was provided in the 1960's by the identification of CHCl_3 (2) and C_2Cl_6 (3) in tissues of animals that were given CCl_4 . More recently, spin-trapping experiments have verified directly

Scheme 1



the production of $\text{CCl}_3\cdot$ from CCl_4 by the rat liver microsomal cytochrome P-450 system (4).

The major reactions of $\text{CCl}_3\cdot$ with liver microsomes (Scheme 1) appear to be covalent binding (5) and H^\cdot abstraction from polyunsaturated fatty acids (PUFA).¹ The H^\cdot abstraction results in CHCl_3 and PUFA $^\cdot$ formation. PUFA $^\cdot$ can enter into cross-linking reactions or lipid peroxidation, depending on the O_2 tension.

In 1978, Packer et al. (6) reported that O_2 reacts very rapidly with $\text{CCl}_3\cdot$ to yield $\text{CCl}_3\text{OO}^\cdot$. Thus, the presence of O_2 when $\text{CCl}_3\cdot$ is formed results in the formation of a second radical species as shown in Scheme 1. Because this reaction is so rapid, O_2 can be expected to compete with microsomal constituents for $\text{CCl}_3\cdot$.

$\text{CCl}_3\text{OO}^\cdot$ is many orders of magnitude more reactive with PUFA than is $\text{CCl}_3\cdot$ (7), and therefore is a potent initiator of lipid peroxidation. It or some of its breakdown products can bind covalently to microsomes as well (8). $\text{CCl}_3\text{OO}^\cdot$ seems unlikely to serve as a source of CHCl_3 (9). Therefore, although both $\text{CCl}_3\cdot$ and $\text{CCl}_3\text{OO}^\cdot$ can cause membrane injury through covalent binding and lipid peroxidation, CHCl_3 arises only from $\text{CCl}_3\cdot$ and may serve as an index of $\text{CCl}_3\cdot$ concentration.

O_2 inhibits the cytochrome P-450-mediated formation of $\text{CCl}_3\cdot$ from CCl_4 . This inhibition is profound, but is not complete even under 21% O_2 (10, 11). Thus, O_2 diminishes $\text{CCl}_3\cdot$ concentration by (1) inhibiting its formation and (2) promoting its conversion to $\text{CCl}_3\text{OO}^\cdot$. The highest concentration of $\text{CCl}_3\cdot$ in a microsomal system will occur then under anaerobic conditions, and its concentration should fall as O_2 tension increases. The effect of O_2 on $\text{CCl}_3\text{OO}^\cdot$ concentration will be more complex. As O_2 tension rises, a greater fraction of the $\text{CCl}_3\cdot$ that is present will be converted to $\text{CCl}_3\text{OO}^\cdot$, but a smaller amount of $\text{CCl}_3\cdot$ will be formed. Thus, the highest $\text{CCl}_3\text{OO}^\cdot$ concentration should be achieved at an intermediate O_2 tension. At higher O_2 tensions, virtually all the CCl_4 -derived free radicals will be $\text{CCl}_3\text{OO}^\cdot$, but the total conversion of CCl_4 to radicals can be expected to be small, so $\text{CCl}_3\text{OO}^\cdot$ concentration will decline.

In an earlier study with NADPH-supplemented rat liver microsomes (11), we found that substantial amounts of CCl_4 disappeared from the flask under anaerobic conditions. CHCl_3 appeared and covalent binding occurred. When the atmosphere in the flask was 21% O_2 , no disappearance of CCl_4 could be detected by the method used. No CHCl_3 production was detected. A small amount of CCl_4 metabolism was occurring, however, because covalent binding and lipid peroxidation were detected.

It seems likely that nearly all the CCl_4 -derived free radicals present under the anaerobic conditions (see Scheme 1) were $\text{CCl}_3\cdot$ and nearly all those present under 21% O_2 were $\text{CCl}_3\text{OO}^\cdot$. This has potential importance because reduced glutathione (GSH) inhibited covalent binding and lipid peroxidation under 21% O_2 , but had no effect on covalent binding or CHCl_3

formation under anaerobic conditions, which suggests that a GSH-dependent mechanism can protect the membrane against $\text{CCl}_3\text{OO}^\cdot$ but not against $\text{CCl}_3\cdot$.

We undertook the present study to investigate the relevance of these observations to the hepatotoxicity of CCl_4 . Experiments were designed to examine the effect of O_2 tensions present in the liver and the effect of GSH on CCl_4 metabolism and the injurious processes of lipid peroxidation and covalent binding. The effect of hyperbaric O_2 in vivo was also assessed as a potential treatment for CCl_4 ingestion.

Methods

Animals. Male Sprague-Dawley strain rats were purchased from Timco Laboratories, Houston, TX. They were given food and water ad libitum and were housed in rooms with alternating 12-h light and dark cycles. Rats used in the experiments depicted in Figs. 3 and 4 were fed standard laboratory rations and rats from which microsomes were prepared (Figs. 1 and 2) were fed a nutritionally adequate semisynthetic diet (12). Experiments were begun between 8 and 10 a.m.

Rats were killed by cervical dislocation and exsanguination. Hepatic microsomes were prepared as described before (13) and were used for experiment on the day of preparation.

Microsomal incubations. The microsomal incubations were carried out in a volume of 5 ml in 25-ml Erlenmeyer flasks that were sealed with rubber septums. They were shaken in a water bath at 37°C, and the buffer used (microsomal incubation buffer) was 50 mM Tris-HCl, pH 7.5, with 140 mM NaCl and 50 μM EDTA. When used, GSH was added to a concentration of 1 mM.

The incubations reported in Figs. 1 and 2 were carried out under O_2 - N_2 mixtures. O_2 concentration is indicated on the figures. The gas mixtures were made with a three-tube flowmeter (Matheson Gas Products, Newark, NJ) and solutions and buffers used were equilibrated by bubbling the buffer and solutions to be added with the gas mixture for 15 min. Then the flask that contained the incubation mixture without CCl_4 and NADPH was flushed with the gas mixture for 5 min at 37°C before the flask was sealed and CCl_4 (10 μl vol in ethanol) was injected. After a further 5 min equilibration, the incubation was started by injection of NADPH. Flask protein concentrations varied from 1.14 to 1.60 mg/ml. NADPH concentration was 400 μM . After 20 min of incubation, flask atmosphere was sampled for CHCl_3 analysis. Then the sealed flask was chilled on ice for several minutes before it was opened. A sample was taken immediately for assay of thiobarbituric acid-reactive substances when the flask was opened.

For the experiment in Fig. 1, 1 mM CCl_4 was used. For the one in Fig. 2, 72 μM ^{14}C -labeled CCl_4 (2.8 Ci/mol, sp act) was used. To determine covalent binding to microsomes, 1 ml of 10% trichloroacetic acid was added to each flask after cooling and the microsomes were pelleted by centrifugation. The pellet was washed three times with 5 ml ethyl acetate to remove ^{14}C -labeled material that was not covalently bound. Then the pellet was digested in a scintillation vial by 0.5 ml Protosol and ^{14}C was determined after addition of 10 ml of Ready-Solv EP (Beckman Instruments, Palo Alto, CA).

In vivo studies. The in vivo studies were carried out using airtight plexiglas metabolism chambers (14). Each chamber contained soda lime to remove carbon dioxide and anhydrous CaSO_4 to remove water vapor. Hyperbaric conditions were maintained by pressurization of the O_2 -filled chambers to two atmospheres using compressed 100% O_2 . An air atmosphere was maintained by attachment of the chambers, sealed with air inside, to a respirometer that contained 100% O_2 . As

1. Abbreviations used in this paper: PUFA, polyunsaturated fatty acids; GSH, reduced glutathione.

O₂ was consumed by the rat, it was replaced by O₂ from the respirometer, which maintained the original O₂ concentration. 3 h after the chamber measurements were begun, the chambers were flushed with room air for 3 min and then resealed for a subsequent 3-h experimental period under an air atmosphere. Chamber atmosphere was sampled through a rubber septum with a gas-tight syringe and needle and was injected directly into the gas chromatograph for analysis.

Rats given phorone (2,6-dimethyl-2,5-heptadien-4-one) received 250 mg/kg i.p. 2 h before CCl₄ was given. The phorone was dissolved in corn oil and 2 ml of the mixture per kilogram was injected. Controls received corn oil injections. Liver glutathione content was reduced to 4% of control 2 h after this phorone administration, as determined in a separate experiment (data not shown).

CCl₄ was mixed with an equal volume of mineral oil and 0.5 ml of the mixture was given by stomach tube per 100 g body weight immediately before the rat was put into the metabolism chamber. CHCl₃ in chamber atmosphere was used to determine its production from CCl₄. Because CHCl₃ can be metabolized by rats, it was necessary to determine the extent to which metabolism would influence chamber CHCl₃ levels. Separate experiments were performed (results not shown). Injection of CHCl₃ into a chamber that contained a rat resulted in rapid disappearance of the CHCl₃ from the chamber atmosphere. CHCl₃ did not disappear, however, when the rat in the chamber was treated with CCl₄ simultaneously. Therefore, CHCl₃ in chamber atmosphere represented production under these conditions. The absence of metabolism was probably due to CCl₄-induced destruction of cytochrome P-450.

Assays. Liver glutathione was determined by the recirculating assay of Tietze (15) as modified by Griffith (16). Thiobarbituric acid-reactive substances were measured as described previously (17). Ethane was determined in chamber atmosphere as described previously (14). CHCl₃ was measured using a Porasil C (80:100) column in a Hewlett-Packard 5880A gas chromatograph with a flame-ionization detector. Oven temperature was 80°C. Chamber atmosphere (1 ml) was injected directly into the instrument. CHCl₃ in flask atmosphere was measured by its direct injection into the gas chromatograph. Standard curves for the chamber studies were made by dilution of CHCl₃ in glass gas vessels. Standard curves for the microsomal experiments were made by injecting CHCl₃ into 25-ml Erlenmeyer flasks that contained 5 ml of buffer.

Materials. CCl₄ and phorone were purchased from Aldrich Chemical Co., Milwaukee, WI and CHCl₃ from American Scientific Products, Houston, TX; Protosol and ¹⁴CCl₄ were supplied by New England Nuclear, Boston, MA; NADPH was purchased from Boehringer-Mannheim Biochemicals, Indianapolis, IN; GSH, disodium EDTA, and Tris were from Sigma Chemical Co., St. Louis, MO.

Statistics. Statistical significance was determined with *t* test.

Results

The O₂ tension in the rat liver varies from the equivalent of 7% of an atmosphere of O₂ to almost 0% (18). The regions around the central veins have the lowest O₂ tensions. For this study of CCl₄ metabolism by liver microsomes, we chose three different O₂ concentrations in the range found in the liver, as well as 0 and 21% O₂ (Figs. 1 and 2).

Figure 1 *A* demonstrates the effect of O₂ tension on CHCl₃




Figure 1. Effect of O₂ concentration and GSH on CHCl₃ production from CCl₄ (*A*) and CCl₄-induced lipid peroxidation (*B*) in rat liver microsomes. Open bars indicate that no GSH was present; shaded bars indicate that 1 mM GSH was present. Incubations were carried out as described in Methods. Values are mean ± 1 SD (*n* = 4 animals).

formation by microsomes. As O₂ tension increased, CHCl₃ formation decreased. Under 21% O₂, no CHCl₃ production was detected. Because CHCl₃ is derived from CCl₃, these results indicate that increasing the O₂ tension decreases CCl₃ concentration. GSH had no detectable effect on CHCl₃ formation. These results support our earlier suggestion that GSH has little or no effect on CCl₃ metabolism (11).

Fig. 1 *B* shows the effect of O₂ and GSH on CCl₄-induced lipid peroxidation. The greatest amount occurred under 1% O₂. Further increments in O₂ were associated with decrements in lipid peroxidation. No significant lipid peroxidation occurred under anaerobic conditions because the process is O₂-dependent. CHCl₃ production (Fig. 1 *A*) and lipid peroxidation (Fig. 1 *B*) were both diminished by increases in O₂ tension above 1%,




Figure 2. Effect of O₂ concentration and GSH on covalent binding of ¹⁴C derived from ¹⁴C-labeled CCl₄. Open bars indicate that no GSH was present; shaded bars indicate that 1 mM GSH was present. Incubations were carried out as described in Methods. Values are mean ± 1 SD (*n* = 4 animals).

but the diminution in CHCl_3 production was the greater. At 21% O_2 , no CHCl_3 production could be detected, yet significant lipid peroxidation occurred.

GSH protected against lipid peroxidation but did so best at the higher O_2 tensions. Lipid peroxidation was inhibited 83–88% by GSH under 3–21% O_2 , but under 1% O_2 the inhibition was only 50%. This result is consistent with the hypothesis that the GSH-dependent protection is effective against $\text{CCl}_3\text{OO}^\bullet$ but not against CCl_3^\bullet .

Fig. 2 demonstrates that covalent binding decreases as O_2 tension increases. It, like lipid peroxidation, did not fall as profoundly as did CHCl_3 production, particularly at the higher O_2 concentrations. This is illustrated in Fig. 3, which shows that the respective decreases in CHCl_3 production and in covalent binding were the same when O_2 tension rose from 0 to 1%. Successive increases in O_2 tension led to progressively more profound drops in CHCl_3 production without comparable drops in covalent binding. As O_2 tension rose from 5 to 21%, CHCl_3 production ceased completely, but covalent binding fell only by 33%. GSH inhibited covalent binding under 3–21% O_2 but had no effect on it at 0 or 1% O_2 (Fig. 2). These results are consistent with the GSH-dependent protection, being effective against $\text{CCl}_3\text{OO}^\bullet$ but not against CCl_3^\bullet .

Based on these results, raising hepatic O_2 tensions could be expected to lessen CCl_4 injury. The higher O_2 tension would inhibit the formation of CCl_3^\bullet by the cytochrome P-450 system and promote the conversion of CCl_3^\bullet formed to $\text{CCl}_3\text{OO}^\bullet$, which can be detoxified by a GSH-dependent mechanism. If this supposition is correct, raising hepatic O_2 tension with hyperbaric O_2 might constitute effective treatment for CCl_4 poisoning.

Using metabolic chambers, we studied the effect of hyperbaric O_2 on CHCl_3 production from CCl_4 in rats, and also measured ethane production as an index of lipid peroxidation. The effect of GSH depletion was examined as well.

Fig. 4 shows that hyperbaric O_2 sharply depressed CHCl_3 production by the rats. GSH depletion had no statistically significant effect on CHCl_3 production, although it appeared to augment the hyperbaric O_2 effect at 2 and 3 h. The inhibition was relieved when hyperbaric O_2 was replaced by air after 3 h. All rats produced the same amount of CHCl_3




Figure 3. Comparison of the O_2 effect on CHCl_3 production and covalent binding. Values used here were calculated from the data presented in Figs. 1 A and 2 without GSH addition. These values are ratios and indicate the effect of the indicated increment in O_2 concentration on CHCl_3 production (solid bars) and on covalent binding (shaded bars).




Figure 4. Effect of hyperbaric O_2 and GSH depletion on CCl_4 metabolism to CHCl_3 in the rat. The experiment was carried out as described in Methods. All rats received 2.5 ml CCl_4/kg at zero time and were put into the metabolism chambers. Treatments were as follows: ●, air atmosphere; ○, hyperbaric O_2 atmosphere from 0–3 h; △, air atmosphere and depleted of GSH; □, hyperbaric O_2 atmosphere from 0–3 h and depleted of GSH. All rats were in a normobaric air atmosphere from 3 to 6 h. Values shown are averages of four rats. Brackets indicate 1 SD.

from 4 to 6 h after CCl_4 administration while breathing air. The apparent differences in CHCl_3 production in the hour after the chambers were flushed were likely due to redistribution of CHCl_3 from the tissues of the animals into the chamber atmosphere. These results strongly suggest that hyperbaric O_2 diminishes CCl_3^\bullet concentration in rat liver after CCl_4 administration. This finding correlates well with the results that were obtained using the microsomal system (Fig. 1 A).

Fig. 5 shows that hyperbaric O_2 sharply depressed ethane production after CCl_4 administration, which indicated that it prevented CCl_4 -induced lipid peroxidation. GSH depletion had a striking effect on lipid peroxidation. It doubled ethane production by animals breathing air. The ethane production by animals in hyperbaric O_2 was so low, however, that no effect of GSH depletion could be seen. The hyperbaric O_2 inhibition of ethane production was not reversed when the animals were returned to an air atmosphere (Fig. 5). These in vivo experiments lend support to the hypothesis that O_2 and GSH are protective against the hepatotoxicity of CCl_4 .

Discussion

Fig. 6 is a proposed scheme of the early free radical portion of CCl_4 metabolism. It serves as a basis for discussing the present results. The conversion of CCl_4 to CCl_3^\bullet is catalyzed by cytochrome P-450 (19). O_2 inhibits this reaction. Once CCl_3^\bullet has been formed, it reacts very rapidly with O_2 to give $\text{CCl}_3\text{OO}^\bullet$ a much more reactive radical than CCl_3^\bullet (6, 7). If little or no O_2 is present, CCl_3^\bullet can bind covalently to lipids and proteins or abstract an H^\bullet , usually from a PUFA. The PUFA $^\bullet$




Figure 5. Effect of hyperbaric O_2 and GSH depletion on lipid peroxidation as measured by ethane production in the rat. These data are from the same experiment shown in Fig. 3 and the symbols used are the same.

can then react with other free radicals, which leads to polymerization, or, if any O_2 is present, PUFA* can undergo lipid peroxidation.

CCl_3OO^* is very reactive and is a good initiator of lipid peroxidation (7). It, or compounds derived from it, can bind covalently also (8). Our earlier work indicated that GSH can affect the metabolism of CCl_3OO^* , preventing membrane damage by it, but that GSH could not protect against CCl_3^* (11). Those conclusions are supported by the experiments shown in Figs. 1 and 2. GSH was more efficient in blocking CCl_4 -induced lipid peroxidation and covalent binding at the higher O_2 tensions, where CCl_3OO^* formation would be favored, than at the lower ones, where CCl_3^* concentration would be higher.

Mico and Pohl (8) have recently presented evidence that further reactions of CCl_3OO^* yield an electrophilic chlorine and phosgene, both of which are potentially toxic metabolites. The present studies were undertaken before that report and did not measure production of those metabolites. Consequently, determination of the role of GSH in their formation and detoxification will require further studies.

This scheme (Fig. 6) can potentially be used to understand the pattern of liver injury by CCl_4 . In the well-oxygenated periportal areas, cytochrome P-450 produces a small amount of CCl_3^* , which is rapidly converted to CCl_3OO^* . CCl_3OO^* is very




Figure 6. Scheme to explain O_2 and GSH effects on CCl_4 metabolism by rat liver microsomes.

reactive and may destroy the activating cytochrome P-450 by direct attack or by damage to the adjacent membrane (11, 19, 20). GSH protects the rest of the cell against CCl_3OO^* .

In the poorly oxygenated centrilobular regions of the liver, CCl_3^* is produced in large quantities by cytochrome P-450. Because O_2 tension is low, little CCl_3^* is converted to CCl_3OO^* . CCl_3^* , being less reactive than CCl_3OO^* , can diffuse farther before reacting and therefore should not destroy the cytochrome P-450 which activates CCl_4 to the extent that CCl_3OO^* does. Thus, there is much greater metabolism of CCl_4 under these conditions (11). CCl_3^* can bind covalently and can cause lipid peroxidation (if some O_2 is present). GSH does not appear to inhibit these events. Because there is an O_2 gradient between the periportal and the centrilobular regions, some areas will have O_2 tensions in which both CCl_3^* and CCl_3OO^* are present. Since the O_2 tension is relatively low, CCl_3^* may diffuse away from the activating cytochrome P-450 before reacting with O_2 , to produce the highly reactive CCl_3OO^* . This may protect the activating cytochrome P-450 and allow more CCl_4 metabolism. In those areas, perhaps analogous to the 1 and 3% O_2 in Figs. 1 and 2, GSH would be expected to exhibit significant protection. This may explain the protective effect of GSH against lipid peroxidation that was observed in air-breathing animals (Fig. 5).

The entire hepatic lobule would be well oxygenated under hyperbaric O_2 . GSH protects the membrane remote from the activating cytochrome P-450 species from CCl_3OO^* , but under well-oxygenated conditions it should have a small effect because of the small CCl_3OO^* production. This is a potential explanation for the failure of GSH depletion to cause increased ethane production in CCl_4 -treated rats under hyperbaric O_2 (Fig. 4). Also contributing to the lowering of ethane production by hyperbaric O_2 could be reaction of O_2 with the ethyl radical, as suggested by Cohen (21).

This scheme (Fig. 6) can explain why hypoxic conditions are associated with increased lipid peroxidation and hepatotoxicity from CCl_4 (22, 23). Likewise, it provides the rationale for the use of hyperbaric O_2 in the treatment of CCl_4 poisoning. European workers have shown that prolonged treatment with hyperbaric O_2 protects rats against CCl_4 -induced liver damage (24, 25), and there have been case reports of its efficacy in human CCl_4 poisoning (26). Hyperbaric O_2 should inhibit the metabolism of CCl_4 to free radicals and promote the formation of CCl_3OO^* in all parts of the liver lobule. That should result in decreased lipid peroxidation and covalent binding, but still allow destruction of the activating cytochrome P-450 with cessation of CCl_4 metabolism (11). Further studies are needed to determine the significance of CCl_4 metabolism to $CHCl_3$, which takes place under normoxic conditions even after a period of hyperbaric O_2 exposure (Fig. 4).

Hyperbaric O_2 would appear to be a logical treatment for CCl_4 poisoning. Available data (24-26) indicate it should be instituted as soon after ingestion as possible and that exposure should be maintained for as long as possible. This will allow

excretion of CCl₄ while inhibiting its conversion to injurious free radical intermediates. Conversely, hypoxia should be avoided.

These findings may have relevance to the hepatotoxicity of other agents. Halothane metabolism has recently been shown to yield a peroxy radical under aerobic conditions (7), and the GSH-dependent microsomal free radical scavenging mechanism might protect against it as well as against CCl₄.

Acknowledgments

The authors are grateful to Dr. M. A. Venkatachalam for reviewing the manuscript, to Drs. K. E. Hill and P. A. Krieter for helpful discussions, and to Mrs. R. E. Ortiz for typing the manuscript.

This work was supported by National Institutes of Health grant ES 02497 and the Robert A. Welch Foundation of Houston, TX, grant AQ-870.

References

1. Recknagel, R. O., E. A. Glende, and A. M. Hruszkewycz. 1977. Chemical mechanisms in carbon tetrachloride toxicity. In *Free Radicals in Biology*. W. A. Pryor, editor. Academic Press, Inc., New York. III:97-132.
2. Butler, T. C. 1961. Reduction of carbon tetrachloride *in vivo* and reduction of carbon tetrachloride and chloroform *in vitro* by tissues and tissue constituents. *J. Pharmacol. Exp. Ther.* 134:311-319.
3. Fowler, J. S. L. 1969. Carbon tetrachloride metabolism in the rabbit. *Br. J. Pharmacol.* 37:733-737.
4. Poyer, J. L., P. B. McCay, E. K. Lai, E. G. Jenzen, and E. R. Davis. 1980. Confirmation of assignment of the trichloromethyl radical spin adduct detected by spin trapping during ¹³C-carbon tetrachloride metabolism *in vitro* and *in vivo*. *Biochem. Biophys. Res. Commun.* 94:1154-1160.
5. Gordis, E. 1969. Lipid metabolites of carbon tetrachloride. *J. Clin. Invest.* 48:203-209.
6. Packer, J. E., T. F. Slater, and R. L. Willson. 1978. Reactions of the carbon tetrachloride-related peroxy free radical with amino acids: pulse radiolysis evidence. *Life Sci.* 23:2617-2620.
7. Forni, L. G., J. E. Packer, T. F. Slater, and R. L. Wilson. 1983. Reaction of the trichloromethyl and halothane-derived peroxy radicals with unsaturated fatty acids: a pulse radiolysis study. *Chem.-Biol. Interact.* 45:171-177.
8. Mico, B. A., and L. R. Pohl. 1983. Reductive oxygenation of carbon tetrachloride: trichloromethylperoxyl radical as a possible intermediate in the conversion of carbon tetrachloride to electrophilic chlorine. *Arch. Biochem. Biophys.* 225:596-609.
9. Kubic, V. L., and M. W. Anders. 1981. Mechanism of the microsomal reduction of carbon tetrachloride and halothane. *Chem.-Biol. Interact.* 34:201-207.
10. Nastainczyk, W., H. J. Ahr., and V. Ullrich. 1982. The reductive metabolism of halogenated alkanes by liver microsomal cytochrome P-450. *Biochem. Pharmacol.* 31:391-396.
11. Burk, R. F., K. Patel, and J. M. Lane. 1983. Reduced glutathione protection against rat liver microsomal injury by carbon tetrachloride: dependence on O₂. *Biochem. J.* 215:441-445.
12. Lawrence, R. A., and R. F. Burk. 1978. Species, tissue, and subcellular distribution of non selenium-dependent glutathione peroxidase activity. *J. Nutr.* 108:211-215.
13. Burk, R. F., and B. S. S. Masters. 1975. Some effects of selenium deficiency on the hepatic microsomal cytochrome P-450 system in the rat. *Arch. Biochem. Biophys.* 170:124-131.
14. Burk, R. F., and J. M. Lane. 1979. Ethane production and liver necrosis in rats after administration of drugs and other chemicals. *Toxicol. Appl. Pharmacol.* 50:467-478.
15. Tietze, F. 1969. Enzymatic method for quantitative determination of nanogram amounts of total and oxidized glutathione: applications to mammalian blood and other tissues. *Anal. Biochem.* 27:502-522.
16. Griffith, O. W. 1980. Determination of glutathione and glutathione disulfide using glutathione reductase and 2-vinylpyridine. *Anal. Biochem.* 106:207-212.
17. Burk, R. F., M. J. Trumble, and R. A. Lawrence. 1980. Rat hepatic cytosolic glutathione-dependent enzyme protection against lipid peroxidation in the NADPH-microsomal lipid peroxidation system. *Biochim. Biophys. Acta.* 618:35-41.
18. Lubbers, D. W. 1977. Quantitative measurement and description of oxygen supply to the tissue. In *Oxygen and Physiological Function*. F. F. Jobsis, editor. Professional Information Library, Dallas. 254-276.
19. Noguchi, T., K.-L. Fong, E. K. Lai, S. S. Alexander, M. M. King, L. Olson, J. L. Poyer, and P. B. McCay. 1982. Specificity of a phenobarbital-induced cytochrome P-450 for metabolism of carbon tetrachloride to the trichloromethyl radical. *Biochem. Pharmacol.* 31:615-624.
20. Frank, H., H. J. Hausemann, and H. Remmer. 1982. Metabolic activation of carbon tetrachloride: induction of cytochrome P-450 with phenobarbital or 3-methylcholanthrene and its effect on covalent binding. *Chem.-Biol. Interact.* 40:193-208.
21. Cohen, G. 1982. Production of ethane and pentane during lipid peroxidation: biphasic effect of oxygen. In *Lipid Peroxides in Biology and Medicine*. Yagi, K. editor. Academic Press, Inc., New York. 199-212.
22. Kieczka, H., and H. Kappus. 1980. Oxygen dependence of CCl₄-induced lipid peroxidation *in vitro* and *in vivo*. *Toxicol. Lett.* 5:191-196.
23. Shen, E. S., V. F. Garry, and M. W. Anders. 1982. Effect of hypoxia on carbon tetrachloride hepatotoxicity. *Biochem. Pharmacol.* 31:3787-3793.
24. Rapin, M., C. Got, J. R. LeGall, and M. Goulon. 1967. Effet de l'oxygene hyperbare sur la toxicite hepatique du tetrachlorure de carbone chez le rat. *Rev. Franc. Etudes Clin. Biol.* 12:594-599.
25. Montani, S., and C. Perret. 1967. Oxygenation hyperbare dans l'intoxication experimentale au tetrachlorure de carbone. *Rev. Franc. Etudes Clin. Biol.* 12:274-278.
26. Truss, C. D., and P. G. Killenberg. 1982. Treatment of carbon tetrachloride poisoning with hyperbaric oxygen. *Gastroenterology.* 82:767-769.