

Bradford C. Berk, Gino Vallega, A. J. Muslin, H. M. Gordon, Mitzy Canessa, and R. Wayne Alexander
The Journal of Clinical Investigation, Volume 83, No. 3, March 1989

Page 823.

The description for part B of Table I should correctly state: B. Cell number 24 h after logarithmic growth no./dish $\times 10^{-4}$, as shown below

Table I. Growth Responses of SHR and WKY VSMC

A. [³ H]Thymidine incorporation (% increase over 0.4% serum control)			
	SHR	WKY	SHR/WKY
≤ Subculture 6	589	200	2.95
	135	56	2.41
	<u>594</u>	<u>317</u>	<u>1.87</u>
	439±264	191±131	2.41±0.54*
> Subculture 7	213	265	0.80
	149	204	0.73
	102	141	0.72
	<u>253</u>	<u>278</u>	<u>0.91</u>
	179±67	222±63	0.79±0.09
B. Cell number 24 h after logarithmic growth (no./dish $\times 10^{-4}$)			
	SHR	WKY	
≤ Subculture 6	97±31 (n = 3)*	30±20 (n = 3)	
> Subculture 7	154±29 (n = 3)	108±35 (n = 3)	

SHR and WKY cells were incubated in six-well dishes (Costar Corp.) at $1-2 \times 10^4$ cells/well. [³H]Thymidine incorporation was determined on cells made quiescent by 48 h incubation in 0.4% serum-DME during the logarithmic growth phase. Cell counts were performed on cells 24 h after the end of logarithmic growth. Values are the mean±SD of three wells and represent data obtained from serial passage (3–23) of one set of primaries. Similar data were observed with one other set of primaries. (**P* < 0.05 vs. WKY.)

Page 824.

In the legend to Figure 1 a sentence is missing that may cause confusion in interpretation of the figure. The correct legend is as follows:

Figure 1. Proliferation of cultured SHR and WKY VSMC stimulated by 10% serum. Cells were replicate plated at 2×10^4 cells per 35-mm dish in 10% calf serum-DME. 2 d later, the medium was changed to DME with 10% calf serum. Cell counts are shown beginning with addition of 10% calf serum. Each data point is the mean±SD from three dishes.