

Disulfide-linked $\gamma\delta$ T Cell Antigen Receptors Expressed on T Cells Derived from Patients with Primary Immunodeficiency Disorders

Hidetoshi Seki,* Masanobu Nanno,* Noorbibi K. Day,† Robert A. Good,† and Chris D. Platsoucas*

*Department of Immunology, The University of Texas M.D. Anderson Cancer Center, Houston, Texas 77030; and

†All Children's Hospital, Department of Pediatrics, University of South Florida, St. Petersburg, Florida 33701

Abstract

T cell lines or clones from two patients, one with a partial DiGeorge syndrome and one with severe common variable immunodeficiency expressed disulfide-linked $\gamma\delta$ T cell antigen receptor (TCR) comprised of a γ -chain polypeptide of 40–43 kD, and a δ -chain polypeptide of 37–40 kD. This $\gamma\delta$ TCR appears to be similar to that found on T cell clones, and lines derived from peripheral blood lymphocytes from normal donors. Previous studies have shown that T cell lines derived from the peripheral blood of patients with immunodeficiency disorders express non-disulfide-linked $\gamma\delta$ TCR. In contrast to the latter and coincident with findings in the present study, the vast majority of T cell lines and clones derived from the peripheral blood of normal donors express disulfide-linked $\gamma\delta$ TCR.

Introduction

The $\gamma\delta$ T cell antigen receptor (TCR)¹ is comprised of a γ - and a δ -chain polypeptide associated either noncovalently (non-disulfide-linked TCR) or by a disulfide bond (disulfide-linked TCR) (1). Although only one δ -chain constant region gene segment has been identified (2–4), three different forms of the γ -chain have been demonstrated (1, 5–12). These are: (a) A disulfide-linked γ -chain polypeptide of 40 kD, coded by the C γ 1 gene segment, which is expressed on T cells derived from the peripheral blood or thymocytes from normal donors (1, 9, 10). (b) A non-disulfide-linked γ -chain polypeptide of 40 kD, coded by a C γ 2 gene segment containing two copies of the CII exon and expressed on T cells derived from the peripheral blood of patients with primary immunodeficiency disorders (8), certain normal donors (1, 11, 12) and on a thymocyte clone (6). (c) A non-disulfide-linked γ -chain polypeptide of 55–60 kD, coded by a C γ 2 gene segment containing three copies of the CII exon, and expressed primarily on T cells derived from patients with immunodeficiency disorders (5, 8), and the Peer (13) and Molt-17 (14) tumor cell lines. Our pre-

vious studies (8) and studies of others (1, 5) have shown that T cell lines derived from the peripheral blood of patients with primary immunodeficiency disorders expressed non-disulfide-linked $\gamma\delta$ TCRs. In contrast, the vast majority of T cell clones or lines derived from the peripheral blood of normal donors expressed disulfide-linked $\gamma\delta$ TCR. We report here that T cell lines and clones derived from the peripheral blood from two patients with partial DiGeorge syndrome or a severe form of common variable immunodeficiency expressed disulfide-linked $\gamma\delta$ TCR.

Methods

Patients. Peripheral blood lymphocytes from a patient with partial DiGeorge syndrome and a patient with severe common variable immunodeficiency (CVI) were used in this study. The patient with partial DiGeorge syndrome had characteristic dysmorphic facial features, no thymic shadow, and only very low levels of thymulin (15). The patient with severe CVI had significant T cell immunologic abnormalities, besides antibody deficiency.

Monoclonal antibodies. OKT4 and OKT8 MAb were obtained from Ortho Diagnostic Systems Inc. (Raritan, NJ). Anti- δ TCS1 MAb, anti-TCR δ 1 MAb, FITC-conjugated anti- δ TCS1 MAb, and FITC-conjugated anti-TCR δ 1 MAb were purchased from T Cell Sciences (Cambridge, MA). Cells stained with the anti- δ TCS1 MAb or with the anti-TCR δ 1 MAb were designated in this study as δ TCS1⁺ or δ 1⁺, respectively. Anti-Leu 4, phycoerythrin-conjugated anti-Leu 4, and FITC-conjugated anti-WT31 MAbs were purchased from Becton-Dickinson (Sunnyvale, CA). An anti- γ chain specific MAb, designated 3D5, was developed in this laboratory by immunizing mice with a hybrid protein containing a large part of the CII exon of the human C γ 2 gene segment (16). This MAb immunoprecipitates all three forms of γ -chain polypeptides from denatured lysates. It recognizes both the C γ 1 and C γ 2 gene segments. It is γ -chain specific and does not immunoprecipitate $\alpha\beta$ TCR (16).

T cell lines and clones. Nonadherent peripheral blood lymphocytes from patients with immunodeficiency, or from normal donors were prepared as previously described (17). The cells were treated with OKT4 and OKT8 MAbs, plus rabbit complement as described (17). CD4⁺CD8[−] lymphocytes were isolated by centrifugation on a Ficoll-Hypaque density cushion and were found to contain 40–60% CD3⁺ cells, ~ 5% WT31⁺ cells, and 3% CD4⁺ or CD8⁺ cells. To generate T cell lines, the cells were cultured at a concentration of 1×10^6 /ml in 24-well tissue culture plates in RPMI 1640 supplemented with 10% fetal calf serum, 25 mM Hepes buffer, 2 mM L-glutamine, 100 μ g/ml streptomycin, and 100 U/ml penicillin (all from Gibco Laboratories, Grand Island, NY), 100 U/ml of highly purified recombinant human interleukin-2 (rIL-2) (Cetus Corp., Emeryville, CA), 5 μ g/ml of PHA, and irradiated (5,000 rad) allogeneic peripheral blood mononuclear cells as feeders (2×10^6 /well). The cultures were fed every 4 d. T cell lines were harvested 4–5 wk later. To generate T cell clones, purified CD4⁺CD8[−] peripheral blood lymphocytes were stimulated under lim-

Address reprint requests to Dr. Chris D. Platsoucas, Department of Immunology, Box 178, M.D. Anderson Cancer Center, 1515 Holcombe Blvd., Houston, TX 77030.

Received for publication 22 June 1989.

1. Abbreviations used in this paper: CVI, common variable immunodeficiency; TCR, T cell antigen receptor.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/89/11/1679/04 \$2.00

Volume 84, November 1989, 1679–1682

iting dilution conditions (1, 3, 10, or 30 cells per well) with irradiated (5,000 rad) allogeneic mononuclear leukocytes from a normal donor (100,000 cells per well), rIL-2 (100 U/ml), and PHA (5 μ g/ml), as previously described (12). Clones derived from these cultures were expanded for analysis.

Immunofluorescence analysis. Cell surface immunofluorescence was carried out as previously described (18).

Cell-mediated cytotoxicity. Cell-mediated cytotoxicity was determined by a 51 Cr-release assay as described (18).

Immunoprecipitation. Cells were labeled with Na 125 I (14.8 mCi/ μ g sp act of iodine) by the lactoperoxidase method and immunoprecipitations were carried out as previously described (12). For immunoprecipitations with the anti- γ chain specific 3D5 MAb, lysates were denatured and alkylated as described previously (12, 16). Samples were analyzed by SDS/PAGE, either under reducing or nonreducing conditions.

Results

Two CD4 $^{+}$ CD8 $^{-}$ T cell lines exhibiting substantial proportions of cells expressing $\gamma\delta$ TCR were developed by expanding in culture purified CD4 $^{+}$ CD8 $^{-}$ peripheral blood lymphocytes from a patient with severe CVI and a patient with partial DiGeorge syndrome. Immunofluorescence analysis of these lines is shown in Fig. 1. The CVI cell line was 99% CD3 $^{+}$, 86% δ 1 $^{+}$, 25% δ TCS1 $^{+}$, and 11% WT31 $^{+}$. The DiGeorge line was 99% CD3 $^{+}$, 51% δ 1 $^{+}$, 43% δ TCS1 $^{+}$, and 39% WT31 $^{+}$ (Fig. 1). Two T cell clones were derived from CD4 $^{+}$ CD8 $^{-}$ cells from the patient with partial DiGeorge Syndrome, and both were CD3 $^{+}$ CD4 $^{+}$ CD8 $^{-}$ WT31 $^{+}$ δ TCS1 $^{+}$ δ 1 $^{+}$. These T cell lines and clones exhibited MHC unrestricted cytotoxicity and lysed cells of the K562 cell line and of the MEL21 melanoma tumor cell line (Table I). Cytotoxicity by the DiGeorge T cell clones against the K562 targets was enhanced by the OKT3 MAb.

Immunoprecipitation of 125 I-labeled lysates of the DiGeorge T cell line by anti-Leu 4 MAb, followed by SDS/PAGE analysis under nonreducing conditions revealed polypeptide bands in the range of 83–93 kD (in addition to the CD3 bands of 20–28 kD). SDS/PAGE analysis under reducing conditions demonstrated two polypeptide chains with molecular masses of 40 and 36 kD, in addition to the CD3 bands (Fig. 2).

Immunoprecipitation, by the γ -chain specific 3D5 MAb, of nonreduced denatured lysates from two T cell clones (CD3 $^{+}$ WT31 $^{+}$ CD4 $^{+}$ CD8 $^{-}$ δ TCS1 $^{+}$ δ 1 $^{+}$) derived from CD4 $^{+}$ CD8 $^{-}$ peripheral blood lymphocytes from the patient with partial DiGeorge syndrome, demonstrated that they also ex-




Figure 1. Immunofluorescence analysis of T cell lines derived from CD4 $^{+}$ CD8 $^{-}$ peripheral blood lymphocytes from a patient with severe CVI and a patient with partial DiGeorge syndrome. The cells were stained with FITC-conjugated WT31; FITC-conjugated anti- δ TCS1; FITC-conjugated anti-TCR δ 1, or phycoerythrin-conjugated anti-Leu 4. Control cells were stained with FITC-conjugated goat anti-mouse IgG antibody.

Table I. Cytolytic Activity of $\gamma\delta$ TCR T Cell Lines or Clones from Patients with Immunodeficiency Diseases

Line or clone	% 51 Cr release (E/T = 20:1)			
	K562		MEL21	
	Medium	Anti-CD3	Medium	Anti-CD3
CVI T cell line	79.4 \pm 5.2	86.5 \pm 7.9	50.7 \pm 2.4	25.4 \pm 2.5
DiGeorge T cell line	53.7 \pm 2.1	ND	19.7 \pm 3.0	ND
DiGeorge T cell clone	28.7 \pm 2.1	85.6 \pm 1.6	2.1 \pm 0.3	6.1 \pm 6.0
DiGeorge T cell clone	15.6 \pm 4.1	63.6 \pm 4.1	0.1 \pm 1.2	4.8 \pm 1.6

ND, not done.

pressed disulfide-linked $\gamma\delta$ TCR. SDS-PAGE analysis under nonreducing conditions revealed on both clones polypeptides of 81 kD, which were resolved to two polypeptide bands of 37 and 40 kD under reducing conditions (Fig. 2). These two T cell clones were δ TCS1 $^{+}$, demonstrating that both disulfide- and non-disulfide-linked $\gamma\delta$ TCR can express the δ TCS1 determinant (12).

Biochemical analysis of the T cell line CD3 $^{+}$ WT31 $^{+}$ CD4 $^{+}$ CD8 $^{-}$ δ 1 $^{+}$ derived from the patient with severe CVI demonstrated that it also expressed a disulfide-linked $\gamma\delta$ TCR. Im-




Figure 2. Cells from a T cell line and two T cell clones (CD3 $^{+}$ WT31 $^{+}$ CD4 $^{+}$ CD8 $^{-}$ δ TCS1 $^{+}$ δ 1 $^{+}$) derived from peripheral blood lymphocytes from a patient with partial DiGeorge syndrome were labeled with 125 I and lysed in 5 mM 3[(3-cholamidopropyl)dimethyl-ammonio]1-propanesulfonate lysis buffer (lanes 1 and 2), or 1% NP-40 lysis buffer (lanes 3–6). Lysates were immunoprecipitated under nonreducing conditions with mouse IgG (lanes 1, 3, and 5), anti-Leu 4 MAb (lane 2), or anti- γ chain 3D5 MAb (lanes 4, and 6). Immunoprecipitations with the anti- γ chain 3D5 MAb were carried out after denaturation of the lysates as described in Methods. Samples were solubilized in SDS sample buffer with β -mercaptoethanol (reducing conditions, R) or without (nonreducing conditions, NR), and subjected to electrophoresis on 12.5% SDS-PAGE gels.




Figure 3. Cells from a T cell line derived from peripheral blood lymphocytes from a patient with severe CVI were labeled with ^{125}I and lysed in 1% NP-40 lysis buffer. Lysates were immunoprecipitated under nonreducing conditions with mouse IgG (lane 1), or anti- γ chain 3D5 MAb (lane 2). Immunoprecipitations with the anti- γ 3D5 MAb were carried out after denaturation of the lysates as described in Methods. Samples were solubilized in SDS sample buffer with β -mercaptoethanol (reducing conditions, R) or without (nonreducing conditions, NR), and subjected to electrophoresis on 12.5% SDS-PAGE gels.

munoprecipitation of denatured nonreduced lysates by the anti- γ 3D5 MAb followed by SDS-PAGE analysis under nonreducing conditions revealed a band of 81 kD. Under reducing conditions, two bands of 40 and 43 kD were observed (Fig. 3).

Discussion

T cell lines and clones derived from peripheral blood lymphocytes from a patient with partial DiGeorge syndrome and a patient with severe CVI expressed disulfide-linked $\gamma\delta$ TCR. These $\gamma\delta$ TCR appeared to use a γ -chain polypeptide of ~ 40 – 43 kD and a δ -chain of 37 – 40 kD and appeared to be similar to the $\gamma\delta$ TCR found on T cell lines or clones derived from normal donors. T cell lines derived from the peripheral blood of three patients with primary immunodeficiency disorders expressed exclusively non-disulfide linked $\gamma\delta$ TCR using either a 40-kD or a 55–60-kD γ -chain polypeptide (8). Similarly, Brenner et al. (1, 5) observed only non-disulfide-linked γ -chain polypeptides on T cell lines derived from the peripheral blood of certain patients with primary immunodeficiency disorders.

Our results suggest that T cell lines derived from peripheral blood lymphocytes from patients with primary immunodeficiency disorders can also express disulfide-linked $\gamma\delta$ TCR.

Our findings (12, and unpublished observations), and those of others (19) suggest that ~ 70 – 80% of the $\gamma\delta$ TCR T cell clones or lines derived from peripheral blood lymphocytes from normal donors expressed disulfide-linked $\gamma\delta$ TCR. The remaining clones expressed non-disulfide linked $\gamma\delta$ TCR, using the 40 kD C γ 2 γ -chain polypeptide (11, 12, 19). $\gamma\delta$ TCR expressing T cells derived from normal donors, and using the 55–60-kD non-disulfide-linked γ -chain polypeptide have been rare. Moretta et al. (19) have reported the utilization of the 55–60-kD γ -chain by 2% of the $\gamma\delta$ TCR T cell clones from normal donors that they examined. Also, they found a correlation of the expression of the CD8 T cell differentiation antigen with the 55–60-kD non-disulfide-linked γ -chain polypeptide. We have also obtained one $\gamma\delta$ TCR expressing T cell clone derived from peripheral blood lymphocytes of a normal donor that used the 55–60-kD non-disulfide-linked γ -chain polypeptide. These cells were CD8 $^{-}$ (manuscript in preparation).

The T cell line and the T cell clones from the patient with the partial DiGeorge syndrome expressed the δ TCS1 determinant. Similarly, $\sim 25\%$ of the $\gamma\delta$ TCR-expressing cells of the T cell line from the patient with severe CVI were δ TCS1 $^{+}$. All these T cell lines and clones expressed disulfide-linked $\gamma\delta$ TCR in agreement with our previous findings (12) and in contrast to those of Bottino et al. (20), who reported that the δ TCS1 determinant is expressed only on non-disulfide-linked $\gamma\delta$ TCR.

The function of the $\gamma\delta$ TCR is not known at the present. $\gamma\delta$ TCR using different constant region gene segments may have different functions in T cell differentiation or may be responsive to different antigenic stimuli. Different γ -chain constant region gene segments may be preferentially associated with different variable (V γ) and joining (J γ) segments. Identification of such a preferential association will indicate different functions for $\gamma\delta$ TCR using C γ 1, and C γ 2. Additional studies are needed to answer these questions.

Acknowledgments

This work was supported in part by grants AI-24669 and AG-05628 from the National Institutes of Health, grant CH-420 from the American Cancer Society, and a grant from the Eleanor Naylor Dana Charitable Trust.

References

1. Brenner, M. B., J. McLean, H. Scheft, J. Riberdy, S.-L. Ang, J. G. Seidman, P. Devlin, and M. S. Krangel. 1987. Two forms of the T-cell receptor γ protein found on peripheral blood cytotoxic T lymphocytes. *Nature (Lond.)*. 325:689–694.
2. Chien, Y.-H., M. Iwashima, K. B. Kaplan, J. F. Elliot, and M. M. Davis. 1987. A new T-cell receptor gene located within the alpha locus and expressed early in T cell differentiation. *Nature (Lond.)*. 327:677–682.
3. Hata, S., M. B. Brenner, and M. S. Krangel. 1987. Identification of putative human T cell receptor δ complementary DNA clones. *Science (Wash. DC)*. 238:678–682.
4. Loh, E. Y., L. L. Lanier, C. W. Turan, D. R. Littman, M. M. Davis, Y.-H. Chien, and A. Weiss. 1987. Identification and sequence of a fourth human T cell antigen receptor chain. *Nature (Lond.)*. 330:569–572.

5. Brenner, M. B., J. McLean, D. P. Dialynas, J. L. Strominger, J. A. Smith, F. L. Owen, J. G. Seidman, S. Ip, F. Rosen, and M. S. Krangel. 1986. Identification of a putative second T-cell receptor. *Nature (Lond.)* 322:145-149.
6. Bank, I., R. A. DePinho, M. B. Brenner, J. Cassimeris, F. W. Alt, and L. Chess. 1986. A functional T3 molecule associated with a novel heterodimer on the surface of immature human thymocytes. *Nature (Lond.)* 322:179-181.
7. Moingeon, P., A. Ythier, G. Goubin, F. Faure, A. Nowill, L. Felmon, M. Rainaud, F. Forestier, F. Daffos, C. Bohuon, and T. Hercend. 1986. A unique T-cell receptor complex expressed on human fetal lymphocytes displaying natural-killer-like activity. *Nature (Lond.)* 323:638-640.
8. Ioannides, C. G., K. Itoh, F. E. Fox, R. Pahwa, R. A. Good, and C. D. Platsoucas. 1987. Identification of a second T-cell antigen receptor in human and mouse by an anti-peptide γ -chain-specific monoclonal antibody. *Proc. Natl. Acad. Sci. USA* 84:4244-4248.
9. Borst, J., R. J. vandeGriend, J. W. Oostveen, S.-L. Ang, C. J. Melief, J. G. Seidman, and R. L. H. Bolhuis. 1987. A T-cell receptor γ /CD3 complex found on cloned functional lymphocytes. *Nature (Lond.)* 325:683-688.
10. Lanier, L. L., N. A. Federspiel, J. J. Ruitenbergh, J. H. Phillips, J. P. Allison, D. Littman, and A. Weiss. 1987. The T cell antigen receptor complex expressed on normal peripheral blood CD4⁺CD8⁻ T lymphocytes. A CD3-associated disulfide-linked γ chain heterodimer. *J. Exp. Med.* 165:1076-1094.
11. Hochstenbach, F., C. Parker, J. McLean, V. Gieselmann, H. Band, I. Bank, L. Chess, H. Spits, J. L. Strominger, J. G. Seidman, and M. B. Brenner. 1988. Characterization of a third form of the human T cell receptor γ/δ . *J. Exp. Med.* 168:761-776.
12. Seki, H., M. Nanno, P.-F. Chen, K. Itoh, C. G. Ioannides, R. A. Good, and C. D. Platsoucas. 1989. Molecular heterogeneity of $\gamma\delta$ T-cell antigen receptors expressed by CD4⁺CD8⁻ T-cell clones from normal donors. *Proc. Natl. Acad. Sci. USA* 86:2326-2330.
13. Weiss, A., M. Newton, and D. Crommie. 1986. Expression of T3 in association with a molecule distinct from the T-cell antigen receptor heterodimer. *Proc. Natl. Acad. Sci. USA* 83:6998-7002.
14. Tighe, L., A. Foster, D. M. Clar, A. W. Boylston, I. Lavenir, and T. H. Rabbitts. 1987. Unusual forms of T cell γ mRNA in a human T cell leukemia cell line: implications for γ gene expression. *Eur. J. Immunol.* 17:1729-1736.
15. Dardenne, M., J. M. Pleau, B. Nabaira, P. Lefrancier, M. Derrien, J. Choay, and J. F. Bach. 1982. Concentration of zinc and other metals to the biological activity of serum thymic factor. *Proc. Natl. Acad. Sci. USA* 79:5370-5374.
16. Nanno, M., H. Seki, Y. Bao, C. Ioannides, J. Morkowski, and C. D. Platsoucas. 1989. Development of a monoclonal antibody specific for the gamma chain of the T-cell antigen receptor using an open reading frame expression vector. *Hybridoma* 8:277-291.
17. Platsoucas, C. D., and R. A. Good. 1981. Inhibition of cell-mediated cytotoxicity by monoclonal antibodies to human T-cell differentiation antigens. *Proc. Natl. Acad. Sci. USA* 78:4500-4504.
18. Platsoucas, C. D. 1984. Human T cell antigens involved in cytotoxicity against allogeneic or autologous chemically modified targets. Association of the Leu 2a/T8 antigen with effector-target cell binding and of the T3/Leu 4 antigen with triggering. *Eur. J. Immunol.* 14:566-577.
19. Moretta, A., C. Bottino, E. Ciccone, G. Tambussi, M. C. Mingari, S. Ferrini, G. Casorati, P. Varese, O. Viale, M. Migone, and L. Moretta. 1988. Human peripheral blood lymphocytes bearing T cell receptor γ/δ . Expression of CD8 differentiation antigen correlates with the expression of the 55-kD, C γ 2-encoded γ chain. *J. Exp. Med.* 168:2349-2354.
20. Bottino, C., G. Tambussi, S. Ferrini, E. Ciccone, P. Varese, M. C. Mingati, L. Moretta, and A. Moretta. 1988. Two subsets of human T lymphocytes expressing γ/δ antigen receptors are identifiable by monoclonal antibodies directed to two distinct molecular forms of the receptor. *J. Exp. Med.* 168:491-505.