Characterization of the Immunoglobulin Heavy Chain Complementarity Determining Region (CDR)-III Sequences from Human B cell Precursor Acute Lymphoblastic Leukemia Cells

Hitoshi Kiyoi,* Tomoki Naoe,* Keizo Horibe,† and Ryuzo Ohno*

*Department of Medicine, the Branch Hospital Nagoya University School of Medicine, Higashi-ku, Nagoya 461, Japan; and †Department of Pediatrics, Nagoya University School of Medicine, Showa-ku, Nagoya 466, Japan

Abstract

Sequence analysis of the immunoglobulin heavy chain complementarity determining region (CDR)-III of B-lineage cells at various stages has provided important insights concerning B cell maturation and selection. Knowledge of human CDR-III sequences has been relatively limited compared with that of the murine system. We analyzed the CDR-III sequences of B cell precursor acute lymphoblastic leukemia (pre-B ALL) cells in 23 newly diagnosed and 10 relapsed patients, in order to elucidate the organization of CDR-III in B cell precursors. We found a very low frequency of somatic mutations in D and J_H regions, preferential use of D_{LR}, D_{XP}, D_{HQ52}, and D_N elements, and of 3' side J_H segments, and no predominant usage of D coding frames. Unusual joinings such as V_H-D-D-J_H and V_H-J_H were observed in three, and one sequences, respectively. We compared the CDR-III sequences derived from 10 patients between diagnosis and relapse. Two of them had three spots of mutated nucleotides at relapse, all of which were found in the N region near the D segments. Our data showed the possibility of somatic mutation at relapse, in addition to developmentally regulated rearrangement of the immunoglobulin gene at the stage of B cell precursors. (*J. Clin. Invest.* 1992; 89:739-746.) **Key words:** immunoglobulin • CDR-III • pre-B ALL • somatic mutation • PCR

Introduction

The immunoglobulin heavy chain (IgH)¹ variable region is encoded by three separate genes: variable (V_H), diversity (D), and joining (J_H) segments. During B lymphocyte differentiation, the IgH gene is assembled in two steps. D and J_H segments are joined first, followed by V_H to D-J_H joining. During these joining steps, antibody diversity is generated by the following mechanisms:

(a) somatic recombination of multiple V_H, D, and J_H segments; (b) variation of joining sites and addition of the N region nucleotides; and (c) somatic mutation (1-3).

The assembled variable region of the Ig gene contains three hypervariable regions surrounded by framework regions that are relatively well conserved. These regions are called complementarity determining regions (CDR)-I, II, and III, and are thought to contact the antigenic epitope directly (1-4). CDR-I and II are encoded by a V_H segment, and CDR-III is encoded by D elements, V_H-D and D-J_H junctional segments. As a result of these mechanisms, enormous diversity of CDR-III is created from a limited number of germline D segments (1-3).

In murine system, recent studies have demonstrated several characteristics of the IgH gene at each stage of B cell differentiation. First, the preferential use of the most 5' D segments, and the most J_H-proximal D segments, is developed during initial D to J_H joining in immature B cells (5). Second, the CDR-III sequences from fetal and neonatal B-lineage cells preferentially use the J_H-proximal V_H genes (6-9). In contrast, adult B-lineage cells show random V_H utilization (9-13). Third, N region nucleotides are thought to be added by the enzyme terminal deoxynucleotidyl transferase (TdT) (14-17), and very few N region nucleotides are observed in most fetal and neonatal V_H-D-J_H junctional sequences. The addition of N regions thus appears to be a developmentally regulated process in B cells (18-20). Fourth, one particular reading frame of the D elements is preferentially used in productively rearranged V_H-D-J_H genes, and other reading frames are used at the stage of D-J_H joining, although the joining border varies in each clone (21, 22). It is suggested that the selection of reading frames is developmentally regulated at the stage of B cell ontogeny.

In the human system, D_{HQ52}-J1 joining was frequently observed in Epstein-Barr virus-transformed fetal B cell lines (23). Another study showed the biased use of J_H segments (J3 > J4 > J5 > J1, J2) and the D_{HQ52} element in fetal liver (24). Recently Yamada et al. reported that 3' side J_H segments (J4, J5, and J6) were preferentially used, and all coding frames of germline D genes were used to generate CDR-IIIs in adult human peripheral B cells (25). Until now, molecular studies of human B-lineage malignancies at various stages have provided many insights into B cell ontogeny. For example, follicular B cell lymphoma has extensive somatic mutations in the V_H regions (26-28), although B cell precursor leukemia has few mutations (29).

In this study, in order to elucidate the characteristics of CDR-III at the stage of B cell precursors, we have analyzed the CDR-III sequences from human B cell precursor acute lymphoblastic leukemia (pre-B ALL) cells using the polymerase chain reaction (PCR) amplification and a fluorescent chain-terminating dideoxy-nucleotide sequencing system. We demonstrate here the structure of the N region, the low frequency of somatic

Address correspondence and reprint requests to Tomoki Naoe, Department of Medicine, the Branch Hospital Nagoya University School of Medicine, 1-1-20, Daiko-Minami, Higashi-ku, Nagoya 461, Japan

Received for publication 19 March 1991 and in revised form 13 September 1991.

1. *Abbreviations used in this paper:* CDR, complementarity determining region; D, diversity segment; dd-NTP-Fs, four dideoxynucleotide terminators; IgH, immunoglobulin heavy chain; J_H, joining segment; P, palindromic; PCR, polymerase chain reaction; pre-B ALL, B cell precursor acute lymphoblastic leukemia; SSC, standard saline citrate; Tdt, terminal deoxynucleotidyl transferase; V_H, variable segment.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/92/03/0739/08 \$2.00

Volume 89, March 1992, 739-746

mutations in the D and J_H regions, and the segments in the D and J_H genes which are preferentially used. In addition, we compared the CDR-III sequences between initial diagnosis and relapse, and found the possibility of somatic mutation in pre-B ALL cells at relapse.

Methods

Leukemia cells. Bone marrow samples were taken from 23 patients (13 children and 10 adults) with pre-B ALL at the time of initial diagnosis or relapse, after informed consent was obtained. All bone marrow samples contained over 90% leukemia cells. The diagnosis of ALL in each patient was confirmed by standard morphological and immunophenotyping studies. All leukemia cells had an L1 or L2 French-American-British (FAB) subtype, expressed HLA-DR, CD10, and CD19 antigens, and lacked immunophenotypic evidence of both T (CD1, CD2, CD3, CD4, CD5, CD7, CD8) and mature B-cell lineage (CD20 and surface Ig). The expression of cytoplasmic μ chain and TdT was examined by an immunofluorescence test. Karyotypic analysis revealed that five patients (patients S.B., S.I., F.K., S.M., and C.S.) had the Philadelphia chromosome (Ph¹). In the 10 relapsed patients, the morphological, immunophenotypical, and karyotypical features of their leukemia cells were identical with those at initial diagnosis. These data are summarized in Table I.

Southern blot analysis. High molecular weight DNA was extracted from leukemia cells according to a previously published method (30). Rearrangements of the IgH gene were analyzed by Southern blot analysis (31), using a probe made of the subcloned 4.3-kb fragment (PstI-HindIII) from the genomic clone containing J_H region. High molecular weight DNA from pre-B ALL was digested with HindIII, EcoRI, or a combination of HindIII and BamHI restriction endonucleases (Boehringer Mannheim Yamanouchi, Tokyo, Japan), electrophoresed in an 0.8% agarose gel, and blotted onto nitrocellulose membranes. Hybridization was carried out with ³²P-labeled nick-translated probes under previously published conditions (31). Membranes were washed twice with 2× standard saline citrate (SSC), containing 0.1% SDS, and twice with 0.1× SSC, containing 0.1% SDS and exposed to x-ray films.

Oligonucleotide primers. Oligonucleotide primers for PCR amplification of CDR-III were synthesized on a DNA synthesizer (model 381-A; Applied Biosystems, Inc., Foster City, CA). The consensus sequences of the V_H and J_H segments were determined according to the published sequence data (32–34). Recently, several PCR primers for the amplification of the human IgH CDR-III have been reported (35–37). We also synthesized similar primers: Vcomm: 5'-GAGT-C-GAC(A/T)C(A/G)GC(G/C)(G/A)TGTA(T/C)T(T/A)CTG-3', and Jcomm: 5'-CCAAG-CTTACCTGAGGAGACGGTGA-3' that was designed to correspond more specifically to the published IgH sequences than those previously reported. These primers contain SalI and HindIII cloning sites, respectively, to allow ligation of the amplified sequences into recombinant vectors (38).

PCR amplification of the CDR-III sequences. PCR was essentially performed as described by Saiki et al. (39). A 50- μ l reaction mixture contained 500 ng of genomic DNA, 50 mM KCl, 10 mM Tris-HCl (pH 8.4), 1.0 to 1.5 mM MgCl₂, 100 μ g/ml gelatin, 0.25 μ M of Vcomm and Jcomm primers, 200 μ M of each deoxynucleotide triphosphate (dATP, dCTP, dGTP, and dTTP; Boehringer Mannheim Yamanouchi). The reaction mixture was first incubated at 95°C for 10 min to denature double-stranded DNA, followed by 1 min at 55–58°C to anneal primer and template. Primer extension was started by the addition of 2.5 U *Taq* polymerase (AmpliTag; Perkin-Elmer Cetus Corp., Norwalk, CT), and allowed to proceed for 1 min at 72°C. Subsequent denaturing, annealing, and extension steps were performed at 92°C for 1 min, at 55 to 58°C for 1 min and at 72°C for 1 min, respectively, for 40 cycles on a program temperature control system (model PC-700; Astec, Fukuoka, Japan).

DNA sequencing of amplified fragments. The amplified fragments were separated through 4% NuSieve GTG agarose gels (FMC BioProd-

ucts, Rockland, ME) and 12% polyacrylamide gels in TBE buffer (0.09 M Tris-borate, 0.002 M EDTA; pH 8.0). Amplified fragments were isolated from the polyacrylamide gels by the crush and soak technique, then digested with SalI and HindIII restriction endonucleases (Boehringer Mannheim Yamanouchi), and ligated to the M13mp18 phage vector (Takara, Kyoto, Japan). The ligated materials were transfected into *Escherichia coli* strain JM109. More than five recombinant plaques per one clone were picked up and cultured in 2 × TY medium (16 g/liter Bacto-tryptone; Difco Laboratories, Inc., Detroit, 10 g/liter Bacto-yeast extract; Difco Laboratories, 5 g/liter NaCl). Single-stranded phage DNA was prepared from these cultures. DNA was sequenced using fluorescent chain-terminating dideoxynucleotides on a Genesis 2000 (DuPont Co., Wilmington, DE) (40). This sequencing system uses a special set of four dideoxynucleotide terminators (dd-NTP-Fs), each having a unique member of the succinyl fluorescein family attached. Single-stranded phage DNA, M13 M1-primer (Takara), and reaction buffer were heated at 90°C for 2 min and then transferred to a 37°C water bath for 10 min to allow the primer to anneal. At this time, a single solution containing a mixture of each deoxynucleotide triphosphate (dCTP, dTTP, 7-deaza dATP, and 7-deaza dGTP), dd-NTP-Fs, DTT, and T-7 DNA polymerase (Takara) was prepared. An aliquot of this solution was added to the annealed mixture, and after a 5-min incubation, the unincorporated nucleotide, proteins, and reaction buffer were removed by ammonium acetate/ethanol precipitation. The DNA was washed with 70% ethanol and dried up, followed by resuspension in the loading solution. The mixture was then applied to the auto sequencer (Genesis 2000; DuPont Co.). The sequence data of each clone was determined if all the sequences from each recombinant plaque were identical.

Results

Southern blot analysis. All leukemia cells analyzed here demonstrated rearranged bands of the J_H region in either HindIII, EcoRI, or the combination of HindIII and BamHI digests. In the 10 relapsed patients, the rearranged patterns were identical with those at initial diagnosis (summarized on Table I).

Analysis of the CDR-III sequences from pre-B ALL. All genomic DNAs from the pre-B ALL cells were well amplified using the Vcomm and Jcomm primers. However, DNA from normal peripheral blood lymphocytes did not result in a significant band, and gave a faint smear after an additional 40-cycle PCR amplification (data not shown). In patient S.I., two amplified bands were observed that presumably corresponded to two rearranged IgH genes. We analyzed both bands and determined their CDR-III sequences.

The CDR-III sequences from the pre-B ALL cells are shown in Fig. 1. In the amplified sequences from all patients except T.E., we could determine D elements followed by J_H sequences. In patient T.E., a V_H gene joined directly to a J_H gene without a D region. Similar joining clones have been reported in peripheral B cells (25). We do not know if this joining results from deletions of the D region during V_H to D-J_H joining, or whether it reflects extensive exonuclease modification of the D region in the V_H-D-J_H gene.

D segment. All D-coding elements were tentatively assigned to germline D segments (32, 41, 42). Somatic mutations of D elements were rare (4/412; 0.97%), although three patients (K.N., N.K., and C.S.) showed mutated D elements: CA to TG in D_{N1}; A to G in D_{HQ52}; and G to C in D_{HQ52}, respectively. The DNAs obtained from these three patients at the time of both initial diagnosis and relapse were independently analyzed. The sequences of the mutated D elements were completely identical in each patient. The D element of patient K.N. may be

Table I. Clinical and Laboratory Data of Patients with Pre-B ALL

Patient	Age	Sex	D/R*	J _H (HindIII) [†]	TdT	C _μ
	yr				%	
MS	1	F	D	G/R/R	NT	NT
			R	G/R/R	NT	NT
YM	2	M	D	G/R	NT	NT
			R	G/R	NT	NT
TH	2	M	D	NT	80	—
MK	3	F	D	G/R/R	NT	—
RF	6	M	D	G/R	NT	—
SB	7	M	D	G/R	60	—
SE	9	F	D	G/R/R	NT	NT
			R	G/R/R	NT	NT
KN	10	M	D	G/R	70	—
			R	G/R	NT	—
HI	10	M	D	G/R/R	70	—
HT	10	M	D	G/R	NT	+
NK	10	M	D	G/R	NT	NT
			R	G/R	NT	NT
TY	11	M	D	G/R	30	—
			R	G/R	30	—
SH	12	F	D	G/R	NT	—
KM	18	F	D	G/R/R	NT	+
			R	G/R/R	70	+
EY	18	F	D	G/R/R	NT	—
			R	G/R/R	NT	—
YI	20	M	D	G/R	70	—
TE	24	M	D	G/R	80	—
MH	24	F	D	G/R/R	60	+
			R	G/R/R	90	+
HM	25	M	D	G/R/R	70	—
SI	47	F	D	G/R/R	80	—
SM	55	M	D	G/R	60	—
FK	57	M	D	G/R/R/R	90	+
CS	57	F	D	G/R/R	NT	—
			R	G/R/R	90	—

* D and R indicate diagnosis and relapse, respectively.

[†] Rearranged pattern of Southern blot hybridization are indicated. G, germline band; R, rearranged band. C_μ, cytoplasmic μ chain. NT, not tested.

encoded by another member of the D_N gene family, since the same sequences were reported as being frequently found in peripheral B cells (25). Similarly, the D_{K4} element in patient S.B. that was matched in a narrow region might be encoded by another member of the D_K gene family.

The frequencies of D segment use in human pre-B ALL cells are shown in Fig. 2. To clarify our data, previously described CDR-III sequences from pre-B ALL cells are also shown in Fig. 2 (29, 36, 37, 43). The D_{LR} family was most frequently used (our data: 41%, published data: 48%). The frequency of the D_{XP} family (our data: 29%, published data: 29%) was a little lower than that of the D_{LR} family. The D segment usage frequency gradient, i.e., D_{LR} > D_{XP} > D_{HQ52}, D_N > D_K, D_A > D_M, is similar to the published data (29, 36, 37, 43).

It is noteworthy that three clones contained D–D joining sequences. In patient T.Y., both sequences at the time of initial diagnosis and relapse had the same D–D joins consisting of

D_{LR4}, D_{HQ52}, and a five-nucleotide N region between these D elements. In patient S.M., D–D join consisted of D_{LR4}, D_{XP4}, and a seven-nucleotide N region between them. Also in patient H.T., D–D join consisted of D_{LR4}, and D_{XP4}, with a 12-nucleotide N region between them (Fig. 1).

Coding frame of D segment. Although all CDR-III sequences shown in Fig. 1 are not necessarily translated, nucleotide sequences corresponding to the D element are typed in triplets to identify the temporary coding frame. The frame sequence in V_H segments were decided to start with TA(T/C)T(T/A)C TGT. In Table II the coding frames of the D segment are shown. All three coding frames were equally used. Of interest was the use of the first coding frame in the D_{LR} family, which contained the stop codon, and was not used by the peripheral B cells.

J_H segment. The J_H segments of the sequenced clones were assigned to J1–J6 according to published data (32).

It could not be determined whether the J_H segment from patient F.K. was assigned to J4 or J5, because several nucleotides were deleted at the 5' coding region of the J_H segment. Somatic mutations of eight nucleotides (two nucleotides deleted, four nucleotides substituted, and two nucleotides inserted) were observed in 848 nucleotides encoded by the J_H segments. This rate (8/848; 0.94%) is almost the same as that (4/412; 0.97%) in the nucleotides encoded by D segments. Furthermore, our sequences of J4, J5, and J6 had a discrepancy of one, two, and one nucleotides, respectively, compared with the published germ line sequences (32). However, since these have been reported as germ line polymorphisms (24), we did not include these discrepancies among the numbers described above.

The frequency of J_H segment use was also examined (Fig. 3). There was marked correlation between the 3' location and J_H usage. This biased use was similar to previous reports in pre-B ALL (29, 36, 37, 43), and appeared equally in the CDR-III of both children and adults.

N region. We determined the 3' end of V_H segment in each clone according to the published V_H sequences (34). Nucleotides which could not be assigned to any V_H, D, nor J_H segments were determined as N region. The N region nucleotides of 1 to 20 bases were observed at the V_H–D junction in all clones. In three patients (T.Y., H.T., and S.M.), the N regions were surrounded by D elements as described above. At the D–J_H junction, two clones (Y.M. and K.N.) lacked a N region, and other clones had 1 to 22 nucleotides. GC nucleotides were predominant (249/432; 58%) in the N region as previously described (14, 44) (Fig. 1).

Although TdT is probably a responsible enzyme that mediates the addition of the N nucleotides (14–17), it was difficult to find a correlation between the expression of TdT in leukemia cells, and the length of the N nucleotides.

It has been proposed that palindromic (P) regions contribute to junctional diversity (45). The P region consists of dinucleotides palindromic to the coding nucleotides adjacent to the signal heptamer. We found P nucleotides in 14 out of 24 clones, as shown in Fig. 1. However, the contribution of P nucleotides to the N region formation of these clones may be difficult to assess, because there was no significant difference between the actual frequency of the inserted P nucleotides (24/96; 21%: number of the P region/number of V_H–N, N–D, D–N, and N–J_H junctions that could potentially be inserted with P nucleotides.) and that of random nucleotide insertion (25%).

Case		3' end of V	N, D, N region	5' end of J	D	J
EY	D	TGCGAGAG	gcttt ggA TTA CGA TTT TTG GAG tgtttcccttcg	TGACTACTGGGGCCAGGGAACCTGG	XP4	4
SI	D	TGCGAGAGA	tcgcggttag ggA TTA CGA TTT TTG GAG TGT cccttcacgaactcgg	TACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	XP4	6
KM	D	TGCAA	aaggg GAT TTT TGG AGT GGT TAT cccgcgcgtccg	TGACTACTGGGGCCAGGGAACCTGG	XP4	4
MS	D	TGCAAGAG	cccggg TAC GAT ATT TTG ACT GGT TAT TAa tc	CTACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	XP1	6
YI	D	TGCGAGAG	gc TAT TAC TAT GGT TCG GCc ccaaagg	AAGTGTTCGACCTCTGGGGCCAGGGAACCTGG	XP1	5
SI	D	TGCGAGA	ctccatccgtctacg gTG ACT ACt gt	TACTACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	A4/1	5
SB	D	TGCG	tccgag ggT GGT TAC cctctaggac	AAGTGTTCGACCTCTGGGGCCAGGGAACCTGG	K4	5
FK	D	TGCGAAA	tggtag GTA TAG CAG CTg gctggtac	GGGGCCAGGGAACCTGG	N4	4 5
KN	D	TGCGAAA	tgg tGG GTA TAG CAG TGG CTG GTA C	GGGGCCAGGGAACCTGG	N1	4
MH	D	T	tccagtccctccggcgt ggA TAG CAG CAG CTG GTg gggcagctgtaatgggtcgg	ACTGTTTCGACCTCTGGGGCCAGGGAACCTGG	N1	5
SE	D	TGCGAGA	ggg gGT AGT ACC AGC TGC ccaatgagcggcgtgg	TACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	LR4	6
HI	D	TGCGAGATA	taatatgtcgggggga ctA GGA TAT TGT AGT AGT ACC AGC TGC TAT Gag ggaaggaa	TTACTACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	LR4	6
MK	D	TGCGAGAG	ttgtttccgttac tAG GAT ATT GTA GTA GTA CCA GCT GCT ATG gagt	ACTACTTTGACTACTGGGGCCAGGGAACCTGG	LR4	4
SH	D	TGCGAGAG	ccctga ATT GTA GTA GTA CCA GCT GCT ATG Cgc ctgata	TTACTACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	LR4	6
HM	D	TGCGAGAGA	c TAT TGT ACT AA g	ACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	LR1	6
RF	D	TGCGAGAGA	gccccggaaggag gaG GTG GTA Ggc	TACTACTACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	LR2	6
YM	D	TGCGAG	gatg ATT GTG GTG GTG ATT GCT A	TACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG	LR3	6
TH	D	TGCGAGAG	ctcagggg TGG GGA gacttt	TACTGTGACTTGTGATCTCTGGGGCCAGGGAACCTGG	HQ52	2
NK	D	TGCGAGAGA	agaaggatttaagtt gTA GCT GGG attatgtagt	TACTTTGACTACTGGGGCCAGGGAACCTGG	HQ52	4
CS	D	TG	gT AAC TCG Gc cgg	CTGGGGCCAGGGAACCTGG	HQ52	4
HT	D	TGCAA	aatatcgaatgag gAC CAG CTG Cca aaatgatag TAG TGG TTA TTA ctacgg	ACTGTTGTGACCTCTGGGGCCAGGGAACCTGG	LR4 & XP4	5
TY	D	TGCAA	aagat aGA TAT TGT AGT AGT ACC AGC TGC TAT ata taA ACT GGG GAa ctaacctttct	TTTGACTACTGGGGCCAGGGAACCTGG	LR4 & HQ52	4
SM	D		gcctcggggagtgcccc tTA TTG TAG TAG TAC CAG CTG CTA Taa atc ggA CGA TTT TTG GAG TGG TTA ggcgcctt	AGTGTGTTGACCTCTGGGGCCAGGGAACCTGG	LR4 & XP4	5
TE	D	T	acta	ACTACTACTACTACGGTATGGAAGTCTGGGGCCAGGGAACCAAGG		6

Figure 1. CDR-III sequences from pre-B ALL cells. D and R indicate diagnosis and relapse, respectively. N region nucleotides are indicated by lower-case letters. D elements which could be tentatively assigned to previously described D genes are underlined. Dots indicate identical nucleotides. * nucleotide discrepancy with previously described germline J_H segment. These nucleotide discrepancies were developed at the same position in each J_H segment, and were thought to be germline polymorphisms. ‡ nucleotide mutation. § nucleotide insertion. || nucleotide deletion. ¶ P region nucleotide.

Figure 2. D element usage among human pre-B ALL cells. Distribution of D element among 24 clones sequenced here (black bars), and among 21 clones previously reported (open bars) (29, 36, 37, 43). The clones which could be examined both at initial diagnosis and relapse were counted as one clone.

Comparison of CDR-III sequences between initial diagnosis and relapse. We compared the CDR-III sequences derived from 10 patients between initial diagnosis and relapse. Sequencing data were determined by examining several recombinant clones from each sample to exclude the possibilities of clonal heterogeneity and sequencing error. Although eight of them had completely matched sequences, the other two (K.M. and C.S.) differed by two or three nucleotides. To elucidate that these sequence differences were not due to Taq polymerase replication errors during early stages of the PCR amplification, we amplified genomic DNA from the initial and relapse specimens three times independently, and sequenced five recombinant clones derived from each amplified sample. Since each sequenced data (15 clones per one specimen) was completely identical, we could rule out the possibility of Taq polymerase replication errors. The mutated clones at relapse were thought to be the same as the original clones, because Southern blot analysis using J_H probe showed similar patterns (Fig. 4). Additionally, in patient K.M., a nucleotide insertion in the J_H segment was observed at diagnosis, and also at relapse. In patient C.S., the D element mutation was observed in both clones. These nucleotide differences at relapse were therefore not due to clonal change but to somatic mutation. All these mutations occurred at the N region. In patient K.M., two nucleotides, GG in the V_H - D_{XP4} junction, and one nucleotide C adjacent to the 3' end of the D_{XP4} element, had changed to CA and T at relapse, respectively. In patient C.S., two nucleotides GG downstream from the D_{HQ52} element had changed to TA at relapse (Fig. 1).

Discussion

Knowledge of the DNA sequences of the human IgH variable region has so far been limited, in contrast to that of the murine system. Recently a rapid cloning and sequencing method for IgH CDR-III using PCR has been developed (25, 35-38, 43). Using this technique, we analyzed the CDR-III sequences from pre-B ALL cells.

Somatic mutation of the CDR-III sequences from pre-B ALL cells. As mentioned in Results, we found several characteristics of the CDR-III sequences from pre-B ALL cells. So-

matic mutation of D and J_H segments was very rare (4/412 in D segments, 8/848 in J_H segments). Furthermore, all of our sequences of J4, J5, and J6 had a discrepancy of one, two, and one nucleotides, respectively, compared with published germline sequences (32). However, since these have been previously pointed out as germline polymorphisms (24), and the majority of sequenced human J_H segments had the same nucleotide sequences as ours (24, 25, 29, 36, 37, 43), we did not include these discrepancies among the numbers described above.

Somatic mutation appears to occur at specific stages of B cell development. Bird et al. reported that the rearranged V_H genes from pre-B ALL showed no evidence of mutation (29). In

Table II. Usage of D coding Frame

	1st	2nd	3rd
DXP4	4 (4)	1 (2)	0 (0)
DXP1	0 (1)	1 (1)	0 (1)
DXP'1	0 (0)	1 (3)	0 (1)
DA1/4	0 (0)	0 (0)	1 (1)
DK4	0 (0)	0 (1)	1 (2)
DK1	0 (0)	0 (0)	0 (0)
DN4	0 (0)	0 (0)	1 (2)
DN1	0 (1)	0 (0)	2 (3)
DM1	0 (0)	0 (0)	0 (0)
DM2	0 (0)	0 (0)	0 (0)
DLR5	0 (0)	0 (0)	0 (0)
DLR4	2 (4)	3 (4)	2 (3)
DLR1	0 (0)	1 (3)	0 (0)
DLR2	0 (1)	0 (0)	1 (3)
DLR3	0 (0)	0 (0)	1 (2)
DHQ52	2 (3)	0 (1)	2 (2)
Total	8 (14)	7 (15)	11 (20)

The coding frame of the D elements was temporary decided to start with TA(T/C) T(T/A)C TGT in the V_H region. All three coding frames were equally used. The total number of our data and previously described data (29, 36, 37, 43) are listed in parentheses.

chronic lymphocytic leukemia and Burkitt's lymphoma, somatic mutation has not been observed (46–48). On the other hand, many cases of follicular lymphoma contained significant somatic mutations in the IgH variable region (26–28). Additionally, Zelenetz et al. reported that transformed diffuse lymphoma cells could mutate somatically (49). Although malignant cells may not necessarily reflect the phenomena of normal B cell development, these data suggest that somatic mutation is activated after B cell maturation.

Utilization and coding frame of the D segment. All CDR-III sequenced here contained one or two D elements as reported by Siebenlist et al. (41), Ravetch et al. (32), Ichihara et al. (42), and Matsuda et al. (50).

The total number of human D genes is predicted to be about 30, since each five 9-Kb repeats contains six D gene families (D_M , D_{LR} , D_{XP} , D_A , D_K , and D_N) (42). Until now, 17 D genes have been sequenced. Four of five repeats are located between the V_H region and the J_H region (5'- D_4 - D_1 - D_2 - D_3 -3'). The D_5 gene is located among V_H genes (50).

There seemed to be a certain biased use of the D gene family in pre-B ALL cells, since the frequency of D segment usage was as follows; $D_{LR} > D_{XP} > D_{HQ52}$, $D_N > D_K$, $D_A > D_M$. In adult human peripheral blood B lymphocytes, D segments were used as follows with low frequency; $D_{XP} > D_N > D_K$, $D_{LR} > D_M$, $D_A > D_{HQ52}$ (25). Of interest is the difference of D segment usage between peripheral B cells and pre-B ALL cells. The low frequency of the D_K and D_A families, and relatively high frequency of the D_{LR} family and D_{HQ52} in pre-B ALL cells were noticeable compared with peripheral B cells. Although the reason for the low frequency of the D_K and D_A families remains unclear, the relatively high frequency of the D_{LR} family and D_{HQ52} seems to be related to the stage of B cell ontogeny. One reason for the low frequency of D_{LR} usage in peripheral B cells is that B cells using the first coding frame of the D_{LR} family contains stop codon and must be eliminated. There is a remarkable discrepancy in the usage of the D_{HQ52} element between human fetal liver and peripheral B cells (23–25). The first step in IgH gene rearrangement is a D to J_H join. The initial join often involves a D_{HQ52} gene (23, 24). In the murine system, the initial D- J_H rearrangement is often deleted and replaced by a secondary D- J_H rearrangement consisting of a more 5' D gene

Figure 3. J_H usage among human pre-B ALL cells. Black and open bars are as described in the legend to Fig. 2.

Figure 4. Southern blot hybridization of DNAs from patients C.S. and K.M. Southern blot hybridization of DNAs from the two patients (C.S. and K.M.) who had mutations in the N region nucleotides between initial diagnosis and relapse probed with J_H , indicates the identical rearranged pattern. (Lane 1) Patient C.S. at diagnosis; (Lane 2) Patient C.S. at relapse; (Lane 3) Patient K.M. at diagnosis; (Lane 4) Patient K.M. at relapse; (Lane 5) Germline. H, Digested with HindIII. B/H, digested with the combination of BamHI and HindIII. E, digested with EcoRI.

and a more 3' J_H gene (5, 51). The pre-B ALL cells using the D_{HQ52} element might have been transformed into malignancy during an early stage of ontogeny.

On the other hand, the D_4 subgroup (D_{XP4} and D_{LR4}) is most frequently used among D_{XP} and D_{LR} families. This finding concurs with Tsukada's report which showed the preferential use of the most 5' D in murine immature B cell lines (5). D_{LR5} was not found in our sequence data, although other D_{LR} families were frequently used. As mentioned above, moderately preferential usage of the D_4 subgroup and D_{HQ52} , and no use of D_{LR5} , might be explained by their location.

We observed three clones in which CDR-III sequences contained unusual D-D joins. The D-D joining model was first proposed by Kurosawa and Tonegawa to explain the origin of the N region of the murine IgH gene (44). They thought that a D segment (D_a) could join to another D segment (D_b), using a 24-mer signal spacer consisting of the end pentamer within the D_a segment, the D_a heptamer, and a 12mer of spacer instead of a 23mer, because intact D segments cannot join among themselves by the 12/23 rule. Meek et al., however, proved that murine germline D segments could directly join with another D segment either directly or invertedly (52). In fact, other investigators also reported the existence of D-D joins which were observed more frequently in adult murine B cells (18, 20). Before our analysis, Jonsson et al. also reported two clones with D-D joins; D_{LR1} - D_{XP1} and D_{K4} - D_{HQ52} from pre-B ALL (37). Furthermore, Yamada et al. reported several clones with D-D joins derived from peripheral B cells (25). Thus, in the CDR-III from human pre-B cells, the D-D joins might occur more frequently than in murine B cell differentiation.

Recent studies have shown that truncated or complete IgH chain molecules are expressed together with surrogate light chain molecules on the surface of the murine pre-B cells (22). Although the human counterpart molecule has not been demonstrated, a selection of D reading frames might be also influenced by a kind of selective pressure. Our DNA study, however, suggests no biased usage of D coding frames. Furthermore, some CDR-III sequences contained a stop codon. The

transcripts should be analyzed to elucidate the selection of coding frames.

Utilization of the J_H segment. We found a clear correlation between J_H utilization and the 3' localization of J_H . At an early stage of B cells in fetal liver, D_{HQS2} - $J1$ joining was frequently observed (23). During B cell differentiation, a second D - J_H or a third D - J_H might occur before V_H - D - J_H rearrangement is accomplished (51). In this case the most 5' D and the most 3' J_H would join at a later stage. This explanation matches our observations, and supports the one-dimensional tracking model of recombination (6, 53). Schroeder et al. reported a striking gradient of J_H usage in human fetal liver as follows: $J3 > J4 > J5 > J2, J6$ (24). Yamada et al. reported a different gradient of J_H usage in peripheral B cells as follows: $J4 > J6 > J5 > J3$ (25). In our findings and previously reported sequences derived from pre-B ALL cells (29, 36, 37, 43), $J4, J5$, and $J6$ segments were equally used. It is not clear at the present time whether these discrepancies are due to the different utilization of B cells in each stage of ontogeny, or the selection resulting from malignant transformation.

Mutated CDR-III sequence at relapse. 2 out of 10 patients had mutated CDR-III sequences at relapse. All of these changes were observed within the N region near the D segments; in addition, the G nucleotide was most frequently substituted. Misincorporations due to the relative lack of fidelity of Taq polymerase might be the important problem with our approach. In order to rule out the possibility of Taq polymerase replication errors, we examined the genomic DNAs from these two patients three times independently, and confirmed the mutations. In the earlier studies, a cumulative error frequency was reported about 0.25% after 30 cycles of PCR (39). However, recent studies using lower dNTP and Mg^{2+} ion concentrations (200 μ M each dNTP, and 1.5 mM $MgCl_2$), which were the same as ours, have reported a significant increase in the fidelity of Taq polymerase. Now it is recognized that the cumulative error frequency is less than 6.6×10^{-5} , and the average mutation rate is 5×10^{-6} errors per nucleotide incorporated per cycle, assuming 25 cycles of doubling (54, 55).

Generally, somatic mutation has been thought to occur in mature B cells after antigenic stimulation. In the murine system, somatic mutation in antibody variable genes appeared early after primary immunization (56). In a pre-B cell line, hypermutation at the IgH locus whose rate was determined to be $0.3-1 \times 10^{-4}$ per cell generation, was reported (57). On the other hand, in human systems it is not known from what stage human B cells mutate, although somatic mutations were frequently observed in follicular B cell lymphoma (26-28, 49). Our data, however, suggests that a primary mutation might occur at specific sequences even at the stage of pre-B cell.

Recently Yamada et al. reported the usefulness of the tumor-specific CDR-III sequence to detect and monitor minimal residual disease at the level of 10^{-5} , using the PCR amplification (43). We also reported the detection of minimal residual disease at the level of 10^{-6} in a patient who had received an allogeneic bone marrow transplantation using two-step PCR amplification (38). However the mutated CDR-III sequence cannot be detected, since these detection systems are based on the enormous diversity of CDR-III. The clinical usefulness of these systems, therefore, must be limited when the leukemia cells have mutated CDR-III sequences at relapse.

Structural analysis of CDR-III sequence is important to elucidate the B cell repertoire. As presented here, the sequential

study of clinical lymphoid tumor samples can serve as a valuable model in B cell differentiation. Additionally, a rapid cloning and sequencing method of CDR-III could be applied to many clinical fields.

Acknowledgments

We thank Dr. Yoshikazu Kurosawa for helpful advice and discussion, and Yujiro Mori and Alan F. Brunner (Aloka Co., Ltd.) for technical support for DNA sequencing.

This work was supported by grants from the Japanese Ministry of Health and Welfare.

References

1. Tonegawa, S. 1983. Somatic generation of antibody diversity. *Nature (Lond.)* 302:575-581.
2. Honjo, T. 1983. Immunoglobulin genes. *Annu. Rev. Immunol.* 1:499-528.
3. Alt, F. W., T. K. Blackwell, R. A. DePinho, M. G. Reth, and G. D. Yancopoulos. 1986. Regulation of genome rearrangement events during lymphocyte differentiation. *Immunol. Rev.* 89:5-30.
4. Wu, T. T., and E. A. Kabat. 1970. An analysis of the sequences of the variable regions of Bence Jones proteins and myeloma light chains and their implications for antibody complementarity. *J. Exp. Med.* 132:211-250.
5. Tsukada, S., H. Sugiyama, Y. Oka, and S. Kishimoto. 1990. Estimation of D segment usage in initial D to J_H joinings in a murine immature B cell line. *J. Immunol.* 144:4053-4059.
6. Yancopoulos, G. D., S. V. Desiderio, M. Paskind, J. F. Kearney, D. Baltimore, and F. W. Alt. 1984. Preferential utilization of the most J_H -proximal V_H gene segments in pre-B-cell lines. *Nature (Lond.)* 311:727-733.
7. Perlmutter, R. M., J. F. Kearney, S. P. Chang, and L. E. Hood. 1985. Developmentally controlled expression of immunoglobulin V_H genes. *Science (Wash. DC)* 227:1597-1600.
8. Jeong, H. D., and J. M. Teale. 1988. Comparison of the fetal and adult functional B cell repertoires by analysis of V_H gene family expression. *J. Exp. Med.* 168:589-603.
9. Dildrop, R., U. Krawinkel, E. Winter, and K. Rajewsky. 1985. V_H -gene expression in murine lipopolysaccharide blasts distributes over the nine known V_H -gene groups and may be random. *Eur. J. Immunol.* 15:1154-1156.
10. Wu, G. E., and C. J. Paige. 1986. V_H gene family utilization in colonies derived from B and pre-B cells detected by the RNA colony blot assay. *EMBO (Eur. Mol. Biol. Organ.) J.* 5:3475-3481.
11. Schulze, D. H., and G. Kelsoe. 1987. Genotypic analysis of B cell colonies by in situ hybridization. Stoichiometric expression of three V_H families in adult C57BL/6 and BALB/c mice. *J. Exp. Med.* 166:163-172.
12. Yancopoulos, G. D., B. A. Malynn, and F. W. Alt. 1988. Developmentally regulated and strain-specific expression of murine V_H gene families. *J. Exp. Med.* 168:417-435.
13. Sheehan, K. M., and P. H. Brodeur. 1989. Molecular cloning of the primary IgH repertoire: A quantitative analysis of V_H gene usage in adult mice. *EMBO (Eur. Mol. Biol. Organ.) J.* 8:2313-2320.
14. Alt, F. W., and D. Baltimore. 1982. Joining of immunoglobulin heavy chain gene segments: implications from a chromosome with evidence of three D - J_H fusions. *Proc. Natl. Acad. Sci. USA* 79:4118-4122.
15. Desiderio, S. V., G. D. Yancopoulos, M. Paskind, E. Thomas, M. A. Boss, N. Landau, F. W. Alt, and D. Baltimore. 1984. Insertion of N regions into heavy-chain genes is correlated with expression of terminal deoxynucleotidyl transferase in B cells. *Nature (Lond.)* 311:752-755.
16. Landau, N. R., D. G. Schatz, M. Rosa, and D. Baltimore. 1987. Increased frequency of N-region insertion in a murine pre-B-cell line infected with a terminal deoxynucleotidyl transferase retroviral expression vector. *Mol. Cell. Biol.* 7:3237-3243.
17. Lieber, M. R., J. E. Hesse, K. Mizuuchi, and M. Gellert. 1988. Lymphoid V(D)J recombination: Nucleotide insertion at signal joints as well as coding joints. *Proc. Natl. Acad. Sci. USA* 85:8588-8592.
18. Feeney, A. J. 1990. Lack of N regions in fetal and neonatal mouse immunoglobulin V-D-J junctional sequences. *J. Exp. Med.* 172:1377-1390.
19. Gu, H., I. Forster, and K. Rajewsky. 1990. Sequence homologies, N sequence insertion and J_H gene utilization in V_HDJ_H joining: implications for the joining mechanism and the ontogenetic timing of Ly1 B cell and B-CLL progenitor generation. *EMBO (Eur. Mol. Biol. Organ.) J.* 9:2133-2140.
20. Bangs, L. A., I. E. Sanz, and J. M. Teale. 1991. Comparison of D, J_H and junctional diversity in the fetal, adult, and aged B cell repertoires. *J. Immunol.* 146:1996-2004.
21. Ichihara, Y., H. Hayashida, S. Miyazawa, and Y. Kurosawa. 1989. Only D_{FL16} , D_{SP2} , and D_{QS2} gene families exist in mouse immunoglobulin heavy chain

diversity gene loci, of which D_{H16} and D_{SP2} originate from the same primordial D_H gene. *Eur. J. Immunol.* 19:1849–1854.

22. Gu, H., D. Kitamura, and K. Rajewsky. 1991. B cell development regulated by gene rearrangement: arrest of maturation by membrane-bound D_H protein and selection of D_H element reading frames. *Cell*. 65:47–54.

23. Nickerson, K. G., J. Berman, E. Glickman, L. Chess, and F. W. Alt. 1989. Early human IgH gene assembly in Epstein-Barr virus-transformed fetal B cell lines. Preferential utilization of the most J_H -proximal D segment (DQ52) and two unusual V_H -related rearrangements. *J. Exp. Med.* 169:1391–1403.

24. Schroeder JR, H. W., and J. Y. Wang. 1990. Preferential utilization of conserved immunoglobulin heavy chain variable gene segments during human fetal life. *Proc. Natl. Acad. Sci. USA*. 87:6146–6150.

25. Yamada, M., R. Wasserman, B. A. Reichard, S. Shane, A. J. Caton, and G. Rovera. 1991. Preferential utilization of specific immunoglobulin heavy chain diversity and joining segments in adult human peripheral blood B lymphocytes. *J. Exp. Med.* 173:395–407.

26. Cleary, M., T. C. Meeker, S. Levy, E. Lee, M. Trela, J. Sklar, and R. Levy. 1986. Clustering of extensive somatic mutations in the variable region of an immunoglobulin heavy chain gene from a human B cell lymphoma. *Cell*. 44:97–106.

27. Levy, R., S. Levy, M. L. Cleary, W. Carroll, S. Kon, J. Bird, and J. Sklar. 1987. Somatic mutation in human B-cell tumors. *Immunol. Rev.* 96:43–58.

28. Levy, S., E. Mendel, S. Kon, Z. Avnur, and R. Levy. 1988. Mutational hot spots in Ig V region genes of human follicular lymphomas. *J. Exp. Med.* 168:475–489.

29. Bird, J., N. Galili, M. Link, D. Stites, and J. Sklar. 1988. Continuing rearrangement but absence of somatic hypermutation in immunoglobulin genes of human B cell precursor leukemia. *J. Exp. Med.* 168:229–245.

30. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular Cloning: a Laboratory Manual. 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

31. Southern, E. 1975. Detection of specific sequence among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98:503–517.

32. Ravetch, J. V., U. Siebenlist, S. Korsmeyer, T. Waldmann, and P. Leder. 1981. Structure of the human immunoglobulin mu locus: Characterization of embryonic and rearranged J and D genes. *Cell*. 27:583–591.

33. Kabat, E. A., T. T. Wu, M. Reid-Miller, H. M. Perry, and K. S. Gottesman. 1987. Sequences of Proteins of Immunological Interest. 4th ed. National Institutes of Health publication, Bethesda, MD.

34. Berman, J. E., S. J. Mellis, R. Pollock, C. L. Smith, H. Suh, B. Heinke, C. Kowal, U. Surti, L. Chess, C. R. Cantor, et al. 1988. Content and organization of human Ig V_H locus: Definition of three new V_H families and linkage to the Ig V_H locus. 1988. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:727–738.

35. Yamada, M., S. Hudson, O. Tournay, S. Bittenbender, S. S. Shane, B. Lange, Y. Tsujimoto, A. J. Caton, and G. Rovera. 1989. Detection of minimal disease in hematopoietic malignancies of the B-cell lineage by using third-complementarity-determining region (CDR-III)-specific probes. *Proc. Natl. Acad. Sci. USA*. 86:5123–5127.

36. Brisco, M. J., L. W. Tan, A. M. Orsborn, and A. A. Morley. 1990. Development of a highly sensitive assay, based on the polymerase chain reaction, for rare B-lymphocyte clones in a polyclonal population. *Br. J. Haematol.* 75:163–167.

37. Jonsson, O. G., R. K. Kitchens, F. C. Scott, and R. G. Smith. 1990. Detection of minimal residual disease in acute lymphoblastic leukemia using immunoglobulin hypervariable region specific oligonucleotide probes. *Blood*. 76:2072–2079.

38. Kiyoi, H., T. Naoe, K. Kitamura, T. Yamauchi, T. Ichihashi, and R. Ohno. 1991. Disappearance of minimal residual lymphoblastic leukemia cells six months after allogeneic bone marrow transplantation without GVHD. *Bone Marrow Transplant.* 8:59–62.

39. Saiki, R. K., D. H. Gelfand, S. Stoffel, S. J. Scharf, R. Higuchi, G. T. Horn,

K. B. Mullis, and H. A. Erlich. 1988. Primer-directed enzymic amplification of DNA with a thermostable DNA polymerase. *Science (Wash. DC)*. 239:487–491.

40. Prober, J. M., G. L. Trainor, R. J. Dam, F. W. Hobbs, C. W. Robertson, R. J. Zagursky, A. J. Cocuzza, M. A. Jensen, and K. Baumeister. 1987. A system for rapid DNA sequencing with fluorescent chain-terminating dideoxynucleotides. *Science (Wash. DC)*. 238:336–341.

41. Siebenlist, U., J. V. Ravetch, S. Korsmeyer, T. Waldmann, and P. Leder. 1981. Human immunoglobulin D segments encoded in tandem multigenic families. *Nature (Lond.)*. 294:631–635.

42. Ichihara, Y., H. Matsuoka, and Y. Kurosawa. 1988. Organization of human immunoglobulin heavy chain diversity gene loci. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:4141–4150.

43. Yamada, M., R. Wasserman, B. Lange, B. A. Reichard, R. B. Womer, and G. Rovera. 1990. Minimal residual disease in childhood B-lineage lymphoblastic leukemia. *N. Engl. J. Med.* 323:448–455.

44. Kurosawa, Y., and S. Tonegawa. 1982. Organization, structure, and assembly of immunoglobulin heavy chain diversity DNA segments. *J. Exp. Med.* 155:201–217.

45. Lafaille, J. L., A. DeCloux, M. Bonneville, Y. Takagaki, and S. Tonegawa. 1989. Junctional sequences of T cell receptor genes: implications for T cell lineages and for a novel intermediate of V-(D)-J joining. *Cell*. 59:859–870.

46. Kipps, T. J., E. Tomhave, P. P. Chen, and D. A. Carson. 1988. Autoantibody-associated kappa light chain variable region gene expressed in chronic lymphocytic leukemia with little or no somatic mutation. Implications for etiology and immunotherapy. *J. Exp. Med.* 167:840–852.

47. Meeker, T. C., J. C. Grimaldi, R. O'Rourke, J. Loeb, G. Juliusson, and S. Einhorn. 1988. Lack of detectable somatic hypermutation in the V region of the Ig H chain gene of a human chronic B lymphocytic leukemia. *J. Immunol.* 141:3994–3998.

48. Caroll, W. L., M. Yu, M. P. Link, and S. J. Korsmeyer. 1989. Absence of Ig V region gene somatic hypermutation in advanced Burkitt's lymphoma. *J. Immunol.* 143:692–698.

49. Zelenetz, A. D., T. T. Chen, and R. Levy. 1991. Histologic transformation of follicular lymphoma to diffuse lymphoma represents tumor progression by a single malignant B cell. *J. Exp. Med.* 173:197–207.

50. Matsuda, F., K. H. Lee, S. Nakai, T. Sato, M. Kodaira, S. Q. Zong, H. Ohno, S. Fukuhara, and T. Honjo. 1988. Dispersed localization of D segments in the human immunoglobulin heavy-chain locus. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:1047–1051.

51. Reth, M. G., S. Jackson, and F. W. Alt. 1986. V_HDJ_H formation and DJ_H replacement during pre-B differentiation: non-random usage of gene segments. *EMBO (Eur. Mol. Biol. Organ.) J.* 5:2131–2138.

52. Meek, K. D., C. A. Hasemann, and J. D. Capra. 1989. Novel rearrangements at the immunoglobulin D locus: Inversions and fusions add to IgH somatic diversity. *J. Exp. Med.* 170:39–57.

53. Wood, C., and S. Tonegawa. 1983. Diversity and joining segments of mouse immunoglobulin heavy chain genes are closely linked and in the same orientation: implications for the joining mechanism. *Proc. Natl. Acad. Sci. USA*. 80:3030–3034.

54. Gelfand, D. H., and T. J. White. 1990. Thermostable DNA polymerase. In PCR Protocols. M. A. Innis, D. H. Gelfand, J. J. Shinsky, and T. J. White, editors. Academic Press, San Diego, CA. 129–141.

55. Erlich, H. A., D. Gelfand, and J. J. Sninsky. 1991. Recent advances in the polymerase chain reaction. *Science (Wash. DC)*. 252:1643–1651.

56. Levy, N. S., U. V. Malipiero, S. G. Lebecque, and P. J. Gearhart. 1989. Early onset of somatic mutation in immunoglobulin V_H genes during the primary immune response. *J. Exp. Med.* 169:2007–2019.

57. Wabl, M., P. D. Burrows, A. V. Gabain, and C. Steinberg. 1985. Hypermutation at the immunoglobulin heavy chain locus in a pre-B-cell line. *Proc. Natl. Acad. Sci. USA*. 82:479–482.