

Mineralocorticoid Modulation of Apical and Basolateral Membrane $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ Transport Processes in the Rabbit Inner Stripe of Outer Medullary Collecting Duct

Steven R. Hays

Department of Internal Medicine, The University of Texas Southwestern Medical Center, Dallas, Texas 75235

Abstract

To examine the mechanism by which mineralocorticoids regulate HCO_3^- absorption in the rabbit inner stripe of the outer medullary collecting duct, we microfluorometrically measured intracellular pH (pH_i) in *in vitro* perfused tubules using 2',7'-bis(carboxyethyl)-5(6)-carboxyfluorescein (BCECF) assaying the apical and basolateral membrane $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ transport processes in three groups of animals: those receiving chronic *in vivo* DOCA treatment (5 mg/kg per d \times 2 wk); those with surgical adrenalectomy (ADX, [chronic \times 2 wk]) on glucocorticoid replacement; and controls. Baseline pH_i was not different in the three groups. Cellular volume (vol/mm) was increased 38% in DOCA tubules versus controls, but unchanged in ADX tubules versus controls. Buffer capacities (B_T) were not different in the three groups. Apical membrane H^+ pump activity, assayed as the Na^+ -independent pH_i recovery from an acid load ($\text{NH}_3/\text{NH}_4^+$ prepulse) and expressed as J_H ($\text{dpH}_i/\text{dt} \cdot \text{vol}/\text{mm} \cdot B_T$) was increased 76% in DOCA tubules versus controls, and decreased 56% in ADX tubules versus controls. Basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchange activity assayed as the pH_i response to basolateral Cl^- addition was increased 73% in DOCA tubules versus controls, and decreased 44% in ADX tubules versus controls. When examined as a function of varying $[\text{Cl}^-]$, the V_{max} of $\text{Cl}^-/\text{HCO}_3^-$ exchange activity was significantly increased in DOCA tubules (control, $72.7 \pm 15.7 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$ vs DOCA, $132.3 \pm 22.5 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P < 0.02$), while the $K_{1/2}$ for Cl^- was unchanged. Basolateral membrane Na^+/H^+ antiporter activity assayed as the Na^+ -dependent pH_i recovery from an acid load was not changed in chronic DOCA tubules versus controls. In conclusion, the apical membrane H^+ pump and basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchanger of the rabbit OMCD_i are regulated in parallel without chronic alterations in pH_i under the conditions of mineralocorticoid excess and deficiency. The parallel changes in these transporters accounts for the alterations in OMCD_i HCO_3^- absorption seen under these condi-

tions. (*J. Clin. Invest.* 1992. 90:180–187.) Key words: proton pump • chloride-bicarbonate exchanger • intracellular pH • adrenalectomy • chronic desoxycorticosterone acetate

Introduction

Mineralocorticoids are known to be an important regulator of net acid excretion by the kidney (1, 2). This action is mediated in part by a direct effect of mineralocorticoids, independent of effects on Na^+ transport (3), on H^+ secretion in the inner stripe of the outer medullary collecting duct (OMCD_i).¹ Several mechanisms exist whereby mineralocorticoids could directly stimulate transepithelial H^+ secretion in the OMCD_i: (a) stimulating the apical membrane H^+ pump; (b) stimulating the basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchanger; (c) inhibiting the basolateral membrane Na^+/H^+ antiporter; or (d) stimulating the apical and basolateral membrane $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ transporters in parallel. Several investigators have found that the *in vivo* mineralocorticoid status of the animal alters the level of NEM-sensitive H^+ -ATPase activity measured in the OMCD_i (4–7). In addition, in OMCD_i's harvested from adrenalectomized animals, acute exposure of aldosterone stimulates NEM-sensitive H^+ -ATPase activity by a process requiring protein synthesis (7). These enzymatic studies suggest that the primary effect of mineralocorticoids may be mediated by a direct effect on the apical membrane H^+ pump. Effects on the other $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ transport mechanisms have not been investigated.

The purpose of the present study was to examine the mechanism(s) by which mineralocorticoids modulate HCO_3^- absorption in the rabbit OMCD_i, using the fluorescent measurement of cell pH (pH_i). The results demonstrate that increases and decreases in mineralocorticoid activity modulate apical and basolateral membrane $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ transporters in parallel without chronically altering pH_i , accounting for changes in OMCD_i HCO_3^- absorption in these conditions.

Methods

The technique of *in vitro* microperfusion of isolated rabbit OMCD_i was used as previously described (8). Briefly, female New Zealand White rabbits weighing 1.5–2.0 kg were maintained on standard laboratory chow and tap water *ad lib*. OMCD_i segments were identified and dissected free from 1-mm coronal slices at 4°C (pH 7.4, solution 1, Table I). To avoid the outer stripe, perfused segments were dissected from the

Portions of this work were presented in 1990 before the National Meeting of the American Society of Nephrology, Washington, DC, and appeared in abstract form (1990. *J. Am. Soc. Nephrol.* 1:650a).

Address correspondence and reprint requests to Dr. Steven R. Hays, Department of Internal Medicine, The University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, TX 75235-8856.

Received for publication 16 September 1991 and in revised form 3 February 1992.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/92/07/0180/08 \$2.00

Volume 90, July 1992, 180–187

1. Abbreviations used in this paper: ADX, adrenalectomy; BCECF, 2',7'-bis(carboxyethyl)-5(6)-carboxyfluorescein; $B_{\text{non-CO}_2\text{HCO}_3^-}$, non- $\text{CO}_2/\text{HCO}_3^-$ buffer capacity; B_T , total buffer capacity; DIDS, 4,4'-diisothiocyano-stilbene-2, 2'-disulfonate; DOCA, desoxycorticosterone acetate; $[\text{NH}_4^+]_o$, concentration of NH_4^+ in the bath; OMCD_i, outer medullary collecting duct; pH_i , intracellular pH.

Table I. Composition of Solutions

	1	2	3	4	5	6	7	8	9	10
Na ⁺	145	135	—	125	135	135	135	135	135	—
K ⁺	5	5	5	5	5	5	5	5	5	5
Mg ²⁺	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Ca ²⁺	1.6	1.6	1.6	1.6	7.0	1.6	6.9	6.7	5.0	1.6
Choline	—	—	120	—	—	—	—	—	—	110
NH ₄ ⁺	—	—	—	10	—	—	—	—	—	10
N-methyl										
D-glucosamine	—	—	15	—	—	—	—	—	—	15
Cl ⁻	125.2	125.2	125.2	125.2	—	103.2	10	20	40	125.2
HCO ₃ ⁻	25	—	—	—	25	25	25	25	25	—
HPO ₄ ²⁻	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
SO ₄ ²⁻	—	—	—	—	1.0	1.0	1.0	1.0	1.0	—
Gluconate	—	—	—	—	114	—	103.8	93.4	70	—
Glucose	8.3	8.3	8.3	8.3	8.3	8.3	8.3	8.3	8.3	8.3
Alanine	5	5	5	5	5	5	5	5	5	5
Hepes	—	25	25	25	25	25	25	25	25	25

All units are millimolar. All solutions were titrated to 290 mOsm with addition of raffinose. Solutions 3 and 10 were titrated to pH 7.40 with 1 N N-methyl-D-glucosamine hydroxide. Solutions 1, 2, and 4–9 were titrated to pH 7.40 with 1 N NaOH. All HCO₃⁻ containing solutions were equilibrated with 94% O₂/6% CO₂; other solutions were equilibrated with 100% O₂.

inner half of the inner stripe. Tubules were transferred into a bath chamber with a vol of ~90 μ l, constructed of black lucite to minimize light reflection. The peritubular fluid was continuously exchanged at ~10 ml/min by hydrostatic pressure, allowing a complete fluid exchange within 1 s. Tubular lumens were perfused at flow rates of 25–50 nl/min. Bath pH was monitored continuously by placing a commercial flexible pH electrode into the bath (MI-5089; Microelectrodes, Inc., Londonderry, NH). Bath solutions were prewarmed at 37°C, continuously equilibrated with either 94% O₂/6% CO₂ or 100% O₂, and passed to the bath chamber through CO₂ impermeable tubing (Clarkson Controls and Equipment Co., Detroit, MI). Bath temperature of 37±0.3°C was maintained by a specially designed water-jacketed glass coiled line.

To minimize motion, the distal end of tubule was sucked gently into a collection pipette. In addition, the average length of the tubule exposed to the bath fluid was limited to ~250–500 μ m. The tubules were loaded with the acetoxymethyl derivative of 2',7'-bis(carboxyethyl)-5(6)-carboxyfluorescein (BCECF-AM) (Molecular Probes, Inc., Eugene, OR), 8 μ M, from the bath using solution 1 (Table I). Loading was continued until signal to background fluorescence at 450 nm excitation was $\geq 10:1$, usually requiring 10–15 min. Tubules were then washed with solution 1, (Table I) at pH 7.4 for a minimum of 10 min, followed by the control solution of each experiment.

Luminal and peritubular solutions used in these studies are listed in Table I. CO₂/HCO₃⁻-free solutions were bubbled with 100% oxygen passed through a 3-N KOH trap. Desoxycorticosterone acetate (DOCA), 4,4'-diisothiocyano-stilbene-2,2'-disulfonate (DIDS), nigericin, amiloride hydrochloride, and all solution salts were purchased from Sigma Chemical Co. (St. Louis, MO).

Cell pH (pH_i) measurement. BCECF fluorescent emission was measured at 530 nm with an inverted epifluorescent microscope (Nikon Diaphot; Nikon Inc., Garden City, NY) attached to a dual excitation microspectrofluorimeter (SPEX CM-1; SPEX Inds., Inc., Edison, NJ) which allows rapid alternation between two excitation wavelengths (500 and 450 nm), as previously described (9). Fluorescence was measured using a 20 \times objective, on an area of the tubule ~125 μ m in length that included the entire width of the tubule. No attempts were made to make measurements on single cells. Although immunocytochemical studies based on the staining with antibodies against the H⁺-ATPase and band 3 Cl⁻/HCO₃⁻ exchanger have found evidence

for two cell types in the inner stripe segment (10), ultrastructural (11), morphologic (12), and electrophysiologic (13, 14) studies strongly support one cell type in the inner half of the OMCD, where the present studies were performed. Wiener and Hamm (15) were able to divide OMCD_i cells into two groups based on ease of BCECF loading, but found both cell types possessed the same apical and basolateral membrane H⁺/OH⁻/HCO₃⁻ transporters. Thus, these studies suggest that the OMCD_i is composed of one cell type with variable degrees of differentiation. Generally, the measured segment started ~50–100 μ m from the perfusion pipette. Background fluorescence at each of the excitation wavelengths was measured on the tubule before loading with BCECF, and the results were subtracted from the measured fluorescence during the experiment. A fluorescent ratio was then calculated as fluorescence with 500 nm excitation divided by that with 450 nm excitation. The initial rate of change in the fluorescence excitation ratio was defined by the slope of a line drawn tangent to the initial deflection ([d(F500/F450)/dt]).

BCECF fluorescence excitation ratios were calibrated intracellularly using the method of Thomas et al. (16). Tubules were bathed and perfused with well-buffered solutions (25 mM Hepes, 60 mM phosphate, and appropriate [HCO₃⁻]) of varying pH containing 7 μ M nigericin (a K⁺/H⁺ antiporter) and 120 mM K⁺ (9). Because of the small amount of variability between tubules, a calibration was generated in each of three experimental groups of animals (chronic DOCA, adrenalectomy, and control) and was used to convert F500/F450 fluorescent ratios to pH units in each experimental group.

Buffer capacity. The buffer capacity was determined using the technique of either rapid NH₃/NH₄⁺ addition or withdrawal, as described by Roos and Boron (17). The non-CO₂/HCO₃⁻ buffer capacity B_{non-CO₂/HCO₃⁻} (mmol · liter⁻¹ · pH unit⁻¹) is given by the formula:

$$B_{\text{non-CO}_2/\text{HCO}_3^-} = [\text{NH}_4^+]_i / \Delta\text{pH}_i \quad (1)$$

where ΔpH_i is the measured pH_i change. [NH₄⁺]_i is calculated from the formula:

$$[\text{NH}_4^+]_i = [\text{NH}_4^+]_o (\text{pH}_o - \text{pH}_i) \quad (2)$$

where [NH₄⁺]_o is the concentration of NH₄⁺ in the bath; pH_o is the pH of the peritubular solution; and pH_i is either the cell pH after NH₃/NH₄⁺ addition, or the cell pH just before NH₃/NH₄⁺ withdrawal. In

studies performed in the absence of $\text{CO}_2/\text{HCO}_3^-$, $B_{\text{non-CO}_2/\text{HCO}_3^-}$ represents the total buffer capacity of the cell (B_T). In studies performed in the presence of $\text{CO}_2/\text{HCO}_3^-$, the CO_2 -buffering capacity, B_{CO_2} ($\text{mmol} \cdot \text{liter}^{-1} \cdot \text{pH unit}^{-1}$) is calculated by the formula:

$$B_{\text{CO}_2} = 2.3[\text{HCO}_3^-]_i \quad (3)$$

where $[\text{HCO}_3^-]_i$ is given by the formula:

$$[\text{HCO}_3^-]_i = \alpha \cdot \text{pCO}_2 \cdot 10(\text{pHi} - \text{pK}) \quad (4)$$

where α is the solubility of CO_2 in water, pHi is the cell pH at the start of the experimental maneuver, and a pK of 6.1 was used. In the presence of $\text{CO}_2/\text{HCO}_3^-$, the total buffer capacity of the cell is equal to:

$$B_T = B_{\text{non-CO}_2/\text{HCO}_3^-} + B_{\text{CO}_2/\text{HCO}_3^-} \quad (5)$$

Calculation of proton fluxes. Transporter activities were expressed as proton fluxes (J_H , $\text{pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$) induced by the experimental maneuvers, calculated using the formula:

$$J_H = \text{dpHi}/\text{dt} \cdot \text{vol}/\text{mm} \cdot B_T \quad (6)$$

where vol/mm is the cellular volume of the tubules expressed per millimeter length. Cellular volumes were determined for each tubule by optical measurement of internal and external diameters of each tubule before experimentation. When transporter activity was measured under acidic pHi conditions (H^+ pump and Na^+/H^+ antiporter), the buffer capacity was determined by $\text{NH}_3/\text{NH}_4^+$ withdrawal. In these studies measuring Na^+/H^+ antiporter and H^+ pump activity, the buffer capacity was determined on the same tubule. When transporter activity was measured under alkaline pHi conditions ($\text{Cl}^-/\text{HCO}_3^-$ exchanger), $B_{\text{non-CO}_2/\text{HCO}_3^-}$ was calculated from $\text{NH}_3/\text{NH}_4^+$ addition made during the Na^+/H^+ antiporter and H^+ pump studies, plus the $B_{\text{CO}_2/\text{HCO}_3^-}$ determined for each $\text{Cl}^-/\text{HCO}_3^-$ exchanger tubule.

Experimental groups. Three experimental groups of animals were used. The first group consisted of rabbits maintained on standard laboratory chow (Teklad, Madison, WI) and tap water ad lib. (control). There were two control groups of animals, one for each experimental group. The second group of rabbits was maintained on standard laboratory chow and tap water ad lib., and received 5 mg/kg body wt DOCA (dissolved in sesame oil) intramuscularly every day for 14 d (DOCA). The third group of rabbits was surgically adrenalectomized and maintained on standard laboratory chow and 0.9% normal saline drinking water ad lib. (ADX). Preoperatively and on the first postoperative day, ADX animals received 50 μg of dexamethasone (dexamethasone sodium phosphate; Elkins-Sinn, Cherry Hill, NJ) subcutaneously. Thereafter, the rabbits were maintained on glucocorticoid replacement therapy (dexamethasone 10 $\mu\text{g}/\text{kg}$ s.c.) from the second postoperative day until death (12–15 d). This dose of glucocorticoid minimizes mineralocorticoid activity, while enhancing food and water intake as well as the general well-being of the animals (18–20).

Statistics. Results are reported as mean \pm standard error. Data are analyzed using the two-tailed Student's t test for paired and unpaired data, as appropriate.

Results

Chronic in vivo DOCA administration

The first set of studies examined the effects of chronic in vivo DOCA administration on the apical and basolateral membrane OMCD_i $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ transport processes. Shown in Table II is a comparison of baseline pHi , cell volume, and buffer capacities obtained in OMCD_i's harvested from chronic DOCA and control rabbits. Baseline pHi values obtained in the presence of 25 mM HCO_3^- and 40 mm Hg pCO_2 (pH 7.4, solution 1, Table I) in 21 DOCA tubules was 6.93 ± 0.02 . This value was not significantly different from the baseline pHi values determined under similar conditions in 23 control tubules

Table II. Baseline Cell pH, Cell Volume, and Buffer Capacities in Chronic DOCA Tubules

	Control	Chronic DOCA
Baseline cell pH	6.92 ± 0.03 $n = 23$	6.93 ± 0.02 $n = 21$
Cell vol (10^{-10} liter/mm)	6.51 ± 0.29 $n = 17$	$8.97 \pm 0.25^{\S}$ $n = 33$
Buffer capacity ($\text{mmol} \cdot \text{liter}^{-1} \cdot \text{pH unit}^{-1}$) (H^+ pump and Na^+/H^+ antiporter studies)*	33.16 ± 3.80 $n = 17$	34.37 ± 4.21 $n = 22$
($\text{Cl}^-/\text{HCO}_3^-$ exchanger studies) [†]	39.02 ± 2.50 $n = 18$	40.16 ± 2.58 $n = 20$

Values are means \pm SEM. n = number of tubules. * Buffer capacity was determined by $\text{NH}_3/\text{NH}_4^+$ withdrawal. [†] Buffer capacity was the mean $B_{\text{non-CO}_2/\text{HCO}_3^-}$ value obtained from $\text{NH}_3/\text{NH}_4^+$ addition in the H^+ pump and Na^+/H^+ antiporter studies plus the $B_{\text{CO}_2/\text{HCO}_3^-}$ for each tubule in the $\text{Cl}^-/\text{HCO}_3^-$ exchanger studies. [§] $P < 0.05$ chronic DOCA versus controls.

(6.92 ± 0.03 , $P = \text{NS}$). However, chronic DOCA treatment led to a 38% increase in cell volume (control, $n = 17$, $6.51 \pm 0.29 \cdot 10^{-10}$ liter/mm; chronic DOCA, $n = 33$, $8.97 \pm 0.25 \cdot 10^{-10}$ liter/mm, $P < 0.05$). There was no significant difference in buffer capacities between the two groups determined by either $\text{NH}_3/\text{NH}_4^+$ addition or withdrawal (Table II).

Apical membrane H^+ pump activity. Apical membrane H^+ pump activity was assessed as the Na^+ -independent pHi recovery from an acid load performed in the absence of ambient $\text{CO}_2/\text{HCO}_3^-$. In previous studies, we have found Na^+ -independent pHi recovery from an acid load to be NEM-sensitive and SCH 28080-insensitive, likely reflecting the presence of a simple vacuolar H^+ pump (21). In addition, we have shown that a sustained increase in cell calcium induced by peritubular Na^+ removal results in inactivation of the H^+ pump (22). To lessen the influence of this variable, we examined H^+ pump activity in the presence of peritubular Na^+ , with Na^+ -dependent pHi recovery inhibited by 1 mM peritubular amiloride. 1 mM peritubular amiloride completely blocks Na^+ -dependent pHi recovery in this segment (21). Shown in Fig. 1 is a typical experimental tracing. 12–15 min before the experiment, tubules were bathed in a 135-mM Na^+ , $\text{CO}_2/\text{HCO}_3^-$ -free solution (pH 7.4, solution 2, Table I) and luminally perfused with a Na^+ -free, $\text{CO}_2/\text{HCO}_3^-$ -free solution (pH 7.4, solution 3, Table I). 1 mM amiloride was then added to the peritubular solution and continued throughout the remainder of the experiment. After 5 min exposure to this solution, the peritubular solution was changed to an identical solution containing 10 mM NH_4Cl (pH 7.4, solution 4, Table I), and then 4–5 min later it was changed back to an identical solution without NH_4Cl (solution 2). The initial rate of pHi recovery from the induced acid load (Fig. 1) was used as an index of apical membrane H^+ pump activity. Shown in Fig. 2 are the results of these studies. In 11 chronic DOCA tubules, J_H was increased 76% when compared to seven control tubules (control, $n = 7$, 2.23 ± 0.48 $\text{pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; DOCA, $n = 11$, 3.93 ± 0.51 $\text{pmol} \cdot$




Figure 1. Assay for apical membrane H⁺ pump activity. Na⁺-independent cell pH recovery after an acid load in the presence of peritubular Na⁺ plus amiloride. Before the experimental tracing shown above, tubules were initially bathed for 15 min with a CO₂/HCO₃⁻-free solution (pH 7.4, solution 2, Table I) and luminally perfused with a Na⁺-free, CO₂/HCO₃⁻-free solution (pH 7.4, solution 3, Table I). At the beginning of the experimental tracing above, 1 mM amiloride was then added to the peritubular solution and continued throughout the remainder of the experiment. After 5 min exposure to this solution, the peritubular solution was changed to an identical solution containing 10 mM NH₄Cl (pH 7.4, solution 4, Table I) for 4–5 min before changing back to an identical solution without NH₄Cl (solution 3).

mm · min⁻¹, $P < 0.05$). Thus, chronic in vivo DOCA treatment leads to a stimulation of the apical membrane H⁺ pump.

Basolateral membrane Cl⁻/HCO₃⁻ exchange activity. Basolateral membrane Cl⁻/HCO₃⁻ exchange activity was assessed as the effect of peritubular Cl⁻ addition on pH_i. Shown in Fig. 3 is a typical experimental tracing. Tubules were bathed and luminally perfused with a Cl⁻-free solution containing 25 mM HCO₃⁻ (pH 7.4, solution 5, Table I) for 15 min before experimentation. The peritubular solution was then changed to an identical solution containing 103 mM Cl⁻ (pH 7.4, solution 6, Table I) for 1 min before changing back to the Cl⁻-free solution (solution 5). The initial rate of cell acidification after peritubular Cl⁻ addition (Fig. 3) was used as an index of basolateral membrane Cl⁻/HCO₃⁻ exchange activity. Shown in Fig. 4 are




Figure 2. Effect of chronic DOCA treatment on apical membrane H⁺ pump activity. Mean J_H ± SEM for control and chronic DOCA tubules is shown.




Figure 3. Assay for basolateral membrane Cl⁻/HCO₃⁻ exchange activity: peritubular Cl⁻ addition-induced acidification. Tubules were initially bathed with a control solution containing 125.2 mM Cl⁻ and 25 mM HCO₃⁻ (pH 7.4, solution 1, Table I) and luminally perfused with a Cl⁻-free, 25 mM HCO₃⁻ solution (pH 7.4, solution 5, Table I). The peritubular solution was then changed to a Cl⁻-free, 25 mM HCO₃⁻ solution (solution 5) for 15 min and continued for the first 1–2 min of the experimental tracing above. Cl⁻ was then added to the peritubular solution (pH 7.4, solution 6, Table I) for 1 min before switching back the Cl⁻-free solution (solution 5).

the results of these studies. In 20 chronic DOCA tubules, J_H was increased 73% when compared to 18 control tubules (control, 43.45 ± 6.34 pmol · mm⁻¹ · min⁻¹; DOCA, 75.24 ± 10.84 pmol · mm⁻¹ · min⁻¹, $P < 0.02$). Thus, chronic in vivo DOCA treatment leads to a stimulation of the basolateral membrane Cl⁻/HCO₃⁻ exchanger.

To assure that the assay was measuring a DOCA-induced increase in basolateral membrane Cl⁻/HCO₃⁻ exchange activity, similar experiments were performed in chronic DOCA tubules in the absence and presence of the disulfonic stilbene inhibitor of Cl⁻/HCO₃⁻ exchange activity, DIDS. After initially determining the rate of peritubular Cl⁻-induced cell acidification in the absence of peritubular DIDS, the peritubular solution was changed to a Cl⁻-free solution containing 100 μM peritubular DIDS for 6–8 min. The rate of peritubular Cl⁻-induced cell acidification was then reexamined. In seven chronic DOCA tubules, the rate of cell acidification induced by peritubular Cl⁻ addition was inhibited 96% in the presence of peritubular




Figure 4. Effect of chronic DOCA treatment on basolateral membrane Cl⁻/HCO₃⁻ exchange activity. Mean J_H ± SEM of control and chronic DOCA tubules is shown.

bular DIDS (baseline $69.18 \pm 23.84 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; DIDS $2.69 \pm 1.24 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P < 0.05$). The 96% inhibition of peritubular Cl^- -induced acidification by peritubular DIDS was similar to the 97% inhibition found in control tubules ($n = 3$, baseline $34.47 \pm 0.51 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; DIDS $1.06 \pm 1.31 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P < 0.01$). These studies indicate that the cell acidification that occurred upon peritubular Cl^- addition in the DOCA tubules was predominantly due to an increase in basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchange activity.

To further examine the nature of this DOCA-induced increase in basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchange activity, the apparent $K_{1/2}$ for peritubular Cl^- of the exchanger was determined in similar studies. Tubules were again bathed and lumenally perfused in a Cl^- -free solution containing 25 mM HCO_3^- (solution 5) for 15 min before experimentation. Peritubular additions of 10, 20, 40 and 103 mM Cl^- (solutions 7, 8, 9, and 6, respectively) were examined. The order of the Cl^- additions was varied from tubule to tubule. The kinetics of this transporter were determined from the rate of change of J_{H} in response to peritubular Cl^- addition. When more than one measurement was made with the same Cl^- concentration on a tubule, these were averaged to provide a result for that tubule. The initial acidification rate (J_{H}) in eight DOCA-treated tubules with addition of 10 mM Cl^- was $15.48 \pm 1.99 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; 20 mM, $27.70 \pm 3.21 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; 40 mM, $44.17 \pm 3.77 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; and 103 mM, $80.65 \pm 9.19 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$. The initial acidification rate (J_{H}) in 11 control tubules with addition of 10 mM Cl^- was $10.22 \pm 2.28 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; 20 mM, $17.05 \pm 3.5 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; 40 mM, $26.78 \pm 4.34 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; and 103 mM, $47.28 \pm 7.41 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$. Shown in Fig. 5 is the Eadie-Hofstee transformation of this data. There was no significant difference between the two groups in the $K_{1/2}$ values (control, $67.72 \pm 16.00 \text{ mM}$; DOCA $75.29 \pm 12.85 \text{ mM}$, $P = \text{NS}$), whereas a significant difference in the V_{max} values was seen (control $72.69 \pm 15.73 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; DOCA $132.31 \pm 22.49 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P < 0.02$). Thus, chronic DOCA treatment leads to no change in the $K_{1/2}$, but increases the V_{max} of the basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchanger.

Basolateral membrane Na^+/H^+ antiporter activity. Basolateral membrane Na^+/H^+ antiporter activity was assessed as the Na^+ -dependent pH_i recovery from an acid load. Shown in Fig. 6 is a typical experimental tracing. Tubules were initially bathed in a HCO_3^- -free solution containing 135 mM Na^+ (pH 7.4, solution 2, Table I) and lumenally perfused with a Na^+ -free, HCO_3^- -free solution (pH 7.4, solution 3, Table I) for 15




Figure 5. Effect of chronic DOCA treatment on the acidification rate as a function of peritubular $[\text{Cl}^-]$: Eadie-Hofstee transformation.

Peritubular $[\text{Na}^+]$, mM	135	0	135
Peritubular $[\text{NH}_4\text{NH}_4^+]$, mM	0	10	0




Figure 6. Assay for basolateral membrane Na^+/H^+ antiporter activity: Na^+ -dependent pH_i recovery from an acid load. Tubules were initially bathed with a control solution containing 145 mM Na^+ and 25 mM HCO_3^- (pH 7.4, solution 1, Table I) and lumenally perfused with a Na^+ -free, $\text{CO}_2/\text{HCO}_3^-$ -free solution (pH 7.4, solution 3, Table I). The peritubular solution was then changed to $\text{CO}_2/\text{HCO}_3^-$ -free solution containing 135 mM Na^+ (pH 7.4, solution 2, Table I) for 15 min and continued for the first 1–2 min of the experimental tracing above. The peritubular solution was then changed to a Na^+ -free, $\text{CO}_2/\text{HCO}_3^-$ -free solution containing 10 mM NH_4Cl (pH 7.4, solution 10, Table I). After 3–4 min, the peritubular solution was changed to an identical solution without NH_4Cl (pH 7.4, solution 3, Table I). After 1 min, the peritubular solution was changed to an identical solution containing 135 mM Na^+ .

min before experimentation. The peritubular solution was then changed to a Na^+ -free solution containing 10 mM NH_4Cl (pH 7.4, solution 10, Table I) for 4–5 min. The peritubular solution was then changed to an identical solution without the NH_4Cl (pH 7.4, solution 3, Table I) for 1 min before changing to an identical solution containing 135 mM Na^+ (solution 2). The initial rate of pH_i recovery from the induced acid load in the presence of Na^+ was used as an index of basolateral membrane Na^+/H^+ antiporter activity (Fig. 6). In addition, the initial rate of pH_i recovery for 1 min in the absence of Na^+ (Fig. 6) is an index of Na^+ -independent H^+ extrusion (apical membrane H^+ pump activity). In 11 DOCA-treated tubules, the initial Na^+ -independent pH_i recovery rate (J_{H}) from an acid load was 71% increased in DOCA-treated tubules when compared to 10 control tubules (control $1.08 \pm 0.20 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; DOCA $1.85 \pm 0.26 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P < 0.05$), thus confirming our previous findings on the apical membrane H^+ pump above. Previously, we have found that the lower rate of apical membrane H^+ pump activity determined by this method is due to inhibition of the H^+ pump by a peritubular Na^+ removal-induced sustained increase in cell calcium (22).

However, as shown in Fig. 7, Na^+ -dependent pH_i recovery from an acid load was not significantly different between the two groups (control $28.66 \pm 6.73 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; DOCA $40.98 \pm 6.31 \text{ pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P = \text{NS}$). For chronic DOCA treatment to enhance transepithelial HCO_3^- absorption, one would expect to see an inhibition in basolateral membrane Na^+/H^+ antiporter activity, but in fact if anything, there was a tendency for an increase in Na^+/H^+ antiporter activity. Thus, it is probable that changes in Na^+/H^+ antiporter activity




Figure 7. Effect of chronic DOCA treatment on basolateral membrane Na^+/H^+ antiporter activity. Mean $J_H \pm \text{SEM}$ for control and chronic DOCA tubules is shown.

do not contribute to the increased OMCD_i transepithelial HCO_3^- absorption seen with chronic DOCA treatment.

Chronic adrenalectomy

To determine whether mineralocorticoid deficiency also results in parallel modulation of apical and basolateral membrane $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ transporters, we next examined the effects of chronic ADX. Shown in Table III is a comparison of baseline pH_i , cell volume, and buffer capacities obtained in OMCD_i's harvested from ADX and control animals. Baseline pH_i values obtained in the presence of 25 mM HCO_3^- and 40 mm Hg pCO_2 (pH 7.4, solution 1, Table I) in 37 tubules from ADX rabbits was 6.98 ± 0.02 , which was not significantly different from the baseline pH_i values obtained in 18 tubules from control rabbits (7.02 ± 0.03 , $P = \text{NS}$). In addition, ADX had no effect on cellular volume (control, $6.45 \pm 0.22 \cdot 10^{-10}$ liter/mm, $n = 17$; ADX $6.55 \pm 0.18 \cdot 10^{-10}$ liter/mm, $n = 41$, $P = \text{NS}$). There was no significant difference in buffer capacities between the two groups determined by either $\text{NH}_3/\text{NH}_4^+$ addition or withdrawal (Table III).

Apical membrane H^+ pump activity. Apical membrane H^+ pump activity was assessed as the Na^+ -independent pH_i recovery from an acid load. Shown in Fig. 1 is a typical experimental tracing. Shown in Fig. 8 are the results of this study. ADX led to a 56% reduction in apical membrane H^+ pump activity when compared to control tubules (control, $n = 6$, 1.47 ± 0.21




Figure 8. Effect of chronic adrenalectomy on apical membrane H^+ pump activity. Mean $J_H \pm \text{SEM}$ for control and chronic adrenalectomy tubules is shown.

$\text{pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; ADX, $n = 9$, 0.65 ± 0.07 $\text{pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P < 0.01$).

Basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchange activity. Basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchange activity was assessed as the effect of peritubular Cl^- addition on pH_i , as shown in Fig. 3. Shown in Fig. 9 are the results of these studies. ADX led to a 44% reduction in basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchange activity when compared to control tubules (control, $n = 7$, 43.95 ± 6.78 $\text{pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$; ADX, $n = 26$, 24.60 ± 3.83 $\text{pmol} \cdot \text{mm}^{-1} \cdot \text{min}^{-1}$, $P < 0.05$).

Discussion

Mineralocorticoid hormone receptors are present in all cells of the OMCD_i (23). There is no net transport of Na^+ or K^+ in this segment under normal conditions and chronic DOCA treatment has no effect on the transport of these two ions in this segment (24). Although mineralocorticoid hormones can influence acidification through such factors as volume status, potassium balance, ammoniogenesis, and distal delivery of nonbicarbonate buffer, Hulter et al. (1, 2) found that mineralocorticoids can independently modulate distal nephron acidification. In addition, Stone et al. (3) found that chronic minera-

Table III. Baseline Cell pH and Cellular Volume in Chronic ADX Tubules

	Control	Chronic ADX
Baseline cell pH	7.02 ± 0.03 $n = 18$	6.98 ± 0.02 $n = 37$
Cell vol (10^{-10} liter/mm)	6.45 ± 0.22 $n = 17$	6.55 ± 0.18 $n = 41$
Buffer capacity (mmol · liter ⁻¹ · pH unit ⁻¹)		
(H^+ pump studies)*	25.07 ± 2.09 $n = 7$	26.23 ± 1.95 $n = 9$
($\text{Cl}^-/\text{HCO}_3^-$ exchanger studies)†	29.80 ± 1.40 $n = 7$	29.50 ± 1.29 $n = 26$

Values are means \pm SEM. n = number of tubules. * Buffer capacity was determined by $\text{NH}_3/\text{NH}_4^+$ withdrawal. † Buffer capacity was the sum of the mean $\text{B}_{\text{non-}\text{CO}_2/\text{HCO}_3^-}$ value obtained from $\text{NH}_3/\text{NH}_4^+$ addition in the H^+ pump studies plus the $\text{B}_{\text{CO}_2/\text{HCO}_3^-}$ for each tubule in the $\text{Cl}^-/\text{HCO}_3^-$ exchanger studies.




Figure 9. Effect of chronic adrenalectomy on basolateral membrane $\text{Cl}^-/\text{HCO}_3^-$ exchange activity. Mean $J_H \pm \text{SEM}$ for control and chronic adrenalectomy tubules is shown.

locorticoid treatment of rabbits in vivo increased, and chronic adrenalectomy decreased net H^+ secretion in the OMCD_i perfused in vitro.

Apical membrane H^+ pump. We found a 71–76% stimulation of apical membrane H^+ pump activity by two different apical membrane H^+ pump assays after chronic DOCA treatment, and a 56% decrease in H^+ pump activity after adrenalectomy. The DOCA results are of a similar magnitude to the increase in net transepithelial HCO_3^- absorption found by Stone et al. (3) in OMCD_is dissected from rabbits chronically treated with DOCA. In addition, our ADX results are of a similar magnitude to the 40% reduction in net transepithelial HCO_3^- absorption we found previously (8). Of note, the measured H^+ flux (J_H) across the apical membrane in our studies was less than rates of net transepithelial H^+ secretion found in control and DOCA tubules perfused in vitro (3, 8). The difference may be a result of the nominal presence of ambient CO_2/HCO_3^- in our studies that was removed to decrease any contribution to pH_i by the Cl^-/HCO_3^- exchanger (9). Jacobsen (25) has found that decreasing peritubular CO_2 from 40 mmHg to 10–14 mmHg results in a 52% reduction in net transepithelial H^+ secretion. Thus, in our experiments J_H may have been less than predicted from flux studies due to the nominal presence of CO_2/HCO_3^- .

Enzymatic studies of rat medullary collecting ducts by Mujaïs (4) and rabbit medullary collecting ducts by Garg and Narang (5) found a twofold increase in NEM-sensitive ATPase activity after chronic aldosterone treatment of either normal rats or adrenalectomized rabbits, respectively. In addition, Khadouri et al. (7) found almost a 65% reduction in medullary collecting duct NEM-sensitive ATPase activity in adrenalectomized rats. These results agree with our transport measurements.

With chronic DOCA treatment, enhanced cortical collecting duct K^+ secretion may lead to a state of relative K^+ deficiency. Recently, Wingo (26) found in K^+ -deficient rabbits that 100 μM luminal omeprazole inhibited both net HCO_3^- and K^+ absorption in the OMCD_i suggesting the presence of an H^+-K^+ -ATPase on the apical membrane under these conditions. Previously, we have found that chronic DOCA treatment does not significantly alter rabbit muscle K^+ content (unpublished observations) and recently, Garg (27) found no increase in H^+-K^+ -ATPase activity in the rabbit medullary collecting duct after chronic DOCA treatment. This is in contrast to the marked elevation of H^+-K^+ -ATPase activity they found in the medullary collecting duct under conditions of K^+ deficiency (28). Thus, Na^+ -independent pH_i recovery from an induced acid load under conditions of mineralocorticoid excess most likely represents a vacuolar H^+ pump.

Basolateral membrane Cl^-/HCO_3^- exchanger. In our studies, we found that chronic DOCA treatment led to a 73% stimulation of basolateral membrane Cl^-/HCO_3^- exchange activity, while adrenalectomy led to a 44% decrease in basolateral membrane Cl^-/HCO_3^- exchange activity. In further studies, we found that chronic DOCA treatment led to no change in the apparent $K_{1/2}$ for Cl^- , while the V_{max} was increased. These results are suggestive that chronic DOCA treatment increases the number of active Cl^-/HCO_3^- exchangers present.

Therefore, for both the apical membrane H^+ pump and basolateral membrane Cl^-/HCO_3^- exchanger, mineralocorticoid excess and deficiency increases and decreases, respectively, the activity of these two $H^+/OH^-/HCO_3^-$ transporters in the OMCD_i. These effects occurred to a degree comparable

to the changes found in OMCD_i net transepithelial HCO_3^- absorption under similar conditions of mineralocorticoid excess and deficiency.

Aldosterone mechanism of action. One possible explanation for our results is an aldosterone-induced increase in H^+ pump and Cl^-/HCO_3^- exchanger number. Aldosterone has been shown to increase Na^+-K^+ -ATPase pump number in the cortical collecting duct (20, 29–31) by: (a) acutely stimulating exocytosis of Na^+-K^+ -ATPase pumps from a latent pool of pumps already present within the cell (20, 31); (b) stimulating Na^+-K^+ -ATPase mRNA and protein biosynthesis (32); and, (c) stimulating Na^+-K^+ -ATPase α and β subunit gene transcription (33). Thus, it is possible that aldosterone has a similar mechanism of action in the OMCD_i to increase H^+ pump and Cl^-/HCO_3^- exchanger number.

Khadouri et al. (7) found NEM-sensitive H^+ -ATPase activity from ADX rats to increase threefold after acute (2 h) aldosterone exposure. In addition, the aldosterone-induced increase in NEM-sensitive ATPase activity was prevented by inhibitors of mRNA and protein synthesis, actinomycin D and cycloheximide, respectively. The early sensitivity to cycloheximide and actinomycin D suggests a requirement for protein synthesis, but the synthesized protein need not be the H^+ pump, but may rather be a regulatory protein.

One interesting observation in our studies was the disparity in alterations of cell volume and $H^+/OH^-/HCO_3^-$ transport activity seen with chronic DOCA treatment and ADX. Previously, in the proximal tubule, conditions associated with hypertrophy (chronic metabolic acidosis, chronic K^+ deficiency, and chronic hyperfiltration) have been found to lead to a parallel increase in apical and basolateral membrane $H^+/OH^-/HCO_3^-$ transport processes (34). In our studies, chronic DOCA treatment resulted in similar findings with an increase in cell volume and a parallel increase in apical membrane H^+ pump and basolateral membrane Cl^-/HCO_3^- activities. However, with ADX, decreases in $H^+/OH^-/HCO_3^-$ transport activity were found without detectable changes in cell volume. These observations suggest a possible disparity between the modulation of $H^+/OH^-/HCO_3^-$ transport processes and cell volume under conditions where $H^+/OH^-/HCO_3^-$ transport processes are decreased in parallel. The mechanism(s) of a DOCA-induced increase in cell volume and $H^+/OH^-/HCO_3^-$ transport processes and a ADX-induced decrease in $H^+/OH^-/HCO_3^-$ transport processes without a change in cell volume remain to be elucidated.

Parallel modulation of apical and basolateral membrane $H^+/OH^-/HCO_3^-$ transporters. Parallel modulation of apical and basolateral membrane $H^+/OH^-/HCO_3^-$ transporters in response to a variety of conditions has been extensively documented in the proximal tubule. Parallel adaptive increases in apical membrane Na^+/H^+ antiporter activity and basolateral membrane $Na^+/3HCO_3^-$ cotransporter activity have been found under the chronic conditions of metabolic acidosis (35, 36) and alkalosis (35), respiratory acidosis (37, 38) and alkalosis (37), hyperfiltration (39), and K^+ depletion (40). In addition, acute exposure of the in vitro microperfused proximal tubule to angiotensin II has been found to increase in parallel the activities of these two transporters (41). Under the conditions of chronic mineralocorticoid excess or deficiency, we found no difference in baseline pH_i measured under identical conditions between experimental and control tubules. Chronic DOCA treatment caused a parallel increase in apical mem-

brane H^+ pump activity (76%) and basolateral membrane Cl^-/HCO_3^- exchange activity (73%). Similarly, chronic adrenalectomy resulted in a parallel decrease in apical membrane H^+ pump activity (56%) and basolateral membrane Cl^-/HCO_3^- exchange activity (44%). Thus, parallel regulation of both apical and basolateral membrane $H^+/OH^-/HCO_3^-$ transporters may be a common mechanism to alter net transepithelial transport of HCO_3^- across the nephron without deleterious effects of prolonged alterations in pH_i .

Acknowledgments

The author is grateful to Robert J. Alpern for his continuing support and advice throughout the performance of these studies. The author also gratefully acknowledges the excellent technical assistance of Rebecca Aricheta and Ebtesam M. Shaway.

These studies were supported by National Institute of Health grant R01 DK-39298, and by funds from the National Kidney Foundation of Texas and from the Ruby B. Hexter estate. Steven Hays is a recipient of a Clinical Investigator Award K08 DK 01859.

References

- Hulter, H. N., L. P. Ilnicki, J. A. Harbottle, and A. Sebastian. 1977. Impaired renal H secretion and NH_3 production in mineralocorticoid-deficient glucocorticoid-replete dogs. *Am. J. Physiol.* 232 (Renal Fluid Electrolyte Physiol. 1):F136-F146.
- Hulter, H. N., J. H. Licht, R. D. Glynn, and A. Sebastian. 1979. Renal acidosis in mineralocorticoid deficiency is not dependent on $NaCl$ depletion or hyperkalemia. *Am. J. Physiol.* 236 (Renal Fluid Electrolyte Physiol. 5):F283-F294.
- Stone, D. K., D. W. Seldin, J. P. Kokko, and H. R. Jacobson. 1983. Mineralocorticoid modulation of rabbit medullary collecting duct acidification: a sodium-independent effect. *J. Clin. Invest.* 72:77-83.
- Mujais, S. K. 1987. Effect of aldosterone on rat collecting tubule NEM-sensitive ATPase. *J. Lab. Clin. Med.* 109:34-39.
- Garg, L. C., and N. Narang. 1988. Effects of aldosterone on NEM-sensitive ATPase in rabbit nephron segments. *Kidney Int.* 34:13-17.
- Khadouri, C., S. Marsy, C. Barlet-Bas, and A. Doucet. 1987. Effect of adrenalectomy on NEM-sensitive ATPase along rat nephron and on urinary acidification. *Am. J. Physiol.* 253 (Renal Fluid Electrolyte Physiol. 22):F495-F499.
- Khadouri, C., S. Marsy, C. Barlet-Bas, and A. Doucet. 1989. Short-term effect of aldosterone on NEM-sensitive ATPase in rat collecting tubule. *Am. J. Physiol.* 257 (Renal Fluid Electrolyte Physiol. 26):F177-F181.
- Hays, S. R., J. P. Kokko, and H. R. Jacobson. 1986. Hormonal regulation of proton secretion in rabbit medullary collecting duct. *J. Clin. Invest.* 78:1279-1286.
- Hays, S. R., and R. J. Alpern. 1990. Basolateral membrane Na^+ -independent Cl^-/HCO_3^- exchange in the inner stripe of the rabbit outer medullary collecting tubule. *J. Gen. Physiol.* 95:347-367.
- Schuster, V. L. 1990. Bicarbonate reabsorption and secretion in the cortical and outer medullary collecting tubule. *Semin. Nephrol.* 10:139-147.
- Ridderstrale, Y., M. Kashgarian, B. Koepfen, G. Giebisch, D. Stetson, T. Ardito, and B. Stanton. 1988. Morphological heterogeneity of the rabbit collecting duct. *Kidney Int.* 34:655-670.
- Madsen, K. M., J. W. Verlander, J. Linser, and C. C. Tisher. 1989. Identification of intercalated cells in rabbit medullary collecting duct. *Kidney Int.* 35:458a. (Abstr.)
- Koepfen, B. M. 1985. Conductive properties of the rabbit outer medullary collecting duct: inner stripe. *Am. J. Physiol.* 248 (Renal Fluid Electrolyte Physiol. 17):F500-F506.
- Muto, S., K. Yasoshima, K. Yoshitomi, M. Imai, and Y. Asano. 1990. Electrophysiological identification of α - and β -intercalated cells and their distribution along the rabbit distal nephron segments. *J. Clin. Invest.* 86:1829-1839.
- Weiner, I. D., and L. L. Hamm. 1990. Differential loading of luminal BCECF in the inner stripe of the rabbit outer medullary collecting duct (OMCD_{iii}). *J. Am. Soc. Nephrol.* 1:649a. (Abstr.)
- Thomas, J. A., R. N. Buchsbaum, A. Simnik, and E. Racker. 1979. Intracellular pH measurements in Ehrlich ascites tumor cells utilizing spectroscopic probes generated in situ. *Biochemistry.* 18:2210-2218.
- Roos, A., and W. F. Boron. 1981. Intracellular pH. *Physiol. Rev.* 61:296-434.
- Cordova, H. R., J. P. Kokko, and D. Marver. 1989. Chronic indomethacin increases rabbit cortical collecting tubule Na^+-K^+ -ATPase activity. *Am. J. Physiol.* 256 (Renal Fluid Electrolyte Physiol. 25):F570-F576.
- Field, M. R., B. A. Stanton, and G. H. Giebisch. 1984. Differential acute effects of aldosterone, dexamethasone, and hyperkalemia on distal tubular potassium secretion in the rat kidney. *J. Clin. Invest.* 74:1792-1802.
- Blot-Chabaud, M., F. Wanstock, J.-P. Bonvalet, and N. Farman. 1990. Cell sodium-induced recruitment of Na^+-K^+ -ATPase pumps in rabbit cortical collecting tubules is aldosterone-dependent. *J. Biol. Chem.* 265:11676-11681.
- Hays, S. R., and R. J. Alpern. 1990. Apical and basolateral membrane H^+ extrusion mechanisms in the inner stripe of the rabbit outer medullary collecting duct. *Am. J. Physiol.* 259 (Renal Fluid Electrolyte Physiol. 28):F628-F635.
- Hays, S. R., and R. J. Alpern. 1991. Inhibition of Na^+ -independent H^+ pump by Na^+ -induced changes in cell Ca^{2+} . *J. Gen. Physiol.* 98:791-813.
- Farman, N., M. E. Oblin, M. Lombes, F. Delahaye, H. M. Westphal, J.-P. Bonvalet, and J. M. Gasc. 1991. Immunolocalization of gluco- and mineralocorticoid receptors in the rabbit kidney. *Am. J. Physiol.* 260 (Cell Physiol. 29):C226-C233.
- Stokes, J. B., M. J. Ingram, A. D. Williams, and D. Ingram. 1981. Heterogeneity of the rabbit collecting tubule: localization of mineralocorticoid hormone action to the cortical portion. *Kidney Int.* 20:340-347.
- Jacobson, H. R. 1984. Medullary collecting duct acidification: effects of potassium, HCO_3^- concentration, and pCO_2 . *J. Clin. Invest.* 74:2107-2114.
- Wingo, C. 1989. Active proton secretion and potassium secretion in the rabbit outer medullary collecting duct. Functional evidence for proton-potassium-activated adenosine triphosphatase. *J. Clin. Invest.* 84:361-365.
- Garg, L. C. 1991. Effects of aldosterone on H-K-ATPase activity in rabbit nephron segments. In Aldosterone: Fundamental Aspects. Proceedings of the International Symposium on Aldosterone, France, 1991. J.-P. Bonvalet, N. Farman, M. Lombes, and M.-E. Rafestin-oblin, editors. Colloques INSERM/John Libbey Eurotext Ltd./Montrouge, France. 328a. (Abstr.)
- Garg, L. C., and N. Narang. 1988. Ouabain-insensitive K-adenosine triphosphatase in distal nephron segments of the rabbit. *J. Clin. Invest.* 81:1204-1208.
- Mujais, S. K., M. A. Chekal, W. J. Jones, J. P. Hayslett, and A. I. Katz. 1985. Modulation of renal sodium-potassium-adenosine triphosphatase by aldosterone: effect of high physiologic levels on enzyme activity in isolated rat and rabbit tubules. *J. Clin. Invest.* 76:170-176.
- Blot-Chabaud, M., N. Coutry, J.-P. Bonvalet, and N. Farman. 1991. Influence of intracellular sodium on aldosterone-induced increase in the functional expression of Na^+-K^+ -ATPase. In Aldosterone: Fundamental Aspects. Proceedings of the International Symposium on Aldosterone, France, 1991. J.-P. Bonvalet, N. Farman, M. Lombes, and M.-E. Rafestin-oblin, editors. Colloques INSERM/John Libbey Eurotext Ltd./Montrouge, France. 151-161.
- Barlet-Bas, C., C. Khadouri, S. Marsy, and A. Doucet. 1990. Enhanced intracellular sodium concentration in kidney cells recruits a latent pool of Na^+-K^+ -ATPase whose size is modulated by corticosteroids. *J. Biol. Chem.* 265:7799-7803.
- Verrey, F., E. Schaefer, P. Zoerkler, M. P. Paccolat, K. Geering, J. P. Kraehenbuhl, and B. C. Rossier. 1987. Regulation by aldosterone of Na^+-K^+ -ATPase mRNAs, protein synthesis, and sodium transport in cultured kidney cells. *J. Cell Biol.* 104:1231-1237.
- Verrey, F., J. P. Kraehenbuhl, and B. C. Rossier. 1989. Aldosterone induces a rapid increase in the rate of Na,K -ATPase gene transcription in cultured kidney cells. *Mol. Endocrinol.* 3:1369-1376.
- Alpern, R. J., S. Horie, O. Moe, A. Tejedor, R. T. Miller, and P. A. Preisig. 1991. Chronic adaptations in proximal tubular H/HCO_3^- transporters. *Kidney Int.* 40(Suppl.):S29-S32.
- Akiba, T., V. K. Rocco, and D. G. Warnock. 1987. Parallel adaptation of the rabbit renal cortical sodium/proton antiporter and sodium/bicarbonate cotransporter in metabolic acidosis and alkalosis. *J. Clin. Invest.* 80:308-315.
- Preisig, P. A., and R. J. Alpern. 1988. Chronic metabolic acidosis causes an adaptation in the apical membrane Na/H antiporter and basolateral membrane $Na(HCO_3)_3$ symporter in the rat proximal convoluted tubule. *J. Clin. Invest.* 82:1445-1453.
- Ruiz, O. S., J. A. L. Arruda, and Z. Tabor. 1989. $Na-HCO_3$ cotransport and $Na-H$ antiporter in chronic respiratory acidosis and alkalosis. *Am. J. Physiol.* 256 (Renal Fluid Electrolyte Physiol. 25):F414-F420.
- Krapf, R. 1989. Mechanisms of adaptation to chronic respiratory acidosis in the rabbit proximal tubule. *J. Clin. Invest.* 83:890-896.
- Preisig, P. A., and R. J. Alpern. 1991. Increased Na/H antiporter and $Na/3HCO_3$ symporter activities in chronic hyperfiltration: a model of cell hypertrophy. *J. Gen. Physiol.* 97:195-217.
- Soleimani, M., J. A. Bergman, M. A. Hosford, and T. D. McKinney. 1989. Potassium depletion increases Na^+/H^+ exchange and $Na^+:CO_3^{2-}:HCO_3^-$ cotransport in rat renal cortex. *J. Clin. Invest.* 86:1076-1083.
- Geibel, J., G. Giebisch, and W. F. Boron. 1990. Angiotensin II stimulates both $Na-H$ exchange and Na/HCO_3 cotransport in the rabbit proximal tubule. *Proc. Natl. Acad. Sci. USA.* 87:7917-7920.