

**Influence of the high-affinity growth hormone (GH)-binding protein on plasma profiles of free and bound GH and on the apparent half-life of GH. Modeling analysis and clinical applications**

*J Clin Invest.* 1993;91(5):2335-2335. <https://doi.org/10.1172/JCI116243C1>.

**Correction**

**Find the latest version:**

<https://jci.me/116243C1/pdf>



*Correction*

---

Veldhuis, J. D., M. L. Johnson, L. M. Faunt, M. Mercado, and G. Baumann.  
*The Journal of Clinical Investigation*, Volume 91, No. 2, February 1993.

Page 629.

Because of a publisher's error, the following changes should be made in the text:

Page 630, left hand column, line 4, should read: “. . . equilibrium  $K_d$  of 1.5 nM as a nominal value . . .”

Page 631, Table I: the column heading that now reads “GH bound to protein” should read “GH bound to BP.”

Page 633, left hand column, line 2, should read: “. . . 18, and 36 min (Fig. 6).”

Page 636, right hand column, line 3, should read: “. . . (12, 13, 38, 39).”

Page 637, legend to Figure 7, line 7, should read: “. . . and an equilibrium  $K_d$  of 1.5 nM.”

Page 637, left hand column, line 3, should read: “. . . (41, and unpublished).”

Page 637, left hand column, line 6, should read: “. . . (42, 43) . . .”

Page 637, left hand column, line 7, should read: “. . . (44, 45) . . .”