

Reduced β_1 Receptor Messenger RNA Abundance in the Failing Human Heart

Michael R. Bristow,* Wayne A. Minobe,* Mary V. Reynolds,* J. David Port,*
Randy Rasmussen,* Phillip E. Ray,[§] and Arthur M. Feldman[§]

*Temple Hoyne Buell Laboratories, Division of Cardiology, University of Colorado Health Sciences Center, Denver, Colorado 80262;

[†]Department of Biology, University of Utah, Salt Lake City, Utah 84112; and [§]The Peter Belfer Laboratory for the Molecular Biology of Heart Failure, The Johns Hopkins University School of Medicine, Baltimore, Maryland 21205

Abstract

Heart failure in humans is characterized by alterations in myocardial adrenergic signal transduction, the most prominent of which is down-regulation of β_1 -adrenergic receptors. We tested the hypothesis that down-regulation of β_1 -adrenergic receptors in the failing human heart is related to decreased steady-state levels of β_1 receptor mRNA. Due to the extremely low abundance of β_1 receptor mRNA, measurements were possible only by quantitative polymerase chain reaction (QPCR) or by RNase protection methods. Because the β_1 receptor gene is intronless and β_1 receptor mRNA abundance is low, QPCR yielded genomic amplification in total RNA, and mRNA measurements had to be performed in poly(A)⁺-enriched RNA. By QPCR the concentration of β_1 receptor mRNA varied from 0.34 to 7.8×10^7 molecules/ μ g poly(A)⁺-enriched RNA, and the assay was sensitive to 16.7 zeptomol. Using 100-mg aliquots of left ventricular myocardium obtained from organ donors (nonfailing ventricles, $n = 12$) or heart transplant recipients (failing ventricles, $n = 13$), the respective β_1 mRNA levels measured by QPCR were $4.2 \pm 0.7 \times 10^7/\mu$ g vs. $2.10 \pm 0.3 \times 10^7/\mu$ g ($P = 0.006$). In these same nonfailing and failing left ventricles the respective β_1 -adrenergic receptor densities were 67.9 ± 6.9 fmol/mg vs. 29.6 ± 3.5 fmol/mg ($P = 0.0001$). Decreased mRNA abundance in the failing ventricles was confirmed by RNase protection assays in total RNA, which also demonstrated a 50% reduction in β_1 message abundance. We conclude that down-regulation of β_1 receptor mRNA contributes to down-regulation of β_1 adrenergic receptors in the failing human heart. (*J. Clin. Invest.* 1993. 92:2737–2745.) **Key words:** β -adrenergic • heart failure • messenger RNA • polymerase chain reaction • receptor

Introduction

The failing human heart is characterized by desensitization to adrenergic stimulation (1–5), which results in attenuation of the ability of the heart to increase contractility during exercise or physiologic stress. The majority of this subsensitivity to adrenergic drive is the result of a selective down-regulation of the myocardial β_1 -adrenergic receptor (2–5). In the failing hu-

man heart ventricular myocardial β_1 -adrenergic receptors appear to be down-regulated in all subcellular fractions (4). Since there is no significant β receptor reserve in the human heart (6, 7), β_1 receptor density is a major control point in the regulation of contractility (5). Less clear are the molecular alterations responsible for myocardial β_1 -adrenergic receptor down-regulation in the failing human heart.

In model systems the steady-state abundance of β_2 -adrenergic receptor mRNA and receptor protein levels correlate closely (8, 9). Further, studies in cultured heart cells (10, 11) suggest that the same relationship between mRNA abundance and receptor expression may also exist for the β_1 -adrenergic receptor subtype. Accordingly, we tested the hypothesis that, in the human heart, the steady-state abundance of β_1 -adrenergic receptor mRNA would correlate with β_1 receptor expression at the protein level. Because of the extremely low abundance of β_1 receptor mRNA, we measured β_1 receptor mRNA levels by quantitative polymerase chain reaction (QPCR)¹ (12–14) or by RNase protection (15) assays. Using QPCR with poly(A)⁺-enriched RNA as the starting material or RNase protection in total RNA, the data support the premise that down-regulation of β_1 -adrenergic receptors in failing human ventricular myocardium is related to decreased β_1 -adrenergic receptor mRNA abundance.

Methods

Tissue procurement

Human ventricular myocardium was obtained from subjects undergoing heart or heart/lung transplant, and from organ donors whose hearts were unsuitable for donation owing to blood type or size incompatibility. 15 nonfailing left ventricles were taken from 14 organ donors and one subject undergoing heart/lung transplantation, with all 15 hearts having normal left ventricular function determined by echocardiography. Failing left ventricular myocardium was taken from 13 subjects with end-stage left ventricular failure due to idiopathic dilated cardiomyopathy who underwent heart transplantation. These subjects were not receiving intravenous β -adrenergic receptor agonists, phosphodiesterase inhibitors, or antagonists before transplantation. Mean age of the nonfailing controls was 28.8 ± 5.1 (SEM) yr, compared with 41.7 ± 4.7 yr for the subjects with heart failure ($P = \text{NS}$). Eight of the nonfailing controls were female, and seven were male. Of the failing subjects, 11 were male and two were female.

All subjects with end-stage heart failure had New York Heart Association class IV symptoms before transplantation. Their hemodynamic data included a left ventricular ejection fraction of 0.16 ± 0.01 , cardiac index of 2.01 ± 0.23 liter/min/m², mean pulmonary capillary wedge pressure of 20.5 ± 2.3 mmHg, mean pulmonary artery pressure of 30.5 ± 2.6 mmHg, and mean right atrial pressure of 6.6 ± 2.3 mmHg.

Tissue aliquots were removed from the still-beating heart immediately upon explantation, and either immersed in liquid nitrogen (see

Address reprint requests to Dr. Michael R. Bristow, Temple Hoyne Buell Laboratories, Division of Cardiology, University of Colorado Health Sciences Center, Denver, CO 80262.

Received for publication 12 November 1992 and in revised form 21 July 1993.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/93/12/2737/09 \$2.00

Volume 92, December 1993, 2737–2745

1. Abbreviation used in this paper: QPCR, quantitative PCR.

below) or placed in ice-cold, oxygenated Tyrode's solution as described previously (2).

RNA extraction

1-g aliquots of human left ventricular mid free wall were removed on explantation, immediately immersed in liquid nitrogen, and stored at -70°C until used. Poly(A)⁺-enriched mRNA was extracted from 50–100-mg aliquots of human left ventricular myocardium using oligo(dT) cellulose (Micro-FastTrackTM mRNA Isolation Kit Ver. 1.2, Invitrogen Corp., San Diego, CA). Total RNA was extracted from separate 50–100-mg aliquots by the acid guanidinium thiocyanate phenol-chloroform method (16) using RNA Stat-60 (TelTest, Inc., Friendswood, TX), as previously described (13).

Quantitative PCR

β_1 receptor mRNA measurement. The basic technique for measuring mRNA abundance by QPCR in small aliquots of human heart has been previously described (16). The ethanol precipitated pellet of extracted poly(A)⁺-enriched RNA (0.8–3.2 μg) was dissolved in 16 μl of diethyl pyrocarbonate-treated H_2O , and a 4- μl aliquot was directly measured by a capillary spectrophotometer (model DU-64, Beckman Instruments, Inc., Fullerton, CA). For RNA extracted from both non-failing and failing hearts 200 ng (1–4 μl) of the remaining 12 μl of RNA was then added to the reverse transcriptase reaction, along with 0.5 pg of a synthetic (84mer) cRNA “internal” standard (17). The reverse transcriptase reaction was carried out according to manufacturer's instructions (Promega Corp., Madison, WI), using a 20- μl reaction volume, RNase H-free Moloney murine leukemia virus reverse transcriptase, and random primers. The synthetic RNA (or cRNA) was generated by an in vitro transcription reaction from synthetic DNA templates having a T7 RNA polymerase promoter sequence. The cRNA was DNase (RNase free) treated, phenol/chloroform extracted, and ethanol precipitated before resuspension and quantification by spectrophotometry. The amount of cRNA added to the reverse transcription reaction was determined experimentally to ultimately yield collinear amplification of the β_1 receptor and internal standard cDNAs.

Extensive characterization of amplification of reverse transcribed β_1 receptor mRNA resulted in selection of the following optimal conditions: 2% DMSO and 2 mM MgCl_2 in the Taq polymerase buffer, the use of RNase H to degrade the RNA strand in the RNA:cDNA duplex after reverse transcription, 50 pmol of each primer, and 4 U of Taq polymerase (Promega Corp.) per 100- μl reaction. Additionally, the reaction mixture was denatured at 94°C for 5 min, then cooled to the annealing temperature of 55°C for 2 min, and brought to the 72°C extension temperature before Taq polymerase was added (18). Typically a 4- μl aliquot of the reverse transcriptase/RNase H-treated reaction was used in the PCR reaction, with a final volume of 100 μl .

The primers utilized in the β_1 receptor PCR reaction spanned a 202-bp segment of the human β_1 cDNA sequence (+150 to +351 [17]) and incorporated restriction enzyme recognition sites at the 5' ends for use in the cloning of this segment for other purposes. The sequences were: 5' · CCGAATTCCGAGCCGCTGTCTCAGCAGTGGACA · 3' for the 5' primer and 5' · CCAAGCTTGGTGGCCCCGAACGGC-ACCACCAGCA · 3' for the 3' primer. For subsequent quantification of PCR product yield after gel electrophoretic separation, an aliquot of the 3' primer was end-labeled with [γ - ^{32}P]ATP using T4 polynucleotide kinase (Promega Corp.). All DNA oligonucleotides were purchased from Operon Technologies, Alameda, CA. As described previously (13), 10- μl aliquots were removed from sequential amplification cycles (15 through 33) and fractionated with molecular weight standards by agarose gel electrophoresis. Specific product bands were located by ethidium bromide staining and ultraviolet irradiation, excised, and quantified by Cerenkov counting.

With this technique, separate amplification curves for the unknown cDNA and the internal standard cDNA were constructed. Since a known amount of synthetic RNA internal standard is carried through the cDNA synthesis and amplification, the original β_1 receptor mRNA

level can be determined by extrapolation from the internal standard cDNA standard curve. It is essential that this extrapolation be performed during the exponential phase of amplification and that the curves for the unknown and internal standard are collinear, which occurred in all cases between cycles 20 and 30. An RNA sample not subjected to reverse transcription is run as a control for contamination by genomic DNA.

β_2 mRNA abundance was measured in total RNA with linear amplification occurring in all cases between cycle numbers 20 and 30. The assay conditions for measuring β_2 receptor mRNA abundance by QPCR have been previously described (13, 19); 1 μg of total RNA from both failing and nonfailing groups plus 30 pg of cRNA was used in the reverse transcriptase reaction. Typically a 1 μl aliquot of the reverse transcriptase/RNase H treated reaction was used in the PCR reaction.

Measurement of β_1 receptor mRNA by RNase protection. The human β_1 -adrenergic receptor cDNA (17) was subcloned into pBluescript II KS (Stratagene, Inc., La Jolla, CA) and digested with either SmaI or EagI. Antisense RNA transcribed from the T3 promoter of the SmaI digested vector comprised ~ 450 bp of the terminal end of the 3' untranslated region (SmaI probe, probe I). Antisense RNA transcribed from the T3 promoter of the EagI digested vector contained ~ 400 bp of the 3' end of the coding region and the entire 3' untranslated region (~ 700 bp) (EagI probe, probe II).

Antisense riboprobes were transcribed from the appropriately digested β_1 receptor cDNA using the Maxiscribe kit (Ambion, Inc., Austin, TX). The riboprobes were labeled by incorporation of [α - ^{32}P]UTP 800 Ci/mM (New England Nuclear, Boston, MA) into the RNA during transcription with T3 DNA-dependent RNA polymerase. Approximately $1\text{--}2 \times 10^5$ cpm of ^{32}P -labeled antisense β_1 RNA was used in each nuclease protection assay.

Labeled riboprobes were hybridized in solution with 2 or 5 μg of total RNA extracted from failing or nonfailing explanted hearts for 18 h at 44°C . Unhybridized single-stranded RNA was digested with RNaseA and RNaseT1. RNA-RNA hybrids were resolved by electrophoresis in an 8% acrylamide denaturing gel and visualized by autoradiography at -80°C . The RPAII kit (Ambion, Inc.) was used for solution hybridization and RNase digestion.

Protected RNA species were quantified by densitometry. In all experiments in vitro transcribed β_1 receptor sense strand RNA generated by the same method from the same vector as the antisense riboprobes served as a positive control. In addition, in order to verify the specificity of the β_1 probes for β_1 mRNA, the synthetic full-length β_2 mRNA (20) was transcribed and 100 pg hybridized with each of the two β_1 antisense riboprobes.

Hybridization intensity for protected RNA species of the appropriate size was measured by densitometry of autoradiograms developed for 2–7 d. Two or three nonfailing control samples were run on each gel and the OD reading averages of these samples gave the 100% or control value. Each sample from failing or nonfailing hearts was then referenced to this 100% value, with results reported as percentage of control.

β -Adrenergic receptor quantification. β_1 and β_2 adrenergic receptors were quantified in a crude membrane fraction prepared from 5-g aliquots of left ventricular free wall, using previously described methods (2). Briefly, the total population of β_1 plus β_2 receptors was measured by the nonselective radioligand [^{125}I]iodocyanopindolol. Maximum binding (B_{max}) and the [^{125}I]iodocyanopindolol dissociation constant (K_d) were determined by nonlinear least-squares computer modeling of the specific binding curve (2). Fractions of β_1 vs. β_2 receptors in the membrane preparations were determined by computer modeling of competition curves using the highly selective β_1 antagonist CGP-20712A (4). Multiplication of the total B_{max} by the fraction of β_1 receptors obtained from CGP-20712A/[^{125}I]iodocyanopindolol competition curves then yielded β_1 receptor density.

Protein concentrations were determined by the Peterson modification of the method of Lowry (21). RNA concentration and purity was determined by measuring absorbance at 260 and 280 nM using a highly sensitive capillary spectrophotometer (Beckmann Instrument model DU-64).




Figure 1. Assessment of genomic DNA contamination levels between poly(A)⁺-enriched RNA (H, I, J, K, P, Q, R, S) and total RNA (L, M, N, O, T, U, V, W) fractions subjected to QPCR. 200 ng of poly(A)⁺-enriched RNA and 1 μ g of total RNA extracted from two failing and two nonfailing hearts were added to duplicate reverse transcription reactions differing only in whether they contained reverse transcriptase or not. Subsequently, 10% of each of the total RNA reactions and 20% of the poly(A)⁺ reactions were amplified by PCR. All the reverse transcription reactions with transcriptase (H, I, J, K, L, M, N, O) show strong amplification of specific 202-bp product (compared with the positive synthetic β_1 receptor mRNA control, (C) but of the reactions without transcriptase (P, Q, R, S, T, U, V, W) only the poly(A)⁺-enriched RNA reactions (P, Q, R, S) do not amplify the 202-bp product. (A) DNA molecular weight markers. (B) PCR reagent blank. (C) Synthetic, full-length β_1 receptor mRNA/positive amplification control. (D, E, F, G) Synthetic β_1 receptor mRNA/no reverse transcriptase controls.

Statistical analysis

Comparison of numerical data between two groups was by an unpaired t-test. Correlation analysis was by simple regression. All statistical computation was performed on the StatView 512 + TM program (Brainpower, Inc., Calabasas, CA). Statistical significance was taken as $P < 0.05$ in a two-tailed distribution.

Results

QPCR

β_1 receptor mRNA abundance. Shown in Fig. 1 is a 4% agarose gel stained with ethidium bromide in which were run PCR products generated from four samples each of total and poly(A)⁺-enriched RNA, extracted from two nonfailing and two failing human ventricles. In the absence of reverse transcriptase [RT(−) condition], genomic material of the expected size for the β_1 receptor PCR product is clearly present in the total RNA samples after 38 cycles. In contrast, poly(A)⁺-enriched RNA shows no amplification under RT(−) conditions. Based on these results we used poly(A)⁺-enriched RNA as the starting material for QPCR quantification of β_1 mRNA abundance.

Fig. 2 shows an ethidium stained 4% agarose gel in which were separated the contents of six PCR reactions using poly(A)⁺-enriched RNA as the starting material. The amplified cDNAs were made by reverse transcribing 36–44 ng of poly(A)⁺-enriched RNA extracted from six samples of human left ventricle (designated A, B, C, D, E, and F) along with 200–500 fg of the internal standard cRNA. Each PCR reaction is represented by a set of nine sequential 10- μ l aliquots which were removed from the reaction tube at (left to right) cycles 14, 17, 19, 21, 24, 25, 27, 29, and 32. The PCR amplification product for β_1 receptor mRNA appears as a 202-bp band, while

the internal standard appears as an 84-bp band. The gel also includes “genomic controls”: that is, poly(A)⁺-enriched RNA from the six samples (labeled, respectively, G–L) not subjected to reverse transcription. There was progressive amplification of the target mRNA and the internal standard cRNA in all six samples (A–F). However, there was no amplification of genomic DNA in any sample (G–L). Identity of the 202-bp band was confirmed by direct PCR sequencing (22).

Fig. 3 shows amplification curves from four individual samples (two failing and two nonfailing) using 1.9–11.4 fg of internal standard cRNA per ng of poly(A)⁺ RNA. It can be seen that there is collinear amplification between 20 and 30 cycles; curves were accepted for analysis only if amplification was linear for at least six cycles.

In order to compare the efficiency of reverse transcription of the internal standard cRNA to that of the β_1 receptor mRNA, we measured by QPCR a dilution series of cDNAs reverse transcribed from known, equal amounts (16.7 zeptomol to 16.7 amol of each; or 10^4 to 10^7 molecules) of the cRNA and full-length β_1 receptor mRNA. The latter was generated by in vitro transcription from a pBluescript IIS phagemid expression vector. An amplification curve was constructed for each concentration of the series, and quantities of mRNA vs. cRNA were calculated from equivalent points in the exponential phase of amplification. As shown in Fig. 4, the incorporation of labeled primers into the PCR products made from the internal standard cRNA was essentially identical to those for the full-length β_1 receptor mRNA. In order to determine if the QPCR assay was linear over the expected range of β_1 receptor mRNA concentrations, we added increasing amounts of in vitro transcribed β_1 receptor mRNA to poly(A)⁺-enriched RNA that contained a β_1 mRNA abundance of 2.5×10^7 molecules/ μ g. Using either 0.25 or 0.50 pg of cRNA as the internal stan-




Figure 2. Agarose gels containing the fractionated contents of six separate PCR reactions using poly(A)⁺-enriched RNA subjected to reverse transcription and PCR. Amplified β_1 receptor mRNA appears as a 202-bp product, and the 84-bp product is the internal standard. MW, molecular weight markers. 10- μ l aliquots were removed from the PCR reaction in cycles 14, 17, 19, 21, 24, 25, 27, 29, and 32, in lanes 1–9 from left to right. G–L are cycles 14, 25, and 32 for each of six samples in which reverse transcriptase was omitted (“genomic control”).

dard, the assay was linear with respect to added mRNA from measured values of 2.5×10^7 to 18×10^7 molecules/ μ g poly(A)⁺ RNA.

Table I gives the QPCR quantification of β_1 mRNA in poly(A)⁺-enriched samples from 13 failing and 12 nonfailing human hearts. Compared to nonfailing controls, failing hearts had a 50% lower abundance of β_1 -adrenergic receptor mRNA. This finding was similar to the 56% lower density of β_1 receptor observed in the failing vs. nonfailing heart and β_2 receptor mRNA abundance.

β_2 receptor mRNA abundance. β_2 receptor mRNA abundance was also measured by QPCR, in total RNA extracted from nonfailing and failing human hearts (Table II). In contrast to β_1 receptor mRNA, collinear amplification of reverse-transcribed β_2 message always occurred in < 30 cycles. Genomic amplification was not observed for the β_2 mRNA assay, presumably because of the lower PCR cycle requirements (data not shown). As can be seen in Table I, there was no difference ($P = \text{NS}$) in β_2 receptor mRNA abundance or β_2 receptor density between nonfailing and failing hearts.

Relation between β_1 receptor mRNA abundance and β_1 or β_2 receptor density. Fig. 5 gives the simple regression relation

between QPCR-measured β_1 receptor mRNA abundance and β_1 or β_2 receptor density in the 25 (13 failing and 12 nonfailing) left ventricles. There is a significant direct relation between β_1 mRNA and β_1 receptor density (*top panel*, $r = 0.52$, $P = 0.008$) but no relation between β_1 mRNA and β_2 receptor density (*bottom panel*, $r = 0.07$, $P = \text{NS}$).

RNase protection

Nuclease protection assays were also used to measure β_1 mRNA abundance, in total RNA extracted from explanted failing and nonfailing hearts. A representative autoradiogram from an RNase protection assay utilizing probe I in total RNA is shown in Fig. 6. In Table II are given the β_1 mRNA abundance measurements for probes I and II, and for the average of both probes when both were used in the same sample. The abundance of β_1 mRNA was less in failing than in nonfailing hearts, confirming the results of quantitative PCR. Using the average data for both probes, β_1 receptor mRNA was decreased by 50.2% and β_1 receptor density by 65.2% in the same hearts (Table II). As shown in Fig. 7, β_1 mRNA measurement by RNase protection was highly correlated ($r = 0.90$, $P < .001$) with QPCR measurements in the same samples, which in-

Table I. β_1 and β_2 Receptor mRNA Abundances Measured by QPCR and β_1 and β_2 Receptor Densities in Nonfailing and Failing Human Left Ventricular Myocardium

Group (n)	β_1 mRNA abundance <i>molecules $\times 10^7/\mu$g poly(A)⁺ RNA</i>	β_1 receptor density <i>fmol/mg</i>	β_2 mRNA abundance <i>molecules $\times 10^7/\mu$g total RNA</i>	β_2 receptor density <i>fmol/mg</i>
Nonfailing (12)	4.2 \pm 0.7	67.9 \pm 6.9	2.2 \pm 0.6	22.1 \pm 1.9
Failing (13)	2.1 \pm 0.3*	29.6 \pm 3.5*	1.6 \pm 0.3	17.7 \pm 1.3

Values are given \pm SEM. * $P < 0.01$ vs. nonfailing controls.




Figure 3. PCR amplification of reverse transcription products (cDNA) of β_1 receptor mRNA from poly(A)⁺-enriched RNA (○, *BETA 1*) and internal standard cRNA from the β_1 construct (●, *CONSTRUCT*).

cluded RNA extracted from four nonfailing and six failing ventricles.

To verify the specificity of the antisense probes full-length sense human β_2 -adrenergic receptor mRNA was transcribed and hybridized to both antisense β_1 probes. The two β_1 antisense probes used were highly specific for β_1 -adrenergic receptor mRNA, as neither the *Sma*I (probe I) nor the *Eag*I β_1 probe (probe II) protected a fragment of β_2 mRNA of the correct size (data not shown).

As the β_1 -adrenergic receptor gene is apparently intronless (Fig. 1), protection of genomic DNA present as a contaminant in the RNA preparations could be a potential problem as it was with QPCR. Accordingly, RNA extracted from a nonfailing heart was treated with DNase I before hybridization with the *Sma*I riboprobe. By densitometry the abundance of β_1 mRNA in DNase I treated total RNA was no different from the abundance of β_1 mRNA in untreated total RNA from the same sample (data not shown).

Discussion

Using QPCR we have previously reported that the failing human heart exhibits marked abnormalities of gene expression, including de novo expression of atrial natriuretic peptide mRNA and decreased levels of phospholamban mRNA (13). Furthermore, these changes in gene expression are potentially reversible with restoration of normal cardiac function (23).

Within the context of the altered β receptor-G-protein-adenylyl cyclase signal transduction pathways in the failing heart, the molecular mechanisms responsible for the various described abnormalities (1–5) remain elusive. The mRNA levels for the β_2 adrenergic receptor (19) and the G proteins, $G_{\alpha s}$ (19) and $G_{\alpha 13}$ (19), are not different in total RNA extracted from failing and nonfailing human ventricular myocardium. This is not totally unexpected, since the respective gene products are not quantitatively altered in the failing human heart (5). Based on previous work in the failing human heart (2, 3)




Figure 4. Simultaneous measurement of identical molar amounts of β_1 receptor mRNA (■, BETA-1) and β_1 cRNA internal standard (standard ○, INT STND) by quantitative PCR.

and in cell culture model systems (10, 11), one element of the receptor-G-protein-adenylyl cyclase complex that is a strong candidate for abnormal gene expression is the β_1 -adrenergic receptor.

By Northern analysis or QPCR β_1 -adrenergic receptor mRNA abundance measurements in total RNA extracted from human heart have thus far been unreliable, due to low abundance of the G + C rich β_1 receptor message (17) and genomic contamination of PCR products. Based on the structures of closely related genes (20, 24) and our PCR data the β_1 -adrenergic receptor gene likely contains no introns, and therefore PCR primers could not be chosen to differentiate the cDNA amplification signal from that of genomic DNA. The need for high cycle numbers (> 35) in detecting this low-abundance message exacerbated the genomic DNA contamination problem. Consequently, it was necessary to use poly(A)⁺-enriched RNA as the starting material. In using these conditions, fewer than 32 cycles of PCR were required to produce satisfactory levels of amplification. Furthermore, under these conditions, amplifications of non-reverse-transcribed [RT(-)] RNA produced no detectable signal, indicating the elimination of genomic DNA contamination.

In using quantitative QPCR it is mandatory that collinear amplification be obtained with the internal standard. This, in turn, requires that approximately equal amounts of internal standard and mRNA are present in the initial reaction mixture. Additionally, it is important that this collinear amplification occur at relatively low cycle numbers (i.e., < 35), to avoid artifacts caused by PCR. These criteria were satisfied by the above described assay conditions.

In failing ventricles, steady-state β_1 receptor mRNA levels measured by QPCR were 50% lower than in nonfailing control hearts. In these same failing hearts β_1 receptor density was 56% lower than in nonfailing controls. Moreover, when all 25 ventricles in which β_1 receptor mRNA abundance was measured were considered, there was a direct relation between β_1 mRNA abundance and β_1 receptor density, but no relation between β_1 mRNA and β_2 receptor density. The degree of down-regulation of mRNA and the degree of down-regulation of β_1 -adrenergic

Table II. β_1 Receptor mRNA Abundance Measured by RNase Protection and β_1 -Adrenergic Receptor Density in Nonfailing and Failing Human Ventricular Myocardium

Group	Probe I (SmaI digest)	Probe II (EagI digest)	Both riboprobes	β_1 receptor density
	% of control			fmol/mg
Nonfailing	100.6±13.3 (n = 7)	99.5±19.1 (n = 4)	99.4±9.5 (n = 7)	83.0±10.3 (n = 7)
Failing	52.3±9.0* (n = 6)	48.8±9.3* (n = 6)	49.7±6.6* (n = 6)	29.9±4.2* (n = 6)

Values are given±SEM. * $P < .05$ vs. nonfailing.

receptors were therefore quite similar considering the vast array of factors which could potentially influence the regulation of both mRNA and proteins. These factors include altered transcriptional and translation frequency, mRNA stability, and receptor protein turnover. Also, the fact that the nonfailing controls were obtained from organ donors who are invariably subjected to very high levels of endogenous sympathetic drive due to brain injury (25) cannot be ignored. Since the kinetics of mRNA turnover are likely to be faster than receptor turnover (i.e., mRNA half-life vs. protein half-life), it may be that mRNA abundance was already beginning to decrease in the nonfailing controls.

Decreased β_1 receptor mRNA in failing left ventricles was confirmed by RNase protection assay, which showed a relative decrease in message levels by 50% in six failing hearts. The presence of β_1 receptor genomic material contaminating the total RNA in these assays did not pose a problem, presumably due to the preference of the riboprobes for RNA-RNA vs. RNA-DNA hybridization (26). Of the two riboprobes, the 3' UTR probe (SmaI digest, probe I) gave the best hybridization signal. Both probes were specific for β_1 receptor mRNA, as in vitro transcribed β_2 mRNA did not cross-protect against RNase degradation. The fact that in two RNA fractions two totally independent methods, QPCR and RNase protection, yielded decreased β_1 receptor mRNA abundance measurements in failing human heart provides strong evidence that transcript steady-state level is one important determinant of β_1 receptor regulation in human ventricular myocardium.

Despite the 50% decrease in β_1 mRNA in poly(A)⁺-enriched or total RNA extracted from failing human hearts β_2 mRNA abundance in total RNA was similar in failing and in nonfailing ventricles. This is in agreement with our previously reported data in total RNA (19). Apparently because β_2 receptor mRNA abundance is greater than β_1 in the human heart, QPCR can be performed in total RNA with relatively low (< 30) cycle numbers and genomic contamination can be avoided (19).

There are two previous reports of QPCR measurements of β_1 receptor mRNA in failing human ventricular myocardium, one reporting an increase compared to nonfailing controls (27) and one reporting a decrease (28). In the previous report showing a decrease (28), 38 cycles of PCR were used on total RNA and no RT(-) control condition was reported. The report of an increase in β_1 mRNA in failing ventricles also used total RNA (27). Again, the combination of low abundance and an





Figure 5. Simple regression plots of β_1 receptor mRNA (y-axis) vs. β_1 receptor density (top panel) and β_2 receptor density (bottom panel) in the 25 left ventricles in which β_1 mRNA abundance was measured by QPCR.




Figure 6. A representative autoradiogram of an RNase protection assay. Total RNA extracted from explanted human hearts was hybridized with an [$\alpha^{32}\text{P}$]UTP-labeled antisense β_1 riboprobe (SmaI digest of β_1 cDNA; 450-bp probe). Fragments protected from RNase digestion were resolved by denaturing gel electrophoresis and visualized by autoradiogram. Lanes 1 and 2: SmaI antisense β_1 riboprobe without (lane 1)

and with (lane 2) RNase treatment; lane 2 contains eight times the amount of riboprobe as lanes 1 or 3–8. The smaller, faster migrating bands present in all lanes are probably self-protected probe fragments. As the probe is very G + C rich (17), higher order structures in the single stranded antisense probe are likely to be present, even under denaturing conditions. These structures will protect portions of the probe from RNase digestion. Lanes 3, 4, and 5: synthetic β_1 mRNA (200, 100, and 50 pg) hybridized with the labeled β_1 riboprobe. The prominent lower bands in the standard lanes are most likely due to premature termination of transcription of the synthetic sense β_1 mRNA. Lanes 6, 7, and 8: Total RNA from three different explanted nonfailing human hearts hybridized with the antisense β_1 riboprobe. Lanes 9, 10, and 11: Total RNA from three different explanted failing human hearts hybridized with the antisense β_1 riboprobe. β_1 mRNA abundance is significantly lower in failing human hearts compared to the abundance in nonfailing human hearts as determined by RNase protection assay.




Figure 7. Correlation of β_1 mRNA abundance measured by QPCR in poly(A)⁺-enriched RNA vs. measurement by RNase protection in total RNA; four nonfailing and six failing human left ventricles.

intronless gene makes measurements of β_1 receptor mRNA levels by QPCR challenging, and unreliable in total RNA using high (> 30) cycle numbers. Depending on the DNA/RNA ratio in myocardium comprising the starting material, β_1 "mRNA" abundance in total RNA could be found to be increased or decreased.

There are two previous reports of RNase protection performed as solution hybridization with the hybrids isolated by filtration (29, 30). Although identical riboprobes were used in these two investigations (29, 30), one study in idiopathic dilated cardiomyopathy found an increase in β_1 receptor "mRNA" (30) and the other in valvular heart disease (30) found a decrease. Based on the number of self-protecting fragments detected on acrylamide gels in our own RNase protection assays, it is obvious that the appropriately sized RNA:RNA hybrid needs to be separated and isolated by gel electrophoresis. Unless it has been documented that a single protected or a very dominant protected product accounts for the overwhelming majority of the measured radioactivity in the solution assay, filtration isolation of hybridized material can lead to spurious results. Circumstances such as these may have contributed to the divergent results in these two previous studies (29, 30).

Although the starting material (left ventricular myocardium) used for mRNA and receptor measurements was multicellular, the majority of cell volume, protein, and RNA in ventricular myocardium is derived from cardiac myocytes. In mammalian ventricles β_1 adrenergic receptors are selectively expressed in myocardial cells (31), and therefore it is likely that the vast majority of β_1 receptor protein and mRNA measured in this investigation was derived from cardiac myocytes. We have previously shown that this is in fact the case for β_1 -adrenergic receptors measured in crude membrane fractions extracted from human ventricles (2).

In summary, β_1 receptor mRNA abundance is lower in the failing human heart than in nonfailing, control hearts. This decrease can fully account for the observed down-regulation of β_1 -adrenergic receptor expression in failing hearts. Further studies will be required to determine the molecular basis for this alteration in gene expression.

Acknowledgments

We wish to thank Dr. Carl T. Wittwer of the University of Utah Medical School for helpful suggestions on optimal conditions for PCR, Dr.

M. Benjamin Perryman for critical review of the manuscript, and Frank Stewart for manuscript preparation.

This work was supported by National Institutes of Health (NIH) grants HL-13408-15 and HL-48013-01A2, awarded to Dr. Michael R. Bristow. Dr. Arthur M. Feldman was supported by NIH Program Grant HL-39719 and The W. W. Smith Charitable Trust.

References

1. Bristow, M. R., R. Ginsburg, W. Minobe, R. S. Cubicciotti, W. S. Sageman, K. Lurie, M. E. Billingham, D. C. Harrison, and E. B. Stinson. 1982. Decreased catecholamine sensitivity and β -adrenergic-receptor density in failing human hearts. *N. Engl. J. Med.* 307:205-211.
2. Bristow, M. R., R. Ginsburg, M. Fowler, W. Minobe, R. Rasmussen, P. Zera, R. Menlove, P. Shah, and E. Stinson. 1986. β_1 and β_2 adrenergic receptor subpopulations in normal and failing human ventricular myocardium: coupling of both receptor subtypes to muscle contraction and selective β_1 receptor down-regulation in heart failure. *Circ. Res.* 59:297-309.
3. Brodde, O. R., S. Schuler, R. Kretsch, M. Brinkmann, H. G. Borst, R. Hetzer, J. C. Reidemeister, W. Warnecke, and H. R. Zerkowski. 1986. Regional distribution of β -adrenoceptors in the human heart: coexistence of function β_1 - and β_2 -adrenoceptors in both atria and ventricles in severe congestive cardiomyopathy. *J. Cardiovasc. Pharmacol.* 8:1235-1242.
4. Bristow, M. R., W. Minobe, R. Rasmussen, P. Larrabee, L. Skerl, J. W. Klein, F. L. Anderson, L. Murray, L. Mestroni, S. V. Karwande, et al. 1992. β -Adrenergic neuroeffector abnormalities in the failing human heart are produced by local, rather than systemic mechanisms. *J. Clin. Invest.* 89:803-815.
5. Bristow, M. R., R. E. Hersberger, J. D. Port, E. M. Gilbert, A. Sandoval, R. Rasmussen, A. E. Cates, and A. M. Feldman. 1990. β -Adrenergic pathways in nonfailing and failing human ventricular myocardium. *Circulation.* 82(Suppl. 1):12-25.
6. Port, J. D., and M. R. Bristow. 1988. Lack of spare β -adrenergic receptors in the human heart. *FASEB J.* 2:A602.
7. Brown, L., N. M. Deighton, S. Bals, W. Sohlmann, H. R. Zerkowski, M. C. Michel, and O. E. Brodde. 1992. Spare receptors for β -adrenoceptor-mediated positive inotropic effects of catecholamines in the human heart. *J. Cardiovasc. Pharmacol.* 19:222-232.
8. Hadcock, J. R., and C. C. Malbon. 1988. Down-regulation of beta-adrenergic receptors: agonist-induced reduction in receptor mRNA levels. *Proc. Natl. Acad. Sci. USA.* 85:5021-5025.
9. Hadcock, J. R., H. Y. Wang, and C. C. Malbon. 1989. Agonist-induced destabilization of β -adrenergic receptor mRNA: attenuation of glucocorticoid-induced up-regulation of β -adrenergic receptors. *J. Biol. Chem.* 264:19928-19933.
10. Bristow, M. R., C. Durham, J. Klein, R. Rasmussen, J. D. Port, R. R. Gesteland, W. H. Barry, and A. M. Feldman. 1991. Down-regulation of β -adrenergic receptors and receptor mRNA in heart cells chronically exposed to norepinephrine. *Clin. Res.* 39:256A. (Abstr.)
11. Lazar-Wesley, E., J. R. Hadcock, C. C. Malbon, G. Kunos, and E. J. Ishac. 1991. Tissue-specific regulation of α_{1B} , β_1 , and β_2 -adrenergic receptor mRNAs by thyroid state in the rat. *Endocrinology.* 129:1116-1118.
12. Wang, A. M., M. V. Doyle, and D. F. Mark. 1989. Quantitation of mRNA by the polymerase chain reaction. *Proc. Natl. Acad. Sci. USA.* 86:9717-9721.
13. Feldman, A. M., P. E. Ray, C. M. Silan, J. A. Mercer, W. Minobe, and M. R. Bristow. 1991. Selective gene expression in failing human heart. *Circulation.* 83:1866-1872.
14. Bouaboula, M., P. Legoux, B. Pessegue, B. Delpech, X. Dumont, M. Piechaczyk, P. Casellas, and D. Shire. Standardization of mRNA titration using a polymerase chain reaction method involving co-amplification with a multispecific internal control. 1992. *J. Biol. Chem.* 267:21830-21838.
15. Lee, J. J., and N. A. Costlow. 1987. A molecular titration assay to measure transcript prevalence levels. *Methods Enzymol.* 152:633-648.
16. Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162:156-159.
17. Frielle, T., S. Collins, K. W. Daniel, M. G. Caron, R. J. Lefkowitz, and B. K. Kobilka. 1987. Cloning of the cDNA for the human beta 1-adrenergic receptor. *Proc. Natl. Acad. Sci. USA.* 84:7920-7924.
18. Kawasaki, E. S. 1990. Amplification of RNA. In PCR Protocols—A Guide to Methods and Applications. M. A. Innis and D. H. Gelfand, editors. Academic Press, Inc., San Diego, CA. 3-12.
19. Feldman, A. M., P. E. Ray, and M. R. Bristow. 1991. Expression of α -subunits of G proteins in failing human heart: a reappraisal utilizing quantitative polymerase chain reaction. *J. Mol. Cell. Cardiol.* 23:1355-1358.
20. Dixon, R. A. F., B. K. Kobilka, D. J. Strader, J. L. Benovic, H. G. Dohman, T. Frielle, M. A. Bolanowski, C. D. Bennett, E. Rands, R. A. Diehl, et al. 1986. Cloning of the gene and cDNA for mammalian β -adrenergic receptor and homology with rhodopsin. *Nature (Lond.).* 321:75-79.

21. Peterson, G. L. 1977. A simplification of the protein assay method of Lowry et al. which is more generally applicable. *Anal. Biochem.* 83:346–356.
22. Gyllenstein, U. B., and H. A. Erlich. 1988. Generation of single-stranded DNA by the polymerase chain reaction and its application to direct sequencing of the HLA-DQA locus. *Proc. Natl. Acad. Sci. USA.* 85:7652–7656.
23. Ladenson, P. W., S. I. Sherman, K. L. Baughman, P. E. Ray, and A. M. Feldman. 1992. Reversible alterations in myocardial gene expression in a young man with dilated cardiomyopathy and hypothyroidism. *Proc. Natl. Acad. Sci. USA.* 89:5251–5255.
24. Shimomura, H., and A. Terada. 1990. Primary structure of the rat beta-1 adrenergic receptor gene. *Nucleic Acids Res.* 18:4591.
25. Hamill, R. W., P. D. Woolf, J. V. McDonald, L. A. Lee, and M. Kelly. 1987. Catecholamines predict outcome in traumatic brain injury. *Ann. Neurol.* 21:438–443.
26. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning, a Laboratory Manual*. 2nd edition. Cold Spring Harbor Press, Cold Spring Harbor, NY.
27. Ihl-Vahl, R., R. Marquetant, and R. H. Strasser. 1992. Subtype-selective regulation of mRNA for β_1 -adrenergic receptor in chronic heart failure. *Circulation.* 86(Suppl. I):I-767. (Abstr.)
28. Ungerer, M., M. Bohm, J. S. Elce, E. Erdmann, and M. J. Lohse. 1993. Altered expression of β -adrenergic receptor kinase and β_1 -adrenergic receptors in the failing human heart. *Circulation.* 87:454–463.
29. Sylven, C., M. Bronnegard, E. Jansson, P. Sotonyi, F. Liang-Xiong, F. Waagstein, and A. Hjalmarsson. 1992. Increased β_1 receptor expression at mRNA level in dilated cardiomyopathy. *J. Am. Coll. Cardiol.* 19:273A. (Abstr.)
30. Sylven, C., P. Arner, L. Hellstrom, E. Jansson, P. Sotonyi, A. Somogyi, and M. Bronnegard. 1991. Left ventricular β_1 and β_2 adrenoceptor mRNA expression in normal and volume overloaded human heart. *Cardiovasc. Res.* 25:737–741.
31. Murphree, S. S., and J. E. Saffitz. 1988. Delineation of the distribution of beta-adrenergic receptor subtypes in canine myocardium. *Circ. Res.* 63:117–125.