



SHP-2/*PTPN11* mediates gliomagenesis driven by *PDGFRA* and *INK4A/ARF* aberrations in mice and humans

Kun-Wei Liu,^{1,2} Haizhong Feng,^{1,2} Robert Bachoo,³ Andrius Kazlauskas,⁴ Erin M. Smith,⁵ Karen Symes,⁵ Ronald L. Hamilton,² Motoo Nagane,⁶ Ryo Nishikawa,⁷ Bo Hu,^{1,8} and Shi-Yuan Cheng^{1,2}

¹University of Pittsburgh Cancer Institute and ²Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

³Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, Texas, USA. ⁴Schepens Eye Research Institute and Harvard Medical School, Boston, Massachusetts, USA. ⁵Department of Biochemistry, Boston University School of Medicine, Boston, Massachusetts, USA.

⁶Department of Neurosurgery, Kyorin University Faculty of Medicine, Tokyo, Japan. ⁷Department of Neurosurgery, Saitama Medical University, Saitama-ken, Japan. ⁸Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

Recent collaborative efforts have subclassified malignant glioblastomas into 4 clinical relevant subtypes based on their signature genetic lesions. Platelet-derived growth factor receptor α (*PDGFRA*) overexpression is concomitant with a loss of cyclin-dependent kinase inhibitor 2A (*CDKN2A*) locus (encoding P16INK4A and P14ARF) in a large number of tumors within one subtype of glioblastomas. Here we report that activation of PDGFR α conferred tumorigenicity to *Ink4a/Arf*-deficient mouse astrocytes and human glioma cells in the brain. Restoration of p16INK4a but not p19ARF suppressed PDGFR α -promoted glioma formation. Mechanistically, abrogation of signaling modules in PDGFR α that lost capacity to bind to SHP-2 or PI3K significantly diminished PDGFR α -promoted tumorigenesis. Furthermore, inhibition of SHP-2 by shRNAs or pharmacological inhibitors disrupted the interaction of PI3K with PDGFR α , suppressed downstream AKT/mTOR activation, and impaired tumorigenesis of *Ink4a/Arf*-null cells, whereas expression of an activated PI3K mutant rescued the effect of SHP-2 inhibition on tumorigenicity. PDGFR α and PDGF-A are co-expressed in clinical glioblastoma specimens, and such co-expression is linked with activation of SHP-2/AKT/mTOR signaling. Together, our data suggest that in glioblastomas with *Ink4a/Arf* deficiency, overexpressed PDGFR α promotes tumorigenesis through the PI3K/AKT/mTOR-mediated pathway regulated by SHP-2 activity. These findings functionally validate the genomic analysis of glioblastomas and identify SHP-2 as a potential target for treatment of glioblastomas.

Introduction

Human gliomas account for the most common and malignant tumors in the CNS (1). Despite intensive treatments including maximal surgical resection, combined with radiotherapy and concurrent or adjuvant chemotherapies, median survival time of patients with high-grade glioblastoma multiforme (GBM) remains 14–16 months after diagnosis (1, 2). Recently, coordinated genomic analyses of a large cohort of clinical GBM specimens identified frequent co-alterations of genes in 3 core pathways – the P53, retinoblastoma (RB), and receptor tyrosine kinase (RTK) pathways – that are crucial in gliomagenesis (3). Notably, the gene encoding platelet-derived growth factor receptor α (PDGFR α) is amplified in approximately 13% of total GBMs analyzed, and deletion of the tumor suppressor cyclin-dependent kinase inhibitor 2A (*CDKN2A*) locus is frequently found in these PDGFR α -amplified GBMs (3, 4). However, to our knowledge, studies have not been conducted to determine whether these genetic aberrations act in concert to promote gliomagenesis, or to determine the underlying mechanisms of PDGFR α -stimulated tumorigenesis.

The homozygous deletion of the *CDKN2A* locus at chromosome 9p21, which eliminates both *INK4A* and *ARF* genes (encoding P16INK4A and P14ARF, respectively), concomitantly deregulates

lates 2 of the major tumor suppressor pathways, the RB and P53 pathways (5). Mice lacking the *Cdkn2a* locus develop spontaneous tumors and are more prone to carcinogenesis (6). Whereas no spontaneous tumors are found in the brain of *Ink4a/Arf*-deficient mice, astrocytes and neural stem cells (NSCs) from these mice form high-grade gliomas in the brain upon EGFR activation (7). Additionally, K-Ras activation has been shown to cooperate with loss of *Ink4a* and *Arf* in mouse astrocytes (mAst) or neural progenitor cells to generate GBMs (8).

PDGFR α is a RTK that elicits a variety of biological activities such as cell proliferation and migration via stimulation by its ligand dimers, PDGF-AA, -AB, -BB, -CC, and -DD. Activated PDGFR α associates via the autophosphorylated tyrosine (p-Y) residues at its cytoplasmic domain to various downstream SH2 domain-containing signaling molecules, including SRC family kinases (SFKs), phosphotyrosine phosphatase SHP-2, PI3K, and PLC γ (9, 10). Mice deficient in PDGFR α or PDGF-A or engineered to separately express PDGFR α with mutations of the individual p-Y sites show alterations in cellular behavior and embryo development (9, 11). In particular, PI3K has been identified as the major effector of PDGFR α signaling in vitro and in vivo (10, 12, 13). SFKs and PLC γ contribute to some but not all PDGFR α functions (10, 12–14), whereas SHP-2 is not required for cell survival during *Xenopus* embryogenesis (12). However, contributions of each of these signaling modules to glioma formation have not to our knowledge been demonstrated.

Conflict of interest: The authors have declared that no conflict of interest exists.

Citation for this article: *J Clin Invest.* 2011;121(3):905–917. doi:10.1172/JCI43690.



Activation of PDGFR α signaling has been observed in human gliomas. In clinical glioma specimens, PDGFR α and PDGF-A are overexpressed in tumor cells, while PDGFR β is only expressed in endothelial and peri-endothelial compartments (15). PDGF-B, which binds to both PDGFR α and PDGFR β , is an oncoprotein that causes glioma formation in the brain (16). Loss of *Ink4a/Arf* in mAst cells enhances PDGF-B-initiated gliomagenesis and tumor progression in the brain (16). Specific activation of PDGFR α signaling by infusion of PDGF-A proteins, which only bind to PDGFR α in PDGFR α -positive type B NSCs in the subventricular zone (SVZ), leads to glioma-like growth of these cells in adult brain (17). However, how the activation of PDGFR α signaling causes glioma formation and whether co-alteration of tumor suppressor pathways is required in PDGFR α -mediated gliomagenesis have not been directly shown. In this study, we determined the synergistic impact of *PDGFRA* overexpression and *INK4A/ARF* deletion on gliomagenesis in mAst cells and human glioma cells. We then used genetic and pharmacological approaches targeting individual downstream signaling enzymes to examine which specific signaling pathway(s) emanating from PDGFR α are critical in tumorigenesis. Furthermore, we confirmed our observations in clinical glioma specimens that co-overexpress PDGFR α and PDGF-A.

Results

Overexpression of PDGFR α and/or PDGF-A confers tumorigenicity to *Ink4a/Arf*^{-/-} mAst cells. To determine whether activation of PDGFR α signaling in *Ink4a/Arf*^{-/-} mAst cells leads to gliomagenesis in the brain, we overexpressed PDGFR α and/or its ligand PDGF-A chain in a retroviral vector containing an IRES-GFP in *Ink4a/Arf*^{-/-} mAst cells (Figure 1A). The impact of PDGFR α /PDGF-A overexpression on the growth and survival of these cells in vitro was moderate (Supplemental Figure 1, A and B; supplemental material available online with this article; doi:10.1172/JCI43690DS1). However, when these cells were separately transplanted into the flanks of mice, significant s.c. tumor growth was evident in mice that separately received *Ink4a/Arf*^{-/-} mAst cells expressing PDGFR α , PDGF-A, or PDGFR α /PDGF-A (with the receptor and ligand co-expressed in the same cell to create an autocrine signaling), whereas minimal or no tumor formation was seen in control mice that received *Ink4a/Arf*^{-/-} mAst cells expressing GFP (Figure 1B). To determine tumorigenicity of these cells in the brain, various mAst cells were separately implanted into the brain of mice. Notably, mice that received GFP mAst cells did not show active tumor growth up to 42 days after implantation, while PDGFR α , PDGF-A, and PDGFR α /PDGF-A mAst cells started to form tumors in the brain as early as 8–11 days and the majority of tumors reached a volume of approximately 25 to 30 mm³ in 25–35 days. Moreover, mice that separately received PDGFR α - or PDGF-A-expressing mAst cells survived up to 2 months after implantation. On the other hand, all mice that received autocrine PDGFR α /PDGF-A-expressing mAst cells developed large and invasive tumors by 20 days, with an average survival time of 25 days after implantation. As shown in Figure 1, C–L, significantly larger and highly invasive gliomas formed in the brains of mice that received *Ink4a/Arf*^{-/-} mAst cells overexpressing PDGFR α , PDGF-A, or PDGFR α /PDGF-A, whereas only small tumor lesions were found in the brain of mice that received control mAst cells. Significantly, an approximately 10-fold increase in the cell proliferation index was found in gliomas derived from PDGFR α -activated mAst cells compared with control tumors (Fig-

ure 1, M–R), whereas an approximately 10-fold decrease of cell apoptosis was seen in PDGFR α /PDGF-A-expressing tumors (Figure 1S). Of note, in established s.c. or brain tumors, exogenous expression of PDGFR α or PDGF-A was maintained at the end of the experiments (Supplemental Figure 1, C–K, and data not shown). Taken together, these results indicate that expression of PDGFR α and/or PDGF-A confers tumorigenicity to *Ink4a/Arf*-deficient mAst cells in the brain.

To further characterize the tumors derived from *Ink4a/Arf*^{-/-} PDGFR α mAst cells, we examined molecular markers of various cell lineages in the CNS development. As shown in Figure 2, tumors derived from PDGFR α -expressing mAst cells were highly positive for the neural progenitor marker nestin, which was distributed along the processes of individual cells (Figure 2, A and B). As expected, most of tumor cells showed expression of the progenitor/mature astrocyte marker GFAP (Figure 2, C and D) but were negative for the neuronal marker class III β -tubulin (TUJ1) (Figure 2, E and F). Significantly, NG2, an oligodendrocyte progenitor cell marker (18), was expressed in a population of tumor cells that were actively invading the surrounding brain parenchyma, whereas the core of the tumor mass showed relatively low NG2 expression (Figure 2, G and H). In contrast, control tumors derived from *Ink4a/Arf*^{-/-} GFP mAst cells lacked nestin or NG2 expression (data not shown). In addition, tumors derived from *Ink4a/Arf*^{-/-} PDGFR α mAst cells showed negative staining for oligodendrocyte type 2 astrocyte progenitor marker A2B5 and an oligodendrocyte marker CNPase (data not shown). Notably, brain tumors derived from *Ink4a/Arf*^{-/-} PDGFR α /PDGF-A mAst cells (data not shown) displayed similar IHC features to tumors derived from *Ink4a/Arf*^{-/-} PDGFR α mAst cells, suggesting that similar dedifferentiation events also occurred in autocrine PDGFR α /PDGF-A-co-expressing tumors.

Exogenous expression of PDGF-A enhances tumorigenicity of *Ink4a/Arf*-null but not WT *Ink4a/Arf* human glioma cells in the brain. Next, we used 4 human glioma cell lines that were either *INK4A/ARF* null (LN444 and LN443) or WT *INK4A/ARF* (LN-Z308 and LN319; Figure 3A) (19) and exogenously expressed PDGF-A in these glioma cells that express endogenous PDGFR α (Figure 3B). In vitro, expression of PDGF-A in *INK4A/ARF*-null LN444 cells significantly enhanced their capacity of anchorage-independent growth in soft agar, whereas a minimal effect was seen in WT *INK4A/ARF* LN-Z308 and LN319 cells (Figure 3C). When various glioma cells were separately implanted into the brain or the flanks of mice, PDGF-A expression markedly enhanced tumor growth and invasion of *INK4A/ARF*-null LN444 and LN443 glioma cells, while minimal impact of PDGF-A expression on tumorigenesis or invasion was seen in WT *INK4A/ARF* LN-Z308 and LN319 cells in both anatomic sites (Figure 3, D–K, and Supplemental Figure 2). Mice received LN444/PDGF-A cells in the brain had tumor onset as early as 35 days, while mice with LN443/PDGF-A cells developed invasive intracranial tumor 25–30 days after implantation. In contrast, LN444 and LN443 parental cells only formed small tumor lesions in the brain up to 2–3 months following implantation. Mice that received LN444/PDGF-A cells survived for 75–80 days, while most of the mice that received LN443/PDGF-A cells lived up to 3 months after implantation. On the other hand, no significant brain tumor growth was found 2–3 months after implantation in mice that received LN-Z308/PDGF-A or LN319/PDGF-A cells. Thus, consistent with our findings in the *Ink4a/Arf*^{-/-} mAst cells, these data indicate that PDGFR α /PDGF-A signaling enhances in vivo tumor growth and invasion of human glioma cells deficient in *INK4A/ARF*.





Figure 1

PDGFRα and/or PDGF-A overexpression promotes tumorigenesis of *Ink4a/Arf*^{-/-} mAst cells. (A) IB analysis of exogenous expression of PDGFRα and/or PDGF-A in mAst cells. Arrows indicate PDGFRα and PDGF-A proteins run as doublet bands. β-Actin was used as a loading control. (B) Tumor growth of various mAst cells s.c. Data are shown as mean ± SD. (C–L) Representative H&E staining images of various brain sections from 2 independent experiments with at least 3–5 mice per group with similar results. Brains were harvested 25–30 days after transplantation for C–F and 20 days after transplantation for G. Scale bars: 1 mm (C–G); 200 μm (H–L). (M–Q) Ki-67 staining of the corresponding brain sections in (C–G). Yellow arrows indicate tumor mass or invading tumor cells. Red arrows indicate Ki-67-positive cells. Scale bars: 50 μm. (R and S) Quantification of Ki-67 staining (R) and TUNEL staining (S) of various brain sections. Data are presented as percentage to parental controls (mean ± SD). n, the number of mice used for each group. *P < 0.05; **P < 0.01; ***P < 0.001.

p16INK4a but not *p19ARF* attenuates the PDGFRα-promoted tumorigenesis of *Ink4a/Arf*-deficient mAst cells and human glioma cells. To investigate whether re-expression *p16INK4a* or *p19ARF* is able to abrogate the enhanced tumorigenicity in *Ink4a/Arf*^{-/-} mAst cells, we separately expressed these 2 tumor suppressors in mAst cells (Figure 4A). Surprisingly, re-expression of *p16INK4a* but not *p19ARF* alone in *Ink4a/Arf*^{-/-} mAst cells suppressed soft agar growth of

PDGFRα-expressing mAst cells stimulated by PDGF-A (Figure 4B). Consistently, enforced restoration of *p16INK4a* in *Ink4a/Arf*^{-/-} mAst cells significantly inhibited tumorigenesis of PDGFRα/PDGF-A-overexpressing mAst cells in the mouse brain (Figure 4C). To further demonstrate the cooperative effect of PDGFRα activation and loss of *Ink4a* on tumorigenesis, we knocked down endogenous *p16INK4a* using shRNAs targeting *CDKN2A* in

**Figure 2**

Tumors derived from *Ink4a/Arf*^{-/-} PDGFR α mAst cells express markers of neural progenitor cells. Representative images of IHC staining using antibodies against nestin (A and B), GFAP (C and D), β III-tubulin (E and F), and NG2 (G and H). Arrows indicate positive staining of various markers. T, tumor mass; P, normal brain parenchyma. Scale bars: 100 μ m (A, C, E, and G); 50 μ m (B, D, F, and H).

WT *INK4A/ARF* LN319 cells. As shown in Figure 4D, significant inhibition of P16INK4A without affecting P14ARF expression by 2 separate shRNAs in LN319 cells resulted in a marked increase in colony formation in soft agar. Since p16INK4a inhibits CDK4/6, which inactivates RB by phosphorylation, and since p19ARF (or P14ARF) targets MDM2, which suppresses p53 (5), we thus examined responses of modulating CDK4/6 and p53 in these cells. As shown in Figure 4E, inhibition of phosphorylation of RB, the direct downstream target of CDK4/6 by CDK4/6 inhibitor PD0332991 (20), markedly reduced PDGF-A-stimulated cell growth of *Ink4a/Arf*-deficient mAst cells and LN444 cells in soft agar, which suggests that the tumorigenic effect of PDGFR α signaling is dependent on CDK4/6 inactivation of RB proteins (p-RB). Since we did not observe any tumor-suppressing effect of p19ARF on *Ink4a/Arf*^{-/-} PDGFR α -expressing mAst cells (Figure 4B), we asked whether the downstream p53 protein was functional in these cells. When *Ink4a/Arf*^{-/-} and *Ink4a/Arf*^{-/-} p19ARF mAst cells were treated with cisplatin (21), p53 expression was strongly induced in *Ink4a/Arf*^{-/-} PDGFR α /p19ARF-expressing mAst cells but only moderately induced in *Ink4a/Arf*^{-/-} PDGFR α -expressing mAst cells (Supplemental Figure 3A). The mAst cells expressing p19ARF were more sensitive to cisplatin inhibition of cell survival than those without p19ARF expression, indicating that p19ARF-upregulated p53 in *Ink4a/Arf*^{-/-} PDGFR α -expressing mAst cells rendered sensitivity to cisplatin inhibition (Supplemental Figure 3B). Collectively, these data show that p16INK4a but not p19ARF suppresses the tumori-

genesis promoted by PDGFR α /PDGF-A signaling, suggesting a cooperative effect of PDGFR α activation and p16INK4a inhibition during gliomagenesis. Additionally, p19ARF loss might be required for tumor survival in certain circumstances such as in the presence of DNA-damaging agents.

PDGFR α -enhanced tumorigenicity is mediated by SHP-2 and PI3K signaling. Previous studies using genetic and biochemical approaches have defined the roles of signaling molecules in PDGFR α -mediated cellular functions by specific tyrosine-to-phenylalanine (Y-to-F) mutations (Figure 5A) (10, 12, 13). To investigate the impact of these PDGFR α mutations on tumorigenesis, we separately expressed WT PDGFR α or various PDGFR α mutants in *Ink4a/Arf*^{-/-} mAst cells. The PDGFR α mutant R627 (PDGFR α -R627) that harbors a lysine-to-arginine (K-to-R) mutation was included as a “receptor kinase-dead” control. As shown in Figure 5B, stimulation of WT PDGFR α by PDGF-A resulted in autophosphorylation of the receptor and promoted phosphorylation of downstream signaling molecules Erk1/2 and Akt, whereas there was an undetectable receptor tyrosine autophosphorylation in PDGFR α -R627 (Figure 5B). PDGFR α Y-to-F mutations at Y572 and Y574 (PDGFR α -F572/74; for SHP-2 binding), Y1018 (PDGFR α -F1018; for PLC γ binding), and Y988 (PDGFR α -F988) did not result in a significant decrease in p-Akt and p-Erk levels in response to PDGF-A stimulation (Figure 5B). Moreover, the mutation at Y731 and Y742 (PDGFR α -F731/42; for PI3K binding) led to a marked decrease in PDGF-A-stimulated p-Akt and p-Erk1/2 levels (Figure 5B). In agreement with a previous report (22), p-Erk1/2 was markedly reduced in PDGF-A-stimulated mAst cells expressing PDGFR α -F720 (Y-to-F mutation at Y720; for SHP-2 binding), compared with WT PDGFR α mAst cells (Figure 5B). Interestingly, p-Akt levels were also attenuated in PDGFR α -F720 cells.

Next, we determined the impact of these Y-to-F mutations on PDGFR α -promoted cell transformation of *Ink4a/Arf*^{-/-} mAst cells in vitro. As shown in Figure 5C, mAst cells expressing WT PDGFR α and PDGFR α -F572/74, PDGFR α -F988, and PDGFR α -F1018 PDGFR α mutants showed similar capability of anchorage-independent growth in soft agar. In contrast, expression of PDGFR α -R627, PDGFR α -F731/42, or PDGFR α -F720 instead of WT PDGFR α significantly abrogated the capacity of mAst cells to form colonies in soft agar (Figure 5, C and D), indicating that PI3K and SHP-2 signaling were critical for cell transformation in vitro. When various *Ink4a/Arf*^{-/-} mAst cells were separately implanted into the brains of mice, PDGFR α -R627, PDGFR α -F731/42, PDGFR α -F720, and PDGFR α -F988 significantly impaired PDGFR α -promoted tumorigenesis and invasion compared with WT PDGFR α (Figure 6, B, C, E, and F, compared with Figure 6A). However, mAst cells expressing PDGFR α -F572/74 or PDGFR α -F1018 displayed tumorigenicity comparable to WT PDGFR α tumors (Figure 6, D and G, compared to 6A), but showed markedly reduced tumor invasion compared with the tumors derived from mAst cells expressing WT PDGFR α (Figure 6, K and N, compared with 6H). Higher-magnification images of PDGFR α -R627, PDGFR α -F731/42, and PDGFR α -F720 mutants (Figure 6, I, J, and L) further illustrated that these mutations significantly negated the enhanced tumorigenicity conferred by PDGFR α activation, leading to the formation of tumors similar in volume and invasiveness to those seen in GFP-expressing mAst tumors (Figure 1, D and I). Moreover, PDGFR α -F731/42 and PDGFR α -F720 tumors showed a significantly decreased cell proliferation (Figure 6, O and P) and increased apoptosis





Figure 3

PDGF-A overexpression enhances tumorigenesis of *INK4A*/ARF-deficient but not WT *INK4A*/ARF human glioma cells. (A) RT-PCR (top) and IB analyses (bottom) of *INK4A*/ARF-deficient LN444 and LN443 and WT *INK4A*/ARF LN-Z308 and LN319 human glioma cells. (B) IB analysis of PDGF-A overexpression in glioma cells with endogenous PDGFR α expression. PDGFR α was phosphorylated at tyrosine residues in PDGF-A-expressing cells, but not in parental cells that have no detectable PDGF-A. P, parental cells; A, PDGF-A-overexpressing cells. β -Actin was used as a loading control in both A and B. (C) Representative images (left) and quantification (right) of anchorage-independent growth of various glioma cells in soft agar. Scale bars: 1 mm. Data are presented as percentage of the respective parental cells (mean \pm SD). * $P < 0.001$, Student's t test. (D–K) Representative H&E-stained images of various brain sections from 2 independent experiments with 3–5 mice per group with similar results. Brains were harvested 50–55 days (D and H), 75–80 days (E and I), and 45–50 days (F, G, J, and K) after transplantation. Arrows indicate gliomas formed in the brain. Scale bar: 1 mm.

(Figure 6, Q and R), whereas PDGFR α -F572/74, PDGFR α -F988, and PDGFR α -F1018 tumors exhibited moderate or minimal impacts on cell proliferation and survival compared with WT PDGFR α tumors. Conversely, retention of any 1 of the 5

signaling modules (Y731/42, Y572/74, Y720, Y988, and Y1018 PDGFR α mutants) (10) was insufficient to rescue the abolished tumorigenesis in the brain (data not shown). Taken together, these data indicate that ablation of PDGFR α association with

**Figure 4**

Re-expression of p16INK4a but not p19ARF suppresses the tumorigenesis of mAst cells expressing PDGFR α /PDGF-A. (A) IB analyses of re-expression of p16INK4a or p19ARF in PDGFR α - or PDGF-A-overexpressing *Ink4a/Arf*^{-/-} mAst cells. (B) Top: Anchorage-independent growth of p16INK4a- or p19ARF-expressing mAst cells. Scale bars: 1 mm. Bottom: Quantification of soft agar assays. Data are presented as mean \pm SD. * P < 0.05. (C) Representative H&E staining images of various brain sections from 2 independent experiments with 3–5 mice per group with similar results. Brains were harvested 18–22 days after implantation. Mice that received either type of cell displayed similar tumor onset and survival times as mice in Figure 1. Scale bars: 1 mm. (D) Anchorage-independent growth of WT *INK4A/ARF* LN319 cells transfected with shCDKN2A in soft agar. Top: IB analysis. Bottom: Quantification of soft agar assays. Two stable cell clones (#1 and #2) with different efficiencies of P16INK4A knockdown were used. Data are mean \pm SD. (E) Soft agar growth of *Ink4a/Arf*-deficient mAst cells and LN444 cells treated with PD0332991. Top: IB analysis. Bottom: Quantification of soft agar assays. β -Actin was used as a loading control in all IB experiments. Data are presented as mean \pm SD and are representative of 2 independent experiments. * P < 0.05; ** P < 0.01; *** P < 0.001.

SHP-2 or PI3K abrogates tumorigenicity of *Ink4a/Arf*^{-/-} mAst cells expressing PDGFR α , whereas individual association of PI3K, SHP-2, SFK, or PLC γ with the RTK was insufficient to elicit the full spectrum of tumor-promoting effects of PDGFR α .

Pharmacological inhibition of SHP-2 or PI3K abrogates PDGFR α -promoted tumorigenesis of Ink4a/Arf-deficient mAst cells and human glioma cells. We further investigated whether inhibition of SHP-2 and PI3K activities by pharmacological approaches suppresses tumorigenicity of PDGFR α /PDGF-A-expressing *Ink4a/Arf*^{-/-} mAst and LN444 cells. To this end, we exploited several pharmacological inhibitors against PI3K (LY294002), SHP-2 (PHPS-1 and NSC87877) (23), and SFKs (SU6656, and PP2). As shown in Figure 7, LY294002 inhibited PDGF-A-induced phosphorylation of Akt at 10 μ M in mAst cells and 5 μ M in LN444 glioma cells, whereas Erk1/2 phosphorylation was unaffected by LY294002 treatment in both types of cells (Figure 7, A and B). Of note, a modest decrease was observed in p-Erk1/2 levels in mAst cells treated with 20 μ M LY294002 (Fig-

ure 7A). At a concentration of 100 μ M (23), both PHPS-1 and NSC87877 significantly inhibited p-Erk1/2, a direct downstream target of SHP-2, in mAst cells and LN444 cells (Figure 7, A and B). Next, we examined the impact of the pharmacological interventions of these signaling enzymes on in vitro cell transformation and found that 10 μ M LY294002, which had no effect on Erk1/2 activation, suppressed soft agar growth of PDGFR α -expressing mAst cells stimulated by PDGF-A (Figure 7A) and PDGF-A-expressing LN444 cells (Figure 7B). Similarly, 100 μ M of either PHPS-1 or NSC87877 ablated PDGFR α -stimulated, anchorage-independent growth in soft agar but only had a minimal effect on the cell survival and caspase-3 activation of both PDGFR α /PDGF-A-expressing mAst cells and LN444 cells in culture (Supplemental Figure 4, A and B). In contrast, 2 SFK inhibitors, SU6656 and PP2 each at 5- μ M concentration, only showed a modest impact on tumorigenesis of these cells (Figure 7, A and B). Additionally, the MEK inhibitor PD98059 at 10 μ M, which suppressed PDGF-A-induced




Figure 5
Impacts of PDGFR α mutations on downstream signaling of PDGFR α and anchorage-independent growth in soft agar of *Ink4a/Arf*^{-/-} mAst. (A) Schematics of various PDGFR α mutants. (B) IB analyses of PDGF-A-stimulated *Ink4a/Arf*^{-/-} mAst overexpressing individual PDGFR α mutants. Corresponding total proteins or β -actin were used as loading controls. Vertical lines in left panels of p-Tyr, PDGFR α , and β -actin indicate that these images were modified by removal of two lanes of samples between WT and R627. The longer white vertical line indicates that the samples were analyzed in separate SDS-PAGE gels due to the limit of sample loading per gel. (C and D) Soft agar assay. (C) Representative images of soft agar colonies. (D) Quantification of soft agar assays. Data are presented as mean \pm SD and are representative of 2 independent experiments. Scale bars: 1 mm. **P* < 0.01.

p-Erk1/2 in mAst (Supplemental Figure 4C), also inhibited soft agar growth of these cells (Figure 5A), validating Erk1/2 as a mediator of PDGFR α -SHP-2 signaling. Taken together, these data demonstrate that the PDGFR α /PDGF-A-enhanced tumorigenicity of *Ink4a/Arf*-deficient mAst and human glioma cells requires intact SHP-2 and PI3K enzymatic activities.

SHP-2 ablation disrupts PI3K/AKT/mTORC1/S6K activation and attenuates the enhanced tumorigenesis of PDGFR α -expressing mAst. We then examined the impact of SHP-2 inhibitors (PHPS-1 and NSC87877) and PDGFR α -F720 on PI3K signaling. As shown in Figure 5B and Figure 8A, PDGFR α -F720 mutant and SHP-2 inhibitors significantly decreased PDGF-A-stimulated p-Akt in PDGFR α -expressing mAst. A nearly complete knockdown of SHP-2 by siRNAs markedly reduced the stimulated p-Akt levels in these cells, possibly through interruption of the association between PI3K (p85 subunit) and PDGFR α (Figure 8B). However, a previous study showed that PDGFR α -F720 mutation did not result in a decrease in PI3K association with the RTK in mouse fibroblasts (10). Similarly, when SHP-2 was knocked down in NIH3T3 cells, a minimal impact of SHP-2 inhibition on PDGF-A-induced PI3K association and Akt phosphorylation was observed in these fibroblasts (Supplemental Figure 5A), suggesting that the impact of SHP-2 on PI3K association with the RTK and p-Akt in astrocytes and glioma cells was specific. To examine whether the effect of

SHP-2 knockdown on p-Akt signaling was specific to PDGFR α signaling, we knocked down SHP-2 by siRNAs in EGFRvIII-expressing *Ink4a/Arf*^{-/-} mAst (WT PTEN) and LN444 and U87MG (both PTEN mutant) cells. We observed a reduction or a modest impact of p-AKT level in LN444/EGFRvIII and U87MG/EGFRvIII cells, respectively (Supplemental Figure 5B), suggesting that SHP-2 also regulates PI3K/AKT activation in other RTK signaling. However, Akt phosphorylation was absent in *Ink4a/Arf*^{-/-} EGFRvIII mAst, possibly due to the presence of WT Pten in these cells.

We next tested whether signaling downstream to Akt was affected by SHP-2 inhibition. We found that SHP-2 inhibitors (Figure 8A), SHP-2 siRNA knockdown (Figure 8B), or PDGFR α -F731/42 and PDGFR α -F720 mutations (Figure 8C) significantly impaired PDGF-A-stimulated phosphorylation of S6 kinase (p-S6K) downstream to the mammalian target of rapamycin (mTOR) pathway. Thus we further determined the impact of the ablation of SHP-2 and its downstream target the mTOR pathway on cell transformation. Strikingly, stable knockdown of SHP-2 by shRNAs markedly reduced PDGFR α -promoted soft agar growth of mAst (Figure 8D). Additionally, treatment with mTOR complex 1 (mTORC1) inhibitor rapamycin led to a dose-dependent decrease in soft agar colonies formed by PDGF-A-stimulated *Ink4a/Arf*^{-/-} mAst expressing PDGFR α or LN444 cells (Supplemental Figure 6, A and B), compared with their respective DMSO-treated controls.




Figure 6 Impacts of individual mutations of PDGFR α in mAst on brain tumorigenesis. (A–N) Representative H&E staining images of various brain sections from 2 independent experiments with at least 5 mice per group with similar results. Mice that received various types of cells displayed similar tumor onset and survival times as described in Figure 1. Brains were harvested 25–30 days (A), 30–35 days (B, C, D, and G), and 35–40 days (E and F) after transplantation. Scale bars: 1 mm (A–G), 200 μ m (H–N). Arrows indicate tumors or invading cells. (O) Ki-67 staining of brain sections. Scale bars: 50 μ m. Arrows indicate Ki-67–positive cells. (P) Quantification of Ki-67 staining. (Q) TUNEL staining images. Scale bars: 100 μ m. (R) Quantification of TUNEL staining. Data in P and R are presented as mean \pm SD. * P < 0.01; ** P < 0.001.

Similarly, rapamycin was able to effectively suppress the cell transformation capacity conferred by EGFRvIII expression in these cells (Supplemental Figure 6C). Previous studies suggested that the limited efficacy of rapamycin in clinical use was due to the capacity of rapamycin to potentiate PI3K/Akt signaling (24). However, we did not observe an increase of p-Akt level in both *Ink4a/Arf*-deficient mAst and LN444 cells treated with rapamycin up to 72 hours (Supplemental Figure 6D). These results suggest that PDGFR α -promoted tumorigenesis necessitates an intact SHP-2 activity that regulates the PI3K/AKT/mTOR pathway. Since, in clinical glioblastomas, constitutively active PI3K mutations are mostly observed in specimens with no *PDGFRA* aberrations (3, 4), we introduced a constitutively active PI3K p110 subunit (p110 α -CAAX) into *Ink4a/Arf*^{−/−} mAst that either expressed PDGFR α -F720 or an SHP-2 shRNA. In both cell lines, we observed a rescue effect of p110 α -CAAX in soft agar colony formation (Figure 8E), suggesting that PI3K/AKT/mTOR acts downstream of SHP-2 in

PDGFR α -overexpressing gliomas, and that activating PI3K mutations in clinical gliomas might be able to bypass SHP-2 activation to promote tumorigenesis. Taken together, our results suggest that SHP-2 regulates the PI3K/AKT/mTOR signaling emanating from PDGFR α activation.

PDGFR α and PDGF-A are co-expressed, and their expression is linked with the activation of the SHP-2 and the PI3K/AKT/mTOR signaling in clinical glioblastomas. To determine whether PDGF-A and PDGFR α are co-expressed and whether there is a link of overexpression of PDGF-A and PDGFR α and activation of the SHP-2 and the PI3K/AKT/mTOR signaling in clinical glioma specimens, we performed IHC staining on a total of 158 paraffin-embedded primary human glioma specimens using anti-PDGFR α , anti-PDGF-A, anti-p-SHP-2 Tyr542, anti-p-AKT Ser473, and anti-p-S6 Ser235/236 antibodies. PDGFR α proteins were detected at medium to high levels in 17 of 87 GBMs (WHO grade IV tumors) and in 9 of 71 grade II and III tumors. Among these PDGFR α -positive gliomas,




Figure 7

Inhibition of PI3K or SHP-2 activities suppresses PDGFR α -stimulated signaling and cell transformation. (A) Inhibitors of PI3K (LY294002), SHP-2 (PHPS-1 and NSC87877), and MEK (PD98059), but not SFK (SU6656 and PP2), abrogate cell transformation of PDGFR α -overexpressing *Ink4a/Arf*^{-/-} mAst. Top and middle: IB analyses. Bottom: Quantification of soft agar assays. Cells were grown in triplicate in soft agar with or without LY294002 (10 μ M), PHPS-1 (100 μ M), NSC87877 (100 μ M), PP2 (5 μ M), SU6656 (5 μ M), or PD98059 (10 μ M). (B) PI3K and SHP-2 inhibitors suppress PDGF-A-promoted soft agar growth of *INK4A/ARF*-deficient LN444 glioma cells. Top and middle: IB analysis. Bottom: Soft agar assays. Concentrations of the inhibitors and experimental conditions were identical to those in A. Data are presented as mean \pm SD. For IB analyses, corresponding total proteins or β -actin were used as loading controls. All data are representative of 2 independent experiments. * P < 0.001; ** P < 0.0001; *** P < 0.01.

PDGF-A was often co-expressed in the same population of tumor cells (Figure 9, B and C, and Supplemental Table 1). Significantly, phosphorylation of SHP-2 Y542 (p-SHP-2, required for activation of SHP-2) (25), AKT (p-AKT), and ribosomal S6 subunit (p-S6) was also often detected on sister sections of the same tumor in many of the PDGF-A/PDGFR α -positive glioma specimens (Figure 9, B–F, and Supplemental Table 1), suggesting a link between activation of PDGF-A/PDGFR α and stimulation of SHP-2, AKT, and mTOR in clinical glioblastoma specimens. To validate these data, we performed IB analyses on a separate cohort of a total of 20 snap-frozen clinical GBM specimens. As shown in Figure 9G, PDGFR α was expressed at high levels in 7 of 20 GBM samples, 5 of which expressed PDGF-A proteins, suggesting an autocrine PDGFR α signaling in these tumors. In 4 of these 5 PDGF-A/PDGFR α -positive tumors, p-AKT, p-SHP-2, and p-S6 were also detected. Of note, expression of PDGF-A, p-AKT, p-SHP-2, and p-S6 at various levels in other tumor samples likely reflects the impact of heterogeneous gene alterations such as mutations of PTEN, TP53, and overexpression of EGFR/EGFRvIII and c-MET that affect the expression or activation (phosphorylation) of these proteins in clinical glioblastomas. Taken together, these results establish a link of PDGF-A/PDGFR α expression with activation of SHP-2 and PI3K/AKT/mTOR signaling in clinical glioblastoma samples.


Discussion

In this study, we report that PDGFR α and/or PDGF-A overexpression is able to drive gliomagenesis of *Ink4a/Arf*-deficient mAst and human glioma LN444 and LN443 cells. Re-introduction of p16INK4a but not p19ARF into *Ink4a/Arf*-null mAst suppresses PDGFR α -promoted tumor growth. In the absence of PI3K or SHP-2 signaling, PDGFR α fails to enhance tumorigenesis in the brain of mice. Additionally, we establish a link between activation of SHP-2 and the PI3K/AKT/mTOR signaling in PDGFR α -stimulated tumorigenesis in vitro, in mice, and in clinical glioblastoma specimens. Therefore, our data demonstrate that co-alteration of the RTK PDGFR α and tumor suppressor p16INK4a is required for gliomagenesis and that SHP-2 is a critical linker between the PI3K/

AKT/mTOR pathway and PDGFR α in the formation of gliomas.

A unique feature of this study is that specific activation of PDGFR α signaling in vivo by PDGF-A, a ligand that binds to PDGFR α but not PDGFR β (14) as an autocrine loop significantly enhanced the tumorigenesis of *Ink4a/Arf*-deficient mAst and human glioma cells in the brain. Early studies of clinical glioma specimens showed that PDGF-A and PDGFR α are overexpressed in tumor cells, while PDGF-B and PDGFR β are expressed in hyperplastic capillaries, suggesting both autocrine and paracrine loops for PDGF/PDGFR activation in gliomas (15). In neonate and adult mice, expression of PDGF-B induces de novo gliomas from GFAP-positive astrocytes and nestin-expressing glial progenitor cells through activation of PDGFR α and PDGFR β in the brain of both WT and *Ink4a/Arf*-deficient animals (26, 27). Moreover, in WT *Ink4a/Arf* mice, infusion of PDGF-A proteins into the lateral ventricle stimulated tumor-like growth of PDGFR α -positive NSCs in the SVZ in the brain (17). Importantly, data from The Cancer Genome Atlas (TCGA) and other studies revealed that PDGFR α is overexpressed and amplified and often co-expressed with PDGF-A in clinical glioblastoma samples (3, 28, 29). Our results not only functionally validated these studies but also further demonstrated the significance of specific activation of PDGFR α signaling by PDGF-A in cooperation with loss of p16INK4a but not p19ARF in promoting gliomagenesis. We found that when PDGFR α signaling is activated in *Ink4a/Arf*^{-/-} mAst or human glioma cells, mice that received these cells developed significantly larger and highly invasive tumors in the brain. In contrast, no enhancement of tumorigenesis was found in mice that received glioma cells with PDGFR α and an intact *CDKN2A* locus (19). Together, our studies indicate that PDGFR α activation together with *Ink4a/Arf* loss results in enhanced tumor growth of both mAst and human glioma cells in the brain.

Tumor suppressor P16INK4A is frequently mutated in clinical glioblastomas (3, 4, 29). In mice, loss of p16INK4a and p19ARF was shown to be indispensable in facilitating tumorigenesis (6). *Ink4a/Arf*-deficient mice were viable and developed spontaneous



The third important aspect in this study is that we functionally assigned signal modules of PDGFR α in promoting gliomagenesis and tumor invasion in the brain of mice. Previous studies by uncoupling individual signaling pathways from PDGFR α using a series of F-to-Y mutants revealed unequal contributions of each signaling pathway emanating from PDGFR α activation (11, 12, 14). Our in vivo and biochemical data corroborate these reports. When compared with WT PDGFR α overexpression, loss of intrinsic tyrosine kinase activity (R627) of the RTK or binding capacity to PI3K (F731/42) or SHP-2 (F720) abrogated PDGFR α -promoted gliomagenesis, thus signifying the central roles of PI3K and SHP-2 signaling in PDGFR α function. However, while disruption of PDGFR α association with SFKs (F572/74) or PLC γ (F1018) only had a moderate impact on tumor formation in the brain, significant inhibition of tumor cell infiltration in the brain was seen in these tumors, as compared with the WT PDGFR α tumors, thus validating the role of SFKs and PLC γ in mediating PDGFR α -stimulated cell invasion (10). Moreover, a separate set of experiments with individual “add-back panel” mutants in




Figure 9

PDGFR α and PDGF-A are co-expressed in clinical glioma specimens with activated SHP-2 and PI3K/AKT/mTOR pathways. (A–F) Images of IHC staining on sister sections of a representative human J212 GBM specimen, using anti-PDGFR α (B), anti-PDGF-A (C), anti-p-SHP-2 (Y542) (D), anti-p-AKT (E), and anti-p-S6 (F) antibodies. (A) No primary antibody. Arrows indicate positive staining for the indicated proteins. Scale bars: 50 μ m. (G) IB analysis of tissue lysates of a separate cohort of 20 human GBM snap-frozen specimens using indicated antibodies. β -Actin was used as a loading control. Data are representative of 2 independent experiments with similar results.

which 1 of 5 signal modules was individually retained as single or double Y residues (Y731/42, Y572/74, Y720, Y988, and Y1018) could not fully restore PDGFR α -promoted tumorigenesis in the brain (data not shown), consistent with the previous findings using an identical set of PDGFR α add-back panel mutants (10). It is plausible that, similar to this previous in vitro study (10), in which retention of a combination of 2 or 3 add-back signaling modules with comparable levels of total RTK protein in WT PDGFR α -expressing cells was able to restore PDGFR α -mediated biological responses, co-activation of PI3K (Y731/42)- and SHP-2 (Y720)-mediated signaling may be required for the full spectrum of WT PDGFR α -promoted gliomagenesis in the brain.

The last (and what we believe is also the most novel) finding in this study is the emergence of SHP-2 as an essential mediator in PDGFR α -promoted gliomagenesis. SHP-2 (encoded by *PTPN11* gene) is a protein tyrosine phosphatase (PTP) identified as a bona fide proto-oncogene that activates Ras/MAPK signaling through a yet-to-be-defined mechanism (33). Additionally, the role of SHP-2 in mediating PDGFR α signaling has not been clear (9, 11, 14). In human cancers including GBMs, mutations of SHP-2 or its binding partners have been reported, leading to sustained Ras/MAPK signaling (34, 35). Recent genomic analysis of TCGA data has designated *PTPN11* as one of the 6 “linker” genes, which are statistically enriched for connections to various GBM altered genes, thus suggesting a critical role for SHP-2/*PTPN11* in modulation of downstream biological signaling in gliomagenesis (36). Our data not only establish the critical role of SHP-2 in mediating PDGFR α activation for glioma formation but also functionally validate this hypothesis. We showed that inhibition of SHP-2 function by removal of its binding module in PDGFR α (F720 mutant), gene knockdown or pharmacological inhibitors significantly impaired PDGFR α stimulation of tumorigenesis in vivo and in vitro and its downstream signaling effectors, Erk1/2 and PI3K/Akt/mTOR in *Ink4a/Arf*-deficient mAst and glioma cells. Significantly, re-expression of a constitutively active p110, the catalytic subunit of PI3K, rescued the inhibition of SHP-2 in *Ink4a/Arf*-deficient mAst. In EGFRvIII-expressing U87MG glioma cells, SHP-2 regulates activities of ERK/2 and CDC2

that modulate cell cycle progression but has a minimal effect on AKT activation (23). On the contrary, we found that knockdown of SHP-2 not only attenuated PDGFR α stimulation of Erk1/2 activity but also impaired PI3K/Akt/mTOR activity in *Ink4a/Arf*-deficient mAst and human glioma cells. The impact of inhibition of SHP-2 on PI3K/Akt signaling appears to be specific in astrocytic tumors, since we did not observe inhibited PDGF-A stimulation of p-Akt or association of PI3K with PDGFR α in SHP-2-knockdown NIH3T3 fibroblasts. On the other hand, the MEK inhibitor PD98059 also reduced the tumorigenicity of *Ink4a/Arf*-deficient mAst, indicating that SHP-2-mediated PDGFR α signaling requires not only PI3K/Akt but also Erk1/2 activation in gliomagenesis. Additionally, SHP-2 was found to either positively or negatively regulate PI3K/AKT activity (34). However, in our model systems, SHP-2 is required for full activation of PI3K/Akt/mTOR, and inhibition of mTOR by rapamycin markedly suppressed PDGFR α -promoted tumorigenesis. These data are significant since a recent proteomic study revealed that mTOR signaling is predominantly activated in “PDGFRA co-cluster” glioblastomas (37), thus corroborating our observations. However, since phosphorylation of Akt Ser473 occurs both upstream and downstream of mTORC2 signaling, our data do not rule out the role of mTORC2 in PDGFR α -activated signaling. Lastly, our data also demonstrated the importance of activation of PI3K/mTOR signaling by SHP-2 in PDGFR α - or EGFRvIII-promoted tumorigenesis. Taken together, our findings suggest SHP-2 as a critical modulator that regulates PDGFR α -mediated PI3K/AKT/mTOR activities in the development of malignant glioblastomas.

In summary, this study provides molecular insights into the mechanisms by which *PDGFRA* amplification together with loss of *INK4A/ARF* promotes gliomagenesis in the brain. Our data identified SHP-2, as well as PI3K, as a pivotal mediator of PDGFR α signaling in glioma formation. These results have direct clinical relevance, since we not only establish a model system to demonstrate the co-operative role of *PDGFRA* overexpression and *INK4A/ARF* loss in clinical glioblastomas, but also provide functional evidence to validate genomic analyses demonstrating that SHP-2/*PTPN11* is an essential “linker” among glioma altered genes (36). Secondly, activated mTOR



signaling is predominantly found within the subclass of glioblastomas with abnormal PDGFR α signaling (37). Finally, constitutively active PI3K mutations (*PIK3CA* and *PIK3R1*) occur mostly in clinical glioblastomas without *PDGFRA* aberrations (4). Consequently, our results strongly suggest SHP-2/*PTPN11* as a potential target for treatments of glioblastomas with *PDGFRA* overexpression.

Methods

Cell lines and reagents. Primary *Ink4a/Arf*^{-/-} mAst were derived and propagated as previously described (7). Human glioma cell lines LN444, LN443, LN-Z308, and LN319 were obtained from ATCC or from our own collection (38). Unless otherwise mentioned, all glioma cell lines and primary mAst were routinely maintained in 5% CO₂ at 37°C, in DMEM (Invitrogen) containing 10% FBS (Hyclone), 100 U/ml penicillin, and 100 µg/ml streptomycin (Invitrogen).

Histology and IHC. Studies using human tissues were reviewed and approved by the Institutional Review Board involving Human Subjects of the University of Pittsburgh. The specimens were de-identified human tissues, and thus no informed consent was required. A total of 158 paraffin-embedded thin sections of primary human glioma specimens were used, including 87 WHO grade IV GBMs, 34 grade III anaplastic astrocytomas, anaplastic oligodendrogliomas, or anaplastic oligodendroastrocytomas, and 37 grade II oligodendrogliomas and diffuse astrocytomas. Thin sections of human glioma specimens and mouse brains with various tumors were analyzed by IHC using indicated (e.g., Figure 8 legend and Results) antibodies or a TUNEL staining kit as previously described (38, 39). Briefly, the 5-µm human tissue sections were deparaffinized in xylene, followed by rehydration in graded ethanol. After washing with TBS, the antigen was retrieved by boiling the sections in a citrate buffer (pH 6.0) twice for 5 minutes. For mouse brain tissues with various gliomas, 5-µm frozen sections were fixed in pre-chilled acetone at -20°C for 5 minutes, rinsed with PBS, and blocked by AquaBlock (East Coast Biologics Inc.) for 1 hour at room temperature. Afterward, various tissue sections were then incubated with a primary antibody overnight at 4°C and blocked by Peroxidase Blocking Reagent (DAKO) for 10 minutes, followed by incubation with a biotinylated secondary antibody for 30 minutes at room temperature. After washing in PBS, stained tissue sections were visualized by diaminobenzidine chromophore and H₂O₂, followed by hematoxylin counterstaining. Sections were then dehydrated by graded ethanol and mounted with Permount Solution (Fisher).

Soft agar colony formation assay. Colony formation assay in soft agar was performed as previously described (40). Briefly, approximately 5,000 cells

were seeded in a 0.5% Noble Agar top layer with a bottom layer of 0.8% Noble Agar in each of the triplicate wells of a 24-well plate. Growth factor-reduced Matrigel (1 mg/ml) was added into the top layer with or without inhibitors as indicated in figure legends (e.g., Figure 7 legend) and described in Results. PDGF-A-expressing cells in 10% of total cells were included as a source of PDGF-A. DMEM containing 10% FBS was added 3 days after plating and changed every 3 days thereafter. Colonies were scored after 2–3 weeks using Olympus SZX12 stereomicroscope, and data were analyzed using GraphPad Software.

Mouse glioma xenografts, IP, IB, tissue image analyses and quantifications, and siRNA. Experiments of mouse glioma xenografts, IP, IB, tissue imaging and analyses, and siRNA knockdown were performed as previously described (39). For details, see Supplemental Experimental Procedures.

Statistics. One-way ANOVA or an unpaired, 2-tailed Student's *t* test followed by Newman-Keuls post-test was performed using GraphPad Prism software. A *P* value of 0.05 or less was considered statistically significant.

Acknowledgments

We would like to thank Carl-Henrik Heldin at Uppsala University for PDGFR α and PDGF-A cDNAs, Russell O. Pieper at University of California San Francisco for pMXI-*gfp* retroviral vector, Lynda Chin at Dana-Farber Cancer Institute for the cDNA of p16INK4a, Yi Zheng at Cincinnati Children's Hospital Medical Center for cDNA of p19ARF, Tao Cheng at University of Pittsburgh for pMSCV retroviral vectors, and N. Balass for proofreading the manuscript. This work was supported by grants from the NIH (CA102011 and CA130966 to S.-Y. Cheng; EY012509 to A. Kazlauskas; CA87375 to K. Symes), American Cancer Society (RSG CSM-107144), and Pennsylvania Department of Health and Innovative Research Scholar Awards of the Hillman Foundation (to S.-Y. Cheng and B. Hu).

Received for publication May 13, 2010, and accepted in revised form December 22, 2010.

Address correspondence to: Bo Hu, University of Pittsburgh Cancer Institute and Department of Medicine, 5117 Centre Avenue, 2.26, Pittsburgh, Pennsylvania 15213, USA. Phone: 412.623.7791; Fax: 412.623.4840; E-mail: hub@upmc.edu. Or to: Shi-Yuan Cheng, University of Pittsburgh Cancer Institute and Department of Pathology, 5117 Centre Avenue, 2.26f, Pittsburgh, Pennsylvania 15213, USA. Phone: 412.623.3261; Fax: 412.623.4840; E-mail: chengs@upmc.edu.

- Wen PY, Kesari S. Malignant gliomas in adults. *N Engl J Med*. 2008;359(5):492–507.
- Furnari FB, et al. Malignant astrocytic glioma: genetics, biology, and paths to treatment. *Genes Dev*. 2007;21(21):2683–2710.
- TCGA. Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature*. 2008;455(7216):1061–1068.
- Verhaak RG, et al. Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in *PDGFRA*, *IDH1*, *EGFR*, and *NF1*. *Cancer Cell*. 2010;17(1):98–110.
- Sherr CJ. The *INK4a/ARF* network in tumour suppression. *Nat Rev Mol Cell Biol*. 2001;2(10):731–737.
- Serrano M, Lee H, Chin L, Cordon-Cardo C, Beach D, DePinho RA. Role of the *INK4a* locus in tumor suppression and cell mortality. *Cell*. 1996;85(1):27–37.
- Bachoo RM, et al. Epidermal growth factor receptor and *Ink4a/Arf*: convergent mechanisms governing terminal differentiation and transformation along the neural stem cell to astrocyte axis. *Cancer Cell*. 2002;1(3):269–277.
- Uhrbom L, Dai C, Celestino JC, Rosenblum MK, Fuller GN, Holland EC. *Ink4a-Arf* loss cooperates with *KRas* activation in astrocytes and neural progenitors to generate glioblastomas of various morphologies depending on activated Akt. *Cancer Res*. 2002;62(19):5551–5558.
- Heldin CH, Ostman A, Ronnstrand L. Signal transduction via platelet-derived growth factor receptors. *Biochim Biophys Acta*. 1998;1378(1):F79–F113.
- Rosenkranz S, DeMali KA, Gelderloos JA, Bazenet C, Kazlauskas A. Identification of the receptor-associated signaling enzymes that are required for platelet-derived growth factor-AA-dependent chemotaxis and DNA synthesis. *J Biol Chem*. 1999;274(40):28335–28343.
- Betsholtz C. Insight into the physiological functions of PDGF through genetic studies in mice. *Cytokine Growth Factor Rev*. 2004;15(4):215–228.
- Van Stry M, Kazlauskas A, Schreiber SL, Symes K. Distinct effectors of platelet-derived growth factor receptor- α signaling are required for cell survival during embryogenesis. *Proc Natl Acad Sci USA*. 2005;102(23):8233–8238.
- Klinghoffer RA, Hamilton TG, Hoch R, Soriano P. An allelic series at the PDGF α R locus indicates unequal contributions of distinct signaling pathways during development. *Dev Cell*. 2002;2(1):103–113.
- Tallquist M, Kazlauskas A. PDGF signaling in cells and mice. *Cytokine Growth Factor Rev*. 2004;15(4):205–213.
- Hermanson M, et al. Platelet-derived growth factor and its receptors in human glioma tissue: expression of messenger RNA and protein suggests the presence of autocrine and paracrine loops. *Cancer Res*. 1992;52(11):3213–3219.
- Dai C, Celestino JC, Okada Y, Louis DN, Fuller GN, Holland EC. PDGF autocrine stimulation dedifferentiates cultured astrocytes and induces oligodendrogliomas and oligoastrocytomas from neural progenitors and astrocytes in vivo. *Genes Dev*. 2001;15(15):1913–1925.
- Jackson EL, et al. PDGFR α -positive B cells are neural stem cells in the adult SVZ that form glioma-like growths in response to increased PDGF signaling. *Neuron*. 2006;51(2):187–199.
- Stallcup WB, Huang FJ. A role for the NG2 proteoglycan in glioma progression. *Cell Adh Migr*. 2008;2(3):192–201.
- Ishii N, et al. Frequent co-alterations of TP53, p16/CDKN2A, p14ARF, PTEN tumor suppress-



- sor genes in human glioma cell lines. *Brain Pathol.* 1999;9(3):469–479.
20. Wiedemeyer WR, et al. Pattern of retinoblastoma pathway inactivation dictates response to CDK4/6 inhibition in GBM. *Proc Natl Acad Sci U S A.* 2010; 107(25):11501–11506.
21. Fraser M, Bai T, Tsang BK. Akt promotes cisplatin resistance in human ovarian cancer cells through inhibition of p53 phosphorylation and nuclear function. *Int J Cancer.* 2008;122(3):534–546.
22. Wu CJ, O'Rourke DM, Feng GS, Johnson GR, Wang Q, Greene MI. The tyrosine phosphatase SHP-2 is required for mediating phosphatidylinositol 3-kinase/Akt activation by growth factors. *Oncogene.* 2001;20(42):6018–6025.
23. Zhan Y, Counelis GJ, O'Rourke DM. The protein tyrosine phosphatase SHP-2 is required for EGFR-vIII oncogenic transformation in human glioblastoma cells. *Exp Cell Res.* 2009;315(14):2343–2357.
24. Wan X, Harkavy B, Shen N, Grohar P, Helman LJ. Rapamycin induces feedback activation of Akt signaling through an IGF-1R-dependent mechanism. *Oncogene.* 2007;26(13):1932–1940.
25. Lu W, Gong D, Bar-Sagi D, Cole PA. Site-specific incorporation of a phosphotyrosine mimetic reveals a role for tyrosine phosphorylation of SHP-2 in cell signaling. *Mol Cell.* 2001;8(4):759–769.
26. Hu X, Holland EC. Applications of mouse glioma models in preclinical trials. *Mutat Res.* 2005; 576(1–2):54–65.
27. Calzolari F, Malatesta P. Recent insights into PDGF-Induced gliomagenesis. *Brain Pathol.* 2010;20(3):527–538.
28. Martinho O, et al. Expression, mutation and copy number analysis of platelet-derived growth factor receptor A (PDGFRA) and its ligand PDGFA in gliomas. *Br J Cancer.* 2009;101(6):973–982.
29. Parsons DW, et al. An integrated genomic analysis of human glioblastoma multiforme. *Science.* 2008;321(5897):1807–1812.
30. Sharpless NE, Ramsey MR, Balasubramanian P, Castrillon DH, DePinho RA. The differential impact of p16(INK4a) or p19(ARF) deficiency on cell growth and tumorigenesis. *Oncogene.* 2004;23(2):379–385.
31. Tchougounova E, Kastemar M, Brasater D, Holland EC, Westermarck B, Uhrbom L. Loss of Arf causes tumor progression of PDGFB-induced oligodendroglioma. *Oncogene.* 2007;26(43):6289–6296.
32. Mayo LD, Donner DB. The PTEN, Mdm2, p53 tumor suppressor-oncoprotein network. *Trends Biochem Sci.* 2002;27(9):462–467.
33. Matozaki T, Murata Y, Saito Y, Okazawa H, Ohnishi H. Protein tyrosine phosphatase SHP-2: a proto-oncogene product that promotes Ras activation. *Cancer Sci.* 2009;100(10):1786–1793.
34. Chan G, Kalaitzidis D, Neel BG. The tyrosine phosphatase Shp2 (PTPN11) in cancer. *Cancer Metastasis Rev.* 2008;27(2):179–192.
35. Navis AC, van den Eijnden M, Schepens JT, Hoof van Huijsduijn R, Wesseling P, Hendriks WJ. Protein tyrosine phosphatases in glioma biology. *Acta Neuropathol.* 2010;119(2):157–175.
36. Cerami E, Demir E, Schultz N, Taylor BS, Sander C. Automated network analysis identifies core pathways in glioblastoma. *PLoS One.* 2010;5(2):e8918.
37. Brennan C, et al. Glioblastoma subclasses can be defined by activity among signal transduction pathways and associated genomic alterations. *PLoS One.* 2009;4(11):e7752.
38. Jarzynka MJ, et al. ELMO1 and Dock180, a bipartite Rac1 guanine nucleotide exchange factor, promote human glioma cell invasion. *Cancer Res.* 2007;67(15):7203–7211.
39. Yiin JJ, et al. Slit2 inhibits glioma cell invasion in the brain by suppression of Cdc42 activity. *Neuro Oncol.* 2009;11(6):779–789.
40. Yoshida S, Shimizu E, Ogura T, Takada M, Sone S. Stimulatory effect of reconstituted basement membrane components (matrigel) on the colony formation of a panel of human lung cancer cell lines in soft agar. *J Cancer Res Clin Oncol.* 1997;123(6):301–309.