

**STUDIES IN DIABETIC ACIDOSIS AND COMA, WITH  
PARTICULAR EMPHASIS ON THE RETENTION OF  
ADMINISTERED POTASSIUM**

T. S. Danowski, ... , J. M. Quashnock, L. Greenman

*J Clin Invest.* 1949;28(1):1-9. <https://doi.org/10.1172/JCI102037>.

Research Article

**Find the latest version:**

<https://jci.me/102037/pdf>



# STUDIES IN DIABETIC ACIDOSIS AND COMA, WITH PARTICULAR EMPHASIS ON THE RETENTION OF ADMINISTERED POTASSIUM

By T. S. DANOWSKI, J. H. PETERS, J. C. RATHBUN,  
J. M. QUASHNOCK, AND L. GREENMAN

*(From the Departments of Research Medicine and of Physiological Chemistry, the Children's and Presbyterian Hospitals, and the Renziehausen Foundation, University of Pittsburgh, School of Medicine, Pittsburgh)*

(Received for publication July 22, 1948)

Abnormally low concentrations of potassium, at times associated with muscular paralysis, have been observed in the serum of diabetic patients while under treatment for acidosis or coma (1-4). Available data indicate that deficits of cell potassium are also present in at least some of these subjects (5). In an attempt to characterize and quantitate the magnitude of these deficits, exchanges of water, electrolytes, carbohydrate, and protein have been investigated in diabetic acidosis and coma prior to, during, and following the administration of potassium salts.

## EXPERIMENTAL PROCEDURE AND METHODS

Eight cases in all have been studied. Of these, two were adults and the remaining six juvenile diabetics 16 years of age or younger. On admission all had Kussmaul breathing, a marked reduction of the serum bicarbonate content, hyperglycemia, glycosuria, and ketonuria.

Studies during treatment and recovery were divided into three periods. The first or pre-KCl period extended from admission to the point where the patients had improved sufficiently to be maintained on an oral intake. During this time insulin, 0.9 per cent saline, and, once the blood sugar began to drop, glucose solutions were administered in amounts summarized in Table I. Under this treatment ketosis diminished, overbreathing ceased, hyperglycemia decreased, and mental clarity returned. This interval lasted 12 to 25 hours in the individual subjects. During the second or KCl period, 22 to 37 hours in length, the patients received insulin, 10.0 to 30.0 grams of KCl *per os* or intravenously, whole milk containing added carbohydrate, and water as desired. The third or post-KCl period began after the final dose of KCl and lasted up to 34 hours. During this time the patients received insulin as needed, together with measured amounts of water and milk.

In all subjects concentrations of the whole blood non-protein nitrogen, blood sugar (6-8), and the levels of serum chloride, bicarbonate, sodium, potassium and water were measured at the beginning and end of each period (6, 9-12). Average values for the electrolyte, carbo-

hydrate, and protein content of fresh milk have been used in calculating the intake (13, 14). Urine excreted during each period was analyzed for nitrogen, sodium, potassium, chloride, and glucose (6, 10, 11, 15, 16). Body weight was determined when possible at the start and end of each period.

## METHOD OF CALCULATION

Alterations in extracellular fluid volume were calculated from changes in the chloride space, based on the external balance of this anion and corrected for changes in the serum concentrations of chloride as described by Elkinton and co-workers, and by Darrow (17-19). In view of the dehydration known to develop in the course of diabetic acidosis and coma, the initial extracellular volume was assumed for purposes of this calculation to be 15 per cent of the body weight rather than the usual value of 18.7 to 23 per cent found in non-dehydrated human subjects (20). In two instances a pre-treatment weight could not be obtained. A reasonable assumption was therefore made as to its magnitude, based upon subsequent weights and the balance of water. The volume of the chloride space and the concentrations of sodium and potassium in extracellular water, corrected for the Donnan effect, were used in the calculation of increments and decrements of these ions in extracellular fluid. The retention of sodium and of potassium by cells was taken to be the difference between the intake of these cations and the amounts retained in extracellular fluid plus those excreted in urine. In the case of potassium the final values recorded include corrections for the balance of cellular nitrogen, using the K:N ratio of 2.38 m.eq.:1.0 gm. N. In determining the balance of cellular nitrogen, changes in the concentration of the nonprotein nitrogen have been taken into account, assuming for this purpose that the nonprotein nitrogen is distributed uniformly through all of the body water, and that 0.65 of the body weight is water. Similar treatment of the sodium data does not, of course, significantly affect the balances of this electrolyte because of the low Na:N ratio in cells. The data on cell base in Table III represent, therefore, balances in excess of the sodium and potassium which move with nitrogen in the anabolism and catabolism of protein.

TABLE I  
Intake and output during therapy of diabetic acidosis and coma

Pt.	Period	Intake							Urine output					
		Insulin	H <sub>2</sub> O	Cl	Na	K	N	CHO	Vol.	Cl	Na	K	N	Glucose
		<i>units</i>	<i>ml.</i>	<i>m. eq.</i>	<i>m. eq.</i>	<i>m. eq.</i>	<i>gms.</i>	<i>gms.</i>	<i>ml.</i>	<i>m. eq.</i>	<i>m. eq.</i>	<i>m. eq.</i>	<i>gms.</i>	
T.W.	I	650	13,335	1,723	1,723	49	0	300	1,675	178	92	37	7.6	11
	II	135	3,333	629	212	490	12.1	270	1,180	192	85	39	7.3	13
	III	75	2,150	71	57	85	11.8	268	2,300	442	262	90	10.5	27
M.S.	I	170	6,255	726	726	8	0.6	132	1,663	79	38	55	10.9	24
	II	5	3,075	182	38	192	8.0	74	2,190	279	133	83	10.4	0
	III	20	1,025	34	29	42	6.8	43	970	121	92	20	5.8	0
J.K.	I	240	5,240	769	769	0	0	132	3,490	257	265	74	8.6	155
	II	120	5,275	488	69	505	14.2	343	4,050	523	187	208	11.1	65
	III	15	800	20	16	24	3.3	98	315	33	10	16	1.2	14
W.S.	I	154	4,000	615	615	0	0	75	1,100	165	116	53	6.1	26
	II	0	4,350	303	289	322	12.0	155	4,200	718	421	183	4.8	59
	III	58	1,550	51	41	61	8.5	74	2,136	33	207	60	8.4	22
L.D.	I	280	6,570	775	775	0	0	143	4,330	566	473	65	7.1	67
	II	20	5,510	457	170	380	15.9	138	2,800	306	106	36	6.1	5
	III	70	4,525	131	106	158	21.9	190	4,510	409	232	51	11.7	2
R.J.	I	500	6,750	998	1,025	3	8.4	115	1,775	139	106	28	10.0	39
	II	105	4,800	288	16	292	3.8	29	818	133	78	30	9.8	1
	III	50	2,500	16	13	20	2.7	24	1,790	249	146	45	12.7	45
D.C.(a)	I	245	6,876	1,020	1,020	0	0	127	3,630	355	275	105	14.1	70
	II	105	2,550	475	58	490	12.0	105	6,580	990	584	251	21.7	167
	III	57	3,200	105	85	126	17.5	152	2,000	168	16	88	16.6	83
D.C.(b)	I	230	6,925	973	973	0	0	275	4,250	73	164	124	5.4	129
	II	90	6,135	495	90	518	18.5	404	6,710	846	475	233	15.7	268
	III	70	3,175	75	61	91	12.6	185	1,770	72	24	60	7.9	60

## RESULTS

### A. Alteration in serum water and in extracellular volume

On admission a variable degree of plasma dehydration, as measured by the water content, was present (Table II). The values ranged from 897 to 924 gms. of water per liter of serum in contrast to levels of 922 to 938 gms. per liter ordinarily found in healthy adults (21). There is no means of deciding how much of the deficit in any particular patient should be assigned to losses of body fluid alone, and how much is to be considered secondary to a disappearance of circulating plasma proteins of the type observed during depletion of body water and solutes (22, 23). At the end of the first period of treatment all serum water values were again in the normal range, or actually above it. No consistent pattern is apparent in the fluctuations in serum water during the second and third treatment periods.

The expansion in the chloride space uniformly

observed in each patient at the end of the first treatment period suggests that considerable deficits of extracellular water as well had been incurred prior to admission (Table III). These increments, as calculated from the chloride space, ranged from +1.0 to +9.5 liters. Since subsequent changes in Periods II and III of each study in no instance entirely cancelled this initial expansion it is reasonable to interpret this persistent positive balance of extracellular water as indirect evidence suggestive of extracellular dehydration on admission. On the other hand, it is likely that in at least some of the subjects the extracellular fluid volume had been overexpanded, even though edema did not develop. This appeared to be true in some of the infants with diarrhea treated by Darrow (19).

### B. Serum electrolyte concentrations prior to and during therapy

1. *Sodium and chloride:* Despite these deficits of body water it is to be noted that the concen-

trations of sodium and chloride, the chief electrolytes of extracellular fluid and serum, were not increased above the usual range of isotonicity (Table II, at zero point of Period I). As a matter of fact, the concentrations of sodium were abnormally *low* in five patients. In all of these and in one other the concentrations of chloride in serum, taking 97 m.eq. per liter as the lower limit of normal, were also decreased (24). In view of the dehydration described above, it is immediately apparent that these patients had lost, in keeping with the findings in other series, considerable amounts of the chief extracellular electrolytes, con-

comitant with, and in most instances, in excess of, the losses of body water (25, 26). Had water alone been lost, the levels of sodium and chloride would have been increased above normal.

During the first period of treatment in which 0.9 per cent saline solution was given, the concentrations of sodium and of chloride rose in all patients (Table II, end of Period I). As a matter of fact in all but one instance, patient J. K., definite hyponatremia and hyperchloremia appeared. In subject R. J., for example, who started with normal concentrations, the sodium in serum increased to 161.6 m.eq. per liter and the chloride

TABLE II  
*Body weight and analyses of blood and serum during treatment of diabetic acidosis and coma*

Patient (age—sex)	Period	Time from start	Body* wtg.	Blood*		Serum*				
				NPN	Sugar	HCO <sub>3</sub>	Cl	Na	K	H <sub>2</sub> O
		hours	kgm.	mgm. %	mgm. %	m. eq./liter	m. eq./liter	m. eq./liter	m. eq./liter	grams/liter
T.W. (26F)	I pre-KCl	0		—†	572	5.5	101.9	142.3	4.0	—†
	II KCl p.o.	0-25		40	112	11.8	127.8	157.6	2.8	951
	III post-KCl	25-48		37	—		124.9	153.1	5.6	948
		48-70	60.8	34	—	16.4	110.2	144.1	5.4	944
M.S. (15F)	I pre-KCl	0		65	667	9.3	77.6	118.6	3.4	897
	II KCl p.o.	0-17		36	104	17.4	107.7	144.6	3.9	943
	III post-KCl	17-39		26	139	24.0	109.8	143.2	5.4	943
		39-61	41.5	26	259	23.4	102.8	144.8	5.6	937
J.K. (18F)	I pre-KCl	0	50.0	32	600	8.1	94.1	135.5	5.4	916
	II KCl p.o.	0-20	53.9	21	145	19.0	97.8	140.5	5.0	934
	III post-KCl	20-67	53.4	27	235	23.4	98.8	144.4	5.4	930
		67-72	53.4	33	302	22.7	98.1	137.8	6.0	932
W.S. (14M)	I pre-KCl	0	36.2	39	470	6.7	96.3	129.9	5.5	912
	II KCl p.o.	0-12		21	70	14.7	114.9	149.6	4.1	943
	III post-KCl	12-44		27	362	22.4	98.9	144.8	5.5	933
		44-68	38.1	30	181	27.2	94.4	151.2	4.4	930
L.D. (15M)	I pre-KCl	0	38.7	33	1189	4.1	96.5	124.2	4.0	924
	II KCl p.o.	0-23	38.3	24	51	11.2	109.3	143.9	1.8	943
	III post-KCl	23-41	40.9	28	328	15.7	101.8	137.3	3.6	948
		41-65	38.4	35	352	24.0	98.0	142.5	3.0	945
R.J. (14M)	I pre-KCl	0	34.9	103	1395	10.4	101.9	142.1	3.7	912
	II KCl p.o.	0-19		79	230	20.6	137.3	161.6	2.0	933
	III post-KCl	19-44		49§	118	21.3	123.5	143.9	4.1	949
		44-68	47.2	41§	288	25.4	108.9	133.5	3.8	945
D.C.(a) (16F)	I pre-KCl	0	52.7	65	582	4.8	89.9	122.7	6.5	899
	II KCl p.o.	0-23	53.4	28	171	15.1	111.5	155.9	3.0	941
	III post-KCl	23-59	52.8	32	140	26.9	95.5	133.2	4.3	926
		59-83	54.0	37	167	29.2	98.1	139.1	4.1	925
D.C.(b) (16F)	I pre-KCl	0	50.0	67	786	6.8	91.4	125.3	6.7	907
	II KCl i.v.	0-19	53.9	28	163	15.4	117.1	145.6	3.3	936
	III post-KCl	19-51	50.9	29	149	26.1	92.1	140.8	5.7	949
		51-84	50.9	34	55	28.7	94.1	141.0	4.7	927

\* Values as recorded refer to body weight, blood, or serum observed on admission (0 hour), and at the end of each individual period.

† Assumed to be 60 for purposes of calculating balances of cell nitrogen.

‡ Not obtained. Average value of the seven other cases used in calculating concentrations in extracellular water, 910.

§ NPN drawn before end of period; lower value therefore assumed in calculations.

to 137.3, levels about 20 and 35 m.eq., respectively, above the usual range in healthy adults. There was no clinical evidence that this hypertonicity was in any way deleterious. The findings do suggest, however, that in most instances a less concentrated saline solution could have been used, or that the patient should have received larger quantities of glucose solution. During the KCl period, and subsequently, the serum chloride tended to return to, or even below, physiological concentrations. The serum sodium levels during Periods II and III in general showed a similar trend, but in no instance, in contrast to the chloride ion, did hyponatremia appear.

2. *Bicarbonate*: The markedly lowered concentrations of serum bicarbonate present on admission rose toward or to normal during therapy in

all patients. The development of hyperchloremia, which has already been noted an almost invariable occurrence, did not prevent these increases in bicarbonate. However, it is not possible to answer unequivocally from these data the question as to whether or not hyperchloremia repressed the rise of serum bicarbonate. It is clear, for example, that the increase during the first period in serum bicarbonate level of 10.9 m.eq. per liter in 20 hours in patient J. K. who did not develop hyperchloremia is quite comparable to the 10.2 m.eq. per liter increase in 19 hours in the patient with the most pronounced hyperchloremia (R. J.). On the other hand, the possibility that hyperchloremia prolonged the acidosis in patient T. W. can by no means be excluded. At 70 hours the bicarbonate concentration was still only

TABLE III  
*Balances of electrolytes, nitrogen, and carbohydrate during treatment of diabetic acidosis and coma\**

Pt.	Period	External balance†					Extracellular balance			Intracellular balance‡		
		Cl	Na	K	N	CHO	Fluid	Na	K	N	Na	K
		m. eq.	m. eq.	m. eq.	gms.	gms.	liters	m. eq.	m. eq.	gms.	m. eq.	m. eq.
T.W.	I	+1533	+1616	+ 11	- 7.6	+289	+9.5	+1561	+15	- 0.5	+ 56	- 3
	II	+ 434	+ 124	+450	+ 4.7	+257	+3.5	+ 463	+69	+ 5.8	-339	+367
	III	- 375	- 209	- 5	+ 1.2	+240	-0.3	- 228	- 6	+ 2.3	+ 19	- 5
M.S.	I	+ 638	+ 674	- 48	-10.4	+108	+3.7	+ 672	+17	- 2.6	+ 2	- 59
	II	- 102	- 101	+108	- 2.5	+ 74	-1.0	+ 165	+10	+ 0.2	- 64	+ 98
	III	- 90	- 68	+ 22	+ 1.0	+ 43	-0.2	- 6	+ 1	+ 1.0	- 62	+ 19
J.K.	I	+ 503	+ 491	- 75	- 8.6	- 16	+4.4	- 650	+19	- 5.0	+159	- 82
	II	- 38	- 122	+297	+ 3.1	+278	-0.5	- 20	+ 2	+ 1.2	-102	+292
	III	- 16	+ 2	+ 8	+ 2.1	+ 84	0	+ 86	+ 8	+ 0.2	- 88	0
W.S.	I	+ 443	+ 489	- 53	- 6.2	+ 49	+2.7	- 494	+ 3	- 1.9	+ 5	- 52
	II	- 418	- 136	+139	+ 7.2	+ 96	-2.5	+ 398	- 3	+ 5.9	-262	+128
	III	+ 15	- 170	+ 1	+ 0.1	+ 52	+0.4	- 100	- 4	- 0.6	+270	+ 7
L.D.	I	+ 195	+ 284	- 67	- 7.1	+ 76	+1.0	+ 249	-12	- 4.8	+ 35	- 44
	II	+ 148	+ 61	+344	+ 9.8	+133	+1.9	+ 205	+19	+ 8.8	+144	+304
	III	- 280	- 130	+107	+10.2	+188	-2.3	- 274	-12	+ 8.4	+144	+ 99
R.J.	I	+ 841	+ 896	- 25	- 1.7	+ 76	+4.2	+ 772	- 1	+ 3.7	+124	- 34
	II	+ 146	- 74	+262	- 6.1	+ 28	+2.3	- 138	+28	+ 2.6	-212	+228
	III	- 236	- 136	- 25	-10.0	- 21	-0.4	- 174	+ 4	- 8.6	+ 38	- 9
D.C.(a)	I	+ 655	+ 730	-106	-14.2	+ 57	+4.1	- 865	-17	- 1.5	+135	- 86
	II	- 517	- 529	+239	- 9.7	- 62	-3.0	+ 657	+ 3	-11.0	-128	+262
	III	- 65	+ 65	+ 38	+ 1.0	+ 69	-0.8	+ 62	- 5	- 0.8	-127	+ 45
D.C.(b)	I	+ 892	+ 798	-125	- 5.4	+146	+5.3	- 909	- 9	+ 7.3	-111	-116
	II	- 355	- 392	+285	+ 2.7	+135	+0.2	- 57	+31	+ 2.4	-335	+248
	III	+ 1	+ 33	+ 31	+ 4.7	+125	-0.6	+ 33	-14	+ 3.1	+ 66	+ 38

\* Balance data are expressed per individual period rather than cumulatively.

† Corrected for the electrolytes and nitrogen withdrawn in serum samples. Stools, when passed, were formed and not analyzed. Small amounts of vomitus in T.W. and 240 cc. in L.D. in the pre-KCl periods were not analyzed.

‡ The K balances represent changes in excess of the transfers associated with the anabolism and catabolism of proteins. Similar treatment of the Na data does not significantly affect the results. Changes in the NPN have been taken into account in calculating the balance of cell nitrogen (17-19).

16.4 m.eq. per liter, even though hyperglycemia and acetonuria had disappeared.

3. *Potassium*: Abnormally high levels of potassium (6.5 and 6.7 m.eq. per liter) were noted on admission twice in the same patient, D. C. (a) and D. C. (b) in Table II. In the other six cases the admission values ranged from 3.4 to 5.5. During the initial period of therapy in which no potassium was given a dramatic fall of serum potassium to abnormally low levels was observed in patients T. W., L. D., R. J., D. C. (a) and D. C. (b). Though the intermediate points are not recorded the values given are actually the lowest concentrations observed in each patient before the start of KCl *per os* or parenterally. Only one of the subjects, R. J., developed obvious muscular paralysis, with inability to move his extremities. As a group, however, the patients were still too ill at the time of most marked hypopotassemia to permit clear differentiation between generalized asthenia and partial paralysis.

The administration of KCl without exception increased the concentration of serum potassium. The highest recorded level, 5.7 m.eq. per liter, at the end of KCl therapy was observed in the one patient, D. C. (b), who received the salt intravenously. However, a comparable or even greater increment from 2.8 to 5.6 m.eq. per liter occurred in patient T. W. who took the KCl *per os*. It is to be noted that the muscular paralysis of subject R. J. cleared promptly following the oral administration of the potassium salt. None of the subjects in this series developed gastrointestinal distress or diarrhea, even though the KCl was administered in capsules or intravenously as rapidly as 1 gm. per hour.

### C. Balances of carbohydrate, nitrogen, and electrolytes

Results of calculations based on the data in Tables I and II have been recorded in Table III. During Period I the external balances indicated that, with but one exception, the patients had utilized or stored about one-half and in one instance as much as 96 per cent of the administered carbohydrate. Since the concentration of blood sugar fell in each patient during this interval, it is obvious that the balance was even more positive. Similarly, in the subsequent two periods of

observation the balances of carbohydrate remained predominately positive (Table III).

Without exception during Period I all patients excreted nitrogen in urine in excess of the nitrogen intake (Table III). A considerable portion of this negative balance, however, is assignable to a decrease in the elevated nonprotein nitrogen usually present on admission. Even when this factor has been included in the calculation of the nitrogen balance, six of the eight patients were losing cell nitrogen in Period I (Table III). During the KCl and post-KCl periods, when the patients were taking milk, the trend was reversed and the balances of cell nitrogen, with but two significant exceptions, became positive.

In so far as the balances of electrolytes are concerned during the pre-KCl period, all patients retained large amounts of chloride and sodium. While KCl was being administered, and in the subsequent interval the external balances of sodium and of chloride were more often negative than positive (Table III). However, not one of the negative balances was of sufficient magnitude to cancel entirely the earlier retention of sodium or chloride. The largest positive balances of sodium and of chloride, more than 1500 m.eq. of each ion for the three periods, were recorded in T.W., even though this patient was admitted with normal concentrations of sodium and chloride. This finding is to be contrasted with the relatively small positive balances of these ions in patient L. D. despite initial hypochloremia and hyponatremia. It should be emphasized, however, that irrespective of the initial concentrations of sodium and chloride in the serum all patients retained these two electrolytes in greater or lesser proportion. Again, as in the interpretation of the positive balances of extracellular water, this can be taken as evidence of replacement of deficits, but it is possible that some of it represents over-treatment. The retained sodium was disparately distributed between the extracellular and cellular fluid. Most of it, and frequently all, remained in the extracellular phase (Table III).

In contrast to the positive balances of sodium, all but one patient lost considerable quantities of body potassium during the first period of treatment. This one patient, T. W., had received, however, a moderate amount of potassium during this interval. During the second period of ther-

apy in which KCl was ingested or injected the patients retained, as recorded in Table III under "External Balance," 43 to 92 per cent of the administered potassium. The largest positive external balance was 450 m.eq. in the first case recorded, an average of 7.4 m.eq. per kgm. of body weight. An even greater retention in proportion to body weight, 9.0 m.eq. per kgm., was observed in patient L. D. It is to be noted that comparable degrees of potassium retention were observed in patient D. C. during two separate admissions in coma irrespective of the route of administration. In the first instance, D. C. (a), the salt was administered orally, and in the second, D. C. (b), intravenously. Furthermore, the per kilogram retention of potassium given parenterally in this subject was of the same order of magnitude as that observed on the average in the six other patients treated *per os*. As might be predicted, the major portion of the retained potassium entered the cellular phase (Table III). The cell balances of potassium during the second period, corrected for the intracellular balances of nitrogen, ranged from +98 to +406 m.eq. In some of the experiments sodium left the cells as potassium entered, but this was by no means an invariable finding. The failure to observe reciprocal exchanges of these two ions more often may have been related in part to the lag in the extrusion of sodium reported by Conway and Hingerty following replacement of experimentally induced cell potassium deficits (27).

Even with these large positive balances of potassium it is not certain that the deficits had been completely corrected in all patients. As a matter of fact, in the third period, during which the intake was limited to milk and water, only one of the eight patients developed negative balances of cell potassium in excess of nitrogen. The others were still retaining a part or all of the potassium available to them in the milk ration.

#### DISCUSSION

The electrolyte, nitrogen, and carbohydrate studies in these patients again emphasize that considerable amounts of body constituents are lost during diabetic coma. The losses of body water, of extracellular electrolytes, of cell nitrogen, and of carbohydrate have been recognized and defined (25, 26). The striking retention of potas-

sium administered to this series of patients points to the existence of concomitant deficits of this cation in the body cells. Such an interpretation is supported by the experiments of Tarail and Elkinton which showed that amounts of potassium as great as those employed in some of our studies were not retained in the body when given by mouth to healthy adults (28). Presumably such subjects do not have potassium deficits and hence the administered potassium is promptly excreted. Diabetic patients in acidosis or coma are, however, depleted of potassium. Several mechanisms appear to be involved in the development of this negative balance. It is obvious, for example, that patients in diabetic acidosis or coma are in negative nitrogen balance. This breakdown of tissue releases potassium bound to protein of cells (17). Similarly, potassium laid down with glycogen in the liver (29) is released from the cells during deglycogenation. These two processes, however, can account for only a small portion of the potassium deficit. Most of the deficit must represent, therefore, a decrease in the cell potassium present in excess of the nitrogen there. Several factors probably contribute to this loss of cell base. First, it is known that extracellular dehydration, *per se*, is associated with movements of potassium and of water out of the cells (17). Secondly, it has been shown that interruption of carbohydrate metabolism in the blood cell *in vitro*, either by refrigeration, exhaustion of glucose stores, or by addition of an inhibitor such as fluoride, is associated with a pouring out of cell potassium (30). Finally, the evidence to date suggests that as long as urine is being elaborated it contains potassium. This appears to be true even though deficits of cell potassium, and even extracellular hypopotassemia, are present (21, 28). Hence, with the diuresis characteristic of almost all patients in coma considerable amounts of potassium are lost. At present it is not clear to which of these mechanisms the major portion of the potassium deficit is to be ascribed.

The hypopotassemia of extracellular fluid which develops during treatment on a potassium-free regimen noted by other workers and by us stems from a number of sources. It is obvious from our data that it results in part from an expansion of body water with the administration of fluids,

and in part from a continued loss of potassium in urine. The negative balances of cell potassium recorded during the first period of therapy in seven of the eight patients exclude the possibility that reentry of this ion into cells played any significant role in the decrease in serum potassium. This does not mean, of course, that no potassium was deposited in the liver with reglycogenation, nor that none entered other cells. It merely indicates that these must have been subsidiary processes. Furthermore, the negative balances of potassium cannot be explained away as manifestations of transfers of chloride into cells. Such a movement of the reference ion could not cancel the negative cell balances, since even large alterations of the chloride space could account for only a few milliequivalents of potassium.

During the administration of potassium salts the major portion of the cation entered the cells. Only a small amount of it was laid down with nitrogen. There was no significant urinary loss of previously retained potassium in the post-KCl periods. As a matter of fact, in five of the experiments more potassium entered the cells. On the whole a smaller proportion of the administered potassium was retained during Period III suggesting that in the majority of the patients cell concentrations of this ion were completely or almost completely restored. It should be emphasized of course that the possibility that the potassium administered orally was incompletely absorbed cannot be excluded. Naturally this would alter the magnitude of the exchanges. However, the findings in control studies (28), and the absence of frequent or diarrheal stools in these patients, suggest that absorption was essentially complete. This is further supported by the finding that quite comparable data were obtained when KCl was given intravenously.

It is to be noted that positive cell balances of both potassium and sodium were recorded in patients who were admitted with essentially normal serum levels of these ions. These instances should again serve to emphasize the fact that the presence of normal *concentrations* of any particular electrolyte cannot be interpreted as evidence that the *total amount* is intact (31). This is of obvious importance in planning replacement therapy.

The physiological significance of these extensive potassium deficits can be as yet partially and

only indirectly evaluated. It is known, of course, that muscular paralysis and abnormally low levels of *extracellular* potassium frequently coexist (1, 2, 32-34). This is not, however, an invariable association, since paralysis can occur with normal serum concentrations of potassium (35) and our own data indicate that pronounced hypopotassemia can occur without evident paralysis.

At present only analogies can be drawn as to possible deleterious effects of these deficits of *intracellular* potassium. It has been shown, for example, that the depletion of cellular potassium produced by over-dosage with desoxycorticosterone acetate or by restriction of potassium intake is associated with degenerative changes in the myocardium (36, 37). In this respect it is of interest that in our series patient T. W., 26 years of age and previously well, had on admission a grossly abnormal electrocardiogram, characterized by markedly inverted T waves. These changes compatible with myocardial damage were still present one month later, and cleared only after an additional month of convalescence. It may have been more than chance, therefore, that this occurred in the patient who apparently had developed the greatest deficits prior to therapy.

No adequate statistics are yet available as to whether replacement of potassium deficits may mitigate the considerable mortality still associated with diabetic acidosis and coma (26, 38). The favorable results of Govan and Darrow in decreasing the fatality rate of infantile diarrhea by means of parenteral and oral potassium therapy suggest that a similar beneficial response may be expected in diabetic coma (39). In testing this form of therapy, however, unremitting care must be taken to avoid deaths from potassium poisoning (40, 41). At present it would seem reasonable, therefore, to limit the administration of potassium salts to the oral route, or to give only dilute solutions intravenously. An adequate urine volume and constant electrocardiographic surveillance are essential.

#### SUMMARY AND CONCLUSIONS

Patients recovering from diabetic acidosis or coma retained considerable amounts of potassium administered as KCl, in addition to water, sodium, chloride, carbohydrate, and nitrogen. Comparable positive balances of this cation were re-



corded in one subject during two separate admissions, in one of which KCl was given by mouth and in the other intravenously. In every study potassium entered cells far in excess of amounts which could be ascribed to changes in cell protein. The possible physiological significance of these findings has been discussed.

#### ACKNOWLEDGMENT

The authors wish to acknowledge their indebtedness to Miss J. Kruman, R.N. and Miss D. Hoehl, R.N. and their assistants, to Miss M. Freedman and her staff of technicians, and to the internes and residents of the two hospitals. Without their help this study would not have been possible.

#### BIBLIOGRAPHY

- Holler, J. W., Potassium deficiency occurring during the treatment of diabetic acidosis. *J. A. M. A.*, 1946, 131, 1186.
- Nicholson, W. M., and Branning, W. S., Potassium deficiency in diabetic acidosis. *J. A. M. A.*, 1947, 134, 1292.
- Martin, H. E., and Wertman, M., Serum potassium, magnesium, and calcium levels in diabetic acidosis. *J. Clin. Invest.*, 1947, 26, 217.
- Guest, G. M., and Rapoport, S., Electrolytes of blood plasma and cells in diabetic acidosis and during recovery. *Proc. Am. Diabet. A.*, 1947, 7, 97.
- Butler, A. M., Talbot, N. B., Burnett, C. H., Stanbury, J. B., and MacLachlan, E. A., Metabolic studies in diabetic coma. *Tr. A. Am. Physicians*, 1947, 60, 102.
- Peters, J. P., and Van Slyke, D. D., *Quantitative Clinical Chemistry*. Vol. II. Methods. Williams and Wilkins, Baltimore, 1932.
- Somogyi, M., A method for the preparation of blood filtrates for the determination of sugar. *J. Biol. Chem.*, 1930, 86, 655.
- Benedict, S. R., The estimation of sugar in blood and normal urine. *J. Biol. Chem.*, 1926, 68, 759.
- Van Slyke, D. D., and Neill, J. M., The determination of gases in blood and other solutions by vacuum extraction and manometric measurement. *I. J. Biol. Chem.*, 1924, 61, 523.
- Hald, P. M., The flame photometer for the measurement of sodium and potassium in biological materials. *J. Biol. Chem.*, 1947, 167, 499.
- Quashnock, J. M., Gas pressure regulation in flame photometry. *Federation Proc.*, 1948, 7, 179.
- Eisenman, A. J., Mackenzie, L. B., and Peters, J. P., Protein and water of serum and cells of human blood, with a note on the measurement of red blood cell volume. *J. Biol. Chem.*, 1936, 116, 33.
- Sodium and Potassium Analyses of Foods and Waters. Mead Johnson and Company, Evansville, Indiana, 1947.
- Shohl, A. T., *Mineral Metabolism*. Reinhold Publishing Corp., New York, 1939.
- Jeffery, W. H., Note on the Volhard-Harvey method for the estimation of chlorides in urine. *J. Lab. & Clin. Med.*, 1927, 13, 687.
- Sumner, J. B., A more specific reagent for the determination of sugar in urine. *J. Biol. Chem.*, 1925, 65, 393.
- Elkinton, J. R., and Winkler, A. W., Transfers of intracellular potassium in experimental dehydration. *J. Clin. Invest.*, 1944, 23, 93.
- Elkinton, J. R., Winkler, A. W., and Danowski, T. S., Transfers of cell sodium and potassium in experimental and clinical conditions. *J. Clin. Invest.*, 1948, 27, 74.
- Darrow, D. C., The retention of electrolyte during recovery from severe dehydration due to diarrhea. *J. Pediat.*, 1946, 28, 515.
- Elkinton, J. R., The volume of distribution of mannitol as a measure of the volume of extracellular fluid, with a study of the mannitol method. *J. Clin. Invest.*, 1947, 26, 1088.
- Danowski, T. S., Unpublished data.
- Elkinton, J. R., Danowski, T. S., and Winkler, A. W., Hemodynamic changes in salt depletion and in dehydration. *J. Clin. Invest.*, 1946, 25, 120.
- Danowski, T. S., Winkler, A. W., and Elkinton, J. R., Biochemical and hemodynamic changes following the subcutaneous injection of glucose solution. *J. Clin. Invest.*, 1947, 26, 887.
- Hald, P. M., Heinsen, A. J., and Peters, J. P., The estimation of serum sodium from bicarbonate plus chloride. *J. Clin. Invest.*, 1947, 26, 983.
- Peters, J. P., Kydd, D. M., Eisenman, A. J., and Hald, P. M., The nature of diabetic acidosis. *J. Clin. Invest.*, 1933, 12, 377.
- Danowski, T. S., Winkler, A. W., and Peters, J. P., Salt depletion, peripheral vascular collapse, and the treatment of diabetic acidosis. *Yale J. Biol. Med.*, 1946, 18, 405.
- Conway, E. J., and Hingerty, D., Relations between potassium and sodium levels in mammalian muscle and blood plasma. *Biochem. J.*, 1948, 42, 372.
- Tarail, R., and Elkinton, J. R., Potassium deficiency and the role of the kidney in its production. *J. Clin. Invest.*, 1948, 27, 557; 1949, 28, 99.
- Fenn, W. O., The deposition of potassium and phosphate with glycogen in rat livers. *J. Biol. Chem.*, 1939, 128, 297.
- Danowski, T. S., The transfer of potassium across the human blood cell membrane. *J. Biol. Chem.*, 1941, 139, 693.
- Elkinton, J. R., Winkler, A. W., and Danowski, T. S., The importance of volume and of tonicity in salt depletion shock. *J. Clin. Invest.*, 1947, 27, 1002.
- Danowski, T. S., Elkinton, J. R., Burrows, B. A., and Winkler, A. W., Exchanges of sodium and potassium in familial periodic paralysis. *J. Clin. Invest.*, 1948, 27, 65.

33. Ferrebee, J. W., Parker, D., Carnes, W. H., Gerity, M. K., Atchley, D. W., and Loeb, R. F., Certain effects of desoxycorticosterone; the development of "diabetes insipidus" and the replacement of muscle potassium by sodium in normal dogs. *Am. J. Physiol.*, 1941, 135, 230.
34. Brown, M. R., Currens, J. H., and Marchand, J. F., Muscular paralysis and electrocardiographic abnormalities resulting from potassium loss in chronic nephritis. *J. A. M. A.*, 1944, 124, 545.
35. Talbott, J. H., Periodic paralysis; a clinical syndrome. *Medicine*, 1941, 20, 85.
36. Darrow, D. C., and Miller, H. C., The production of cardiac lesions by repeated injections of desoxycorticosterone acetate. *J. Clin. Invest.*, 1942, 21, 601.
37. Miller, H. C., and Darrow, D. C., Relation of muscle electrolyte to alterations in serum potassium and to the toxic effects of injected potassium chloride. *Am. J. Physiol.*, 1940, 130, 747.
38. Joslin, E. P., Root, H. F., White, P., Marble, A., and Bailey, C. C., *Treatment of Diabetes Mellitus*. Lea & Febiger, Philadelphia. 8th edition, 1946.
39. Govan, C. D., Jr., and Darrow, D. C., The use of potassium chloride in the treatment of dehydration of diarrhea in infants. *J. Pediat.*, 1946, 28, 541.
40. Winkler, A. W., Hoff, H. E., and Smith, P. K., Electrocardiographic changes and concentration of potassium in serum following intravenous injection of potassium chloride. *Am. J. Physiol.*, 1938, 124, 478.
41. Govan, C. D., Jr., and Weiseth, W. M., Potassium intoxication. Report of an infant surviving a serum potassium level of 12.27 millimoles per liter. *J. Pediat.*, 1946, 28, 550.