THE URINARY EXCRETION OF SODIUM AND POTASSIUM DURING THE PYROGENIC REACTION IN MAN^{1,2}

By WILLOUGHBY LATHEM 8

(From the Veterans Administration Hospital, West Haven, Connecticut, and The Department of Internal Medicine, Yale University School of Medicine, New Haven, Conn.)

(Submitted for publication December 7, 1955; accepted May 10, 1956)

Among the mechanisms regulating the renal tubular transport of electrolytes the renal circulation plays an important but as yet imperfectly un-An interrelation between renal derstood role. hemodynamics and sodium reabsorption has been demonstrated (2) in a variety of experimental circumstances associated with renal vasoconstriction (infusions of adrenal medullary hormones [3], neosynephrine [4], and other drugs [5]; during orthostasis [4], hemorrhage [6]). Whether such an interrelation exists during renal hyperemia is, however, not known. The present study was undertaken in order to examine this problem further. For this purpose a study was made of urinary sodium and potassium excretion during the pyrogenic reaction, a physiologic alteration accompanied by both renal vasoconstriction and vasodilatation (7).

METHOD

Thirty-two studies were made in 27 convalescent adult male subjects. All subjects were free from cardiovascular or renal disease, none was recovering from a febrile disorder, and all were maintained on a normal hospital diet containing approximately 10 grams of sodium chloride a day. All studies were made in fasting, recumbent subjects during the morning hours, from approximately 8 A.M. until noon, or later as noted.

The effects of the pyrogenic reaction were examined in 16 subjects. In eleven of these studies the clearances of inulin, PAH, sodium and potassium were determined before and following administration of typhoid-paratyphoid vaccine. All 11 subjects were given 500 ml. of water by mouth one-half to one hour prior to the test and received an infusion of 5 per cent dextrose in water intravenously at a rate of 5 ml. per minute throughout the

¹ A preliminary report of this study has appeared in abstract form (1).

² Supported by grants from The American Heart Association and the Greenwich Health Association and the National Heart Institute, National Institutes of Health, Public Health Service.

⁸ Present address: Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania. test. Urine was collected every 10 to 30 minutes over a period of approximately 3 hours. Typhoid vaccine (75,-000,000 organisms) was administered intravenously at the conclusion of the first 10 to 20-minute collection period. Blood was collected without stasis at approximately half-hour intervals throughout the test. In order to prevent a rise in body temperature 8 of these subjects were given Aminopyrine on the day prior to the test in a dosage of 0.6 gram every four hours. Three subjects (J. P., E. V. and G. R.) did not receive this medication.

In five subjects the effects of the pyrogenic reaction on sodium and potassium excretion were observed over a five-hour period and measurements of renal hemodynamics were not made. These subjects ingested 500 ml. of water one-half to one hour prior to the test and approximately 100 to 200 ml. every hour thereafter in order to maintain an adequate urine flow. No intravenous fluids were given. Urine was collected every hour by spontaneous voiding which usually required that the subject stand briefly by the side of the bed. Typhoidparatyphoid vaccine (75,000,000 organisms) was administered intravenously at the end of the first hourly collection period. All five subjects were premedicated with Aminopyrine.

Control observations of urinary sodium and potassium excretion were made in 16 studies under conditions similar to those employed in the experimental (pyrogen) studies except that typhoid vaccine was not administered. None of these subjects received Aminopyrine.

In all studies the body temperature was measured orally and the cardiac rate and arterial blood pressure (sphygmomanometer) were determined at frequent intervals. Sodium and potassium content of urine and plasma were determined by an internal standard flame photometer. Inulin was determined by the method of Harrison (8) and PAH by the method of Smith, Finkelstein, Aliminosa, Crawford, and Graber (9).

RESULTS

Body temperature and subjective response

In eleven of thirteen subjects premedicated with Aminopyrine the body temperature remained normal following the injection of typhoid vaccine (Tables I and II). These subjects remained asymptomatic throughout the test. In two (J. McN. and R. B.) premedicated and in three (J. P., E. V. and G. R.) untreated subjects a chill occurred approximately one to two hours after the administration of vaccine and the oral temperature increased to approximately 100° F. (Table I). These subjects complained of headache, backache and malaise at the time of the chill.

The arterial blood pressure increased 5-10/5-10 mm. Hg in most subjects, along with an acceleration of cardiac rate (10 to 20 beats per minute), during the initial phase of the pyrogenic reaction (one to two hours) and remained elevated throughout the study (Table I).

Renal hemodynamics

The renal circulatory adjustments accompanying the pyrogenic reaction were similar to those reported in previous studies (10, 11). Renal plasma flow (PAH clearance) increased (on the average, 40 per cent) one and one-half to two hours following administration of typhoid vaccine in all eleven subjects in whom renal hemodynamics were studied (Table I). Glomerular filtration rate (inulin clearance) did not change from resting values during hyperemia (except in A. R. in whom there was a slight decrement and in E. V. in whom

TABLE I

Renal function and urinary electrolyte excretion in eleven normal subjects during the pyrogenic reaction *

Sub	ject	Elapsed	T	B.P.	Plasma Na	Plasma K	Cin	Cpah	FF	v	UnaV	U⊧V
Age	S.A.	time min.	Temp. F .	mm. Hg	mEq./L.	mEq./L.	ml./min.	ml./min.	%	ml./min.	μEq./min.	μEq./min
W. G.	. W.	46	974	104/62	137.8	4.42	115	545	21	4.3	63.4	67.0
45	1.83	79					105	468	22	1.5	59.2	56.8
		104					108	507	21	2.0	22.4	39.3
		127			138.1	3.83	110	572	19	3.5	50.4	31.9
		147					111	769	14	2.1	80.2	34.5
		170	978	106/50			108	813	13	3.2	54.8	51.1
I. D)a.	44	97 2	110/80	136.0	4.12	150	735	20	7.1	142.8	101.0
32	2.16	75		•			134	664	20	4.2	114.2	80.0
		102	974	112/90	136.4	4.20	134	599	22	0.6	71.6	57.8
		127		134/90			154	752	20	0.6	108.2	42.4
		151		,	134.5	3.65	161	1,042	15	2.5	191.4	45.4
		169					162	1,173	14	5.8	158.5	68.8
		188	976	146/90	136.8	3.60	149	1,052	14	8.9	134.8	87.0
W.R	. W.	48	97²	110/72	137.4	4.30	132	723	18	11.2	96.7	96.9
33	1.91	82	970	110/80			105	540	19	3.8	77.7	62.0
		<u>9</u> 9		,			119	776	15	0.8	57.2	61.4
		114					111	1,059	11	0.9	100.6	95.2
		131			139.6	4.20	134	1,256	11	1.4	81.2	56.9
		143					135	1,225	11	1.2	102.3	58.0
		158	986	120/80	136.5	4.05	140	1,232	11	1.5	113.9	61.7
S . 1	P.	54	97º	90/58	135.5	4.06	161	752	21	0.8	29.0	99.5
24	1.95	86		90/58			153	668	23	2.1	15.0	89.0
		109		,	137.2	3.80	170	777	22	2.0	12.1	77.6
		154	97²				168	903	18	6.8	65.0	60.1
		168					161	1.041	15	9.7	137.5	43.8
		181	97²	120/70	136.5	3.63	168	1,206	14	9.8	161.5	46.9
Т. M	cN.	72	98º	102/50	133.4	4.25	108	756	14	0.6	11.1	73.0
J. M 37	1.64	102	100 ^s	106/58	130.0	3.92	159	1,101	14	0.7	22.2	101.5
-		123		,			116	1,080	15	0.9	10.2	70.9
		138		108/58	131.2	3.70	110	1,036	11	1.3	6.2	70.9
		152		,			108	1,055	10	1.8	6.4	79.9
		168					96	968	10	2.3	5.5	76.4
		183	1004	120/60	132.0	3.45	103	1,016	10	3.1	5.5	82.1

* In each subject the initial values for renal hemodynamics and electrolyte excretion are the averages of three periods. All other values represent single determinations. The elapsed times are in reference to the beginning of the first collection period. Typhoid vaccine was administered intravenously at the end of the first collection period (10 to 20 minutes after the beginning of the test). All subjects except J. P., E. V., and G. R. were premedicated with Amino-pyrine. Abbreviations are as follows: S.A. = body surface area (M³), Temp. = body temperature (oral), B.P. = arterial blood pressure, C_{in} = inulin clearance—glomerular filtration rate (ml./min.), C_{pah} = p-aminohippurate clearance—renal plasma flow (ml./min.), FF = filtration fraction— C_{in}/C_{pah} (%), V = urine flow (ml./min.), $U_{na}V$ = urinary sodium output (μ Eq./min.), U_kV = urinary potassium output (μ Eq./min.).

					T.	ABLE I-C	continued					
Sub Age	S.A.	Elapsed time <i>min</i> .	Temp. °F.	B.P. mm. Hg	Plasma N. mEq./L.	Plasma K mEq./L.	Cin ml./min.	Cpah ml./min.	FF %	V ml./min.	UnsV µEq./min.	UkV µEq./min.
G.	Р.	56	976	118/74	132.5	4.45	143	629	23	6.1	26.0	62.1
27	2.01	84		132/88			133	696	19	4.8	11.9	49.7
		116	984		135.7	3.90	132	788	17	8.2	27.9	44.4
		138					126	856	15	10.2	40.0	39.6
		152					135	816	17	9.6	35.6	44.5
		168	986	130/50	135.9	3.95	125	734	17	7.8	24.5	49.6
R.	в.	45	974	128/78	135.5	3.85	161	797	20	10.9	45.3	52.1
26	1.90	71		-			133	640	21	4.7	44.9	28.0
		92	99 2	140/82			171	822	21	0.6	45.6	8.6
		118			135.3	3.68	199	998	20	0.9	79.4	11.5
		141					180	1,061	17	4.0	70.3	55.2
		159					178	1,294	14	2.7	57.0	62.0
		179	998	140/72	136.4	3.65	158	1,279	12	10.6	45.3	72.5
A.	R.	26	98º	130/90	137.1	3.98	112	549	20	10.8	118.8	75.4
30	1.82	43		•			84	434	20	8.3	95.0	69.1
		73					97	475	19	4.2	63.8	50.8
		95			138.6	3.78	98	513	19	2.2	30.5	39.0
		117					101	525	19	5.4	55.6	27.3
		143	980	140/90			96	708	14	7.4	106.5	24.1
		164					96	817	12	6.9	158.0	31.7
		183	98º	140/90	139.2	3.70	90	740	12	2.5	181.0	46.5
Ŀ	P.	35	974	108/68	136.4	3.97	151	813	19	6.0	113.8	44.5
33	2.24	59		•			141	819	17	9.7	143.5	46.6
		83					129	700	18	3.4	140.0	36.6
		113	98²	118/74	137.8	3.80	140	741	19	1.2	141.6	32.8
		144					145	1,004	14	2.8	325.0	34.5
		162					131	1,146	11	3.5	384.0	48.6
		183	99²	120/80	137.4	3.70	115	1,122	10	2.3	208.0	70.8
E.		36	98º	118/72	137.2	4.21	137	640	21	7.5	220.2	109.1
31	1.87	66		-			122	623	20	9.2	271.2	91.6
		98					128	608	19	3.5	289.0	78.4
		123	998	120/60	139.0	4.10	119	618	19	1.4	220.0	69.6
		140		-			119	656	18	1.4	204.6	63.0
		157					140	812	17	2.2	287.5	65.8
		172	1004		138.3	3.85	149	909	15	2.3	317.0	70.8

TITT Continued

it increased). In nine subjects the hyperemic phase was preceded by a transient reduction in renal plasma flow which occurred one to two hours following injection of the vaccine (average change, 13 per cent). In seven of these subjects a transient decrement (11 to 21 per cent; average change, 12 per cent) in glomerular filtration rate also occurred at this time; in the remaining two (W. G. W. and S. P.) the filtration rate did not change. The FF tended to remain the same or to increase slightly at this time. In two subjects (J. McN. and G. P.) neither glomerular filtration nor renal plasma flow changed during the latent period prior to hyperemia.

978

994

100²

124/74

120/64

130/70

141

159 175

136.6

136.3

136.5

3.98

3.90

3.90

G. R. 2.01

Urine flow

118

62

100

117

156

112

118

Urine flow decreased one to two hours following administration of the vaccine in eleven of sixteen subjects (V, Tables I and II). In the remainder (S. P., J. McN., G. P., W. L. and F. W.) urine volume was sustained or increased at this time. The reduction in urine flow was always accompanied by a fall in glomerular filtration rate in those studies in which renal hemodynamics were measured (Table I). The decrement in urine flow exceeded the change in filtration in all. In three studies (S. P., J. McN., and G. P.) glomerular filtration rate did not change prior to hyperemia and

18

10

17

22

15

13

12

12.1

7.6 1.2

1.1

1.2

0.7

1.1

215.2

217.2

104.1

116.0

144.3

41.6

71.0

645

648

595

534

840

967

1.018

141.8

133 5

107.0

80.6

89.9

85.8

113.0

urine volume increased or remained constant. During the later or hyperemic phase of the pyrogenic reaction (2 to 5 hours) no systematic change in urine volume was observed, although flow returned towards resting levels if previously depressed. At this time glomerular filtration returned to initial levels. No difference was noted between afebrile and febrile subjects during either the vasoconstrictive or hyperemic phases.

Sodium excretion

The changes in urinary sodium output were variable. During the initial phase of the pyrogenic reaction (one to two hours following administration of the vaccine) sodium excretion diminished transiently (32 to 93 per cent) in 9 of 16 subjects ($U_{na}V$, Tables I and II). In other studies sodium

output increased (four) or remained unchanged (three) at this time. These alterations were not significant when compared to control subjects and could not be correlated with changes in renal hemodynamics. Although a reduction in glomerular filtration rate accompanied the decrement in sodium output in four of seven studies (J. Da., W. R. W., A. R., and G. R.), sodium excretion increased in two subjects (E. V. and J. P.) and remained unchanged in one (R. B.) despite a fall in glomerular filtration rate. Moreover. in W. G. W., G. P. and S. P. sodium output fell 72, 54, and 48 per cent, respectively, without a significant change in glomerular filtration. There was no correlation between the changes in sodium excretion and renal plasma flow.

During the second to fifth hour following ad-

TABLE II	
rinary electrolyte excretion in five normal subjects during the pyrogenic reaction *	*

Subject	Elapsed time	Temp.	Plasma Na	Plasma K	, v	UnaV	UkV
Age S.A.	min.	• F.	mEq./L.	mEq./L.	ml./min.	µEq./min.	µEq./min
W. L.	56	976	137.8	4.50	0.7	33.0	27.0
18 1.92	118				0.7	40.6	40.8
•••	181				0.6	37.3	45.9
	237	98			1.0	81.7	73.7
	299				2.3	85.6	143.9
	361				1.5	91.2	110.9
	417	98*	137.6	4.50	1.5	94.6	93.7
	(0)	071	100 5	3.95	0.7	28.1	59.1
F. W.	60	978	129.5	3.95	1.1	66.7	80.8
32 1.83	124				2.0	40.1	52.0
	183				2.0	50.2	76.6
	241				3.2	50.2 62.8	112.4
	316				2.0		88.3
	371				1.3	91.1	
	423	98º	139.5	3.87	3.6	56.9	65.8
н. н.	65	974	137.6	4.15	3.2	143.6	107.6
46 1.86	121				4.8	212.0	132.1
40 1.00	182				2.0	97.4	78.6
	262				4.1	236.5	65.0
	303				4.6	153.6	131.4
	366	98²	131.8	3.75	5.1	107.3	249.0
D. B.	66	974			8.8	68. 4	35.9
	145	91-			7.1	55.4	52.0
20 2.14	143				0.3	4.8	10.5
	252				5.5 12.7	32.1	79.9
	306				12.7	48.0	137.5
	372	98º			7.2	54.4	100.8
		974			5.5	101.4	221.5
L. G.	68	97			5.5 7.1	77.4	176.7
25 2.20	139				3.0	31.2	48.5
	198 252				10.9	85.5	84.2
	252				9.4	108.7	73.1
	309 369	98²			8.5	49.4	117.9

* Each value represents a single determination. Typhoid vaccine was administered at the end of the first period. All subjects were premedicated with Aminopyrine. Abbreviations are as in Table I.

	Plas	sma	Urine			
Hours	Na mEq./L.	$\frac{K}{mEq./L.}$	UnaV µEq./min.	U⊧V µEq./min.		
		Control subjects				
0–1	138.0 ± 1.0	4.29 ± 0.13	103.3 ± 18.8	72.7 ± 8.1		
2–3	139.5 ± 0.7	4.17 ± 0.14	153.0 ± 21.4	77.0 ± 6.0		
	E	xperimental (pyrogen) su	ıbjects			
0-1	136.0 ± 0.7	4.12 ± 0.05	91.6 ± 16.4	85.9 ± 11.8		
2–3	136.7 ± 0.6	3.76 ± 0.05	119.0 ± 24.4	64.1 ± 3.7		

TABLE III Mean plasma and urinary electrolyte values in sixteen control subjects and in sixteen subjects before and during the pyrogenic reaction *

* Each value is the mean and standard error of the mean for all subjects in the control and experimental groups, respectively. The time in hours is in reference to the beginning of the first collection period. The experimental subjects receiving typhoid vaccine are those listed in Tables I and II. The mean values for this group during the first hour represent resting values prior to an effect of the pyrogenic reaction. The 2- to 3-hour values are those obtained at the time of intrarenal hyperemia. Abbreviations are as shown in Table I.

ministration of the vaccine, at the time of renal hyperemia, sodium output did not change consistently. The mean excretion at this time exceeded resting levels (by more than 10 per cent) in 10 of 16 subjects and was less than or equal to these values in the remainder. The mean change for the entire group of 16 subjects during the third hour ⁴ was not significant when compared to resting values (t = 1.55; < .50 > .10) or when compared to the mean spontaneous changes observed over a similar time interval in control subjects (t = 0.88; p < .50 > .10) (Table III). No difference was noted between febrile and afebrile subjects. The sodium content of plasma did not change significantly (t = 1.64; p < .50 > .10) (Table III).

Potassium excretion decreased (20 to 46 per cent) one to two hours following administration of typhoid vaccine in nine of eleven subjects in whom renal hemodynamics were measured (U_kV , Table I). Potassium output increased in J. McN. and remained unchanged in S. P. at this time. Although the reduction in potassium excretion was usually accompanied by intrarenal vasoconstriction, this change occurred independently of alterations in glomerular filtration rate and in one instance (G. P.) occurred in the absence of a change in renal plasma flow. During hyperemia potassium output remained essentially constant except in J. P., in whom the excretion of this ion increased and in S. P. and G. R., in whom it diminished. A similar response was observed in five subjects in whom renal hemodynamics were not studied (Table II). Potassium output fell in three (H. H., D. B., and L. G.) of these subjects, remained unchanged in one (F. W.) and increased in one (W. L.) during the first two to three hours following administration of the vaccine. During the fourth and fifth hours the urinary excretion of this ion increased above resting levels in four of five studies. No difference was observed between febrile and afebrile subjects in either group of subjects.

The mean reduction in potassium output from resting levels in all 16 subjects during the first one to two hours of the pyrogenic reaction was of borderline significance (t = 2.70; p < .02 > .01). This change was not significant when compared to the control group (t = 1.53; p < .50 > .10). During hyperemia the mean potassium output did not differ significantly from resting levels (t = 2.01; p < .10 > .05) (Table III), nor was the mean difference significant when compared to spontaneous changes observed in control subjects (t = 1.53; p < .50 > .10) (Table III).

The concentration of potassium in plasma decreased during the pyrogenic reaction in 8 of 14 subjects (plasma electrolytes were not measured in two). In the remainder no change was noted. The mean reduction for all 14 subjects, $0.35 \,\mu\text{Eq}$. per L. (range, 0 to 0.80 μEq . per L.), was significant when compared with resting levels (t = 7.0; p < .01) (Table III), but not when compared

⁴ This time was selected in order to analyze statistically the changes during hyperemia in all 16 subjects. The subjects listed in Table I were not studied beyond the third hour.

with the mean spontaneous change in control subjects studied during a similar interval of time (t = 1.97; p < .10 > .05) (Table III).

DISCUSSION

In the present study no definite interrelation was demonstrated between urinary electrolyte excretion and the renal circulatory adjustments attending the pyrogenic reaction. During the initial vasoconstrictive phase of this reaction, when renal plasma flow and glomerular filtration rate were transiently reduced, sodium and potassium output were not altered systematically, nor was there a significant change in the excretion of these ions during the later and more sustained phase of intrarenal hyperemia.

Previous evidence of a relationship between renal hemodynamics and urinary sodium excretion has been obtained in a variety of experimental and clinical conditions which have in common intrarenal vasoconstriction (2, 4, 5). The mechanism by which sodium excretion is reduced under these conditions is not clear but changes in glomerular filtration rate (2, 7) and perhaps in the intrarenal distribution of blood (6) may play a role. Whether changes in filtration are necessary has not been established. It has been suggested that electrolyte reabsorption may be influenced by the rate of blood flow through the kidney (4, 5). Although the present study does not support this view, since sodium and potassium excretion remained unchanged during both renal vasoconstriction and hyperemia, further studies are needed in order to evaluate this problem in view of the many physiologic and metabolic alterations elicited during the pyrogenic reaction (12) which might influence electrolyte metabolism and reabsorption independently of renal hemodynamics. The results of this study cannot be interpreted with certainty in terms of the relationship between the renal circulation and tubular function. Whether an interrelation between electrolyte excretion and hyperemia exists under circumstances other than the pyrogenic reaction must be determined. During renal vasodilatation as elicited by intravenous l-hydrazinophthalazine, urinary sodium excretion is unaltered (13).

Although alterations in renal hemodynamics did not affect electrolyte excretion in the present study, the reduction in glomerular filtration rate during the vasoconstrictive phase of the pyrogenic reaction appeared to initiate a fall in urine volume, as has been demonstrated previously in man (7), and the urine became more concentrated with respect to sodium and potassium ions. That factors other than a change in filtration contributed to the reduced urine output is suggested by the fact that, once initiated, the antidiuresis tended to persist despite a return of glomerular filtration rate to control levels. These observations differ from those reported recently by Brandt, Ruskin, Zumoff, Castleman, and Zuckerman (14) who describe an increase in the free water clearance of dogs following administration of a bacterial pyrogen derived from pseudomonas.

SUMMARY

A study was made of the effect of the pyrogenic reaction as elicited by triple typhoid vaccine on the urinary excretion of sodium and potassium in 16 normal human subjects. No systematic change in the excretion of these ions was observed in febrile or afebrile subjects during either the early, transient period of renal vasoconstriction or the late, sustained period of renal hyperemia. A reduction in urine volume occurred during the vasoconstrictive phase in association with a decrement in glomerular filtration rate and the urine became more concentrated with respect to sodium and potassium ions.

ACKNOWLEDGMENT

The technical assistance of Miss Marion Brazel and Mrs. Frances Kennedy is gratefully acknowledged.

REFERENCES

- Lathem, W., Urinary excretion of sodium and potassium during the pyrogenic reaction. J. Clin. Invest., 1954, 33, 951.
- Selkurt, E. E., Sodium excretion by the mammalian kidney. Physiol. Rev., 1954, 34, 287.
- Smythe, C. McC., Nickel, J. F., and Bradley, S. E., The effect of epinephrine (USP), *l*-epinephrine, and *l*-norepinephrine on glomerular filtration rate, renal plasma flow, and the urinary excretion of sodium, potassium, and water in normal man. J. Clin. Invest., 1952, 31, 499.
- Elkinton, J. R., Crosley, A. P., Jr., Barker, H. G., and Clark, J. K., Alterations in renal hemodynamics and excretion of electrolytes. Federation Proc., 1950, 9, 37.

- Nickel, J. F., Smythe, C. McC., Papper, E. M., and Bradley, S. E., A study of the mode of action of the adrenal medullary hormones on sodium, potassium and water excretion in man. J. Clin. Invest., 1954, 33, 1687.
- Goodyer, A. V. N., and Jaeger, C. A., Renal response to nonshocking hemorrhage. Role of the autonomic nervous system and of the renal circulation. Am. J. Physiol., 1955, 180, 69.
- Smith, H. W., The Kidney: Structure and Function in Health and Disease. New York, Oxford Univ. Press, 1951.
- Harrison, H. E., A modification of the diphenylamine method for determination of inulin. Proc. Soc. Exper. Biol. & Med., 1942, 49, 111.
- Smith, H. W., Finkelstein, N., Aliminosa, L., Crawford, B., and Graber, M., The renal clearances of substituted hippuric acid derivatives and other aromatic acids in dog and man. J. Clin. Invest., 1945, 24, 388.

- Bradley, S. E., Chasis, H., Goldring, W., and Smith, H. W., Hemodynamic alterations in normotensive and hypertensive subjects during the pyrogenic reaction. J. Clin. Invest., 1945, 24, 749.
- Chasis, H., Ranges, H. A., Goldring, W., and Smith, H. W., The control of renal blood flow and glomerular filtration in normal man. J. Clin. Invest., 1938, 17, 683.
- Bennett, I. L., Jr., and Beeson, P. B., The properties and biologic effects of bacterial pyrogens. Medicine, 1950, 29, 365.
- 13. Judson, W. E., Hollander, W., Friedman, I. H., Hatcher, J. D., and Halperin, M. H., Studies on congestive failure: The effects of exercise, salt loading and of intravenous apresoline on cardiorenal hemodynamics and the excretion of sodium and water. J. Clin. Invest., 1953, 32, 579.
- Brandt, J. L., Ruskin, H. D., Zumoff, B., Castleman, L., and Zuckerman, S., Inhibition of renal tubular responsiveness to antidiuretic hormone by pyrogens. Proc. Soc. Exper. Biol. & Med., 1955, 88, 451.