

Sites of *in vivo* extraction and interconversion of testosterone and androstenedione in dogs

Alcide Chapdelaine

J Clin Invest. 1969;48(11):2063-2073. <https://doi.org/10.1172/JCI106172>.

Research Article

The interconversion and extraction of testosterone and androstenedione across and within different tissues or areas have been studied by the constant infusion technique. The results were calculated using the $^3\text{H}/^{14}\text{C}$ ratios and radioactive concentrations of testosterone and androstenedione obtained from afferent and efferent blood and tissues at equilibrium. In each tissue studied, the interconversion between testosterone and androstenedione inside the tissue was significantly higher than the corresponding interconversion across the tissue. The pulmonary contribution to the total interconversion between testosterone and androstenedione was far more important than that of any of the other tissues studied. The hepatic metabolic clearance rates of testosterone and androstenedione were not different from their metabolic clearance rates in the mesenteric area. The extraction of each of these compounds, although not negligible, was lower in the kidney and the femoral bed compared with the extraction in the liver and the mesenteric area. Finally, with the possible exception of the liver, testosterone and androstenedione were more completely metabolized when they originated from the cells than from afferent blood.

The evaluation of these different tissue transfer constants provides more precise information concerning the relative importance of different sites in the metabolism of these interconverting hormones.

Find the latest version:

<https://jci.me/106172/pdf>



Sites of In Vivo Extraction and Interconversion of Testosterone and Androstenedione in Dogs

ALCIDE CHAPDELAINE

From the Departments of Laboratories and Medicine, Maisonneuve Hospital and the University of Montreal, Montreal, Canada

ABSTRACT The interconversion and extraction of testosterone and androstenedione across and within different tissues or areas have been studied by the constant infusion technique. The results were calculated using the $^3\text{H}/^4\text{C}$ ratios and radioactive concentrations of testosterone and androstenedione obtained from afferent and efferent blood and tissues at equilibrium. In each tissue studied, the interconversion between testosterone and androstenedione inside the tissue was significantly higher than the corresponding interconversion across the tissue. The pulmonary contribution to the total interconversion between testosterone and androstenedione was far more important than that of any of the other tissues studied. The hepatic metabolic clearance rates of testosterone and androstenedione were not different from their metabolic clearance rates in the mesenteric area. The extraction of each of these compounds, although not negligible, was lower in the kidney and the femoral bed compared with the extraction in the liver and the mesenteric area. Finally, with the possible exception of the liver, testosterone and androstenedione were more completely metabolized when they originated from the cells than from afferent blood.

The evaluation of these different tissue transfer constants provides more precise information concerning the relative importance of different sites in the metabolism of these interconverting hormones.

INTRODUCTION

In recent years, many problems related to the secretion and metabolism of androgens, principally testosterone (T) and androstenedione (Δ) have been studied and partly elucidated (1-9). For these two steroids, blood production rates, contribution of other precursors to blood production rates, and their metabolic clearance

rates are well documented (2, 5, 7, 8, 10). It has also been clearly demonstrated (5) that, in the normal female, 60% of the circulating testosterone is derived from androstenedione by peripheral conversion.

Many aspects, however, such as the anatomical sites of their interconversion and metabolism are uncertain. In a study of the role of the splanchnic organs in the metabolism of T and Δ , Horton and Tait (5) have indirectly calculated that the conversion of Δ to T was mostly extrahepatic. However, their calculations depend on four assumptions which have not yet been completely verified experimentally. Thus, it is difficult to evaluate to what extent their calculations will be influenced if their assumptions are not completely applicable. Using a more direct approach, Rivarola, Singleton, and Migeon (9) have also found that the interconversion between T and Δ was mostly extrahepatic.

What also remains to be elucidated are the tissue metabolic clearance rates of T and Δ and their relative contribution to the total metabolic clearance rates. In humans, Rivarola et al. (9) did not find any extraction of T and Δ by the renal tissue. However, in the splanchnic system, 44% of T and 83% of Δ were extracted after one passage.

On the other hand, several tissues (as studied *in vitro*) can interconvert and, in some cases, metabolize testosterone and androstenedione, as shown for the muscle (11), the skin (12, 13), the prostate (14), the kidney (15, 16), and the digestive tract (17).

To study these different aspects, we have designed experiments which allow us to measure the interconversion between T and Δ and their extraction across different tissues or areas. The relative contribution of each of these tissues to the total interconversion between T and Δ and their metabolic clearance rates have also been calculated.

METHODS

Reagents. Methylene chloride, ethyl acetate, benzene, acetone, ligroin, methylcyclohexane, toluene, dichloromethane,

A portion of this work was presented at the International Congress on Hormonal Steroids, Milan, 1966.

Received for publication 15 April 1969.

cyclohexane, and hexane were all of spectral quality¹ and used without further purification. Methanol was redistilled twice. Ethanol was purified over *m*-phenylenediamine dihydrochloride and redistilled four times.

Chromatography. Chromatography paper was washed with redistilled methanol in a Friedrichs condenser² for 12 hr. Eastman chromatogram sheets, type K-301R2, coated with silica gel³ were used for thin-layer chromatography. The plates were heated to 100°C for 1 hr and stored in a desiccator before use. Chromatographic systems used in this work are described later in the text. Gas-liquid chromatography was done on a gas chromatograph, Packard model 871, at 240°C, on a Chromosorb-2 (80-120 mesh) column, coated with 3% SE-30.

Labeled steroids. Androstenedione-1,2-³H (50 mc/m mole) and testosterone-4-¹⁴C (58 mc/m mole) were obtained commercially from New England Nuclear Corp., Boston, Mass. These stock solutions were evaporated under nitrogen, diluted with purified ethanol, and stored at 0°C. To test for purity of the labeled steroids, a portion of each was added to 300 µg of authentic steroid⁴ and the mixture was carried through three successive chromatographies (systems 4, 5, and 2, see text). The specific activity was estimated before and after each chromatography by measuring the mass of steroid by absorption at 240 mµ in ethanol, or by gas chromatography, and the radioactivity was assayed by liquid scintillation spectrometry. The specific activity remained constant for each steroid.

Measurement of radioactivity. Each sample was transferred to a counting vial with absolute ethanol, dried, and counted in 10 ml of toluene containing 0.4% 2,5-diphenyloxazole and 0.005% 1,4-bis-2-(5-phenyloxazolyl)benzene, using a Packard liquid scintillation spectrometer, model 3375. All counting was corrected for quenching with an external standard; discriminator and gain settings were such to give an efficiency of 61.6% for ¹⁴C (³H then 0.1%) and 31.2% for ³H (¹⁴C then 8.2%) with backgrounds of 9.8 and 9.7 cpm respectively. Samples were counted three or four times for 20 min or longer so that the SE was less than 5%.

Subjects. 11 normal male and 2 normal female (dogs 1-2 and 1-5) mongrel dogs, weighing between 17 and 38 kg were used in this study. All experiments were started between 8 and 9 a.m. Dogs were anesthetized with pentobarbital. 2.4-5.5 µc of androstenedione-1,2-³H and 0.31-1.0 µc of testosterone-4-¹⁴C were injected into a foreleg vein in 10 ml of 10% ethanol in saline as a priming dose. 30 min later, a mixture of the same radioactive steroids diluted in 25 ml of 10% ethanol in saline was given as a constant infusion using a 30 ml Yale glass syringe fitted to a Harvard infusion pump⁴ model 600-910/920. Rates of infusion varied from 0.546 to 1.36 µc/min for androstenedione-1,2-³H and from 0.05 to 2.2 µc/min for testosterone-4-¹⁴C. Perfusion was performed via a Teflon⁵ catheter to avoid adsorption of radioactive steroid (18). In most cases, infusion lasted approximately 180 min or more since it was previously established, in two dogs not included in this study, that equilibrium had not been reached after 135 min of perfusion. With the exception of the first three dogs, the arterial-1 blood samples were taken 140 min or more after the begin-

ning of the infusion and the other blood samples 160-196 min after the beginning of the infusion. The amounts of blood taken varied between 50 and 115 ml.

Sampling of blood and tissue. At the beginning of each experiment, Teflon catheters were placed into the femoral artery and vein. During the first 30 min of the infusion, a laparotomy was performed and polyethylene catheters were introduced into the portal and renal veins, as described by Herd and Barger (19) in order to avoid disturbance of tissue blood flow. In each instance, the left spermatic vein was ligatured. At the end of the perfusion period, blood samples were withdrawn through these catheters (Table III). Blood from the right heart was taken by means of a catheter introduced into the right ventricle through a jugular vein. The hepatic vein sample was withdrawn through a Teflon catheter placed into the inferior vena cava up to the level of the hepatic veins. Immediately before sampling the blood, the inferior vena cava was tightly ligatured below and above the hepatic veins; the first 5 ml of blood were discarded. All blood samples were withdrawn into 50 ml heparinized syringes and immediately transferred into an Erlenmeyer flask containing 300 µg of authentic T and Δ and 100 ml of 50% methanol in order to stop any further metabolism of T and Δ by the blood (20). Immediately after, different tissues were taken, washed under running water, and rapidly homogenized in 35% methanol with a Virtis homogenizer¹.

Purification of samples. Each blood and tissue sample was further diluted to a final concentration of 33% methanol and dialyzed four times against 40% methanol (21). The dialysates, after the evaporation of methanol, were extracted three times with equal volumes of chloroform. The aqueous phase was then adjusted to pH 4.5 with hydrochloric acid and incubated at 37°C for 24 hr with β-glucuronidase enzyme⁶ (300 U/ml); after the addition of 100 µg of authentic testosterone, the mixture was extracted three times with equal volumes of chloroform. The free and conjugated steroid extracts were purified separately in the following manner:

- (a) paper chromatography in system 3 (ligroin: methanol: water, 10:9:1, at room temperature)
- (b) paper chromatography in system 4 (methylcyclohexane: toluene: methanol: water, 4:1:4:1)
- (c) chromatography on thin-layer in system 5 (dichloromethane: benzene: methanol, 160:160:15) or on paper in system 1 (petroleum ether: methanol: water, 10:7:3)
- (d) In some instances, each compound was rechromatographed in two dimensions on thin-layer in system 2 (ethyl acetate: benzene, 1:1) and in system 7 (ethyl acetate: methanol, 9:1).
- (e) In some instances, testosterone was acetylated and rechromatographed in system 6 (cyclohexane: methanol: water, 10:10:1) or in system 8 (ligroin: methanol: water, 10:9:1 at 4°C). After saponification (22) or as the acetate, it was then crystallized two or three times to constant (³H/¹⁴C) ratio.
- (f) In a few cases, testosterone was oxidized to androstenedione (23), rechromatographed in system 2, and the material was then crystallized.
- (g) Androstenedione was, in some cases, reduced to testosterone (24), rechromatographed in system 3, and then crystallized two or three times to constant ³H/¹⁴C ratio.

¹ Obtained from Warner-Chilcott, Scarborough, Ontario.

² Fisher Scientific Company, Fair Lawn, N. J.

³ Canlab Co., 8655 Delmeade Road, Montreal.

⁴ Obtained from Steraloids Inc., Pawling, N. Y.

⁵ Harvard Apparatus Co., Inc., Dover, Mass.

⁶ Johnston Industrial Plastics, 103 Smith, Ville Lasalle, Québec, Canada.

TABLE I
 $^3\text{H}/^{14}\text{C}$ Ratios and Radioactivity Concentrations of Testosterone (T) and
 Androstenedione (Δ) during the Purification Procedure* in Dog 1-4

Procedures	Blood samples										
	Arterial-1		Arterial-2		Femoral vein		Portal vein		Hepatic vein		
	T	Δ	T	Δ	T	Δ	T	Δ	T	Δ	
C* on system 3 followed by C on system 4	Ratios.....	0.42	13.56	0.48	13.60	0.50	13.21	0.43	13.17	0.79	7.30
	^3H	12,400	126,720	12,510	124,270	11,335	90,900	6,490	36,810	7,510	37,280
	^{14}C	29,345	9,340	26,060	9,137	22,845	6,880	15,105	2,795	9,530	5,105
C on system 5	Ratios.....		13.76		13.52		13.37		12.88		7.43
	^3H		129,015		125,150		87,915		29,825		32,130
	^{14}C		9,376		9,256		6,575		2,315		4,325
Acetylation and C on system 8	Ratios.....	0.46		0.41		0.51		0.47		0.82	
	^3H	11,795		12,010		10,300		5,765		7,730	
	^{14}C	25,645		29,270		20,390		12,265		9,420	
Saponification and C on system 3	Ratios.....									0.79	
	^3H									6,705	
	^{14}C									8,490	
C on system 1	Ratios.....		13.38		13.55		13.34		13.26		7.54
	^3H		121,665		119,730		107,050		30,090		34,770
	^{14}C		9,095		8,836		8,025		2,270		4,610
C on system 5	Ratios.....	0.44		0.43		0.53		0.45			
	^3H	11,820		12,980		10,180		5,491			
	^{14}C	26,863		29,900		19,155		12,070			
Crystallization, 1st	Crystals.....	0.44	13.49	0.45	13.66	0.51	13.19	0.46	13.04	0.80	7.26
Crystallization, 2nd	Crystals.....	0.43	13.65	0.41	13.46		13.24	0.44	13.16	0.80	7.50
	ml 		13.41								7.33
Crystallization, 3rd	Crystals.....		13.59	0.46	13.57		13.52	0.44	13.10		7.46
	ml			0.48				0.45			7.40
Mean	Ratios.....	0.44	13.55	0.44	13.56	0.51	13.31	0.45	13.10	0.80	7.43
	^3H	12,005	125,800	12,500	123,050	10,605	95,290	5,915	32,240	7,315	34,725
	^{14}C	27,285	9,270	28,410	9,076	20,795	7,160	13,145	2,460	9,145	4,680

* Chromatography.

† ^3H concentration in DPM/100 ml of whole blood corrected for losses.

‡ ^{14}C concentration in DPM/100 ml of whole blood corrected for losses.

|| Mother liquor.

After each purification step, the recovery of added carrier was measured by absorption at 240 $m\mu$ in ethanol and by gas chromatography as previously described, and radioactivity was estimated by liquid scintillation spectrometry. Over-all recoveries of authentic T and Δ added to whole blood varied between 30% and 50%. All radioactive concentrations are expressed in dpm per 100 ml of whole blood and are corrected for procedural losses.

A typical example (results from dog 1-4, Table I) shows that the radioactive concentrations and the $^3\text{H}/^{14}\text{C}$ ratios of T and Δ isolated from different blood samples are quite constant throughout the purification procedures. Comparison of T and Δ ratios and concentrations between the two arterial blood samples taken at 15- to 20-min intervals gave evidence that equilibrium in blood was obtained during the continuous infusion. If the first arterial $^3\text{H}/^{14}\text{C}$ ratios are compared, the mean arterial-2 ratios were 100% for T and Δ ; if we compare radioactive concentrations, the mean arterial-2 values were 100 ± 2.0 SE for testosterone- ^3H , 100 ± 2.0 SE for testosterone- ^{14}C , 100 ± 1.1 SE for androstenedione- ^3H , and 100 ± 1.0 SE for androstenedione- ^{14}C . It is evident that there is no trend in the values. All data used in the calculations are shown in Tables II and III.

Calculations and definition of terms. Metabolic clearance rates were calculated as described by Tait (25), blood transfer constants as described by Gurpide, Mann, and Lieberman (26) (see Appendix), and transfer constants through each organ were determined according to formulas (see Appendix) derived by Gurpide⁷, using the model shown in Fig. 1. This model represents the kidney (K) with its afferent arterial (A) blood and its efferent venous (V) blood. The same model can be applied to any other organ.

Definition of terms used subsequently are as follows: $R_{T^{14}\text{C}}$ and $R_{\Delta^{14}\text{C}}$, perfusion rate of T- ^{14}C and Δ - ^{14}C in dpm per day. Radioactive concentrations (c) in dpm per 100 ml of blood are symbolized as follows: $c_{TA}^{^3\text{H}}$ and $c_{TA}^{^{14}\text{C}}$, concentration of testosterone- ^3H and ^{14}C , isolated from arterial blood. $^3\text{H}/^{14}\text{C}$ ratios are symbolized similarly: example, $(^3\text{H}/^{14}\text{C})_{TK}$ is the $^3\text{H}/^{14}\text{C}$ ratio in testosterone isolated from the kidney. MCR^T and MCR^Δ , blood metabolic clearance rate of T and Δ . ρ_{BB}^{TA} and $\rho_{BB}^{\Delta T}$, blood transfer constants (ρ) between T and Δ .

⁷ Gurpide, E. 1962. Personal communication.

Transfer constants (ρ) across or inside each tissue or area, using the kidney as an example are symbolized as follows:
 ρ_{AV}^{Δ} transfer constant from arterial T to renal venous Δ

(transtissue interconversion). ρ_{KK}^{Δ} transfer constant from T to Δ inside the renal cells (intratissue interconversion). ρ_{AV}^{TT} transfer constant from arterial T to renal venous T. The

percentage of extraction of T across the kidney will then be $\left[1 - \rho_{AV}^{TT}\right] \times 100$. ρ_{KV}^{TT} transfer constant from renal T to corresponding venous T. The percentage of extraction of T formed in the kidney will then be $\left[1 - \rho_{KV}^{TT}\right] \times 100$.

The androstenedione transfer constants are expressed in the same way.

TABLE II
*Data for the Determination of Blood MCR^T MCR^A (ρ), BB, and all Transtissular (ρ)**

Dog No.	Blood samples											
	Arterial		Femoral vein		Renal vein		Portal vein		Hepatic vein		Right heart	
	T	Δ	T	Δ	T	Δ	T	Δ	T	Δ	T	Δ
1	Ratios†.....	1.78	23.08					1.81	21.19			
	³ H§.....	16,816	102,040					6,944	37,591			
	¹⁴ C 	9,290	4,526					4,233	1,756			
3	Ratios.....	2.42	26.1			21.3	2.80	17.51	3.43			
	³ H.....	54,823	220,807			170,403	41,696	119,544	9,784			
	¹⁴ C.....	22,743	8,377			7,640	14,524	6,844	2,871			
4	Ratios.....	1.44	37.23			1.59	22.61	1.48	34.23	1.98	20.20	
	³ H.....	28,787	184,747			27,272	144,999	19,908	114,160	1,505	8,225	
	¹⁴ C.....	19,948	4,928			17,076	6,403	13,466	3,330	756	419	
5	Ratios.....	2.25	30.43	2.76	28.95	2.79	27.5	2.88	26.2	3.05	27.1	
	³ H.....	27,525	224,600	23,242	93,105	32,474	11,121	14,150	92,727	7,120	60,112	
	¹⁴ C.....	12,065	7,403	8,200	3,216	11,680	4,025	4,810	3,410	2,370	2,229	
6	Ratios.....	0.64	9.1	0.71	6.96	1.1	6.8	0.81	7.9	1.0	8.4	
	³ H.....	46,265	304,660	40,085	125,060	71,445	217,200	36,655	110,265	8,450	26,660	
	¹⁴ C.....	72,370	33,479	56,455	17,970	62,070	31,560	47,155	13,760	7,460	3,175	
7	Ratios.....	0.67	10.53	0.95	9.9	0.68	8.0	0.67	8.6	1.06	7.3	
	³ H.....	43,442	273,160	48,425	173,055	31,422	256,900	32,257	244,380	11,530	72,990	
	¹⁴ C.....	64,840	25,940	50,987	16,270	46,210	30,125	48,145	28,420	10,035	9,588	
9	Ratios.....	0.66	11.80	0.87	9.35	0.79	8.90	0.85	10.12	1.29	7.50	
	³ H.....	33,627	190,464	23,826	76,472	31,853	149,624	17,625	91,075	18,177	53,800	
	¹⁴ C.....	50,837	16,140	27,615	8,180	39,350	16,854	21,144	9,000	14,102	7,254	
1-0	Ratios.....	0.62	15.47	1.16	12.62	1.24	9.77	1.01	11.35			
	³ H.....	21,930	145,750	36,855	129,605	33,600	92,225	26,660	83,455			
	¹⁴ C.....	35,375	9,420	31,770	10,270	27,100	9,440	26,395	7,350			
1-1	Ratios.....	0.215	11.05	0.27	9.65			0.82	7.95	0.95	7.01	
	³ H.....	9,147	193,252	9,955	130,195			23,820	94,910	6,945	18,865	
	¹⁴ C.....	42,547	17,490	36,865	13,490			29,050	11,940	7,305	2,690	
1-2	Ratios.....	0.58	8.77							0.25	24.47	
	³ H.....	36,560	145,320							13,850	222,450	
	¹⁴ C.....	63,040	16,570							55,395	9,090	
1-3	Ratios.....	0.40	17.68	0.61	15.66					0.28	26.56	
	³ H.....	16,542	259,755	24,035	217,805					11,070	270,300	
	¹⁴ C.....	41,355	14,690	39,400	13,910					39,535	10,175	
1-4	Ratios.....	0.44	13.58	0.51	13.31			0.45	13.10	0.80	7.42	
	³ H.....	12,252	124,425	10,605	95,290			5,915	32,240	7,315	34,725	
	¹⁴ C.....	27,847	9,173	20,795	7,160			13,145	2,460	9,145	4,680	
1-5	Ratios.....	0.642	8.18							0.399	13.08	
	³ H.....	29,685	150,445							25,744	177,993	
	¹⁴ C.....	46,238	18,391							64,522	13,608	

* MCR^T MCR^A (ρ) BB and all transtissular (ρ) are defined in the text.

† ³H/¹⁴C ratios.

‡ ³H concentration in DPM/100 ml of whole blood corrected for losses.

|| ¹⁴C concentration in DPM/100 ml of whole blood corrected for losses.

TABLE III
Data for the Determination of all Intratissular (ρ)*

No.		Tissue samples									
		Liver		Mesenteric area		Kidney		Muscle		Lung	
		T	Δ	T	Δ	T	Δ	T	Δ	T	Δ
4	Ratios‡	5.51	11.81					0.9	7.8		
6	Ratios			1.4	6.3					3.0	4.5
7	Ratios	1.46	6.5			7.86		1.1	9.7	2.96	3.78
9	Ratios	1.51	7.65	1.02	8.80			2.10	5.31		
1-0	Ratios					2.15	5.1	0.72	11.22		
1-1	Ratios							0.65	9.82		
1-2	Ratios									2.37	5.38
1-3	Ratios									12.73	2.50
1-4	Ratios			6.11	0.53	7.99		0.47	10.51		
1-5	Ratios										4.38

* All intratissular (ρ) are defined in the text.

‡ $^3\text{H}/^{14}\text{C}$ ratios.

RESULTS

Metabolic clearance rate (Table IV). The metabolic clearance rate of testosterone (MCR^T) was 1502 ± 113 (SE) liters/day (13 subjects). There was no significant difference between male and female dogs. The metabolic clearance rate of androstenedione (MCR^A) was also identical in male and female dogs; mean value of 1741 ± 140 (SE) liters/day (13 subjects). Furthermore, there appears to be no significant difference between the metabolic clearance rates of T and Δ in dogs ($P > 0.05$, *t* test).

Blood transfer constants (Table IV). The blood transfer constants ($\rho_{BB}^{T\Delta}$ and $\rho_{BB}^{\Delta T}$) are calculated from the $^3\text{H}/^{14}\text{C}$ ratios of the infused mixture and of the androstenedione and testosterone isolated from the arterial blood (26). The transfer constant from blood T to blood Δ was $41.6 \pm 2.9\%$ SE in males (11 subjects), and 45% in females (2 subjects); these values are not statistically different. The transfer constant from blood Δ to blood T was $13.6 \pm 1.2\%$ SE in 11 male dogs and 10.5 ± 6.5 SE in 2 female dogs; these values are also not statistically different. The combined values were respectively $42 \pm 2.5\%$ SE and $13 \pm 1.3\%$ SE; the $\rho_{BB}^{T\Delta}$ is significantly higher than the $\rho_{BB}^{\Delta T}$ ($P < 0.01$).

Tissue transfer constants (Table V). $^3\text{H}/^{14}\text{C}$ ratios and radioactive concentrations of T and Δ , used in the equations as hepatic arterial values, were calculated assuming that, in the anesthetized dog, 80% of the blood entering the liver comes from the portal vein and 20% from the hepatic arteries (27). The mesenteric area includes all tissues between the artery and the portal

vein, in other words, the digestive tract, its mesentery and the intestinal adipose tissue. Similarly, the posterior limb includes all tissues between the femoral artery and vein, principally the bone, muscle, skin, and subcutaneous tissue.

The interconversion between T and Δ , across and inside the liver, mesenteric area, posterior limb, and kidney, does not significantly favor the formation of an-




FIGURE 1 Model describing the renal (K) transfer constants (ρ). Each arrow, from 1 to 8, represents a specific transfer constant, respectively $\rho_{AV}^{T\Delta}$, $\rho_{AV}^{\Delta T}$, $\rho_{KK}^{T\Delta}$, $\rho_{KK}^{\Delta T}$, ρ_{AV}^{TT} , $\rho_{AV}^{\Delta\Delta}$, ρ_{KV}^{TT} , and $\rho_{KV}^{\Delta\Delta}$.

drostenedione (in each case, $P > 0.1$ for the interconversion across and $P > 0.3$ for the interconversion inside). Across and inside the lung, the interconversion seems to favor the formation of androstenedione, but the P values could not be calculated from so few data. In each tissue studied, the interconversion between T and Δ inside the tissue was significantly higher than the corresponding interconversion across the tissue (in all cases, $P < 0.05$).

Across the liver, mesenteric area, kidney, and posterior limb, the extraction of T was not significantly different from the extraction of Δ (in each case, $P > 0.1$). The extraction of T and Δ was similar across the kidney and the posterior limb, but significantly higher across the liver and the mesenteric area. Except for the posterior limb, the extraction of T and Δ preformed in each tissue was significantly higher than the corresponding transtissue extraction (in all cases, $P < 0.01$). The transpulmonary extraction of T and Δ was highly variable (see Discussion) and these values were not used in the calculations of the tissue metabolic clearance rates.

Tissue metabolic clearance rates (Table VI). The tissue metabolic clearance rate (liters/day) is defined as the product of tissue blood flow multiplied by the corresponding transtissue extraction, divided by 100. It should be pointed out that the metabolic clearance

rate of each tissue is also expressed in liters/day of whole blood to avoid possible error due to different partition coefficients between plasma and red cells (28). If we assume the cardiac output to be 125 ml/min per kg in the anesthetized dog (29), each tissue blood flow can be estimated as a percentage of the total cardiac output. The values used in our calculations were as follows: liver blood flow, 25% of the cardiac output (30, 31); mesenteric area blood flow or portal blood flow, 20% (29, 32, 33); renal blood flow, 20% (34); and blood flow in the femoral bed, approximately 12%. It is also important to realize that these different blood flows can vary widely.

The hepatic metabolic clearance rates were corrected for the fact that 80% of the liver blood flow, corresponding to the portal blood flow, has already crossed the mesenteric area which extracts 42% and 51% of the arterial testosterone and androstenedione. The hepatic MCR^T is higher than the hepatic MCR^Δ ($P < 0.01$). In the kidneys and mesenteric area, MCR^T is not significantly different from MCR^Δ ($P > 0.3$ in both tissues). In the femoral bed, MCR^Δ is higher than MCR^T ($P < 0.02$).

The hepatic MCR^T is not significantly different from the mesenteric MCR^T ($P > 0.2$). However, the testosterone metabolic clearance rates in the liver and in the mesenteric area are significantly higher than MCR^T in the renal and femoral bed ($P < 0.01$ in each case). The hepatic MCR^T is not significantly different from the sum of the MCR^T in the mesenteric area, kidneys, and posterior limbs ($P > 0.3$).

The hepatic and mesenteric MCR^Δ are not different ($P > 0.8$), but are definitely higher than the renal and femoral MCR^Δ ($P < 0.05$ in each instance). Also the renal MCR^Δ is not different from the femoral MCR^Δ ($P > 0.8$). The sum of the MCR^Δ in the mesenteric area, kidneys, and posterior limbs appears higher than the hepatic MCR^Δ but the difference is not significant ($P > 0.1$).

Contribution of each tissue to the total conversion between T and Δ (Table VI). To evaluate the relative contribution of each tissue to the total interconversion between T and Δ , each transtissue transfer constant was compared with the total blood transfer constant between T and Δ , using the formula shown in the second footnote (#) of Table VI. The pulmonary contribution to the total interconversion between T and Δ is far more important than any of the other tissues studied, while the renal contribution to the total interconversion between T and Δ was slightly higher than the contribution of the mesenteric area and posterior limbs ($P < 0.05$), but is not different from the hepatic contribution ($P > 0.1$).

TABLE IV
Metabolic Clearance Rates and Total in Vivo Interconversion of Testosterone and Androstenedione

Dog	MCR^T	MCR^Δ	ρ_{BB}^{TA*}	ρ_{BB}^{PT*}
	liters/day‡	liters/day‡		
1	1719	1440	46	17
2	1208	1334	41	23
4	1329	1859	29	13
5	1774	1281	43	17
6	970	1163	55	13
7	1084	1298	48	13
9	1382	1861	42	13
1-0	1896	2390	32	12
1-1	1576	1802	45	4
1-2	1064	2397	57	12
1-3	1622	1340	28	8
1-4	2459	2769	37	9
1-5	1451	1699	45	17
Mean	1502	1741	42	13
SE	±113	±140	±2.5	±1.3

* ρ_{BB}^{TA} and $\rho_{BB}^{\Delta T}$ were calculated from the perfused $^3H/^{14}C$ ratios and the $^3H/^{14}C$ ratios in testosterone and androstenedione isolated from arterial blood.

‡ MCR^T and MCR^Δ are expressed in liters/day of whole blood.

TABLE V
Tissue Transfer Constants (ρ)* and Extractions†

Tissue or area	Transtissue (ρ)		Intratissue (ρ)		Transtissue extraction‡		Intratissue extraction‡	
	$\rho_{T\Delta}$	$\rho_{\Delta T}$	$\rho_{T\Delta}$	$\rho_{\Delta T}$	T $(1 - \rho_{AV}^{TT})$	Δ $(1 - \rho_{AV}^{\Delta\Delta})$	T $(1 - \rho_{KV}^{TT})$	Δ $(1 - \rho_{KV}^{\Delta\Delta})$
Liver	Mean \pm SE§	4.3 \pm 1.8	2.7 \pm 0.9	31.5 \pm 8.6	31.7 \pm 11.7	72.6 \pm 6.4	66.8 \pm 8.2	85.0 \pm 9.9
	SD	4.8	2.6	17.2	20.2	18.0	21.7	17.2
	No. of dogs	(7)	(8)	(4)	(3)	(8)	(7)	(3)
Mesenteric area	Mean \pm SE	4.6 \pm 1.1	2.6 \pm 1.0	19.0 \pm 2.9	11.0 \pm 5.5	42.0 \pm 4.1	51.0 \pm 5.4	87.0 \pm 4.9
	SE	3.5	3.2	5.9	9.5	13.1	17.2	8.5
	No. of dogs	(10)	(10)	(4)	(3)	(10)	(10)	(3)
Kidney	Mean \pm SE	9.4 \pm 1.2	4.9 \pm 2.0	54.0 \pm 6.6	42.0 \pm 3.6	20.0 \pm 3.6	28.0 \pm 5.1	86.0 \pm 10.6
	SD	3.4	4.9	11.3	6.2	8.8	13.6	18.3
	No. of dogs	(7)	(6)	(3)	(3)	(6)	(7)	(3)
Posterior limb	Mean \pm SE	3.3 \pm 0.7	3.7 \pm 1.3	8.5 \pm 1.5¶	9.4 \pm 2.8¶	23.0 \pm 4.5	37.6 \pm 7.0	
	SD	2.1	3.7	3.7	6.3	12.8	19.9	
	No. of dogs	(8)	(8)	(6)	(5)	(8)	(8)	
Lung	Mean \pm SE	14.3 \pm 2.6	5.9 \pm 2.2	53.5 \pm 7.5	47.2 \pm 11.2		19.3 \pm 9.0	89.5 \pm 5.5
	SD	4.5	3.8	10.6	15.9		15.6	7.8
	No. of dogs	(3)	(3)	(2)	(2)		(3)	(2)

* All (ρ) values are expressed in percentages.

† Transtissue and intratissue extractions are calculated as $[1 - \text{corresponding } (\rho)] \times 100$.

‡ Standard deviation of the mean.

§ Standard deviation from the range.

¶ Transfer constants inside the muscle cells.

$^3\text{H}/^{14}\text{C}$ ratios in testosterone glucuronide. In six subjects, the $^3\text{H}/^{14}\text{C}$ ratios of arterial testosterone glucuronide (TG) were all significantly higher than those observed in the corresponding arterial free testosterone, indicating that testosterone glucuronide is not uniquely derived from blood free testosterone. In three subjects

studied, the $^3\text{H}/^{14}\text{C}$ ratios of the renal TG was 2, 2.5, and 3 times higher than the corresponding ratios in arterial TG and the $^3\text{H}/^{14}\text{C}$ ratios of mesenteric TG, 1.8, 2.0, and 2.1 times higher than the ratios in arterial TG, indicating that testosterone was conjugated in the renal and mesenteric cells. Furthermore, the $^3\text{H}/^{14}\text{C}$ ratios of renal

TABLE VI
Tissue Metabolic Clearance Rates* (Liters/Day and % of Total MCR) and Contribution‡ of Each Tissue to the Total Conversion between T and Δ §

Tissue or area	MCR T^*		MCR Δ^*		% of total‡		% of total‡	
	liters/day	%	liters/day	%	$\rho_{BB}^{T\Delta}$	$\rho_{BB}^{\Delta T}$	$\rho_{BB}^{T\Delta}$	$\rho_{BB}^{\Delta T}$
Liver	552 \pm 47(8)	40 \pm 7.0	463 \pm 52(7)¶	29 \pm 4.7	2.7 \pm 1.2(7)		6.7 \pm 2.4(8)	
Mesenteric area	384 \pm 60(10)	25 \pm 3.8	451 \pm 68(10)	26 \pm 3.2	2.2 \pm 0.5(10)		3.1 \pm 1.4(10)	
Kidneys	208 \pm 52(6)	16 \pm 4.0	252 \pm 47(7)	16 \pm 2.8	4.8 \pm 0.9(7)		7.6 \pm 3.3(6)	
Posterior limbs	140 \pm 39(8)	10 \pm 3.0	211 \pm 49(8)	14 \pm 3.3	1.0 \pm 0.2(8)		3.9 \pm 1.3(8)	
Lungs	**		**		34.0 \pm 6.1(3)		45.4 \pm 16.4(3)	

* Tissue metabolic clearance rates (liters/day of whole blood calculated as tissue blood flow \times transtissue extraction/100.

† % of total $\rho_{BB}^{T\Delta}$ or ΔT calculated as $\frac{\rho_{AV}^{T\Delta} \text{ or } \Delta T}{\rho_{BB}^{T\Delta} \text{ or } \Delta T} \times \frac{\text{tissue blood flow}}{\text{cardiac output}} \times 100$.

‡ All values are mean \pm SE.

¶ Values corrected for the extraction across the mesenteric area.

|| Number of dogs in parenthesis.

** Lung metabolic clearance rates were not calculated (see Discussion).

TG was 25% higher than the ratios of the free renal testosterone, indicating that in the renal cells TG did not originate uniquely from free T.

DISCUSSION

In these experiments, it should be pointed out that the transfer constants across or inside each tissue are quite variable in the dogs studied and several important points should be mentioned to explain these apparent discrepancies: (a) Differences in metabolism could exist from dog to dog, apart from the other following factors. (b) In spite of an extensive purification of the plasma steroids, the variations in the blood concentrations of labeled T and Δ were $\pm 10\%$ in some cases. In these instances, the errors in the calculated results could be appreciable. (c) In acute experiments on anesthetized dogs, the blood flow across any tissue can vary (31), and furthermore, anaesthesia can alter certain functions, particularly those of the liver. Theoretically, our catheters should not influence tissue blood flows (19), but this possibility is not definitely excluded. (d) The trans-pulmonary extraction of T and Δ were quite variable and for this reason pulmonary MCR^T and MCR ^{Δ} were not calculated.

Unlike the human (6-9), the mean values for blood MCR of testosterone and androstenedione are not significantly different in dogs. The MCR of testosterone, however, is higher and the MCR of androstenedione is lower than that found in humans. Furthermore, there is no difference in male and female dogs for both the MCR of testosterone and androstenedione. These species differences and the greater conversion of blood T to blood Δ strongly suggest that, unlike the human, there is a much lower binding of testosterone to protein in dogs.

If, as has been done in the human (5), the MCR of testosterone and androstenedione are compared with the mean hepatic blood flow in our dogs (1053 liters/day), at least 30% of the total testosterone metabolism and 40% of the total androstenedione metabolism occurs extrahepatically. It is evident from the individual tissue MCR that these values are well underestimated because the hepatic MCR^T and MCR ^{Δ} represent only 40% and 29% of the total MCR^T and MCR ^{Δ} . It should also be pointed out that the MCR^T and MCR ^{Δ} in the mesenteric area are not significantly different from their respective hepatic MCR's. If T and Δ were secreted into the bile and reabsorbed by the intestine, our mesenteric metabolic clearance rates would be underestimated.

The blood transfer constants between T and Δ are greater in dogs than in humans (5). The greater conversion of blood T to blood Δ may be due to a lower testosterone protein binding affinity and may also explain, at least in part, the greater testosterone MCR in dogs. In agreement with the values found for the inter-

conversion of estrone and estradiol in humans (28), the over-all interconversion between T and Δ in the dog is in favor of the 17-ketone, androstenedione. However, when examined individually, the transtissue transfer constants between T and Δ , while tending to favor the formation of the 17-ketone compound, are not statistically different, with the exception of the lung. Furthermore, the tissue MCR^A is not significantly higher than the corresponding tissue MCR^T, except in the posterior limb.

Rivarola et al. (9) have reported that in humans not more than 28% of the total conversion of blood T to blood Δ , and not more than 16% of the total conversion of blood Δ to blood T occurs in the splanchnic system. These values, as well as the values obtained by the same authors for the extraction of T and Δ , are very difficult to interpret because they are calculated from radioactive concentrations obtained from a peripheral vein. These data are not necessarily equal to the radioactive concentrations entering the splanchnic circulation, particularly if, as in dogs, T and Δ are extracted and interconverted peripherally and in the mesenteric area. Furthermore, when compared with the total interconversion, the splanchnic interconversion should be corrected by a factor, splanchnic blood flow over total cardiac output. When corrected for their respective blood flow, the liver, mesenteric area, kidneys, and posterior limbs are each responsible for less than 10% of the blood interconversion between T and Δ . In other words, more than 90% of the total interconversion occurs extrahepatically. The lung is by far the most important contributor to the blood interconversion of T and Δ in the dog, undoubtedly due primarily to its higher blood flow.

In all of the organs or areas studied, the interconversion between T and Δ inside the tissue was definitely higher than the corresponding interconversion across the tissue. Furthermore, the intratissue interconversion could be underestimated if the tissue extracts were contaminated by T and Δ present in the interstitial water. This means that part of the T or Δ formed from its precursor inside the cells is further metabolized and does not appear in the efferent vein. It is possible that this "cellular" compound has an important local metabolic effect before being metabolized, and this action would not be reflected in the plasma or blood concentration.

The canine hepatic extraction of testosterone is higher than in humans (5, 9), which is another possible explanation for the high testosterone metabolic clearance rate found in dogs. On the contrary, the hepatic extraction of androstenedione is lower in dogs than in the humans but it is possible that the value found in humans by Rivarola et al. (9) is overestimated for reasons already mentioned. The transhepatic extraction of T was

not significantly lower than the extraction of T formed in the liver. Similar values calculated by Horton and Tait (5), in the human, gave 60% for the maximal extraction of T across the liver and 98% for the extraction of T formed in the hepatic cells; however, the calculations used to evaluate the extraction of T formed in the liver could have been overestimated by the extrahepatic production of testosterone glucuronide (see below), by the conversion of Δ to T in the mesenteric area, and also by the extrasplanchnic metabolism of testosterone.

In the mesenteric area, the kidney, and the lung, the extraction of T and Δ formed within the cells was significantly higher than the extraction of T presented to the cells via its afferent blood supply. This could be explained by the possibility that the compound circulating in blood could not reach the cells due to a rapid flow rate or to protein binding, or it could mean that the compound made in the tissue from its precursor is metabolized differently from the preformed compound entering the tissue.

As already observed in humans (6), testosterone glucuronide is not uniquely derived from blood testosterone. Furthermore, testosterone glucuronide is not solely made in the liver but also in the mesenteric area and in the kidney, a finding which is in agreement with *in vitro* (17) and *in vivo* studies (35). Inside the liver and the kidney, testosterone glucuronide does not seem to derive uniquely from tissue testosterone.

The important interconversion and extraction both inside and across the mesenteric area emphasizes the importance, at least in dogs, of comparing portal vein blood with hepatic vein blood during the study of the role of the liver in steroid metabolism.

In agreement with other studies (15, 16), the canine renal tissue very actively metabolizes testosterone and androstenedione. In one human subject, Rivarola et al. (9) did not find any extraction and interconversion of T and Δ by the kidney.

It is impossible to conclude from our study whether some fraction of free and conjugated testosterone or free androstenedione, formed in the kidney, is directly secreted into the urine without equilibration with the corresponding blood compound.

APPENDIX

1. The subsequent equations show the derivation of formulas used to calculate all tissue transfer constants. These mathematical formulas were derived using the kidney as a model (Fig. 1); this model can be applied to any other tissue.

From Fick's principle and assuming that testosterone is not formed in the renal tissue, the extraction of T across the kidney (KE) will be:

$$KE = \frac{c_{TA}^{3H} - c_{TV}^{3H}}{c_{TA}^{3H}} = 1 - \frac{c_{TV}^{3H}}{c_{TA}^{3H}}$$

Since the transfer constant from arterial T to venous T across the kidney (ρ_{AV}^{TT}) is equal to $(1 - KE)$, the equation 1 becomes:

$$\rho_{AV}^{TT} = \frac{c_{TV}^{3H}}{c_{TA}^{3H}}. \quad (2)$$

However, in this study, the above equation is not entirely correct, since a fraction of Δ -³H entering the kidney will leave this tissue as T-³H, it must therefore be subtracted from T_V-³H:

$$\rho_{AV}^{TT} = \frac{c_{TV}^{3H} - \rho_{AV}^{\Delta T} \times c_{\Delta A}^{3H}}{c_{TA}^{3H}}. \quad (3)$$

The same relation being true for the other label, ¹⁴C:

$$\rho_{AV}^{TT} = \frac{c_{TV}^{3H} - \rho_{AV}^{\Delta T} \times c_{\Delta A}^{3H}}{c_{TA}^{3H}} = \frac{c_{TV}^{14C} - \rho_{AV}^{\Delta T} \times c_{\Delta A}^{14C}}{c_{TA}^{14C}}. \quad (4)$$

Isolating $\rho_{AV}^{\Delta T}$ from equation 4:

$$\rho_{AV}^{\Delta T} = \frac{c_{TV}^{3H} c_{TA}^{14C} - c_{TV}^{14C} c_{TA}^{3H}}{c_{\Delta A}^{3H} c_{TA}^{14C} - c_{\Delta A}^{14C} c_{TA}^{3H}}. \quad (5)$$

Multiplying equation 5 by $c_{TA}^{14C} c_{TV}^{14C} / c_{TA}^{14C} c_{TV}^{14C} = 1$:

$$\rho_{AV}^{\Delta T} = c_{TV}^{14C} \frac{(^3H/^{14}C)_{TV} - (^3H/^{14}C)_{TA}}{c_{\Delta A}^{3H} - (^3H/^{14}C)_{TA} c_{\Delta A}^{14C}}. \quad (6)$$

Multiplying equation 6 by $c_{\Delta A}^{14C} / c_{\Delta A}^{3H} = 1$:

$$\rho_{AV}^{\Delta T} = \frac{c_{TV}^{14C} (^3H/^{14}C)_{TV} - (^3H/^{14}C)_{TA}}{c_{\Delta A}^{14C} (^3H/^{14}C)_{\Delta A} - (^3H/^{14}C)_{TA}}. \quad (7)$$

Equation for $\rho_{AV}^{\Delta A}$ can be similarly derived:

$$\rho_{AV}^{\Delta A} = \frac{c_{\Delta V}^{14C} (^3H/^{14}C)_{\Delta V} - (^3H/^{14}C)_{\Delta A}}{c_{TA}^{14C} (^3H/^{14}C)_{TA} - (^3H/^{14}C)_{\Delta A}}. \quad (8)$$

Substituting the $\rho_{AV}^{\Delta T}$ value, calculated from equation 7, in equation 3 ρ_{AV}^{TT} can now be calculated and similarly the $\rho_{AV}^{\Delta \Delta}$ is equal to:

$$\rho_{AV}^{\Delta \Delta} = \frac{c_{\Delta V}^{3H} - \rho_{AV}^{\Delta T} \times c_{TA}^{3H}}{c_{\Delta A}^{3H}}. \quad (9)$$

To calculate transfer constants inside the renal cells, we must assume that all arterial T or Δ enters the cellular pool and then:

$$(^3H/^{14}C)_{TK} = \frac{c_{TA}^{3H} + \rho_{KK}^{\Delta T} c_{\Delta A}^{3H}}{c_{TA}^{14C} + \rho_{KK}^{\Delta T} c_{\Delta A}^{14C}}. \quad (10)$$

Solving equation 10 for $\rho_{KK}^{\Delta T}$:

$$\rho_{KK}^{\Delta T} = \frac{c_{TA}^{3H} - (^3H/^{14}C)_{TK} c_{TA}^{14C}}{(^3H/^{14}C)_{TK} c_{\Delta A}^{14C} - c_{\Delta A}^{3H}}. \quad (11)$$

Similarly $\rho_{KK}^{T\Delta}$ is equal to:

$$\rho_{KK}^{T\Delta} = \frac{c_{\Delta_A}^{3H} - (3H/14C)_{\Delta_K} c_{\Delta_A}^{14C}}{(3H/14C)_{\Delta_K} c_{\Delta_A}^{14C} - c_{T_A}^{3H}} \quad (12)$$

At equilibrium: $\rho_{AV}^{\Delta T} = \rho_{AK}^{\Delta\Delta} \times \rho_{KK}^{\Delta T} \times \rho_{KV}^{TT}$.

And then:

$$\rho_{KV}^{TT} = \frac{\rho_{AV}^{\Delta T}}{\rho_{AK}^{\Delta\Delta} \times \rho_{KK}^{\Delta T}} \quad (13)$$

Assuming again that $\rho_{AK}^{\Delta\Delta}$ is equal to 1, and $\rho_{AV}^{\Delta T}$ and $\rho_{KK}^{\Delta T}$ being known from equations 7 and 11, ρ_{KV}^{TT} can be calculated; if some arterial Δ does not enter the kidney cellular pool ($\rho_{AK}^{\Delta\Delta} < 1$) the ρ_{KV}^{TT} value will be underestimated. In a similar manner:

$$\rho_{KV}^{\Delta\Delta} = \frac{\rho_{AV}^{T\Delta}}{\rho_{AK}^{TT} \times \rho_{KK}^{T\Delta}} \quad (14)$$

2. Blood transfer constants were calculated according to the formulas derived by Gurgide et al. (26) they were:

$$\rho_{BB}^{T\Delta} = \frac{(3H/14C) \text{ infused}}{(3H/14C)_{\Delta_A}}$$

$$\rho_{BB}^{\Delta T} = \frac{(3H/14C)_{T_A}}{(3H/14C) \text{ infused}}$$

ACKNOWLEDGMENTS

The assistance given by Dr. Serge Carriere of Maisonneuve Hospital for the surgical preparation of the dogs is gratefully acknowledged. The author is also indebted to Dr. Erlio Gurgide of Columbia University for his assistance with the mathematical portion of this paper. Thanks are also due to Miss Lise Heroux for technical assistance and to Miss Micheline Caron for secretarial assistance.

This work was supported by Grant MA-1859 from the Medical Research Council of Canada.

REFERENCES

1. Vande Wiele, R. L., P. C. MacDonald, E. Gurgide, and S. Lieberman. 1963. Studies on the secretion and interconversion of the androgens. *Recent Progr. Hormone Res.* **19**: 275.
2. Horton, R., J. Shinsako, and P. H. Forsham. 1965. Testosterone production and metabolic clearance rates with volumes of distribution in normal adult men and women. *Acta Endocrinol.* **48**: 446.
3. Tait, J. F., and S. Burstein. 1964. In vivo studies of steroid dynamics in man. In *The Hormones*. G. Pincus, K. Thimann, and E. B. Astwood, editors. Academic Press Inc., New York. **5**: 441.
4. Tait, J. F., and R. Horton. 1964. Some theoretical considerations on the significance of the discrepancy in urinary and blood production rate estimates of steroid hormones. *Steroids*. **14**: 365.
5. Horton, R., and J. F. Tait. 1966. Androstenedione production and interconversion rates measured in peripheral blood and studies on the possible site of its conversion to testosterone. *J. Clin. Invest.* **45**: 301.
6. Korenman, S. G., and M. B. Lipsett. 1964. Is testosterone glucuronide uniquely derived from plasma testosterone? *J. Clin. Invest.* **43**: 2125.
7. Southren, A. L., G. G. Gordon, S. Tochimoto, G. Pinzon, D. R. Lane, and W. Stypulkowski. 1967. Mean plasma concentration, metabolic clearance and basal plasma production rates of testosterone in normal young men and women using a constant infusion procedure: effect of time of day and plasma concentration on the metabolic clearance rate of testosterone. *J. Clin. Endocrinol. Metab.* **27**: 686.
8. Bardin, C. W., and M. B. Lipsett. 1967. Testosterone and androstenedione blood production rates in normal women and women with idiopathic hirsutism or polycystic ovaries. *J. Clin. Invest.* **46**: 891.
9. Rivarola, M. A., R. T. Singleton, and C. J. Migeon. 1967. Splanchnic extraction and interconversion of testosterone and androstenedione in man. *J. Clin. Invest.* **46**: 2095.
10. Horton, R., and J. F. Tait. 1967. In vivo conversion of dehydroisoandrosterone to plasma androstenedione and testosterone in man. *J. Clin. Endocrinol. Metab.* **27**: 79.
11. Thomas, P. Z., and R. I. Dorfman. 1964. Metabolism in vitro of androst-4-ene-3,17-dione-4-¹⁴C by rabbit skeletal muscle strips. *J. Biol. Chem.* **239**: 762.
12. Wotiz, H. H., H. Mescon, H. Doppel, and H. M. Lemon. 1956. The in vitro metabolism of testosterone by human skin. *J. Invest. Dermatol.* **26**: 113.
13. Rongone, E. L. 1966. Testosterone metabolism by human male mammary skin. *Steroids*. **7**: 489.
14. Acevedo, H. F., and J. W. Goldzieher. 1964. The metabolism of 4-¹⁴C-4-androstene-3,17-dione by normal human prostatic tissue. *Biochim. Biophys. Acta*. **82**: 118.
15. West, C. D., and L. T. Samuels. 1951. The metabolism of testosterone and related steroids by kidney tissue. *J. Biol. Chem.* **190**: 827.
16. Kochakian, C. D., V. S. Raut, and G. Kinson. 1961. Use of coenzymes I and II in interconversion of testosterone and androstenedione. *Amer. J. Physiol.* **200**: 348.
17. Kreek, M. J., F. G. Guggenheim, J. E. Ross, and D. F. Tapley. 1963. Glucuronide formation in the transport of testosterone and androstenedione by rat intestine. *Biochim. Biophys. Acta*. **74**: 418.
18. Levin, J., E. H. Friedrich, and J. Lobotsky. 1965. Steroid adsorption with polyethylene tubing. *J. Clin. Endocrinol. Metab.* **25**: 1519.
19. Herd, J. A., and A. C. Barger. 1964. Simplified technique for chronic catheterization of blood vessels. *J. Appl. Physiol.* **19**: 791.
20. Blaquier, J., E. Forchielli, and R. I. Dorfman. 1967. In vitro metabolism of androgens in whole human blood. *Acta Endocrinol.* **55**: 697.
21. Nowaczynski, W., and J. Genest. 1955. Recovery of aldosterone, hydrocortisone and cortisone following dialysis. *J. Clin. Endocrinol. Metab.* **15**: 1310.
22. Simpson, S. A. von, J. F. Tait, A. Wettstein, R. Neher, J. von Euw, O. Schindler, and T. Reichstein. 1954. Die konstitution des aldosterons über bestandteile der Nebennierenrinde und verwandte stoffe. *Helv. Chim. Acta*. **37**: 1200.

23. Poos, G. I., G. E. Arth, R. E. Beyler, and C. H. Sarett. 1953. Approaches to the total synthesis of adrenal steroids. V. 4 β -Methyl-7-ethylene-dioxy-1,2,3,4,4A α ,4b,5,6,7,8, 10,10 α - β -dodecahydrophenanthrene-4 β -ol-1-one and related tricyclic derivatives. *J. Amer. Chem. Soc.* **75**: 422.
24. Sandor, T., and A. Lanthier. 1960. Biosynthesis of two steroid glycols from 17 α -hydroxyprogesterone by surviving human ovarian slices. *Can. J. Biochem. Physiol.* **38**: 1167.
25. Tait, J. F. 1963. Review: the use of isotopic steroids for the measurement of production rates in vivo. *J. Clin. Endocrinol. Metab.* **23**: 1285.
26. Gurpide, E., J. Mann, and S. Lieberman. 1963. Analysis of open systems of multiple pools by administration of tracers at a constant rate or as a single dose as illustrated by problems involving steroid hormones. *J. Clin. Endocrinol. Metab.* **23**: 1155.
27. Green, H. D., L. S. Hall, J. Sexton, and C. P. Deal. 1959. Autonomic vasomotor responses in the canine hepatic arterial and venous beds. *Amer. J. Physiol.* **196**: 196.
28. Longcope, C., D. S. Layne, and J. F. Tait. 1968. Metabolic clearance rates and interconversions of estrone and 17 β -estradiol in normal males and females. *J. Clin. Invest.* **47**: 93.
29. Grim, E. 1963. The flow of blood in the mesenteric vessels in circulation. In *Handbook of Physiology*. W. F. Hamilton and P. Dow, editors. The American Physiological Society, Washington, D. C. **2**: 1439.
30. Hollenberg, M., and J. Dougherty. 1966. Liver blood flow measured by portal venous and hepatic arterial routes with Kr⁸⁶. *Amer. J. Physiol.* **210**: 926.
31. Birth, A. G., B. H. Casey, and R. M. Zakheim. 1967. Hepatic blood flow measured by the Krypton-85 clearance technique. *Surgery* **62**: 174.
32. Blalock, A., and M. F. Mason. 1936. Observations on the blood flow and gaseous metabolism of the liver of unanesthetized dogs. *Amer. J. Physiol.* **117**: 328.
33. Stewart, J. D., J. G. Stephens, M. B. Leslie, B. A. Portin, and W. G. Schenkje. 1958. Portal hemodynamics under varying experimental conditions. *Ann. Surg.* **147**: 868.
34. Selkurt, E. E. 1963. The renal circulations in Circulation. In *Handbook of Physiology*. W. F. Hamilton and P. Dow, editors. The American Physiological Society, Washington, D. C. **2**: 1457.
35. Cohn, G. L., M. Hume, and P. K. Bondy. 1960. The in vivo glucuronide conjugation of 17-ketosteroids by dog kidney. *Acta Endocrinol. Suppl.* **51**: 729.