

## Cerebrovascular response to acute hypocapnic and eucapnic hypoxia in normal man

William Shapiro, ... , James P. Baker, John L. Patterson Jr.

*J Clin Invest.* 1970;49(12):2362-2368. <https://doi.org/10.1172/JCI106455>.

### Research Article

Alterations in human cerebral blood flow and related blood constituents were studied during exposure to acute hypoxia. Observations were made during serial inhalation of decreasing O<sub>2</sub> concentrations with and without maintenance of normocarbida, during 8 min inhalation of 10% O<sub>2</sub>, and after hyperventilation at an arterial P<sub>O<sub>2</sub></sub> of about 40 mm Hg. In the range of hypoxemia studied, from normal down to arterial P<sub>O<sub>2</sub></sub> of about 40 mm Hg, the magnitude of the cerebral vasodilator response to hypoxia appeared to be largely dependent upon the coexisting arterial CO<sub>2</sub> tension. The mean slope of the increase in cerebral blood flow with decreasing arterial O<sub>2</sub> tension rose more quickly ( $P < 0.05$ ) when eucapnia was maintained when compared with the slope derived under similar hypoxic conditions without maintenance of eucapnia. When 12 subjects inhaled 10% oxygen, cerebral blood flow rose to more than 135% of control in four whose mean decrease in arterial CO<sub>2</sub> tension was - 2.0 mm Hg. The remaining eight had flows ranging from 97 to 120% of control, and their mean decrease in CO<sub>2</sub> tension was - 5.1 mm Hg. When mean arterial P<sub>O<sub>2</sub></sub> was 37 mm Hg, hyperventilation was carried out in 10 subjects. Arterial P<sub>O<sub>2</sub></sub> increased insignificantly, arterial P<sub>CO<sub>2</sub></sub> declined from 34 to 27 mm Hg ( $P < 0.05$ ), and cerebral blood flow which had been [...]

Find the latest version:

<https://jci.me/106455/pdf>



# Cerebrovascular Response to Acute Hypocapnic and Eucapnic Hypoxia in Normal Man

WILLIAM SHAPIRO, ALBERT J. WASSERMAN, JAMES P. BAKER, and JOHN L. PATTERSON, JR.

*From the Cardiopulmonary Laboratory, Department of Medicine, Medical College of Virginia, Health Sciences Division of Virginia Commonwealth University, Richmond, Virginia 23219, and from the Department of Medicine, The University of Texas Southwestern Medical School and Cardiovascular Section, Medical Service, Veterans Administration Hospital, Dallas, Texas 75216*

**ABSTRACT** Alterations in human cerebral blood flow and related blood constituents were studied during exposure to acute hypoxia. Observations were made during serial inhalation of decreasing  $O_2$  concentrations with and without maintenance of normocarbia, during 8 min inhalation of 10%  $O_2$ , and after hyperventilation at an arterial  $P_{O_2}$  of about 40 mm Hg. In the range of hypoxemia studied, from normal down to arterial  $P_{O_2}$  of about 40 mm Hg, the magnitude of the cerebral vasodilator response to hypoxia appeared to be largely dependent upon the coexisting arterial  $CO_2$  tension. The mean slope of the increase in cerebral blood flow with decreasing arterial  $O_2$  tension rose more quickly ( $P < 0.05$ ) when eucapnia was maintained when compared with the slope derived under similar hypoxic conditions without maintenance of eucapnia. When 12 subjects inhaled 10% oxygen, cerebral blood flow rose to more than 135% of control in four whose mean decrease in arterial  $CO_2$  tension was  $-2.0$  mm Hg. The remaining eight had flows ranging from 97 to 120% of control, and their mean decrease in  $CO_2$  tension was  $-5.1$  mm Hg. When mean arterial  $P_{O_2}$  was 37 mm Hg, hyperventilation was carried out in 10 subjects. Arterial  $P_{O_2}$  increased insignificantly, arterial  $P_{CO_2}$  declined from 34 to 27 mm Hg ( $P < 0.05$ ), and cerebral blood flow which had been 143% of control decreased to 109%, a figure not significantly different from control.

These data demonstrate the powerful counterbalancing constrictor effects of modest reductions in  $CO_2$  tension on the vasodilator influence of hypoxia represented by

arterial  $P_{O_2}$  reductions to about 40 mm Hg. Indeed, mild hyperventilation completely overcame the vasodilator effect provided by an arterial  $O_2$  tension as low as 40 mm Hg. The effects of hypoxia on the control of the cerebral circulation must be analyzed in terms of the effects of any associated changes in  $CO_2$  tension.

## INTRODUCTION

Lowering the arterial or cerebral tissue  $P_{O_2}$  in normal conscious subjects results in cerebral vasodilatation, increased cerebral blood flow (CBF) (1), and, in humans breathing at least 7% oxygen, no change in the cerebral rate of  $O_2$  utilization (2). The magnitude of the increases in cerebral blood flow after moderately severe hypoxia appear to be less than the response to a comparable amount of hypercapnia (3). Simultaneously induced hypoxia and hypercapnia result in additive rather than synergistic effects on cerebral blood flow (4).

There is little information available concerning the separate and possibly antagonistic effects on the cerebral vasculature of the hypocapnia frequently associated with exposure to acute hypoxia. Although cerebral vasodilatation was reportedly enhanced during acute hypoxia with normocarbia, the reduction of arterial  $P_{O_2}$  was marked, and control observations without added  $CO_2$  were omitted (2). Hyperventilation has been considered to attenuate the hypoxic response to high altitude, but cerebrovascular data before 6 hr exposure to altitude are unavailable (5).

The purpose of the present study was to provide a description of the alterations in cerebral blood flow and related blood constituents immediately following achievement of the steady state after induction of various de-

Preliminary reports on this work appeared in abstract form in 1964 *Circulation* 30: III-175, and 1965 *Clin. Res.* 13: 76, and 1966 14: 38.

Received for publication 1 April 1970.

**TABLE I**  
*Effects of Graded Hypoxia on Cerebral Blood Flow (CBF), Cerebrovascular Resistance (CVR), Mean Arterial Pressure (MABP), Arterial and Jugular Venous O<sub>2</sub> and CO<sub>2</sub> Tensions, and Cerebral O<sub>2</sub> Delivery\**

| Inspired gas                | CBF       | CVR       | MABP    | Pao <sub>2</sub> | Pvo <sub>2</sub> | Paco <sub>2</sub> | Pvco <sub>2</sub> | O <sub>2</sub> delivery |
|-----------------------------|-----------|-----------|---------|------------------|------------------|-------------------|-------------------|-------------------------|
|                             | % control | % control | mm Hg   | mm Hg            | mm Hg            | mm Hg             | mm Hg             | % control               |
| Air                         | 100       | 100       | 98 ±4.3 | 91 ±3.1          | 36 ±1.4          | 38 ±1.7           | 46 ±2.3           | 100                     |
| 18% Oxygen                  | 99 ±4.7   | 99 ±5.3   | 97 ±2.1 | 74 ±1.7          | 33 ±1.2          | 37 ±2.0           | 46 ±2.3           | 98 ±4.7                 |
| 16% Oxygen                  | 105 ±2.8  | 92 ±3.0‡  | 96 ±4.3 | 64 ±1.9          | 32 ±1.1          | 36 ±1.9           | 45 ±2.1           | 101 ±2.5                |
| 14% Oxygen                  | 105 ±3.5  | 92 ±1.4§  | 96 ±5.1 | 55 ±1.5          | 30 ±1.1          | 36 ±1.9           | 44 ±2.0           | 98 ±2.7                 |
| 12% Oxygen                  | 115 ±4.5  | 87 ±3.4   | 98 ±5.3 | 47 ±0.9          | 29 ±1.2          | 35 ±1.8           | 43 ±2.1           | 100 ±3.3                |
| 10% Oxygen                  | 135 ±9.7  | 72 ±4.3§  | 94 ±4.8 | 40 ±1.3          | 27 ±2.0          | 33 ±1.8           | 40 ±2.2           | 107 ±5.7                |
| 10% Oxygen +CO <sub>2</sub> | 143 ±8.4§ | 68 ±3.0§  | 95 ±5.0 | 41 ±2.4          | 29 ±2.0          | 36 ±2.0           | 43 ±2.5           | 115 ±5.1§               |

\* Values are mean ± standard error of the mean for six subjects. Symbols represent results of paired *t* test analysis.

‡ 0.05 > *P* > 0.02.

§ *P* < 0.01.

|| 0.02 > *P* > 0.01.

**TABLE IA**  
*Relationship of Statistical Similarities and Differences between the Data in Table I according to Duncan's Multiple Range Test (11)*

Mean values represented by and connected by the lines are not significantly dissimilar. Mean values on separated lines are different at the 5% level. Overlapping lines are caused by means which represent intermediate values not statistically different from other means which are separated by *P* < 0.05.\*

| % Oxygen                | 21    | 18    | 16    | 14    | 12    | 10    | 10 + CO <sub>2</sub> |
|-------------------------|-------|-------|-------|-------|-------|-------|----------------------|
| CBF and CVR             | _____ |       | _____ |       |       | _____ |                      |
| Pao <sub>2</sub>        | _____ | _____ | _____ | _____ | _____ | _____ | _____                |
| Pvo <sub>2</sub>        | _____ | _____ | _____ | _____ | _____ | _____ | _____                |
| Paco <sub>2</sub>       | _____ | _____ | _____ | _____ | _____ | _____ | _____                |
| Pvco <sub>2</sub>       | _____ | _____ | _____ | _____ | _____ | _____ | _____                |
| O <sub>2</sub> delivery | _____ |       |       | _____ |       | _____ | _____                |

\* Example: The values for O<sub>2</sub> delivery obtained at the 18, 16, 14, 12, and 10% levels were not significantly different. The value at 10% was sufficiently greater than the others so that it was not significantly different from the 10% + CO<sub>2</sub> value. This latter mean value, however, was *P* < 0.05 greater than the 18, 16, 14, and 12% values.

TABLE II  
*Effects of Controlled and Uncontrolled Arterial CO<sub>2</sub> Tension and Hyperventilation during Hypoxia on Cerebral Blood Flow, Cerebrovascular Resistance, Mean Arterial Pressure, and Arterial and Jugular Venous O<sub>2</sub> and CO<sub>2</sub> Tensions\**

| Inspired gas                        | CBF         | CVR        | MABP     | PaO <sub>2</sub> | PvO <sub>2</sub> | Paco <sub>2</sub> | Pvco <sub>2</sub> |
|-------------------------------------|-------------|------------|----------|------------------|------------------|-------------------|-------------------|
|                                     | % control   | % control  | mm Hg    | mm Hg            | mm Hg            | mm Hg             | mm Hg             |
| Air                                 | 100         | 100        | 97 ± 2.4 | 95 ± 1.8         | 37 ± 1.2         | 38 ± 0.7          | 47 ± 0.7          |
| 18% Oxygen<br>+ CO <sub>2</sub>     | 105 ± 1.8 † | 95 ± 1.6 § | 96 ± 2.1 | 82 ± 1.7         | 36 ± 1.2         | 38 ± 0.8          | 48 ± 0.8          |
| 14% Oxygen<br>+ CO <sub>2</sub>     | 107 ± 1.3   | 94 ± 1.4   | 98 ± 2.7 | 68 ± 2.3         | 35 ± 1.2         | 39 ± 0.8          | 37 ± 0.8          |
| 9.5% Oxygen<br>+ CO <sub>2</sub>    | 128 ± 4.8   | 81 ± 4.0   | 99 ± 3.0 | 49 ± 1.5         | 32 ± 1.5         | 39 ± 0.8          | 47 ± 1.2          |
| 10.4% Oxygen,<br>no CO <sub>2</sub> | 143 ± 11.4  | 71 ± 6.4   | 92 ± 3.4 | 37 ± 1.0         | 26 ± 1.3         | 34 ± 0.7          | 40 ± 0.8          |
| 9.5% Oxygen<br>hypervent.           | 109 ± 7.5   | 93 ± 5.9   | 95 ± 2.9 | 42 ± 2.2         | 25 ± 1.5         | 27 ± 1.4          | 35 ± 1.1          |

\* Values are mean ± standard error of the mean for 10 subjects.

† 0.05 > P > 0.02. Results of paired *t* test analysis.

§ 0.02 > P > 0.01. Results of paired *t* test analysis.

|| P < 0.01. Results of paired *t* test analysis.

grees of acute hypoxia. By this means, attempts were made to define the threshold of the cerebrovascular response to hypoxia and to expose the magnitude of the potentially counterbalancing effects of the hypocapnia often associated with significant hypoxia.

#### METHODS

The subjects of 28 studies were 13 healthy males, aged 25-40 yr (average age 32) in the postabsorptive resting

state. The planned procedures were discussed fully, and all gave their informed consent. After local procaine infiltration, cannulae were placed in the brachial artery and in the internal jugular bulb. Oxygen saturations were determined by the Nahas spectrophotometric method (6), and blood O<sub>2</sub> and CO<sub>2</sub> tensions and pH were determined with an electrode system (7).<sup>1</sup> Methods for expired CO<sub>2</sub> monitoring,

<sup>1</sup> Epsco Blood Parameter Analyzer, Epsco, Inc., Westwood, Mass.

TABLE IIA  
*Relationship of Statistical Similarities and Differences between the Data in Table II According to Duncan's Multiple Range Test*

Mean values represented by and connected by the lines are not statistically dissimilar. Mean values on separated lines are different at the 5% level. Overlapping lines are caused by means which represent intermediate values not statistically different from other means which are separated by P < 0.05.\*

| % Oxygen                                | 21    | 18 + CO <sub>2</sub> | 14 + CO <sub>2</sub> | 9.5 + CO <sub>2</sub> | 10.4  | 9.5 + Hypervent. |
|---|-------|----------------------|----------------------|-----------------------|-------|------------------|
| CBF and CVR                             | _____ |                      | _____                |                       | _____ |                  |
| PaO <sub>2</sub>                        | _____ | _____                | _____                | _____                 | _____ | _____            |
| PvO <sub>2</sub>                        | _____ | _____                | _____                | _____                 | _____ | _____            |
| Paco <sub>2</sub> and Pvco <sub>2</sub> | _____ | _____                | _____                | _____                 | _____ | _____            |

\* See footnote on Table IA.

delivery of gas mixtures, and recording have been described in detail (8-10).

**Graded hypoxemia.** In six studies arterial and jugular venous blood samples were obtained after serial 5-min inhalations of air, 18, 16, 14, 12, and 10% O<sub>2</sub>, and 10% O<sub>2</sub> with CO<sub>2</sub> added to the inspired gas line to restore the end-tidal CO<sub>2</sub> concentration as close as possible to the control level.

**8 min of 10% oxygen inhalation.** In 12 studies, arterial and internal jugular venous blood samples were obtained during the control state and at 1 min intervals during an 8 min period of 10% inhalation.

**Eucapnic hypoxia.** In 10 studies, arterial and jugular venous blood was obtained after the subjects breathed the following for serial 10-min periods: air, 18, 14, 9.5, 10.4, and 9.5% O<sub>2</sub>. End-tidal CO<sub>2</sub> content was maintained at the control level by adding CO<sub>2</sub> to the inspired gas line during the inhalation of 18, 14, and the first 9.5% O<sub>2</sub> periods. No CO<sub>2</sub> was added while breathing 10.4% O<sub>2</sub>, and the subjects hyperventilated during the second period of 9.5% O<sub>2</sub> inhalation.

Since inspired O<sub>2</sub> concentrations as low as 7% do not effect the rate of cerebral oxygen consumption (2), it was valid to apply the 1/(A-V)O<sub>2</sub> method for estimation of cerebral blood flow. The validity of this method and the necessary calculations have been discussed in detail (5, 8-10).

The data (Tables I and II) were subjected to an analysis of variance, and where significant differences ( $P < 0.05$ ) were found, Duncan's multiple range test was applied to discover which treatments differed by  $P < 0.05$  (Tables IA and IIA) (11). Since one of the critical assumptions for analysis of variance is the equality of variances within treatments, and since there was by definition no variability in the control values for cerebral blood flow, cerebrovascular resistance, and cerebral oxygen delivery, paired *t* tests were also applied to these data.

The technics for regression analysis and for testing the two regression lines derived for CBF vs. PaO<sub>2</sub> followed the methods of Ostle (11), and the multiple regression analyses followed the methods of Draper and Smith (12).




FIGURE 1 The mean alterations of cerebral blood flow (CBF), arterial O<sub>2</sub> and CO<sub>2</sub> tensions during serial reductions in inspired oxygen concentration. See text for discussion.




FIGURE 2 Mean slope and 95% confidence limits of cerebral blood flow response during serial reductions in arterial O<sub>2</sub> tension without maintenance of arterial CO<sub>2</sub> tension at control level.

## RESULTS

**Graded hypoxemia.** Table I and Fig. 1 present the data obtained following 5-min serial inhalations of decreasing concentrations of inspired O<sub>2</sub> without added CO<sub>2</sub>. Significant increases in cerebral blood flow followed 12 and 10% O<sub>2</sub> inhalation, although cerebrovascular resistance was reduced after less marked reductions in inspired O<sub>2</sub>. Addition of CO<sub>2</sub> when 10% O<sub>2</sub> was inspired resulted in a further rise in flow and reduction in vascular resistance. Table I A demonstrates the significant changes wrought by the entire series of interventions. It was apparent that estimated total O<sub>2</sub> delivery to the brain was significantly increased during inhalations of 10% O<sub>2</sub> and 10% O<sub>2</sub> + added CO<sub>2</sub>.

The decreases in arterial O<sub>2</sub> and CO<sub>2</sub> tensions are seen in Tables I and IA and in Fig. 1. The jugular venous O<sub>2</sub> tensions decreased less than did the arterial Po<sub>2</sub>. Changes in pH of the arterial and venous blood paralleled the changes in CO<sub>2</sub> tension and are not tabulated.

Fig. 2 shows the relationship between cerebral blood flow and arterial O<sub>2</sub> tension when hypocapnia was not corrected as derived from the regression equation, CBF (as per cent of control) = 274 - 40.1 log<sub>e</sub> PaO<sub>2</sub>. The mean curve and the 95% confidence limits are shown.

**8 min of 10% oxygen inhalation.** The individual results are shown in Table III in order of decreasing maximum change in cerebral blood flow. The time to maximum cerebral blood flow was estimated to be approximately 6-7 min.

TABLE III  
Maximum Alterations in Cerebral Blood Flow and Associated Arterial O<sub>2</sub> and CO<sub>2</sub> Tensions in 12 Subjects Breathing 10% O<sub>2</sub> for 8 Min without Added CO<sub>2</sub>

| Maximum CBF | Δ PaCO <sub>2</sub> at maximum CBF | PaO <sub>2</sub> at maximum CBF | Time to maximum CBF |
|-------------|------------------------------------|---------------------------------|---------------------|
| % control   | mm Hg                              | mm Hg                           | min                 |
| 158         | -4.3                               | 39                              | 8+*                 |
| 147         | -1.8                               | 37                              | 7                   |
| 140         | +0.3                               | 46                              | 7                   |
| 135         | -2.2                               | 40                              | 5                   |
| 120         | -7.4                               | 50                              | 5                   |
| 120         | -4.2                               | 38                              | 6                   |
| 118         | -4.9                               | 45                              | 7                   |
| 108         | -4.7                               | 42                              | 8+*                 |
| 100         | -1.9                               | 52                              | 7                   |
| 98          | -9.1                               | 44                              | 8+*                 |
| 97          | -3.7                               | 50                              | 5                   |
| 97          | -4.7                               | 40                              | 6                   |

\* Plateau in response of cerebral blood flow not achieved during observation period.

Multiple regression analysis revealed that time and the arterial CO<sub>2</sub> tension were statistically significant correlates of cerebral blood flow during this period of 10% O<sub>2</sub> inhalation, CBF = 51.8 + 1.9 × time (min) + 1.56 PaCO<sub>2</sub> - 0.19 PaO<sub>2</sub>.

Fig. 3 presents mean values for the four studies with the largest rises in cerebral blood flow (135% or more of the control). An increase in cerebral blood flow was evident after 2-3 min of 10% O<sub>2</sub> inhalation, and CBF




FIGURE 3 Mean cerebral blood flow and arterial CO<sub>2</sub> tensions during 8 min of 10% O<sub>2</sub> inhalation in four subjects with largest increases in blood flow.




FIGURE 4 Mean cerebral blood flow and arterial CO<sub>2</sub> tensions during 8 min of 10% O<sub>2</sub> inhalation in eight subjects with maximal flow increases of 120% or less of control.

increased further thereafter. The mean decrease in arterial PCO<sub>2</sub> was 2.0 mm Hg. In the remaining eight subjects, the mean maximum cerebral blood flow was 7% above control, and the mean decrease in arterial PCO<sub>2</sub> at the time of peak cerebral blood flow was 5.1 mm Hg (Fig. 4). The differences between the arterial O<sub>2</sub> tensions for these groups at the times of peak cerebral blood flow (40 vs. 45 mm Hg) as well as their lowest arterial O<sub>2</sub> tensions (40 vs. 43 mm Hg) were not significant, but the difference between the mean decreases in CO<sub>2</sub> tensions was significant (Δ -2.0 vs. Δ -5.1 mm Hg, P < 0.05). Alterations in mean arterial blood pressure were insignificant in both groups.

*Eucapnic hypoxemia.* Tables II and IIA and Fig. 5 show that when eucapnia was maintained, significant increases in cerebral blood flow occurred after slight reductions in inspired O<sub>2</sub> concentration. During inhalation of 10.4% O<sub>2</sub> without added CO<sub>2</sub> the average arterial Po<sub>2</sub> was less than that during the preceding period when 9.5% O<sub>2</sub> with added CO<sub>2</sub> was inspired (37 vs. 49 mm Hg, respectively), and arterial PCO<sub>2</sub> declined. These simultaneous decreases in both arterial Po<sub>2</sub> and PCO<sub>2</sub> were associated with rises in cerebral blood flow in six subjects and decreases in four subjects. Mean cerebral blood flow did not change materially despite more intense hypoxia in the presence of this degree of acute hypocapnia. After this intervention each subject hyperventilated while breathing 9.5% oxygen. This resulted in significant decrease in arterial PCO<sub>2</sub> and no material changes in Po<sub>2</sub> when both were compared with the preceding intervention (10.4% O<sub>2</sub> without added CO<sub>2</sub>). Hyperventilation under these circumstances was associated with profound reductions in mean cerebral blood flow which became insignificantly different from control.

Fig. 6 shows the relationship of cerebral blood flow and arterial oxygen tension during eucapnic hypoxemia as derived from the regression equation, CBF = 274 - 38.6 log<sub>e</sub> PaO<sub>2</sub>. The mean curve and 95% confidence limits are shown. The CBF vs. PaO<sub>2</sub> curves during graded hypoxemia (Fig. 2) and during eucapnic hypoxemia (Fig. 6) were significantly different from one

another, ( $P < 0.05$ ), the latter showing a steeper response of the cerebral blood flow with decreasing arterial  $O_2$  tension.

## DISCUSSION

The important studies of Gibbs, Gibbs, Lennox, and Nims (13) demonstrated that during inhalation of 2 and 4% oxygen, additions of  $CO_2$  to the inspired gas line increased the resulting cerebral blood flow and improved mental function as measured by the electroencephalogram, simple arithmetic tests, and time to unconsciousness. The potential importance of the level of the  $CO_2$  tension in modulating the effects of any level of hypoxia was suggested in studies of cerebral blood flow in man at high altitude but studied only after prolonged exposure to high altitude (5). Acute exposure to hypoxia and the effects of correction of the associated hypocapnia were discussed but were not studied by these authors. Steady-state studies of cerebral blood flow at an even more severe level of hypoxia in the presence of normocarbica (9) appeared to show greater rises in flow with normocarbica than those previously reported when  $CO_2$  tension was uncontrolled (2). The level of hypoxia was not precisely comparable to previous studies, and observations during uncontrolled respiration at the level of hypoxia studied were not made.

The present data showed that during reduction in inspired  $PO_2$  without correction of hypocapnia, the level of the cerebral blood flow was largely dependent upon the response of the respiratory center as reflected by the arterial  $CO_2$  tension level. Those subjects who had the least reduction in arterial  $CO_2$  tension during inhalation of 10%  $O_2$  exhibited the largest increases in cerebral blood flow (Fig. 3). Subjects with greater respiratory sensitivity showed little or no increase in cerebral blood flow despite reductions in arterial  $O_2$  tension to 45–50 mm Hg (Fig. 4). Also, restoration of the




FIGURE 5 The mean effects of eucapnic and hypocapnic hypoxia and of hyperventilation on mean cerebral blood flow and arterial  $O_2$  and  $CO_2$  tensions in 10 subjects. See text for discussion.




FIGURE 6 Mean slope and 95% confidence limits of cerebral blood flow response to eucapnic hypoxia. This slope was steeper ( $P < 0.05$ ) than that derived during hypocapnic hypoxia (Fig. 2).

slightly reduced arterial  $CO_2$  tension to the normal range resulted in increases in cerebral blood flow and decreases in cerebrovascular resistance (Fig. 1). The data uncovered a sensitive cerebrovascular vasodilator response to very mild hypoxemia when eucapnia was maintained (compare Figs. 2 and 6).

Thus, hypocapnia of mild to moderate degree may blunt or abolish the vasodilator stimulus of rather severe hypoxia. We are not aware of a previous demonstration of this phenomenon. Although hypoxia is commonly considered the most important stimulus to vasodilatation (5), it was apparent that, at the levels studied, moderate hypocapnia can overcome the vasodilatation of the hypoxic stimulus. Through the levels of hypoxia and arterial  $PCO_2$  alteration studied, alterations in  $PCO_2$  exert greater effects on the cerebral circulation than comparable alterations of arterial  $PO_2$  (3, 4, 8, 9, 14). Additional evidence of more sensitivity to  $CO_2$  than  $O_2$  tension may be seen in the observed approximate mean time to peak flow response. During  $CO_2$  inhalations the increase in cerebral blood flow achieved a plateau in about  $2\frac{1}{2}$  min (9), whereas the time to peak flow during 10%  $O_2$  inhalation in the present series was approximately 6 min. These observations suggest that respiratory center sensitivity to any induced change in blood gas tension is crucial in determining the rapidity and extent of cerebrovascular response. In studies of the effects of simultaneously applied hypoxia and hypercapnia (10%  $O_2$  + 5%  $CO_2$  inhalation) (4), the respiratory response was marked with rapid changes in

blood gases, and the maximal level of cerebral blood flow occurred sooner than when either stimulus was applied separately.

While not exactly comparable because of the conditions of each study, the present data may be considered to complement certain aspects of the altitude studies of Severinghaus, Chiodi, Eger, Brandstater, and Hornbein (5). Their attempted prediction of the early response to hypoxia may now be modified and amplified with the data provided herein. In the main, their predictions underestimated the likely degree of hyperventilation present and its power to blunt the early vasodilator response to hypoxia.

The mechanism of the vascular changes in acute studies such as those described herein must be mediated through the effects of the acute changes in gas tensions on the respiratory center and the cerebral vasculature rather than by alterations in slowly diffusing substances as has been shown during acclimatization to the chronic hypocapnia and hypoxia at high altitude (5). Whether arterial or cerebral venous gas tensions are best correlated with changes in the cerebrovascular resistance has been the subject of conflicting data (9, 10, 15), but most investigators agree that the sites of ultimate regulatory importance are likely to be found in or near the cerebral vascular cells themselves (5). Any explanation of the effects of acute hypoxia must consider the counterbalancing effects on cerebral blood flow secondary to the associated rapid reductions of arterial  $PCO_2$  as well as other factors which might be pertinent to specific clinical situations.

#### ACKNOWLEDGMENTS

We acknowledge with appreciation the technical assistance of Mrs. Robert T. Dance, Jr., and Miss Amy Cramer, and the statistical assistance of Dr. Walter H. Carter, Jr.

This work was supported by a grant (156-61) from the National Aeronautics and Space Administration and by Public Health Research Grant RR 00016 from the National Institutes of Health. This work was also supported by contract NONR 1134(05) from the Navy.

#### REFERENCES

1. Kety, S. S. 1960. The cerebral circulation. *In* Handbook of Physiology. Neurophysiology. H. W. Magoun, editor. American Physiological Society, Washington, D. C. 3: 1751-1760.
2. Cohen, P. J., S. C. Alexander, T. C. Smith, M. Reivich, and H. Wollman. 1967. Effect of hypoxia and normocarbica on cerebral flow and metabolism in conscious man. *J. Appl. Physiol.* 23: 183.
3. Reivich, M. 1964. Arterial  $PCO_2$  and cerebral hemodynamics. *Amer. J. Physiol.* 206: 25.
4. Shapiro, W., A. J. Wasserman, and J. L. Patterson, Jr. 1966. Human cerebrovascular response to combined hypoxia and hypercapnia. *Circ. Res.* 19: 903.
5. Severinghaus, J. W., H. Chiodi, E. I. Eger II, B. Brandstater, and T. F. Hornbein. 1966. Cerebral blood flow in man at high altitude. *Circ. Res.* 19: 274.
6. Nahas, G. G. 1958. A simplified cuvette for the spectrophotometric measurement of hemoglobin and oxyhemoglobin. *J. Appl. Physiol.* 13: 147.
7. Severinghaus, G. W., and A. F. Bradley. 1958. Electrodes for blood  $PO_2$  and  $PCO_2$  determination. *J. Appl. Physiol.* 13: 515.
8. Wasserman, A. J., and J. L. Patterson, Jr. 1961. The cerebral vascular response to reduction in arterial carbon dioxide tension. *J. Clin. Invest.* 40: 1297.
9. Shapiro, W., A. J. Wasserman, and J. L. Patterson, Jr. 1965. Human cerebrovascular response time to elevation of arterial carbon dioxide tension. *Arch. Neurol. (Chicago)*. 13: 130.
10. Shapiro, W., A. J. Wasserman, and J. L. Patterson, Jr. 1966. Mechanism and pattern of human cerebrovascular regulation after rapid changes in blood  $CO_2$  tension. *J. Clin. Invest.* 45: 913.
11. Ostle, B. 1963. Statistics in Research. Iowa State University Press, Ames, Iowa. 2nd edition. 170-174.
12. Draper, N. R., and H. Smith. 1968. Applied Regression Analysis. John Wiley & Sons, Inc., New York. 243-262.
13. Gibbs, F. A., E. L. Gibbs, W. G. Lennox, and L. F. Nims. 1943. The value of carbon dioxide in counteracting the effects of low oxygen. *J. Aviat. Med.* 14: 250.
14. Kety, S. S., and C. F. Schmidt. 1948. The effects of altered arterial tensions of carbon dioxide and oxygen on cerebral blood flow and cerebral oxygen consumption of normal young men. *J. Clin. Invest.* 27: 484.
15. Severinghaus, J. W., and N. Lassen. 1967. Step hypocapnia to separate arterial from tissue  $PCO_2$  in the regulation of cerebral blood flow *Circ. Res.* 20: 272.