

Thyrotropin Increases Prostaglandin Levels in Isolated Thyroid Cells

S.-C. Yu, ... , L. Chang, G. Burke

J Clin Invest. 1972;51(4):1038-1042. <https://doi.org/10.1172/JCI106864>.

Concise Publication

We have shown that two unrelated prostaglandin antagonists block both thyrotropin (TSH) and prostaglandins E (PGE₁, PGE₂) stimulation of thyroidal adenylyl cyclase activation and cyclic 3',5'-adenosine monophosphate (cAMP) formation, suggesting that prostaglandins play an important role in regulating thyroid function. To further explore this postulate, we measured prostaglandin content by radioimmunoassay in homogeneous bovine thyroid cell preparations in the presence and absence of TSH. Antibodies to albumin-conjugated PGE₁ and PGF_{2α} showed specificity for prostaglandins E and F, respectively, but reacted, albeit far less effectively, with heterologous prostaglandins. A double antibody system was used to separate free from antibody-bound PGE₁-³H and PGF_{2α}-³H. Thyroid cells were extracted with ethanol/ethyl acetate and the various prostaglandins separated on silicic acid columns. Recoveries of added PGE₁-³H and PGF_{2α}-³H through the extraction and separation procedures ranged from 50-80%. The sensitivity of the method was 10-50 pg. Basal thyroid cell content of PGE₁ and PGF_{2α} "equivalents" varied between cell preparations (range = 2-6 ng/0.2 ml cell suspension) but, in each instance, remained constant during 5-30-min incubations at 37°C. TSH, 10-100 mU/ml, increased the levels of cell PGE₁ and PGF_{2α} "equivalents" 30-80% above basal during 5-15-min incubations. The stimulatory effect was specific for TSH, no increase in PGE₁ or PGF_{2α} "equivalent" levels being seen with luteinizing hormone (LH), human growth hormone (HGH), adrenocorticotrophic hormone (ACTH), or glucagon. These [...]

Find the latest version:

<https://jci.me/106864/pdf>



Thyrotropin Increases Prostaglandin Levels in Isolated Thyroid Cells

S.-C. YU, L. CHANG, and G. BURKE

From the Division of Endocrinology and Nuclear Medicine, Department of Medicine, Michael Reese Hospital, and the University of Chicago Pritzker School of Medicine, Chicago, Illinois 60616

ABSTRACT We have shown that two unrelated prostaglandin antagonists block both thyrotropin (TSH) and prostaglandins E (PGE_1 , PGE_2) stimulation of thyroidal adenyl cyclase activation and cyclic 3',5'-adenosine monophosphate (cAMP) formation, suggesting that prostaglandins play an important role in regulating thyroid function. To further explore this postulate, we measured prostaglandin content by radioimmunoassay in homogeneous bovine thyroid cell preparations in the presence and absence of TSH. Antibodies to albumin-conjugated PGE_1 and $\text{PGF}_{2\alpha}$ showed specificity for prostaglandins E and F, respectively, but reacted, albeit far less effectively, with heterologous prostaglandins. A double antibody system was used to separate free from antibody-bound PGE_1 - ^3H and $\text{PGF}_{2\alpha}$ - ^3H . Thyroid cells were extracted with ethanol/ethyl acetate and the various prostaglandins separated on silicic acid columns. Recoveries of added PGE_1 - ^3H and $\text{PGF}_{2\alpha}$ - ^3H through the extraction and separation procedures ranged from 50–80%. The sensitivity of the method was 10–50 pg. Basal thyroid cell content of PGE_1 and $\text{PGF}_{2\alpha}$ "equivalents" varied between cell preparations (range = 2–6 ng/0.2 ml cell suspension) but, in each instance, remained constant during 5-30-min incubations at 37°C. TSH, 10-100 mU/ml, increased the levels of cell PGE_1 and $\text{PGF}_{2\alpha}$ "equivalents" 30-80% above basal during 5-15-min incubations. The stimulatory effect was specific for TSH, no increase in PGE_1 or $\text{PGF}_{2\alpha}$ "equivalent" levels being seen with luteinizing hormone (LH), human growth hormone (HGH), adrenocorticotrophic hormone (ACTH), or glucagon. These data support the thesis that prostaglandins may mediate TSH effects on thyroid.

Received for publication 7 January 1972.

INTRODUCTION

We have recently shown (1,2) that two unrelated prostaglandin (PG)¹ antagonists block both thyrotropin (TSH) and PGE_1 effects on thyroid adenyl cyclase and cyclic 3',5'-adenosine monophosphate (cAMP) formation, suggesting that PGs may play an important role in regulating thyroid function. In an attempt to validate this thesis, we measured PG content in homogeneous thyroid cell preparations in the presence and absence of TSH.

METHODS

Antisera used for PG immunoassay were obtained by immunization of rabbits with PG-protein conjugates. PGE_1 and $\text{PGF}_{2\alpha}$ ² were each conjugated to human serum albumin (HSA) (3). To estimate the number of PG residues incorporated into a molecule of HSA, 4 ng 5,6- PGE_1 - ^3H (88 Ci/mmmole)³ or 0.025 μg 9- $\text{PGF}_{2\alpha}$ - ^3H (14 Ci/mmmole)³ was mixed with 4 mg of the appropriate unlabeled PG before conjugation with HSA. Analysis of the conjugate for tritium content showed that approximately 13 molecules of PGE_1 and 15 molecules of $\text{PGF}_{2\alpha}$, respectively, were coupled to each molecule of HSA.

Dialyzed conjugates were emulsified with equal volumes of complete Freund's adjuvant. Initially the rabbits received 1 mg immunogen subcutaneously (0.25 mg/footpad); this was repeated 1 wk later by subcutaneous injection at multiple sites along the dosal surface. Booster injections were given at 2- and 4-month intervals. The animals were bled periodically and the relative titer of antisera was measured using a double antibody system to separate free from bound PG as described below. Generally the highest titer of anti-

¹ *Abbreviations used in this paper:* ACTH, adrenocorticotrophic hormone; cAMP, cyclic 3',5'-adenosine monophosphate; HGH, human growth hormone; HSA, human serum albumin; LH, luteinizing hormone; PG, prostaglandin; TSH, thyrotropin.

² PGs used in this study were supplied by Dr. John Pike, the Upjohn Co., Kalamazoo, Mich.

³ New England Nuclear Corp., Boston, Mass.

sera was obtained by bleeding 10-14 days after booster injections at bimonthly intervals.

Bovine thyroid cells were isolated by an intermittent trypsinization technique (4). For PG determinations, 0.2 ml cell suspension ($12-20 \times 10^6$ cells/ml) in 5 ml Krebs Ringer bicarbonate buffer, pH 7.4, was incubated in 80×16 mm covered sterile polypropylene tubes with or without added TSH (Thyropar®, Armour Pharmaceutical Co., Kankakee, Ill.) at 37°C for time periods specified below. After incubation the tubes were quickly transferred to Duall #22 homogenizer tubes, and after a 2 min centrifugation (1000 g) at 0°C, the sedimented cells were frozen in dry ice. To the sedimented frozen cells was then added approximately 7000 cpm each PGE_1-^3H and $PGF_{2\alpha}-^3H$ in 1.5 ml H_2O . The cells were homogenized for 15 sec at 23°C with a Potter-Elvehjem homogenizer, 8 ml 100% ethanol was added, and the contents were transferred to a #23 Duall homogenizer tube. The homogenization was repeated. The tubes were centrifuged for 10 min at 1000 g, the ethanol supernate was removed by pipetting, and the cell residue was washed with 4 ml 80% ethanol. The combined ethanol supernatant solutions were allowed to stand overnight at 4°C. They were centrifuged for 10 min at 1000 g to remove residual sediment, and the supernatant was evaporated to about 2.5 ml volume at reduced pressure. Residual volume was uniformly adjusted to 3 ml with H_2O , acidified to pH 3.0 with 0.1 N HCl and extracted three times with half its volume of ethyl acetate (5). The aqueous phase was then discarded; the combined ethyl acetate phases were washed with H_2O to neutral pH and evaporated to dryness under reduced pressure. The efficiency of the extraction procedure was > 90%. The dried residue was dissolved in 3 ml benzene-ethyl acetate (60/40, v/v) immediately before chromatography.




FIGURE 1 Standard curve for PGE_1 and cross-reactions of various prostaglandins with antiserum 2-5 at a dilution of 1/4000. Each value represents the mean of closely agreeing (within 5%) triplicate determinations.




FIGURE 2 Standard curve for $PGF_{2\alpha}$ and cross-reactions of various prostaglandins with antiserum 14-4 at a dilution of 1/2000. Each value represents the mean of closely agreeing (within 5%) triplicate determinations.

Silicic acid chromatography was based on methods previously reported (5). All solvents were spectrograde quality and redistilled before use. Silicic acid (100 mesh, Mallinckrodt Chemical Works, St. Louis, Mo.) was activated at 115°C for 1 hr before use, slurred in benzene-ethyl acetate (9/1, v/v), poured into a 0.7×10 cm glass column, and allowed to settle to a height of 3.3 cm. The cell extract in 3 ml benzene-ethyl acetate (60/40, v/v) was then introduced onto the column; residual cell extract in the tube was washed onto the column with an additional 1 ml benzene-ethyl acetate. "A + E group" PGs were further eluted from the column with an additional 8 ml benzene-ethyl acetate-methanol (60/40/2, v/v). Thereafter, 4 ml of this solvent was passed through the column, the eluate discarded, and the "F group" PGs eluted with 6 ml benzene-ethyl acetate-methanol (60/40/15). The respective "A + E group" (12 ml) and "F group" (6 ml) eluates were evaporated at reduced pressure and dissolved in 800 μ l of 0.005 M phosphate-buffered saline, pH 7.45, for radioimmunoassay. Recoveries of PGE_1-^3H and $PGF_{2\alpha}-^3H$ with this chromatographic technique ranged from 60-80% and 50-70%, respectively.

Radioimmunoassay was performed in 10×56 mm polypropylene tubes; 100 μ l antiserum at a dilution which gave 25-35% binding of labeled PG in the absence of unlabeled PG (1/4000 dilution of anti- PGE_1 , 1/2000 dilution of anti- $PGF_{2\alpha}$) was added to an 100 μ l portion (in triplicate) of PG standard or cell extract. 100 μ l PGE_1-^3H or $PGF_{2\alpha}-^3H$ (approximately 6000-8000 cpm) was added to bring total volume to 300 μ l. The tubes were incubated for 2 hr at 23°C, followed by addition of 100 μ l 5% normal rabbit serum and an excess of goat anti-rabbit gamma globulin (100 μ l, 50%) and overnight incubation at 4°C. The tubes were then centrifuged at 1000 g for 40 min at 4°C, the precipitates dissolved




FIGURE 3 Binding of ^3H -labeled PGs A_1 , E_1 , B_1 , and $\text{F}_{2\alpha}$ by the serum of a rabbit immunized with PGE_1 -HSA (antiserum 2-5). PGA_1 - ^3H (approximately 1.18 pmoles), PGE_1 - ^3H (approximately 1.5 pmoles), PGB_1 - ^3H (approximately 1.5 pmoles), and $\text{PGF}_{2\alpha}$ - ^3H (approximately 0.74 pmoles), each about 7000 cpm per reaction mixture, were incubated with dilutions of antiserum (final dilutions) in a 0.3 ml reaction volume for 1 hr at 37°C . After addition of 5% carrier normal rabbit serum in a 0.1 ml volume, goat anti-rabbit gamma globulin (50%, 0.1 ml) was added and immune precipitation was allowed to proceed overnight at 4°C . Results are means of closely agreeing (within 5%) triplicate determinations.

in 300 μl solubilizer (NCS, Amersham Searle, Des Plaines, Ill.), and counted in toluene scintillation solution.

RESULTS

Radioimmunoassay standard curves showed a significant difference between 10 and 50 pg for PGE_1 (Fig. 1) and

between 50 and 100 pg for $\text{PGF}_{2\alpha}$ (Fig. 2). When 0.2 ml H_2O was carried through the entire procedure, virtually no displacement ($\leq 2\%$) was recorded on the curve for either PGE_1 or $\text{PGF}_{2\alpha}$. Although there was only a 1-2% relative cross-reaction between PGs E and F with the antisera to PGE_1 -HSA and $\text{PGF}_{2\alpha}$ -HSA

TABLE I
Radioimmunoassay of Known Amounts of PGE_1 and $\text{PGF}_{2\alpha}$ Added to Isolated Thyroid Cells

Thyroid cell volume	Prostaglandin added		Level of PGE_1 * measured†			Level of $\text{PGF}_{2\alpha}$ § measured‡		
	PGE_1	$\text{PGF}_{2\alpha}$	Mean \pm SD	No.¶	Recovery	Mean \pm SD	No.¶	Recovery
ml	ng		ng			ng		
0.1	0	0	2.959 ± 0.453	9	—	2.128 ± 0.216	9	—
0.1	1.0	1.0	4.055 ± 0.229	6	102.4	3.196 ± 0.396	6	102.2
0.1	10.0	10.0	12.918 ± 0.679	9	99.7	13.065 ± 0.524	6	107.7

* PGE_1 "equivalents".

† Corrected for extraction and separation recoveries.

§ $\text{PGF}_{2\alpha}$ "equivalents".

¶ No. = number of experimental observations, each experimental determination performed in triplicate.

TABLE II
Radioimmunoassay of PGE₁ and PGF_{2α} in Varying
Volumes of Thyroid Cell Suspension

Cell suspension volume	PGE ₁ "equivalents"		PGF _{2α} "equivalents"	
	Mean ±SD	No.*	Mean ±SD	No.*
ml	ng		ng	
0.05	1.64 ±0.28	8	1.48 ±0.29	9
0.1	3.44 ±0.32	5	2.92 ±0.30	9
0.2	6.06 ±0.47	7	5.70 ±0.56	8

* No. = number of experimental observations, each experimental determination performed in triplicate.

(based on mass of PGF_{1α} or PGF_{2α} required to displace 50% of bound PGE₁-³H [Fig. 1] and mass of PGE₁ or PGE₂ required to displace 50% of bound PGF_{2α}-³H [Fig. 2]), PGE₁ antiserum was unable to effectively distinguish between PGE₁, PGE₂, and PGA₁ (Fig. 1) while PGF_{2α} antiserum recognized both PGF_{1α} and PGF_{2α} (Fig. 2).⁴ Thus, while we are reporting PGE₁ and PGF_{2α} levels, these are in fact PGE₁ and PGF_{2α} "equivalents" (6).

While these studies were in progress, Levine, Gutierrez-Cernosek, and Van Vunakis (7) reported that immunization with PGE₁ produced antibodies directed to PGB₁. We therefore carried out similar studies with our PGE₁ antiserum and PGB₁. PGB₁-³H was obtained by heating PGE₁-³H or PGA₁-³H in 0.02 N NaOH at 100°C for 5 min; Andersen (8) has demonstrated that under these conditions PGE₁ and PGA₁ are quantitatively converted to PGB₁. Fig. 3 illustrates the binding capacity of our PGE₁ antiserum. Of the four labeled antigens used, the greatest binding occurred with PGA₁ and PGE₁. Although cross-reaction with PGB₁-³H was observed, binding was much less effective.

Table I shows the correlation between PG added to 0.1 ml cell suspension and that recovered in the radioimmunoassay. When either 0.05, 0.1, or 0.2 ml thyroid cells was extracted, the levels recorded in the radioimmunoassay were proportional to the volume of cell suspension used (Table II).

During a 15 min incubation at 37°C, TSH effected an increase in thyroid cell levels of both PGE₁ and PGF_{2α} "equivalents" (Fig. 4). Although the increase in PG levels was dose-related, the TSH effect on PGE₁ was biphasic, and both the PGE₁ and PGF_{2α} "TSH dose-response" curves were relatively flat. Thus, the maximum TSH-induced increase in PGE₁ levels was approximately

⁴ It should be noted, however, that presence of a double bond at position 5,6 reduced ability of PGE₂ to cause displacement to one-third that of PGE₁ (Fig. 1) and that loss of a double bond at position 5,6 reduced ability of PGF_{1α} to cause displacement to 70% that of PGF_{2α} (Fig. 2).




FIGURE 4 Effects of TSH on prostaglandin content of isolated bovine thyroid cells during 15-min incubation at 37°C. Results are mean ± SD of 4-9 experimental determinations (number of experimental observations given in parentheses), each experimental determination performed in triplicate.

80% over control and that in PGF_{2α} levels approximately 40% over control. The increases were, however, highly reproducible and specific for TSH (Table III).

The stimulatory effect of TSH on cell PG content was evident within 5 min of incubation at 37°C and increased through 30 min of incubation. Basal levels of PGs did not change during this period. TSH, incubated with thyroid cells at 0°C for periods up to 30 min did not influence cell PG content (Table IV).

TABLE III
Effect of TSH and Other Hormones on Thyroid Cell
Content of Prostaglandins*

Hormone added	PGE ₁ "equivalents"		PGF _{2α} "equivalents"	
	Mean ±SD	No. §	Mean ±SD	No.*
None	4.717 ±0.549	9	3.044 ±0.375	9
TSH, 50 mU/ml	7.918 ±0.688	9	5.652 ±0.519	6
Heat inactivated TSH, 50 mU/ml	4.444 ±0.856	9	3.484 ±0.411	8
LH, 0.5 mg/ml	4.016 ±0.753	6	3.018 ±0.474	5
ACTH, 0.5 mg/ml	3.785 ±0.184	9	2.981 ±0.744	5
HGH, 0.5 mg/ml	3.467 ±0.570	7	3.048 ±0.218	6
Glucagon, 10 ⁻⁶ M	3.909 ±0.769	9	2.970 ±0.218	5

* 0.2 ml thyroid cell suspension, 15 min incubation at 37°C.

† ng/0.2 ml cell suspension.

§ No. = number of experimental observations, each experimental determination performed in triplicate.

|| 100°C for 2 hr prior to addition to thyroid cells.

TABLE IV
Time-Course of TSH-Induced Increase in Thyroid Cell Prostaglandin Content

Incubation time and temperature	PGE ₁ "equivalents"					PGF _{2α} "equivalents"				
	Control		TSH, 50 mU/ml			Control		TSH, 50 mU/ml		
	Mean ±SD	No.‡	Mean ±SD	No.‡	Percent increase	Mean ±SD	No.‡	Mean ±SD	No.‡	Percent increase
	ng*		ng*			ng*		ng*		
5 min, 37°C	2.33 ±0.23	9	2.92 ±0.22	8	25	2.60 ±0.62	6	3.68 ±0.65	8	42
15 min, 37°C	2.28 ±0.27	9	3.35 ±0.60	8	47	2.89 ±0.53	7	4.02 ±0.48	6	39
30 min, 37°C	2.55 ±0.23	9	4.21 ±0.50	9	65	2.40 ±0.40	6	4.33 ±0.91	9	80
30 min, 0°C	2.41 ±0.25	6	2.29 ±0.31	6	—	2.53 ±0.48	5	2.68 ±0.55	5	—

* ng/0.2 ml cell suspension.

‡ No. = number of experimental observations, each experimental determination performed in triplicate.

DISCUSSION

Several laboratories have recently reported the production of antibodies to PGs (3, 6, 9). It has been suggested (3) that the inability of antibody to PGE₁ to distinguish between PGA₁ and PGE₁ may be due to conversion of protein-bound PGE₁ to PGA₁ in vivo or in vitro. Levine et al. (7) noted that methodologic variables might influence the apparent specificity of PG antisera, so that divergent findings with the various PG radioimmunoassays were to be anticipated. The inability of our anti-PGE₁ antiserum to effectively distinguish between PGE₁ and PGE₂ or of our anti-PGF_{2α} antiserum to distinguish between PGF_{2α} and PGF_{1α}, or the ability of our PGE₁ antiserum to recognize PGE₁ but not PGB₁ (contrasting sharply with the findings of Levine et al. (7)) should thus be viewed in this light.

PGs E and F simulate a variety of TSH effects on thyroid and are considerably more potent in this regard than other PGs (10). Demonstration that TSH increases thyroid cell content of these "thyrotropic" PGs supports our earlier contention that PGs may play an important role in mediating TSH effects on thyroid (1, 2, 11). It should be noted in this regard that the thyroid cell preparations used in this study respond functionally (e.g., adenylyl cyclase activation, iodide trapping (1, 2, 11)) in similar fashion to both TSH and PGE₁. Moreover, the TSH-induced increase in cell PGE₁ and PGF_{2α} occurred promptly after addition of TSH, paralleling in this respect the temporal character of TSH- (and PGE₁-) induced activation of thyroid adenylyl cyclase and increase in cAMP (1, 2). (The variation in basal cell PG content is consistent with the considerable variability in other aspects of cell function (12).) Although one cannot, from these findings, determine whether the TSH-induced increase in cell PG content reflects increased synthesis, decreased degradation, or both, this study represents, to our knowledge, the first reported instance of hormone-induced increase in tissue PGs.

ACKNOWLEDGMENTS

This work was supported by U. S. Public Health Service Grant AM-11136.

REFERENCES

- Burke, G., and S. Sato. 1971. Effects of long-acting thyroid stimulator and prostaglandin antagonists on adenylyl cyclase activity in isolated bovine thyroid cells. *Life Sci.* **10**: 969.
- Sato, S., M. Szabo, K. Kowalski, and G. Burke. 1972. Role of prostaglandin in thyrotropin action on thyroid. *Endocrinology.* **90**: 343.
- Jaffe, B. M., J. W. Smith, W. T. Newton, and C. W. Parker. 1971. Radioimmunoassay for prostaglandins. *Science (Washington).* **171**: 494.
- Edmonds, M. W., V. V. Row, and R. Volpé. 1970. Action of globulin and lymphocytes from peripheral blood of patients with Graves' disease on isolated bovine thyroid cells. *J. Clin. Endocrinol.* **31**: 480.
- Anggard, E. 1965. The isolation and determination of prostaglandins in lungs of sheep, guinea pig, monkey and man. *Biochem. Pharmacol.* **14**: 1507.
- Caldwell, B. V., S. Burstein, W. A. Brock, and L. Speroff. 1971. Radioimmunoassay of the F prostaglandins. *J. Clin. Endocrinol.* **33**: 171.
- Levine, L., R. M. Gutierrez-Cernosek, and H. Vanakis. 1971. Specificities of prostaglandins B₁, F_{1α}, and F_{2α} antigen-antibody reactions. *J. Biol. Chem.* **246**: 6782.
- Andersen, N. H. 1969. Dehydration of prostaglandins: study by spectroscopic method. *J. Lipid Res.* **10**: 320.
- Levine, L., and H. Van Vunakis. 1970. Antigenic activity of prostaglandins. *Biochem. Biophys. Res. Commun.* **41**: 1171.
- Field, J., A. Dekker, U. Zor, and T. Kaneko. 1971. *In vitro* effects of prostaglandins on thyroid gland metabolism. *Ann. N. Y. Acad. Sci.* **180**: 278.
- Burke, G., K. Kowalski, and D. Babiarz. 1971. Effects of thyrotropin, prostaglandin E₁, and a prostaglandin antagonist on iodide trapping in isolated thyroid cells. *Life Sci.* **10**: 513.
- Wilson, B., E. Raghupathy, T. Tonoue, and W. Tong. 1968. TSH-like actions of dibutyryl-cAMP on isolated bovine thyroid cells. *Endocrinology.* **83**: 877.