

## Triiodothyronine and Thyroxine in the Serum and Thyroid Glands of Iodine-Deficient Rats

G. M. Abrams, P. R. Larsen

*J Clin Invest.* 1973;52(10):2522-2531. <https://doi.org/10.1172/JCI107443>.

### Research Article

Triiodothyronine ( $T_3$ ) and thyroxine ( $T_4$ ) were measured by immunoassay in the serum and thyroid hydrolysates of control (group A), mildly iodine-deficient (group B), and severely iodine-deficient rats (group C). These results were correlated with changes in thyroïdal weight,  $^{131}\text{I}$  uptake and  $^{127}\text{I}$  content as well as with the distribution of  $^{131}\text{I}$  in Pronase digests of the thyroid. There was a progressive increase in thyroid weight and  $^{131}\text{I}$  uptake at 24 h with decrease in iodine intake. The  $^{127}\text{I}$  content of the thyroids of the group B animals was 44% and that of the group C animals 2% of that in group A. The mean labeled monoiodotyrosine/diiodotyrosine (MIT/DIT) and  $T_3/T_4$  ratios in group A were  $0.42\pm 0.07$  (SD) and  $0.12\pm 0.01$ ,  $0.59\pm 0.06$  and  $0.11\pm 0.03$  in group B, and  $2.0\pm 0.3$  and  $1.8\pm 0.9$  in the group thyroid digests.

Mean serum  $T_4$  concentration in the control rats was  $4.2\pm 0.6$  (SD)  $\mu\text{g } T_4/100 \text{ ml}$ ,  $4.5\pm 0.3 \mu\text{g}/100 \text{ ml}$  in group B animals, and undetectable ( $<0.5 \mu\text{g}/100 \text{ ml}$ ) in group C animals. There was no effect of iodine deficiency on serum  $T_3$  concentrations, which were  $44\pm 9$  (Mean $\pm$ SD)  $\text{ng}/100 \text{ ml}$  in A animals,  $48\pm 6 \text{ ng}/100 \text{ ml}$  in B animals, and  $43\pm 6 \text{ ng}/100 \text{ ml}$  in the C group. Thyroïdal digest  $T_3$  and  $T_4$  concentrations were 39 and 400  $\text{ng}/\text{mg}$  in group A animals and were reduced to 5 and 1% of [...]

Find the latest version:

<https://jci.me/107443/pdf>



# Triiodothyronine and Thyroxine in the Serum and Thyroid Glands of Iodine-Deficient Rats

G. M. ABRAMS and P. R. LARSEN

*From the Division of Endocrinology and Metabolism, Department of Medicine, School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania 15261*

**ABSTRACT** Triiodothyronine ( $T_3$ ) and thyroxine ( $T_4$ ) were measured by immunoassay in the serum and thyroid hydrolysates of control (group A), mildly iodine-deficient (group B), and severely iodine-deficient rats (group C). These results were correlated with changes in thyroidal weight,  $^{131}\text{I}$  uptake and  $^{127}\text{I}$  content as well as with the distribution of  $^{131}\text{I}$  in Pronase digests of the thyroid. There was a progressive increase in thyroid weight and  $^{131}\text{I}$  uptake at 24 h with decrease in iodine intake. The  $^{127}\text{I}$  content of the thyroids of the group B animals was 44% and that of the group C animals 2% of that in group A. The mean labeled monoiodotyrosine/diiodotyrosine (MIT/DIT) and  $T_3/T_4$  ratios in group A were  $0.42 \pm 0.07$  (SD) and  $0.12 \pm 0.01$ ,  $0.59 \pm 0.06$  and  $0.11 \pm 0.03$  in group B, and  $2.0 \pm 0.3$  and  $1.8 \pm 0.9$  in the group thyroid digests.

Mean serum  $T_4$  concentration in the control rats was  $4.2 \pm 0.6$  (SD)  $\mu\text{g } T_4/100 \text{ ml}$ ,  $4.5 \pm 0.3$   $\mu\text{g}/100 \text{ ml}$  in group B animals, and undetectable ( $<0.5 \mu\text{g}/100 \text{ ml}$ ) in group C animals. There was no effect of iodine deficiency on serum  $T_3$  concentrations, which were  $44 \pm 9$  (Mean  $\pm$  SD)  $\text{ng}/100 \text{ ml}$  in A animals,  $48 \pm 6 \text{ ng}/100 \text{ ml}$  in B animals, and  $43 \pm 6 \text{ ng}/100 \text{ ml}$  in the C group. Thyroidal digest  $T_3$  and  $T_4$  concentrations were 39 and 400  $\text{ng}/\text{mg}$  in group A animals and were reduced to 5 and 1% of this, respectively, in group C. The molar ratio of  $T_3/T_4$  in the thyroid digests of the groups A and B animals was identical to the ratio of labeled  $T_3/T_4$  and was slightly less ( $1.0 \pm 0.9$ ) than the labeled  $T_3/T_4$  ratio in the group C animals.

The mean ratio of labeled  $T_4$  to labeled  $T_3$  in the serum of the severely iodine-deficient animals 24 h after isotope injection was  $11 \pm 1$  (SEM). With previously published values, it was possible to correlate the ratio of labeled  $T_4/T_3$  in the thyroid digest with the labeled

$T_4/T_3$  ratio in the serum of each iodine-deficient animal. This analysis suggested that the labeled thyroid hormones in the severely iodine-deficient rat were secreted in the ratio in which they are present in the gland.

Kinetic analysis of total iodothyronine turnover indicated that two-thirds of the  $T_3$  utilized per day by the iodine-sufficient rat arises from  $T_4$ . If the  $T_4$ - $T_3$  conversion ratio remains the same in iodine deficiency, then the analysis suggests that about 90% of the  $T_3$  arises directly from the thyroid. Therefore, it would appear that absolute  $T_3$  secretion by the thyroid increases severalfold during iodine deficiency. The fact that serum  $T_3$  remains constant and  $T_4$  decreases to extremely low levels, combined with previous observations that iodine-deficient animals appear to be euthyroid, is compatible with the hypothesis that  $T_4$  in the normal rat serves primarily as a precursor of  $T_3$ .

## INTRODUCTION

The thyroidal response to iodine deficiency has been an area of active investigation for many years. Previous studies in the rat reviewed by Studer and Greer indicate that when iodine intake is severely restricted there is an increase in thyroid weight, a decrease in protein-bound iodine (PBI)<sup>1</sup> and an altered pattern of tracer iodine distribution in the thyroid gland (1). The last-mentioned changes include an increase in labeled monoiodotyrosine (MIT) and decrease in labeled diiodotyrosine (DIT) as well as a progressive increase in the ratio of labeled triiodothyronine ( $T_3$ ) to labeled thyroxine ( $T_4$ ). Although Studer and Greer, and Greer, Grimm, and Studer have demonstrated that serum PBI decreases to very low levels within 1 mo of initiation of a low-iodine diet, rats maintained on this regimen for even 1 yr re-

P. R. Larsen is a Career Development Awardee of the U. S. Public Health Service, Award #5 KO4 AM 70401.

Received for publication 13 April 1973 and in revised form 4 June 1973.

<sup>1</sup> Abbreviations used in this paper: DIT, diiodotyrosine; MIT, monoiodotyrosine; PBI, protein-bound iodine;  $T_3$ , triiodothyronine;  $T_4$ , thyroxine; TAA, tertiary-amyloalcohol/hexane/ammonia; TSH, thyroid-stimulating hormone.

main healthy and grow at normal rates (2, 3). Recent studies by Silva have shown that  $O_2$  consumption and body temperature regulation of rats under severe iodine restriction for 120 days is not different from that of control animals (4).

The shift of the predominant intrathyroidal-labeled iodothyronine from  $T_4$  to  $T_3$  in response to iodine restriction has led to speculation that the iodine-deficient rat maintains its apparent euthyroid status by preferential synthesis and release of  $T_3$  (3). Since it has not been possible previously to measure serum  $T_3$  directly under these circumstances, this hypothesis has not been substantiated. While this study was in progress, Volpert and Werner presented data suggesting that there was a decrease in the immunoassayable serum  $T_3$  in the iodine-deficient rat though the ratio of  $T_3$  to PBI was increased (5). However, the small group of animals examined necessarily limited the scope of their conclusions. In the following study data are presented that correlate the changes in serum immunoassayable  $T_3$  and  $T_4$  with changes in thyroidal  $T_3$ ,  $T_4$ , and iodine content in mildly and severely iodine-deficient rats.

## METHODS

*Animals and diets.* Sprague-Dawley male rats weighing 150–200 g were maintained on Remington low-iodine test diet (Diet I, Nutritional Biochemicals Corporation, Cleveland, Ohio) for a period of 3 mo. Half of this group received distilled water and the other half distilled water containing 1.3  $\mu\text{g}$  iodide/ml (estimated intake 20–30 ml  $H_2O$ /day). After 3 mo the supplier of the low-iodine diet was changed (Diet II, General Biochemicals Div., Mogul Corp., Chagrin Falls, Ohio) since the anticipated elevation in MIT/DIT ratios was not observed up to this time. The level of iodine supplementation in the control animals remained the same. However, the substitution of a different commercial low-iodine diet after 3 mo of this study made it impossible to examine the longitudinal effects of sustained iodine restriction under constant conditions. Therefore, we have not attempted an analysis of the data as related to duration of iodine deprivation. The diets were analyzed by the Boston Medical Laboratory, and both contained non-detectable amounts of iodine ( $< 0.165 \mu\text{g/g}$  diet [6]). The fact that the thyroidal changes in animals receiving diet II were substantially greater than those in animals receiving diet I is best explained by the assumption that it contained less iodine. However, without verification of this by actual measurement, the theoretical possibility that diet II contains a goitrogen that accelerated the appearance of the changes in the low-iodine animals cannot be entirely discounted.

*Administration of  $^{131}\text{I}$  and calculation of uptake.* 5–40  $\mu\text{Ci}$  of carrier-free  $^{131}\text{I}$  was injected intraperitoneally 24 h before sacrifice. Groups of control and iodine deficient-rats were killed at various intervals after starting the diets and thyroids were dissected from the trachea and weighed. Radioactivity in thyroids and suitable dilutions of the injected dose were measured with an end window Geiger counter under conditions of constant geometry. Blood was obtained from each animal by cardiac puncture.

*Digestion and extraction.* One or two thyroid glands were digested with Pronase (Calbiochem, Los Angeles, Calif.), by the method of Inoue and Taugog (7). This procedure was slightly modified as we have previously described (8). Labeled  $^{125}\text{I}$  and  $T_4$  (Abbott Laboratories, North Chicago, Ill.) were added at the start of digestion to monitor recovery and identify  $^{131}\text{I}$ iodothyronines. The labeled iodothyronines contained less than 3%  $\text{I}^-$  and no other significant contaminants. In addition, after digestion the hydrolysate was extracted twice with 0.4 ml of methanol in 7.4 N ammonia (1:1, vol:vol). Residual  $^{131}\text{I}$  and  $^{125}\text{I}$  in the pellet after this procedure were 6% or less of the total. Analysis of the pellet  $^{127}\text{I}$  in the animals on the high-iodine intake showed similar losses of  $^{127}\text{I}$ . Digestion of extracts was nearly complete as estimated by chromatography of the extracts in collidine—2 N ammonia (3:1, vol:vol). The amount of  $^{127}\text{I}$  and  $^{131}\text{I}$  remaining at the origin was less than 4%. The identification of  $^{131}\text{I}$ -labeled  $\text{I}^-$ ,<sup>\*</sup> MIT\* and DIT\* was carried out as previously described by chromatography in butanol/acetic acid (8). Separation of  $T_4^*$  and  $T_3^*$  was obtained in tertiary-amyl-alcohol hexane/ammonia (TAA) also as previously described (8). Briefly, 25–75  $\mu\text{l}$  of the extract was chromatographed in both systems, and 1–2 cm segments of the strips were subsequently counted for both  $^{125}\text{I}$  and  $^{131}\text{I}$  with suitable correction. In the TAA system, the percent  $T_3^*$  was corrected for the 0.34% artifactual deiodination of  $T_4^*$  during the chromatography as we have previously described (9).  $T_3^*$  and  $T_4^*$  were corrected for losses during the procedure by reference to the original amounts of  $^{125}\text{I}$  and  $T_4$  added. Results were expressed as a fraction of the total  $^{131}\text{I}$  in the extract.

$^{127}\text{I}$  content in 0.1 ml of the extract (about 0.8 ml total volume) was determined by Boston Medical Laboratory (6). This was then corrected for the total extraction volume as well as the small losses occurring during extraction of the Pronase hydrolysate. Results were expressed as ng  $^{127}\text{I}$  per mg wet weight of thyroid.

*Serum and thyroidal  $T_3$  and  $T_4$  content.*  $T_3$  and  $T_4$  content in the thyroid hydrolysates and serum were determined by radioimmunoassays as previously described (10, 11). All assays included samples from paired control and iodine-deficient rats and were performed in duplicate at two dilutions. In analyzing the extract for iodothyronines, suitable dilutions were made in  $\text{MeOH}/\text{NH}_4\text{OH}$  (99:1, vol:vol). 5 or 10  $\mu\text{l}$  of this dilution was then added to tubes containing either  $T_3$ -free or  $T_4$ -free human serum in 1 ml total volume. This concentration of  $\text{MeOH}/\text{NH}_4\text{OH}$  does not affect the assays. In measuring  $T_4$  and  $T_3$  in thyroid hydrolysates from iodine sufficient rats, the  $\text{MeOH}/\text{NH}_4\text{OH}$  dilutions used were 1/100 to 1/500. In both assays, curves parallel to the standard were obtained with increasing quantities of extract. Thus, the duplicate determinations at two dilutions were in excellent agreement. In the extracts of thyroids from the group C animals, dilutions as low as  $\frac{1}{2}$  and  $\frac{1}{3}$  were sometimes necessary for accurate quantitation, indicating that there was no artifactual contribution to either the calculated  $T_3$  or  $T_4$  value from non-iodine-containing substances in the thyroid extracts. Again, excellent agreement was obtained in the estimated iodothyronine content with two different dilutions. There is no significant displacement of either labeled iodothyronine from its antibody by 1000-fold excesses (by weight) of  $\text{I}^-$ , MIT, or DIT. Since there is essentially no cross-reaction of the  $T_3$  antibody with  $T_4$ , prior separation of  $T_3$  and  $T_4$  is

\* An asterisk will be used to denote  $^{131}\text{I}$ -labeled compounds present in thyroid glands or in serum.

not required for this method. Total thyroid  $T_3$  and  $T_4$  were corrected for losses in extraction and expressed as ng  $T_3$  or  $T_4$  per mg wet weight of thyroid.

Analysis of serum  $T_3$  was performed by using a modification of the  $T_3$ -free human serum system that employs sodium salicylate to block  $T_3$ -protein binding (10). Preliminary studies indicated that as much as 100  $\mu$ l of rat serum could be substituted for human serum in the assay without disturbing the parallelism of the rat serum curve with the standard curve or changing the percentage of tracer bound in the absence of antibody. Thus, unknown samples contained either 100 or 50  $\mu$ l unknown rat serum plus 100 or 150  $\mu$ l of  $T_3$ -free human serum added to make the serum concentration 20% in all tubes. Duplicates of the two dilutions were in excellent agreement.  $T_3$ -free rat serum prepared as we have previously described for human serum contained no  $T_3$  (10). Recovery of  $T_3$  from pooled rat sera was  $96 \pm 8\%$  (mean  $\pm$  SD) for 100 ng  $T_3$ /100 ml and  $99 \pm 8\%$  (mean  $\pm$  SD) for 200 ng  $T_3$ /100 ml. Similar recoveries were obtained using serum from iodine-deficient rats in a smaller series of studies. Likewise, the  $T_4$  determinations could be performed in the system containing  $T_4$ -free human serum as long as the quantity of rat serum per tube did not exceed 5  $\mu$ l (total serum per tube, 10  $\mu$ l).

*Distribution of labeled compounds in serum 24 h after  $^{125}I$  injection.* In the severely iodine-deficient animals the

distribution of radioactivity in the serum was analyzed. The value for  $I^*$  was determined as the percent radioactivity migrating from the origin during 90-min electrophoresis of whole serum in glycine-acetate buffer, pH 8.6, at 150 V.  $T_3^*$  and  $T_4^*$  were determined by the method of Sterling, Dellabarba, Newman, and Brenner, as modified in our laboratory (9, 12).  $^{125}I$   $T_3$  and  $T_4$  were added before chromatography to allow correction for losses.  $T_3^*$  was corrected for the artifactual increases due to deiodination of  $T_4^*$  as described above. Results were expressed as a fraction of the total serum radioactivity.

## RESULTS

*Effect of iodine deprivation on thyroidal weight,  $^{127}I$  content,  $^{125}I$  uptake and the distribution of  $^{125}I$  in thyroidal iodoamino acids.* Since previous pulse-labeling studies have shown that the ratio MIT\*/DIT\* is a sensitive index of iodine deficiency (13, 14), animals were grouped on the basis of this parameter. The animals were classified as follows: A, control (MIT\*/DIT\*,  $< 0.5$ ); B, mildly iodine-deficient (MIT\*/DIT\*, 0.51–0.75); and C, severely iodine-deficient (MIT\*/DIT\*,  $> 1.5$ ). All animals that received iodine supplementation had MIT\*/DIT\* ratios less than 0.5, regardless of diet, and were in group A. Group B consisted of animals that had received diet I and no supplement for 4–8 wk. All group C animals had received Diet II for 4 wk (two animals), 8 wk (four animals), or 12 wk (four animals). The body weights of control and iodine-deficient animals studied at each interval were not different.

In Fig. 1 the correlation between the MIT\*/DIT\* and thyroidal iodine content is illustrated. There is a progressive increase in the MIT/DIT ratio as the mean iodine content of the thyroid gland decreases. The mean thyroidal  $^{127}I$  in the thyroids of the various groups is presented in Table I, and for group A was 880 ng/mg; group B, 390 ng/mg; and group C, 21 ng/mg. Thus, the iodine concentration of the thyroids of the group C animals was reduced to about 2% of the iodine present in the A group.

The validity of this classification was further substantiated by the significant differences among the various groups with respect to two other characteristics of the thyroidal response to iodine deficiency. The thyroid weight was moderately increased in the B group, 5.1 mg/100 g body wt as opposed to 4.1 mg in controls ( $P < 0.005$ ). It was increased to 12 mg/100 g body wt in the C group, approximately three times the weight in the control group ( $P < 0.001$ ). The thyroidal uptake of  $^{125}I$  at 24 h increased progressively with iodine deficiency. However, since the maximal uptake of  $^{125}I$  occurs at time intervals earlier than 24 h in the severely iodine-deficient rat, the increase from control values of 15% in group A to 42% in group C ( $P < 0.001$ ) may underestimate the difference in maximal  $^{125}I$  uptake.




FIGURE 1 Relationship between iodine content and the ratio of  $^{125}I$ -labeled MIT to DIT in the pronase hydrolysates of rat thyroid glands. The letters A, B, and C denote the grouping of the animals: A, control; B, mild iodine deficiency; and C, severe iodine deficiency.

TABLE I  
<sup>127</sup>I Content, Gland Weight, and Uptake and Distribution of <sup>131</sup>I in Pronase Hydrolysates of Thyroid Glands  
 from Control and Iodine-Deficient Rats (Mean ± SD)

Group	n	<sup>127</sup> I	Thyroid weight	RAIU‡	I*	MIT*	DIT*	T <sub>3</sub> *	T <sub>4</sub> *	MIT*/DIT*	T <sub>3</sub> */T <sub>4</sub> *
		ng/mg thyroid	mg/100 g body wt								
A	12	880 ± 180	4.1 ± 0.7	15 ± 10	4.8 ± 1.0	19 ± 3	46 ± 3	2.6 ± 0.3	21 ± 1	0.42 ± 0.07	0.12 ± 0.01
B	9	390 ± 100	5.1 ± 0.6	27 ± 11	3.8 ± 1.1	26 ± 3	44 ± 3	2.5 ± 0.8	23 ± 3	0.59 ± 0.06	0.11 ± 0.03
C	1	7.7	10	55	5.0	44	17	20	7.3	2.7	2.7
	2	4.4	13	49	3.5	41	16	19	5.2	2.5	3.5
	3	6.9	12	36	3.3	38	18	17	6.7	2.1	2.4
	4	6.4	14	43	5.5	41	20	16	12	2.1	1.3
	5	6.1	13	37	6.0	38	20	16	14	1.9	1.1
	6	9.5	7	48	5.1	36	21	23	17	1.8	1.4
	7	7.7	12	37	4.3	33	19	22	17	1.7	1.3
	8	7.2	16	54	2.6	32	21	23	17	1.5	1.4
	9§	55	11	26	3.6	37	22	18	12	1.7	1.5
	10§	97	7.1	33	2.7	38	23	18	14	1.7	1.3
Mean											
±SD		21 ± 30	12 ± 3	42 ± 9	4.2 ± 1.2	38 ± 3	20 ± 3	19 ± 3	12 ± 3	2.0 ± 0.3	1.8 ± 0.9

‡ Uptake of <sup>131</sup>I in thyroid 24 h after isotope injection.

§ Animals on iodine restriction for 4 wk.

|| In group C, data are provided for individual animals.

The distribution of radioactivity in the Pronase hydrolysates of the thyroids is presented in Table I. The I\* percentage was not different in the three groups. The percent of MIT\* increases progressively as the iodine supply is further restricted, the mean being 19 in group A, 26 in group B, and 38 in group C. There is no difference in the DIT\* percentage in groups A and B, but the value of 20% in group C is significantly lower than that observed in the control group ( $P < 0.001$ ). In addition, there is also no significant difference between the A and the B animals with regard to the relative distribution of isotope in T<sub>3</sub> and T<sub>4</sub>, while in group C there is a significant increase in the percent T<sub>3</sub>\* from a mean of 2.6 in groups A and B to 19 in group C ( $P < 0.001$ ). In addition, there is a substantial decrease in the percent T<sub>4</sub>\* from a mean of 22 in groups A and B, to 12 in group C ( $P < 0.001$ ). Approximately 90–100% of the label present in the thyroid hydrolysate is accounted for by these five components. The remainder of the labeled material appeared to be distributed evenly along chromatographic paper strips between the identifiable peaks.

As previously mentioned, there is a progressive increase in the ratio of MIT\*/DIT\* with increasing severity of iodine restriction. The ratio of T<sub>3</sub>\*/T<sub>4</sub>\* is 0.12 in group A, and not significantly different (0.11) in group B. However in group C the ratio is reversed, T<sub>3</sub>\* being 1.8 times T<sub>4</sub>\*. The distribution of the isotope in these normal and iodine-deficient glands is substan-

tially the same as that observed by previous investigators (1).

*Serum T<sub>3</sub> and T<sub>4</sub> levels in control and iodine-deficient rats.* Mean levels of serum T<sub>3</sub> and T<sub>4</sub> are presented in Table II. Where thyroid glands from two animals were pooled, the mean of the values for serum T<sub>3</sub> and T<sub>4</sub> were used. Serum T<sub>3</sub> levels in the control animals ranged from 29 to 54 ng/100 ml with a mean of 44. T<sub>3</sub> levels were not significantly different in group B with a range of from 42 to 61 ng/ml with the mean 48 ng/100 ml. In group C, the mean serum T<sub>3</sub> concentration was unchanged from the group A animals, being 43 ng/100 ml. Thus there was no change in the serum T<sub>3</sub> concentration in the iodine-deficient rat.

In the second column are given the concentrations of immunoassayable T<sub>4</sub>. In control animals, the mean was 4.2 μg/100 with a range of 3.6–5.3 μg/100 ml. Again there was no difference in the values obtained in the group B animals, where the mean was 4.5 ± 0.1 (mean ± SD). However, in the group C animals, T<sub>4</sub> was undetectable. With current methods, we were able to quantitate T<sub>4</sub> in rat serum to a minimum of 0.5 μg/100 ml. In parenthesis the values for PBI are shown, as determined by Boston Medical Laboratory in the serum of most of the group C animals. While it is not known precisely what quantities of noniodothyronine iodine are present in severely iodine-deficient rats, these low PBI values confirm the low serum T<sub>4</sub> estimates obtained by immunoassay. Inspection of the group C

animals shows that in animals 9 and 10, which had been on the diet for a period of only 4 wk, T<sub>4</sub> had already disappeared. These animals were examined at 10 days after institution of the low-iodine diet and still had normal T<sub>4</sub> levels at that time.

*T<sub>3</sub> and T<sub>4</sub> in the pronase hydrolysates of control and iodine-deficient thyroid glands.* The T<sub>3</sub> and T<sub>4</sub> concentrations of the rat thyroid hydrolysates are also shown in Table II. In the thyroid glands of the control rats, the T<sub>3</sub> concentration varied from 27 to 58 ng/mg wet wt with a mean of 39 ng/mg. In mildly deficient animals, T<sub>3</sub> content was moderately reduced to a mean of 26 ng/mg ( $P < 0.005$ ). An even more substantial reduction, to less than 5% of the control group, was observed in the severely iodine-deficient thyroids. The mean T<sub>3</sub> content in this group was 1.7 ng/mg. While the T<sub>3</sub> concentration was markedly reduced in the group C animals, the larger size of the thyroid gland resulted in an overall decrease of the gland T<sub>3</sub> content of approximately 85%, from a mean of 0.64 μg to about 0.10 μg/total gland.

T<sub>4</sub> content of the normal rat thyroid gland was 400 ng/mg wet wt. Like T<sub>3</sub>, the concentration of this iodothyronine was reduced in mild iodine deficiency to a mean of 270 ng/mg, significantly less than that of the A group ( $P < 0.001$ ). The mean thyroidal T<sub>4</sub> content in the C animals was 3.2 ng/mg wet wt, less than 1% of the control T<sub>4</sub> concentration. This is similar in magnitude to the reduction in the <sup>125</sup>I content of the gland. The total T<sub>4</sub> content of the goitrous glands was only 3% of the control value.

Of considerable interest is the fact that the molar ratio of T<sub>3</sub> to T<sub>4</sub> in these thyroid glands is similar to the ratio of T<sub>3</sub>\*/T<sub>4</sub>\* given in Table I. This ratio is precisely the same for groups A and B. However, in group C there appears to be a slight difference, the mean molar ratio of T<sub>3</sub>/T<sub>4</sub> of 1.0 being less than the ratio of 1.8 of the labeled hormones. However, the ratio is markedly increased over the control value by both methods. Thus, it would appear that the T<sub>3</sub>\*/T<sub>4</sub>\* ratio even as soon as 24 h after labeling is an accurate reflection of the absolute ratio of T<sub>3</sub> to T<sub>4</sub> present in the iodine-sufficient gland. This agreement occurs despite the fact that prolonged periods are required for complete equilibration of labeled iodine in the thyroid gland, and it will be discussed at greater length below.

*Distribution of <sup>131</sup>I in the serum of severely iodine-deficient rats.* Analysis of the labeled components in the serum in iodine-deficient rats was performed 24 h after isotope injection. Accurate estimates of T<sub>3</sub>\* and T<sub>4</sub>\* were possible only in this group, due to the high ratio of T<sub>4</sub>\* to T<sub>3</sub>\* present in normal animals. Since there is artifactual deiodination of a variable fraction of T<sub>4</sub>\* during chromatographic separation, it is difficult to determine accurately how much T<sub>3</sub>\* present in normal serum arises in vivo.

Direct chromatography of serum in collidine/NH<sub>4</sub>OH indicated that the only labeled components in serum of iodine-deficient rats were I<sup>-\*</sup>, T<sub>3</sub>\*, and T<sub>4</sub>\*, as has been previously reported (3). As can be seen in Table III, the quantity of I<sup>-\*</sup> present at 24 h was quite small, in all cases 6% or less of the total. The percentage of

TABLE II  
T<sub>3</sub> and T<sub>4</sub> Concentrations in Serum and Thyroids of Control and Iodine-Deficient Rats (Mean ± SD)

Group	n	Serum		Thyroid				Molar ratio T <sub>3</sub> /T <sub>4</sub>
		T <sub>3</sub>	T <sub>4</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>3</sub>	T <sub>4</sub>	
		ng/100 ml	μg/100 ml	ng/mg	μg/gland	ng/mg	μg/gland	
A	12	44 ± 9	4.2 ± 0.6	39 ± 10	0.64 ± 0.27	400 ± 100	6.7 ± 3.0	0.12 ± 0.03
B	9	48 ± 6	4.5 ± 0.3	26 ± 6	0.44 ± 0.14	270 ± 85	4.4 ± 0.8	0.12 ± 0.03
C§	1	41	<0.50 (0.4)‡	0.79	0.042	0.82	0.044	1.1
	2	45	" (0.6)	0.55	0.039	0.18	0.013	3.6
	3	40	"	0.88	0.065	1.9	0.14	0.55
	4	48	" (0.6)	0.97	0.081	2.2	0.19	0.53
	5	41	"	0.89	0.079	1.1	0.097	0.97
	6	33	" (0.6)	2.9	0.14	3.7	0.18	0.93
	7	35	" (0.6)	2.0	0.16	2.8	0.22	0.85
	8	58	" (0.6)	2.9	0.30	2.5	0.25	0.83
	9	40	" (0.8)	1.8	0.070	6.4	0.38	0.33
	10	53	" (0.6)	3.6	0.14	9.9	0.38	0.43
Mean ± SD		43 ± 6	<0.50 (0.6 ± 0.1)	1.7 ± 0.9	0.099 ± 0.078	3.2 ± 2.8	0.19 ± 0.12	1.01 ± 0.90

‡ PBI, μg I/100 ml.

§ In Group c, data are provided for individual animals.

|| Animals on iodine restriction for 4 wk.

label as  $T_3$  varied from 5 up to 17% at this time, whereas  $T_4^*$  ranged from 70 to 102% of the total. Since  $T_3^*$  and  $T_4^*$  content of the original serum was determined independently by  $^{125}\text{I}$ -labeled recovery standards, it was possible that the total percent recovery could be greater than 100. The mean recovery of labeled compounds totaled 96.5%, quite close to the anticipated figure of 100%. The ratio of  $T_4^*/T_3^*$  in the serum of the iodine-deficient animals varied from a low of 4.1 to as high as 18 with a mean of 11. In the last column, the predicted ratio of  $T_4^*/T_3^*$  is presented based on the gland  $T_4^*/T_3^*$ . The derivation of these ratios is discussed subsequently.

*Estimation of the respective contributions of the thyroid and peripheral  $T_4$ -to- $T_3$  conversion to the peripheral  $T_3$  pool in the rat.* It has been previously shown that peripheral  $T_4$ -to- $T_3$  conversion contributes substantially to the circulating  $T_3$  pool in man (15). A similar pathway for  $T_4$  metabolism has been shown in the rat by Schwartz, Surks, and Oppenheimer (16). If the clearances of  $T_3$  and  $T_4$  do not change in the iodine-deficient rat, it should be possible to calculate the relative contributions of the thyroid and peripheral  $T_4$ -to- $T_3$  conversion to the serum  $T_3$  pool in this situation. If plasma  $T_3$  or  $T_4$  clearance did change in iodine deficiency, then the knowledge of the serum concentrations alone would not suffice to calculate the total production rate of each hormone. Silva has recently demonstrated that there are no changes in the clearance of labeled  $T_3$  and  $T_4$  in rats deprived of iodine for 3 mo (4). Using the kinetic data for  $T_3$  and  $T_4$  clearance in the rat, previously published by Oppenheimer et al. (17), and Silva's conclusions (4), it is possible to calculate total  $T_3$  and  $T_4$  utilization in the control and iodine-deficient rats by using our values for serum  $T_3$  and  $T_4$ . It is then possible to estimate the relative proportion of  $T_3$  coming from  $T_4$  and that coming from the thyroid gland with the  $T_4$ -to- $T_3$  conversion ratio of 0.17 determined by Schwartz et al. in the rat (16). The estimates of the quantities of  $T_3$  and  $T_4$  metabolized by these rats are presented in Table IV (see

TABLE III  
Labeled Compounds in the Serum of Severely Iodine-Deficient Rats 24 h After  $^{131}\text{I}$  Administration

	% Total				Predicted† $T_4^*/T_3^*$
	$T_3^*$	$T_4^*$	$T_4^*/T_3^*$	% Total	
1	4.5	10	74	7.4	7.0
2	6.3	17	70	4.1	5.5
3	4.4	10	74	7.4	7.7
4	2.6	5.6	84	15	13
5	2.4	5.0	92	18	16
6	4.1	5.6	89	16	13
7	3.2	7.6	83	11	14
8	4.0	7.5	90	12	12
9	4.4	8.6	83	9.7	12
10	4.6	7.4	102	14	13
Mean	4.1	8.4	84	11	11
SEM	0.4	1.1	3	1	1

† Predicted on the basis of the gland  $T_4^*/T_3^*$  (see Discussion).

Appendix for formulae). Total  $T_4$  degradation is 710 and 760 ng/100 g body wt/day in groups A and B. Total  $T_3$  metabolism is 150 and 160 ng/100 g/day with approximately 67% of this  $T_3$  arising from  $T_4$  to  $T_3$  deiodination. In group C, the total  $T_4$  metabolism is reduced to approximately 11% of that present in the A and B groups even if the maximum  $T_4$  value of 0.5  $\mu\text{g}/100$  ml is used as the estimate of the  $T_4$  level. Nevertheless,  $T_3$  utilization is essentially unchanged. Since there is little  $T_3$  arising from  $T_4$ , over 90% of this  $T_3$  must originate from the thyroid gland if the conversion rate of 0.17 remains constant. While these data are obviously approximations, it would appear that a two-to-threefold increase in the absolute thyroidal  $T_3$  secretion rate must occur in the iodine-deficient rat. Even if the  $T_4$  to  $T_3$  conversion rate increased to 100% in these animals, a maximum of 71 ng of  $T_3/100$  g body wt/day could be generated. This is still less than one-half of the  $T_3$  utilized per day.

TABLE IV  
Analysis of the Quantity and Origin of  $T_3$  and  $T_4$  Utilized by the Rat†

Group	$T_4$ ng/100 g day <sup>-1</sup>	$T_3$ ng/100 g day <sup>-1</sup>	Source of $T_3$			
			$T_4$		Thyroid	
			ng	% Total	ng	% Total
A	710	150	101	67	49	33
B	760	160	108	68	52	32
C	85§	150	12	8	138	92

† Values based on mean serum iodothyronine concentrations in the three groups.

§ Calculated if maximum serum  $T_4$  (0.5  $\mu\text{g}/100$  ml) is assumed.




FIGURE 2 Correlation of predicted and observed ratios of <sup>131</sup>I-labeled T<sub>4</sub>\*/T<sub>3</sub>\* in serum and thyroid glands of severely iodine-deficient rats based on the assumption that the labeled hormones are secreted in the same ratio present in the gland. The solid line is constructed from the equation:

$$\frac{\text{Serum } T_4^*}{\text{Serum } T_3^*} = \frac{20.2 \text{ Gland } T_4^*}{\text{Gland } T_3^* + 0.17 \text{ Gland } T_4^*}$$

The points indicate the observed ratios in severely iodine-deficient animals.

Inspection of Table IV also reveals that the calculated molar ratio of secreted T<sub>3</sub> to secreted T<sub>4</sub> in the group C animals is about 2:1, similar to the ratio of T<sub>3</sub>\*/T<sub>4</sub>\* (1.8) in the thyroid glands of this group. The similarity of these ratios in the iodine-deficient rat suggested that these hormones might be secreted in the ratio in which they exist in the gland. Previous investigators have made similar speculations (3). Since both gland and serum T<sub>4</sub>\*/T<sub>3</sub>\* ratios were determined in the iodine-deficient animals, it was possible to test this hypothesis rigorously by attempting to predict the serum T<sub>4</sub>\*/T<sub>3</sub>\* ratio based on the gland T<sub>4</sub>\*/T<sub>3</sub>\*. This requires the assumption that there is relative equilibrium of the T<sub>3</sub>\*/T<sub>4</sub>\* ratios in the gland and serum at 24 h. This has been verified experimentally for the gland T<sub>3</sub>\*/T<sub>4</sub>\* by previous investigators (1). Since the half-lives of T<sub>3</sub> and T<sub>4</sub> in the rat are 6 and 12 h., respectively, it seems likely that the various differences in the distribution volume and fractional clearance rates of T<sub>3</sub> and T<sub>4</sub> have already taken place 1 day after injection.

Based on these assumptions, the ratio of secreted hormones can be calculated by substituting the percent label in gland T<sub>3</sub> and T<sub>4</sub> in the appropriate equations, which are derived in the Appendix. In Fig. 2, the ideal line for this relationship has been constructed. The

equation yields a hyperbolic curve, though at the lower extreme it approximates a straight line. Also plotted in this figure are the actual ratios of gland T<sub>4</sub>\*/T<sub>3</sub>\* and serum T<sub>4</sub>\*/T<sub>3</sub>\* in the group C animals where accurate estimates of serum T<sub>3</sub>\* are possible. There is an excellent agreement of these points with the predicted curve over a fourfold range of gland T<sub>4</sub>\*/T<sub>3</sub>\* ratios. Furthermore, in the last column of Table III, the individual predicted ratios of serum T<sub>4</sub>\* to T<sub>3</sub>\* are given, based on these assumptions. It is apparent that there is excellent correlation between these figures when these animals are examined individually. The mean predicted T<sub>4</sub>\*/T<sub>3</sub>\* ratio of 11 is precisely that observed.

## DISCUSSION

The changes in the pattern of radioiodine distribution in the thyroid glands of the mildly and severely iodine-deficient rats in these studies are similar to those changes observed by others (1). In addition, the decrease of serum T<sub>4</sub> to undetectable levels within 4 wk of iodine restriction is similar to the effect of iodine deficiency on the PBI (2). Despite the profound decrease in serum T<sub>4</sub>, no changes were detected in the levels of serum immunoassayable T<sub>3</sub> at either 1, 2, or 3 mo after restriction of iodine intake.

These data appear to be at variance with the results of a previous systematic study of this problem. Henger and Albright calculated serum  $T_3$  and  $T_4$  values in control and iodine-deficient rats using equilibrium labeling and iodine specific activity (18). In normal rats, the serum  $T_3$  was estimated to be 43 ng/100 ml, similar to our control values. In animals receiving iodine-deficient diet for 2 mo, a serum  $T_3$  concentration of 101 ng/100 ml was calculated. However, determination of the iodine specific activity in the low-iodine animals depended on accurate estimates of iodine at very low levels in the diet. Whether or not the different results obtained might be due to an overestimation of the dietary iodine remains to be determined.

Volpert and Werner reported values of 19–41 ng/100 ml in seven samples of serum from rats on low iodine diet for 4–5 wk as opposed to levels of 55 to 60 ng/100 ml in animals receiving a normal diet (5). Nejad, Bollinger, Mitnick, and Reichlin have recently reported no significant change in  $T_3$  values in rats fed a low iodine diet for an unspecified period of time while  $T_4$  levels decreased to less than 25% of control (19). However, the estimates of normal  $T_3$  concentrations in rat serum in these studies (16 ng/100 ml) were substantially lower than those we have obtained, for reasons not immediately apparent.

It has been previously speculated by Gross and Pitt-Rivers that  $T_4$  might have to undergo deiodination to  $T_3$  in order to be metabolically active and that, therefore,  $T_4$  might act primarily as a precursor of the active thyroid hormone,  $T_3$  (20). Oppenheimer, Schwartz, and Surks have demonstrated that  $T_4$ -to- $T_3$  conversion occurs in the rat and that this process can be inhibited by propylthiouracil (16, 21). In order to maintain hepatic mitochondrial alpha glycerophosphate dehydrogenase at normal levels in the propylthiouracil-treated animals, 2.5 times the usual replacement dose of  $T_4$  had to be given. This also restored the net  $T_3$  production from  $T_4$  to normal levels. If, as these studies suggest,  $T_4$  must be converted to  $T_3$  to exert its metabolic effect, feedback regulation of the thyroid-pituitary-hypothalamic axis would most likely be dependent on tissue (pituitary)  $T_3$  levels. The presence of specific binding sites for  $T_3$  but not for  $T_4$  in the nuclei of pituitary cells provides further support for this concept (22). The  $T_3$  present in the pituitary could be derived either from  $T_4$  or from the thyroid gland via direct secretion. One would anticipate under these circumstances that the normal homeostatic mechanism in the rat would then operate to maintain  $T_3$  production constant regardless of the source of the hormone.

Analysis of data in Tables III and IV suggests that these animals compensate for low iodine uptake by a shift from  $T_4$  as the primary source of  $T_3$ , to the thyroid

gland itself. Our data further indicate that this marked change in the ratio of secreted hormone results from increases in the molar ratio of  $T_3/T_4$  in iodine-deficient thyroglobulin. Whether this change in thyroglobulin composition is a result of increased thyroid-stimulating hormone (TSH) stimulation or decreased iodination, or both, remains to be determined. However, our analysis of the labeled hormone ratios in the serum is consistent with the simple assumption that they are secreted in the proportion in which they exist in the gland. A similar process may also occur in the normal rat, though the calculated molar ratio of secreted  $T_4$  to  $T_3$  is 12:1 (Table IV) and the ratio in the thyroid gland is 8-9/1.

As indicated above, iodine-deficient rats have elevated TSH levels that lead to the goiter and increased radioactive iodine uptake characteristic of this condition. It may be argued that such animals are, by definition, hypothyroid. Nevertheless, as previously pointed out, they appear to be "clinically" euthyroid (2-4). Whether, in fact, these animals are euthyroid in all respects awaits the results of more careful in vitro studies of tissue thyroid status. If it can be demonstrated that such animals are euthyroid and  $T_3$  levels remain normal with markedly decreased serum  $T_4$ , the results would be consistent with the interpretation that the TSH elevation in these animals is a compensatory response designed to maintain the animal in a euthyroid state. In addition, it would support the above-mentioned hypothesis that  $T_4$ , in the normal rat, is primarily a precursor of  $T_3$ . Other investigators, as well as ourselves, have observed that normal serum  $T_3$  concentrations may be associated with substantial TSH elevations and low serum  $T_4$  levels in patients with primary thyroid disease or after radioactive iodine treatment for hyperthyroidism (23-25). It is possible that some of these patients may have "intrathyroidal" iodine deficiency due to acquired defects in iodide trapping or organification with a similar compensatory response.

Loewenstein and Wollman have shown that prolonged periods (> 99 days) are required for complete equilibration of isotopic iodine into the total body iodine pool in the rat (26). For this reason accurate measurement of thyroidal and serum  $T_3$  and  $T_4$  by equilibrium labeling techniques is tedious. Inspection of Tables I and II shows that in groups A and B, multiplication of the fraction of  $T_4^*$  by the  $^{127}\text{I}$  content results in calculated  $T_4$  concentrations that are considerably less than the immunoassayable estimates. This presumably results largely from the incomplete equilibration of the isotope with the iodine pool and is in agreement with earlier studies (14). Nevertheless, immunoassayable thyroid  $T_4$  content in the control group agrees well with the values of about 350 ng  $T_4/\text{mg}$  wet wt in rat thyroid

that can be calculated from recent [ $^{127}\text{I}$ ]T<sub>4</sub> measurements (14). The fact that acute labeling with  $^{127}\text{I}$  results in the same T<sub>3</sub>/T<sub>4</sub> ratios as the immunoassay determination indicates that, in spite of the functional heterogeneity of thyroid gland iodine previously shown by many investigators, the T<sub>3</sub>/T<sub>4</sub> ratio of acutely formed hormone and total thyroidal iodothyronine are essentially the same. Since the immunoassay does not require separation of T<sub>3</sub> and T<sub>4</sub> with its attendant artifacts, it has obvious advantages in studies of this type and even more in the study of human thyroid tissue that cannot be chronically labeled.

Whether or not the rat has a mechanism for conservation of iodine efficient enough to allow a normal T<sub>3</sub> to be maintained indefinitely is not known. Our preliminary evidence suggests that after prolonged restriction of iodine intake, T<sub>3</sub> levels begin to decrease. More extensive studies are currently under way to confirm this observation.

## APPENDIX

The T<sub>4</sub> or T<sub>3</sub> metabolized in 24 h can be expressed as follows:

$$\text{Total } Tx = [Tx] \times V_{Tx} \times K_{Tx}$$

where [Tx] is the concentration of iodothyronine (moles per milliliter), V<sub>Tx</sub> is the volume of distribution per 100 g body wt and K<sub>Tx</sub> is the mean daily fractional removal rate of T<sub>3</sub> or T<sub>4</sub>.

Under steady-state conditions, the T<sub>4</sub> cleared daily equals the T<sub>4</sub> secreted by the thyroid. It has been shown in the rat that approximately 17% of the T<sub>4</sub> is metabolized via conversion to T<sub>3</sub> (16). Thus, thyroidal T<sub>3</sub> secretion can be expressed as the difference between the metabolized T<sub>3</sub> and the quantity derived from T<sub>4</sub> or:

Secreted T<sub>3</sub>

$$= ([T_3] \times V_{T_3} \times K_{T_3}) - 0.17([T_4] \times V_{T_4} \times K_{T_4})$$

Substitution of experimentally measured values for kinetic constants gives (17):

$$\begin{aligned} \text{Secreted T}_4/100 \text{ g/day} \\ = [T_4] \times 16.4 \times 1.032 \end{aligned}$$

$$\begin{aligned} \text{Secreted T}_3/100 \text{ g/day} \\ = ([T_3] \times 165 \times 2.07) - 0.17([T_4] \times 16.4 \times 1.032) \end{aligned}$$

If T<sub>4</sub>\* and T<sub>3</sub>\* are secreted in the same ratio as they exist within the gland and the peripheral pool is subsequently labeled with  $^{127}\text{I}$ , then when the ratios of T<sub>4</sub>\* to T<sub>3</sub>\* in the gland and in the serum are in relative equilibrium:

$$\begin{aligned} \frac{\text{Secreted T}_4^*}{\text{Secreted T}_3^*} &= \frac{\text{Gland T}_4^*}{\text{Gland T}_3^*} \\ &= \frac{(T_4^* \times 16.4 \times 1.032)}{T_3^* \times 165 \times 2.07 - 0.17} \\ &\quad \times (T_4^* \times 16.4 \times 1.032). \quad (1) \end{aligned}$$

Eq. (1) reduces to

$$\frac{\text{Gland T}_4^*}{\text{Gland T}_3^*} = \frac{16.9 T_4^*}{340.96 T_3^* - 2.88 T_4^*} \quad (2)$$

Eq. (2) can be solved for the ratio of labeled iodothyronines in the serum

$$\frac{T_4^*}{T_3^*} = \frac{20.2 \text{ Gland T}_4^*}{\text{Gland T}_3^* + 0.17 \text{ Gland T}_3^*} \quad (3)$$

## ACKNOWLEDGMENTS

The authors would like to acknowledge the careful technical assistance of Ms. Darina Sipula and Jitka Dockalova and the expert secretarial help of Ms. Barbara Brenneman.

This work was supported by Grant AM14283 from the U. S. Public Health Service and Grant 0-20 from the Health Research and Services Foundation of Pittsburgh.

## REFERENCES

1. Studer, H., and M. A. Greer. 1966. The Regulation of Thyroid Function in Iodine Deficiency. Hans Huber, Bern, Switzerland.
2. Studer, H., and M. A. Greer. 1965. A study of the mechanisms involved in the production of iodine-deficiency goiter. *Acta Endocrinol.* **49**: 610.
3. Greer, M. A., Y. Grimm, and H. Studer. 1968. Qualitative changes in the secretion of thyroid hormones induced by iodine deficiency. *Endocrinology.* **83**: 1193.
4. Silva, E. 1972. Disposal rates of thyroxine and triiodothyronine in iodine-deficient rats. *Endocrinology.* **91**: 1430.
5. Volpert, E. M., and S. C. Werner. 1972. Serum triiodothyronine concentration in the iodine-deficient rat. *Am. J. Anat.* **135**: 187.
6. Benotti, J., N. Benotti, S. Pino, and J. Gardyna. 1965. Determination of total iodine in urine, stool, diets, and tissue. *Clin. Chem.* **11**: 932.
7. Inoue, K., and A. Taurog. 1967. Digestion of  $^{127}\text{I}$ -labeled thyroid tissue with maximum recovery of  $^{127}\text{I}$ -iodothyronines. *Endocrinology.* **81**: 319.
8. Larsen, P. R., K. Yamashita, A. Dekker, and J. B. Field. 1973. Biochemical observations in functioning human thyroid adenomas. *J. Clin. Endocrinol. Metab.* **36**: 1109.
9. Larsen, P. R. 1971. Technical aspects of the estimation of triiodothyronine in human serum: evidence of conversion of thyroxine to triiodothyronine during assay. *Metabolism.* **20**: 609.
10. Larsen, P. R. 1972. Direct immunoassay of triiodothyronine in human serum. *J. Clin. Invest.* **51**: 1939.
11. Larsen, P. R., J. Dockalova, D. Sipula, and F. M. Wu. 1973. Immunoassay of thyroxine in unextracted human serum. *J. Clin. Endocrinol. Metab.* In press.
12. Sterling, K., D. Bellabarba, E. S. Newman, and M. A. Brenner. 1969. Determination of triiodothyronine concentration in human serum. *J. Clin. Invest.* **48**: 1150.
13. Inoue, K., and Taurog, A. 1968. Acute and chronic effects of iodide on thyroid radioiodine metabolism in iodine-deficient rats. *Endocrinology.* **83**: 279.
14. Lamas, L., and G. M. de Escobar. 1972. Iodoamino acid distribution in the thyroids of rats on different iodine intakes and with normal plasma protein-bound iodine. *Acta Endocrinol.* **69**: 473.

15. Braverman, L. E., S. H. Ingbar, and K. Sterling. 1970. Conversion of thyroxine ( $T_4$ ) to triiodothyronine ( $T_3$ ) in athyreotic human subjects. *J. Clin. Invest.* **49**: 855.
16. Schwartz, H. L., M. I. Surks, and J. H. Oppenheimer. 1971. Quantitation of extrathyroidal conversion of L-thyroxine to 3,5,3'-triiodo-L-thyronine in the rat. *J. Clin. Invest.* **50**: 1124.
17. Oppenheimer, J. H., H. L. Schwartz, H. Shapiro, G. Bernstein, M. Martinez, and M. I. Surks. 1970. Differences in primary cellular factors in influencing the metabolism and distribution of 3,5,3'-triiodothyronine and L-thyroxine. *J. Clin. Invest.* **49**: 1016.
18. Heninger, R. W., and E. C. Albright. 1966. Effect of iodine deficiency on iodine-containing compounds of rat tissues. *Endocrinology*. **79**: 309.
19. Nejad, I. F., J. A. Bollinger, M. Mitnick, and S. Reichlin. 1973. Importance of  $T_3$  secretion in altered states of thyroid function in the rat: cold exposure, subtotal thyroidectomy, and hypophysectomy. *Trans. Assoc. Am. Physicians Phila.* **85**: 295.
20. Gross, J., and R. Pitt-Rivers. 1954. Thyroid hormone physiology and biochemistry. II. Triiodothyronine in relation to thyroid physiology. *Recent Prog. Horm. Res.* **10**: 109.
21. Oppenheimer, J. H., H. L. Schwartz, and M. I. Surks. 1973. Propylthiouracil inhibits the conversion of L-thyroxine to L-triiodothyronine. *J. Clin. Invest.* **51**: 2493.
22. Schadow, A. R., M. I. Surks, H. L. Schwartz, and J. H. Oppenheimer. 1972. Specific triiodothyronine-binding sites in the anterior pituitary of the rat. *Science (Wash. D. C.)*. **176**: 1252.
23. Sterling, K., M. A. Brenner, E. S. Newman, W. D. Odell, and D. Bellabarba. 1971. The significance of triiodothyronine ( $T_3$ ) in maintenance of euthyroid status after treatment of hyperthyroidism. *J. Clin. Endocrinol.* **33**: 729.
24. Lieblich, J., and R. D. Utiger. 1972. Triiodothyronine radioimmunoassay. *J. Clin. Invest.* **51**: 157.
25. Larsen, P. R. 1972. Triiodothyronine: review of recent studies of its physiology and pathophysiology in man. *Metab. (Clin. Exp.)*. **21**: 1073.
26. Loewenstein, J. E., and S. H. Wollman. 1967. Kinetics of equilibrium-labeling of the rat thyroid gland with  $^{125}I$ . *Endocrinology*. **81**: 1063.