

Promotion of human adipocyte precursor replication by 17beta-estradiol in culture.

D A Roncari, R L Van

J Clin Invest. 1978;**62**(3):503-508. <https://doi.org/10.1172/JCI109153>.

Research Article

The influence of 17beta-estradiol and 17alpha-estradiol on adult human omental adipocyte precursors grown in a propagating culture system was studied. Cells were grown in subculture in the presence or absence of hormone. 17beta-estradiol resulted in significant promotion of adipocyte precursor replication, as determined by cell counting and incorporation of radioactive thymidine into DNA. The hormone stimulated cell multiplication in the concentration range 0.5-500 ng/ml growth medium. The highest level tested was 500 ng/ml. The maximal effects were obtained at 50 ng/ml (P less than 0.001 by paired t test, 48 h after hormone addition). All 10 cell strains (five were derived from men and five from women) that were tested responded similarly to the hormone. 17beta-estradiol did not affect cell size. 17alpha-estradiol did not promote the replication of adipocyte precursors, nor did it influence cell size. Thus, 17beta-estradiol, which is the active isomer in known target tissues, stimulates the multiplication of human adipocyte precursors in culture.

Find the latest version:

<https://jci.me/109153/pdf>



Promotion of Human Adipocyte Precursor Replication by 17β -Estradiol in Culture

DANIEL A. K. RONCARI and ROBIN L. R. VAN, *Institute of Medical Science and the Toronto Western Hospital, Department of Medicine, University of Toronto, Toronto, Canada M5S 1A8*

ABSTRACT The influence of 17β -estradiol and 17α -estradiol on adult human omental adipocyte precursors grown in a propagating culture system was studied. Cells were grown in subculture in the presence or absence of hormone. 17β -estradiol resulted in significant promotion of adipocyte precursor replication, as determined by cell counting and incorporation of radioactive thymidine into DNA. The hormone stimulated cell multiplication in the concentration range 0.5–500 ng/ml growth medium. The highest level tested was 500 ng/ml. The maximal effects were obtained at 50 ng/ml ($P < 0.001$ by paired t test, 48 h after hormone addition). All 10 cell strains (five were derived from men and five from women) that were tested responded similarly to the hormone. 17β -estradiol did not affect cell size. 17α -estradiol did not promote the replication of adipocyte precursors, nor did it influence cell size. Thus, 17β -estradiol, which is the active isomer in known target tissues, stimulates the multiplication of human adipocyte precursors in culture.

INTRODUCTION

The factors that control adipocyte precursor replication are unknown. Recently, a propagating culture system has been elaborated for these cells (1). It is suitable for studies of regulatory mechanisms operative in adipocyte precursor replication and differentiation.

The total body fat content of girls and boys is the same until puberty; it then increases in girls and decreases in boys (2). The fat content of women is nor-

mally higher than that of men (2, 3). The distribution of adipose tissue also becomes different at the time of puberty (4). Subcutaneous adipose tissue is thicker in women than in men (3). The greater thickness is a result of an increased number of adipocytes in several subcutaneous regions; in the gluteal area, however, the difference is mainly accounted for by the larger fat cells present in women (3). These facts suggest that estrogens may have the potential to influence adipose tissue growth.

It is known that estrogens are taken up by adipose tissue and isolated fat cells from mammals including humans (5–7). However, the significance of this uptake has not been established.

We herein report the promotion of cultured adult human adipocyte precursor replication by 17β -estradiol.

METHODS

Source, preparation and culture of adipocyte precursors. Tissue from 10 patients 20- to 60-yr-old (5 males and 5 females) was obtained at the time of abdominal surgery; the patients were not obese according to Metropolitan Life Insurance Company tables. The cells were isolated from omental adipose tissue as reported (1). After centrifugation, the digested stromal fraction was suspended in medium 199 (K. C. Biological, Inc., Lenexa, Kans.) supplemented with 10% (by volume) fetal calf serum (Flow Laboratories Inc., Rockville, Md.), penicillin (100 U/ml), and streptomycin (100 μ g/ml). Cell culture and preparation for subculture were conducted as described (1). The inocula (0.1 – 0.7×10^5 cells) were plated on 25-cm² Falcon tissue culture flasks (Falcon Plastics, Div. of BioQuest, Oxnard, Calif.) and grown in 5 ml of medium. All studies were conducted in first or subsequent subculture.

Cell counting and sizing. Cells were resuspended in 0.15 M NaCl, aliquots were diluted with Isoton (Coulter Electronics, Inc., Hialeah, Fla.) and counted in a Coulter counter model Z_F. The size of cells was determined at the same time in a Coulter Channelyzer II.

Incorporation of [³H]thymidine into DNA. At the beginning of first subculture, 10 μ Ci of [³H]thymidine (sp act = 45.5 Ci/mmol, New England Nuclear, Boston, Mass.) were added to each flask. 17β -estradiol (Ikapharm, Ramat-Gan, Israel, recrystallized from and dissolved in methanol) or 17α -estradiol (Steraloids, Wilton, N. H., dissolved in methanol) were added to one-half of the flasks. The final concentrations

This work was presented in part at the Annual Meeting of the Canadian Society for Clinical Investigation, Toronto, Canada, 25 January 1977 (1976. *Clin. Res.* 24: 662A. [Abstr.]).

Address reprint requests to Dr. Roncari at the Medical Sciences Building, Room 7238, University of Toronto, Toronto, Canada M5S 1A8.

Received for publication 29 July 1977 and in revised form 24 March 1978.

were 0.05–500 ng/ml growth medium for 17 β -estradiol and 50 ng/ml for 17 α -estradiol. The same volume of methanol as that which contained estrogen, was added to the control flasks.

Cells were disrupted by freezing and thawing in water five times. Portions of aqueous broken cell suspensions were then taken for protein determination and for measurement of [3 H]thymidine incorporation into DNA. The samples were rendered 5% (vol/vol) with respect to perchloric acid. After centrifugation, the pellets were washed and spun three times with 5% perchloric acid. The pellets were finally dissolved with dilute NaOH (pH of water adjusted to 9.5) and portions were counted with Bray's solution at an efficiency of 25–30%.

Analytical methods. Protein concentrations were determined by a microbiuret procedure (8). 17 β -estradiol levels were measured by a radioimmunoassay method (9). The proportion of free 17 β -estradiol was determined by equilibrium dialysis.

RESULTS

Characteristics of adipocyte precursors. The human omental adipocyte precursors, which were isolated and cultured as previously described, had the same morphological and enzymological characteristics as reported earlier (1). Under the experimental conditions used in this investigation, these characteristics and the doubling time of all the cell strains were similar in both first and subsequent subculture.

Analysis of 17 β -estradiol in growth medium. The mean concentration of 17 β -estradiol was 15 pg/ml in assays of four fetal calf serum samples (each performed in triplicate). The same stock of serum was used in all studies. When 17 β -estradiol was added to a final concentration of 50 ng/ml (in medium 199 supplemented with 10% by volume fetal calf serum), 40% of the hormone was in its free form.

Effect of 17 β -estradiol on cell counts. 17 β -estradiol resulted in a significant ($P < 0.001$ by paired t test) increase in the number of adipocyte precursors in culture (Fig. 1). As will be confirmed by the data on [3 H]-thymidine incorporation into DNA, these results indicate promotion of cell replication. The maximal effect was observed at a hormone concentration of 50 ng/ml growth medium. Because the lowest level of 17 β -estradiol that resulted in an appreciable increase in cell number was 0.5 ng/ml, the contribution of hormone present in fetal calf serum (15 pg/ml) was negligible.

Fig. 2 illustrates the effect of 17 β -estradiol on cell counts at different times in first subculture. The data are representative of time-courses using cell strains from 10 patients. At a 17 β -estradiol concentration of 50 ng/ml, the greatest promotion of replication occurred 48 h after addition of hormone. The photomicrographs in Fig. 3 show adipocyte precursors in first subculture. After 48 h of growth, the flask that did not contain 17 β -estradiol had a smaller number of adipocyte precursors (Fig. 3A) than the one with 17 β -estradiol at 50 ng/ml (Fig. 3B).




FIGURE 1 Effect of 17 β -estradiol on the number of adipocyte precursors. Cell strains were isolated from the omental adipose tissue stromal fraction of five men and five women. At the beginning of first subculture, 17 β -estradiol was added to one-half of the flasks at a final concentration of 50 ng/ml growth medium. The means of cell counts and their SE are shown at 48 h in culture. The effect of the hormone was significant by the paired t test at $P < 0.001$.

When the adipocyte precursors were suspended for cell counting and sizing, they became spherical enabling determination of cell diameter. 17 β -estradiol did not alter cell size, the diameter being in the range 17–22 μ m in each case.

In contrast to its 17 β -isomer, 17 α -estradiol at a concentration of 50 ng/ml growth medium did not alter the rate of adipocyte precursor replication (Fig. 4).




FIGURE 2 Cell counts of an adipocyte precursor strain in the presence and absence of 17 β -estradiol. At the beginning of first subculture, the hormone was added to one-half of the flasks at a final concentration of 50 ng/ml growth medium. The data are representative of time-courses using cell strains derived from 10 patients (five men and five women).

The data are representative of time-courses using cell strains from two patients, one a man, the other a woman. For comparison, the same strains were grown within the same week, in the same subculture, in the presence of 17 β -estradiol, which promoted replication as described. As was the case for 17 β -estradiol, its 17 α -isomer did not influence cell size.

Effect of 17 β -estradiol on radioactive thymidine incorporation into DNA. 17 β -estradiol resulted in a significant ($P < 0.001$) promotion of [^3H]thymidine incorporation into DNA (Fig. 5). This finding, which corroborated the evidence obtained by direct cell counting, indicated that the hormone enhanced the replication of adipocyte precursors in culture. The concentration of 17 β -estradiol required to obtain the maximal effect and the time when it occurred were the same as described for cell enumeration. As was the case for cell counts, all 10 cell strains responded similarly. 17 α -estradiol did not alter the degree of [^3H]thymidine incorporation into DNA.

In other experiments, when the same concentration of 17 β -estradiol was added again after 48 h in culture, it resulted in renewed promotion of adipocyte precursor replication, as indicated by cell counting and measurement of [^3H]thymidine incorporation into DNA. The same degree of promotion was observed under the following two experimental conditions: first, 17 β -estradiol was added again to cells grown in culture for 48 h; second, the adipocyte precursors were subcultured at 48 h and, using a similar inoculum size as in the previous passage, were grown again in fresh medium containing 17 β -estradiol. Cell replication was promoted to the same extent as observed after the first addition of hormone. Inoculum size, in the range $0.1\text{--}0.7 \times 10^5$ cells, did not influence the magnitude of the 17 β -estradiol effect. Decreasing availability of free, biologically active hormone is a possible reason for the diminishing effect after 48 h in culture.

In other experiments, 3 μCi of [^3H]thymidine were added at two-hourly intervals from 8 to 16 h in culture, and again at 24 and 48 h. The results were consistent with those illustrated in Fig. 2, that is, 17 β -estradiol did not result in an appreciably greater rate of [^3H]thymidine incorporation into DNA during the earlier periods of time, when compared to the 24–48-h interval. It was not feasible to conduct studies before 8 h of culture because an insufficient number of cells adhered to the substratum during this period of time.

For the sake of consistency and to allow statistical analysis on data obtained with cells in the same passage, most of the studies were done with precursors in first subculture. In addition, each cell strain was tested in at least one subsequent subculture. 17 β -estradiol resulted in a similar degree of promotion of adipocyte precursor replication up to the last (fourth) passage that was examined in this regard.

DISCUSSION



The uterus, vagina, corpus luteum, and the mammary glands have been considered to be the major target organs for estrogens (10, 11). Other sites of female sex steroid action include the hypothalamus and the pituitary (11). In the current study, 17 β -estradiol was found to promote the replication of adipocyte precursors in culture. 17 α -estradiol, the isomer inactive in known target tissues, was also without effect on fat cell precursors. Information obtained using cell culture systems must be interpreted with caution because it does not always reflect *in vivo* events. Our findings, however, do suggest that adipocyte precursors may be target cells for estrogens.

17 β -estradiol promotes the replication of human endometrial cells in culture at similar concentrations and time periods as observed for adipocyte precursors in our studies (12, 13). As is the case for target tissues *in vivo*, 17 α -estradiol is also biologically inactive on endometrial cells in culture (13).

Appreciable uptake of estrogens by fat cells occurs *in vitro* and *in vivo* (5–7). A recently reported study (14) has analyzed the effects of two dosage levels of the long-acting estrogen preparation, polyestradiol phosphate (Estradurin), in rats. After 10 wk, relatively high doses resulted in decreases of both total body weight and mass of subcutaneous and parametrial adipose tissue. The effects were ascribed to the anorexic properties of the hormonal preparation. After 4 wk, relatively smaller doses led to an increased weight of perirenal and parametrial fat depots. Body weight was not affected. The alterations in adipose tissue weight appeared to be a result of changes in fat cell size, rather than number. These results are difficult to interpret because of methodological reasons. Adipocyte precursors and more mature fat cells containing relatively small quantities of triglyceride, would not have been detected. Thus, an influence of the estrogen preparation on precursor cell replication may have been missed. In addition, demonstration of such an effect *in vivo* might require a longer treatment period.

It remains to be proven *in vivo* whether precursor cells take up estrogens and then replicate at an accelerated rate. If these events occurred under physiological conditions, augmented multiplication could account, at least partly, for the differences with respect to adipose tissue cellularity between men and women (3). These dissimilarities become apparent at puberty, a stage when differences in the synthesis and secretion of sex steroids become manifest.

The effect of 17 β -estradiol in culture is reproducible and statistically significant, but not dramatic. Because adipocyte precursors of women are exposed to a different (as compared to men) hormonal environment for several years, a “dramatic” effect might not be






FIGURE 4 Cell counts of an adipocyte precursor strain in the presence and absence of 17α -estradiol. At the beginning of first subculture, 17α -estradiol was added to one-half of the flasks at a final concentration of 50 ng/ml growth medium. The data are representative of time-courses with cell strains derived from two patients (one man and one woman).

expected. A relatively small effect exerted over a rather prolonged period of time could account, at least partly, for the fact that several adipose tissue depots of women contain a greater number of adipocytes than those of men (3). Direct proof of this proposal is of course required before a conclusion can be drawn.

Estrogens not only enhance the growth of such known target tissues as the reproductive organs, but also mediate their differentiation (10, 11). These hormones increase the number of initiation sites for transcription (15). Consequently, specific proteins which participate in target cell differentiation are synthesized (11, 16–20). It is not known whether these biochemical events are related to the stimulation of cell multiplication. While our studies indicate that 17β -estradiol promotes the replication of cultured adipocyte precursors, it is possible that estrogens also bring about specific biochemical changes.

Although the physiological significance of findings in cell culture requires confirmation by *in vivo* studies, the adipocyte precursor culture system provides a useful approach to the investigations of factors potentially operative in fat tissue growth and adipocyte differentiation.

FIGURE 3 Photomicrographs of live adipocyte precursors. They were taken after 48 h in first subculture. The control was devoid of added 17β -estradiol (A), whereas the other flask was supplemented with hormone at a final concentration of 50 ng/ml growth medium (B). Original magnification of 160.




FIGURE 5 Effect of 17β -estradiol on $[^3H]$ thymidine incorporation into DNA of adipocyte precursors. Cell strains were isolated from the adipose tissue of the 10 patients referred to in Fig. 1. At the beginning of first subculture, 10 μ Ci of $[^3H]$ thymidine (45.5 Ci/mmol sp act) were added to 5 ml growth medium contained in each flask; at the same time, 17β -estradiol was added to one-half of the flasks at a final concentration of 50 ng/ml growth medium. The means of the disintegrations per minute incorporated and their SE are shown at 48 h in culture. The effect of 17β -estradiol was significant at $P < 0.001$.

ACKNOWLEDGMENTS

The authors are grateful to Dr. D. W. Killinger and Mrs. E. Perel for valuable advice and for determinations of total and free 17β -estradiol. They are also thankful to Doctors B. R. Taylor, R. E. Mathews, and I. H. Koven for their interest and help. The skilled technical assistance of Mr. D. K. Yip is acknowledged.

This work was supported by Ontario Heart Foundation grant I-46, Medical Research Council of Canada grant MA-5827, and The Atkinson Charitable Foundation.

REFERENCES

1. Van, R. L. R., C. E. Bayliss, and D. A. K. Roncari. 1976. Cytological and enzymological characterization of adult human adipocyte precursors in culture. *J. Clin. Invest.* **58**: 699–704.
2. Friis-Hansen, B. 1965. Hydrometry of growth and aging. *In Human Body Composition*. J. Brozek, editor. Pergamon Press Ltd., Oxford. 1st edition. 191–209.
3. Sjöström, L., U. Smith, M. Krotkiewski, and P. Björntorp.

1972. Cellularity in different regions of adipose tissue in young men and women. *Metab. Clin. Exp.* **21**: 1143–1153.
4. Garn, S. M., and J. A. Haskell. 1959. Fat changes during adolescence. *Science (Wash. D. C.)*. **129**: 1615–1616.
 5. Roncari, D. A. K. 1967. Metabolism of estrogens in rat adipose tissue. *Endocrinology*. **80**: 1160–1163.
 6. Bleau, G., K. D. Roberts, and A. Chapdelaine. 1974. The in vitro and in vivo uptake and metabolism of steroids in human adipose tissue. *J. Clin. Endocrinol. Metab.* **39**: 236–246.
 7. Longcope, C., J. H. Pratt, S. H. Schneider, and S. E. Fineberg. 1976. In vivo studies on the metabolism of estrogens by muscle and adipose tissue of normal males. *J. Clin. Endocrinol. Metab.* **43**: 1134–1145.
 8. Munkres, K. D., and F. M. Richards. 1965. The purification and properties of *Neurospora* malate dehydrogenase. *Arch. Biochem. Biophys.* **109**: 466–479.
 9. Lindner, H. R., E. Perel, A. Friedlander, and A. Zeitlin. 1972. Specificity of antibodies to ovarian hormones in relation to the site of attachment of the steroid hapten to the peptide carrier. *Steroids*. **19**: 357–375.
 10. Jensen, E. V., and H. I. Jacobson. 1962. Basic guides to the mechanisms of estrogen action. *Recent Prog. Horm. Res.* **18**: 387–414.
 11. Chan, L., and B. W. O'Malley. 1976. Mechanisms of action of the sex steroid hormones (first part). *N. Engl. J. Med.* **294**: 1322–1328.
 12. Hiratsu, T. 1968. In vitro cultivation of human endometrium and the influence of steroid hormones on a cell line derived from the endometrium. *Kobe J. Med. Sci.* **14**: 29–48.
 13. Chen, L., H. R. Lindner, and M. Lancet. 1973. Mitogenic action of oestradiol-17 β on human myometrial and endometrial cells in long-term tissue cultures. *J. Endocrinol.* **59**: 87–97.
 14. Krotkiewski, M. 1976. The effects of estrogens on regional adipose tissue cellularity in the rat. *Acta Physiol. Scand.* **96**: 128–133.
 15. Kalimi, M., S. Y. Tsai, M.-J. Tsai, J. H. Clark, and B. W. O'Malley. 1976. Effect of estrogen on gene expression in the chick oviduct. Correlation between nuclear-bound estrogen receptor and chromatin initiation sites for transcription. *J. Biol. Chem.* **251**: 516–523.
 16. Oka, R., and R. T. Schimke. 1969. Interaction of estrogen and progesterone in chick oviduct development. II. Effects of estrogen and progesterone on tubular gland cell function. *J. Cell Biol.* **43**: 123–127.
 17. Katzenellenbogen, B. S., and J. Gorski. 1972. Estrogen action in vitro. Induction of the synthesis of a specific uterine protein. *J. Biol. Chem.* **247**: 1299–1305.
 18. O'Malley, B. W., and A. R. Means. 1974. Female steroid hormones and target cell nuclei. *Science (Wash. D. C.)*. **183**: 610–620.
 19. Luskey, K. L., M. S. Brown, and J. L. Goldstein. 1974. Stimulation of the synthesis of very low density lipoproteins in rooster liver by estradiol. *J. Biol. Chem.* **249**: 5939–5947.
 20. Chan, L., R. L. Jackson, B. W. O'Malley, and A. R. Means. 1976. Synthesis of very low density lipoproteins in the cockerel. Effects of estrogen. *J. Clin. Invest.* **58**: 368–379.