# **JCI** The Journal of Clinical Investigation

# Recognition of Hapten-Modified Cells In Vitro by Human T Lymphocytes

## Michael F. Seldin, ..., Robert R. Rich, Stuart L. Abramson

J Clin Invest. 1979;64(4):967-976. https://doi.org/10.1172/JCI109563.

#### Research Article

Clearer definition of the recognitive structures of human T lymphocytes for antigens will be required to elucidate the molecular basis of diseases and immunological responses induced or regulated by normal or abnormal T-cell function. For this purpose we have investigated the cellular requirements for immune responses in vitro to trinitrophenyl-conjugated peripheral blood mononuclear cells. The responding cell was characterized as a T cell on the basis of rosetting with sheep erythrocytes. T-cell recognition of hapten in proliferative responses depended upon presentation of antigen in an appropriate stimulator-cell context. Neither autologous hapten-modified erythrocytes nor T cells restimulated responses of in vitro-primed lymphocytes. Moreover, hapten-conjugated non-T cells were more effective than modified unfractionated cells in restimulating proliferative responses. Both macrophages and non-T lymphocytes effectively restimulated hapten-conjugate responses.

Cell-mixing experiments indicated that the failure of haptenated T cells to stimulate proliferative responses was not because of a lack of fresh macrophages; these experiments suggested instead that T cells do not express appropriate structures necessary to present haptenic determinants in an immunogenic form. Hapten-modified T cells, however, were capable of inducing primed lymphocytes to become efficient cytotoxic effector cells, indicating that T-cell recognitive units for stimulation of proliferative and cytotoxic responses are different. These data support the concept that for induction of proliferative responses, human T cells recognize conventional antigens in association with *HLA-D*-region-encoded [...]



### Find the latest version:

https://jci.me/109563/pdf

## Recognition of Hapten-Modified Cells In Vitro by Human T Lymphocytes

MICHAEL F. SELDIN, ROBERT R. RICH, and STUART L. ABRAMSON, The Howard Hughes Medical Institute Laboratory, and Departments of Microbiology and Immunology and of Medicine, Baylor College of Medicine, and The Methodist and Veterans Administration Hospitals, Houston, Texas 77030

ABSTRACT Clearer definition of the recognitive structures of human T lymphocytes for antigens will be required to elucidate the molecular basis of diseases and immunological responses induced or regulated by normal or abnormal T-cell function. For this purpose we have investigated the cellular requirements for immune responses in vitro to trinitrophenyl-conjugated peripheral blood mononuclear cells. The responding cell was characterized as a T cell on the basis of rosetting with sheep erythrocytes. T-cell recognition of hapten in proliferative responses depended upon presentation of antigen in an appropriate stimulator-cell context. Neither autologous hapten-modified erythrocytes nor T cells restimulated responses of in vitro-primed lymphocytes. Moreover, hapten-conjugated non-T cells were more effective than modified unfractionated cells in restimulating proliferative responses. Both macrophages and non-T lymphocytes effectively restimulated hapten-conjugate responses.

Cell-mixing experiments indicated that the failure of haptenated T cells to stimulate proliferative responses was not because of a lack of fresh macrophages; these experiments suggested instead that T cells do not express appropriate structures necessary to present haptenic determinants in an immunogenic form. Hapten-modified T cells, however, were capable of inducing primed lymphocytes to become efficient cytotoxic effector cells, indicating that T-cell recognitive units for stimulation of proliferative and cytotoxic responses are different. These data support the concept that for induction of proliferative responses, human T cells recognize conventional antigens in association with *HLA-D*-region-encoded Ia-like molecules.

#### INTRODUCTION

Specific abnormalities of human T-cell activation or control of immune responses seem critically important in the pathogenesis of numerous diseases. The increasing recognition of the association of HLA antigens with a predisposition to certain diseases (1, 2) has focused urgent attention on elucidation of the function of major histocompatibility complex (MHC)<sup>1</sup> genes in the pathogenesis of such disorders. Furthermore, the development and course of disease processes induced by immune responses to modifications in autologous cell surfaces may be controlled by genes within the MHC. An understanding, however, will first require a definition of the role of histocompatibility determinants in T-cell recognition. Examination of human in vitro proliferative and cytotoxic responses to hapten-modified autologous cells may allow a functional characterization of the autologous determinants involved in both normal and abnormal immune responses.

In the mouse, recognition of viruses (3), minor histocompatibility antigens (4) and haptenic determinants (5–7) by T cells is dependent upon the expression of autologous determinants controlled by the murine MHC, *H*-2. Thus, in vitro T-cell function is restricted in cytotoxic responses by H-2K and H-2D antigens, and in proliferative responses by H-2Iregion-encoded Ia antigens. In vivo, immune T cells can only transfer resistance to virus-infected hosts if *H*-2K or *H*-2*D* homology requirements are met (8). On the other hand, resistance to mycobacterial infections are only conferred by T cells which share H-2I-region determinants (thought to be analogous to the HLA-D region) with the host (9). In addition, the expression of Ia determinants on guinea pig macrophage subpopula-

This work was presented in part at the Annual Meeting of the American Society for Clinical Investigation, Washington, D. C., 7 May 1979.

Received for publication 11 April 1979 and in revised form 25 May 1979.

<sup>&</sup>lt;sup>1</sup>Abbreviations used in this paper:  $\Delta$ cpm, net counts per minute; MHC, major histocompatibility complex; PBMC, peripheral blood mononuclear cell(s); TNP, trinitrophenyl.

tions has been shown to be critical in the presentation of both protein antigens and hapten-modified cell surface determinants to responder T cells (10-12).

Recent studies in humans have demonstrated that cell-mediated cytotoxic responses to the Y male antigen (13), influenza-infected cells (14, 15), and haptenmodified cells (16, 17) are constrained by HLA-A- or -Blocus antigens. In contrast, our previous data (18, 19), recently confirmed by Thorsby and Nousiainen (20) indicated that secondary proliferative responses of human peripheral blood lymphocytes to haptenmodified cells were associated with the HLA-D region. Nevertheless, a substantial portion of the proliferative response to hapten-modified human cells was not dependent upon the expression of typed histocompatibility antigens. Thus, hapten-modified allogeneic stimulator cells that did not share major histocompatibility antigens with sensitized responders often induced substantial proliferative responses. To better understand the role of autologous determinants in the recognition of antigen by human lymphoid cells, we have characterized the cellular components involved in responses to hapten-modified cells. Human lymphocytes were primed in vitro to autologous trinitrophenyl (TNP) -conjugated stimulator cells and restimulated with various hapten-modified cell populations. The data strongly suggest that human T-cell recognitive units for stimulation of secondary proliferative responses, in contrast with those involved in restimulation of primed cytotoxic responses, are dependent upon the presentation of hapten in the context of macrophage or non-T-lymphocyte surface antigens.

#### **METHODS**

Cell culture technique. Heparinized venous blood was obtained from healthy human subjects and peripheral blood mononuclear cells (PBMC) were isolated and conjugated to TNP with trinitrobenzene sulfonic acid (ICN Nutritional Biochemicals, Cleveland, Ohio) as previously described (18). Cells were cultured in RPMI 1640 (Grand Island Biological Co., Grand Island, N. Y.) supplemented with 20% ABpositive, heat-inactivated, pooled human plasma.

For primary stimulation, cells were cultured in roundbottofm  $17 \times 100$ -mm tubes (2001, Falcon Labware, Oxnard, Calif.). Responder and stimulator cells were each cultured at a final concentration of  $7.5 \times 10^5$  cells/ml in 2-ml cultures for 20-30 d as previously described (18). Stimulator cells were inactivated with 500 rads of <sup>60</sup>Co (500 rads/min).

For secondary stimulation, primed cells were harvested, washed, and cocultured in round-bottom microculture wells (IS-MRC-96; Linbro Chemical Co., Hamden, Conn.) for assessment of proliferative responses, or in  $17 \times 100$ -mm round-bottom tubes for induction of cytotoxic effector cells. Responder cells ( $1.5 \times 10^6$ /microculture well or  $1.5 \times 10^6$ / tube) were cultured with an equal number of 2,000 rad-inactivated fresh TNP-conjugated or unconjugated stimulators.

Glutaraldehyde-fixed stimulator cells were prepared according to the method of Forman (21). TNP-conjugated or unconjugated cells suspended in phosphate-buffered saline (pH 7.4) were added to an equal volume of glutaraldehyde (Polysciences, Inc., Warrington, Pa.) in phosphate-buffered saline. The cells were treated for 10 s before addition of a 50-fold excess of Hanks' balanced salt solution (Microbiological Associates, Walkersville, Md.).

Proliferative responses to TNP-conjugated cells. Proliferative responses were assessed on days 3 or 4 after secondary stimulation by addition of 1.0  $\mu$ Ci of tritiated thymidine (2.0 Ci/mM sp act, New England Nuclear Corp., Boston, Mass.) to cultures for the final 20 h of the incubation period.

Data from separate experiments are expressed as mean counts per minute of three to four replicate cultures with the SEM. Net counts per minute ( $\Delta$  cpm) were calculated by subtracting the counts per minute of responses to unconjugated stimulators from the counts per minute of cultures with TNP-conjugated stimulator cells. Errors for  $\Delta$  cpm calculations were determined by the formula for propagation of error. Percent control responses were calculated by determining the mean response to stimulators and employing the following formula: ([(cpm experimental TNP-conjugated) – (cpm experimental unconjugated)]/[(cpm unfractionated TNP-conjugated)] × 100.

Cytotoxic responses to TNP-conjugated cells. Cytotoxic responses were assessed using a <sup>51</sup>Cr-release assay. Primed effector cells were harvested from tubes 4-5 d after secondary stimulation, washed, and adjusted to appropriate concentrations in RPMI 1640 supplemented with 10% heat-inactivated fetal calf serum (Grand Island Biological Co.). The target cells for the cell-mediated lympholysis assay were PBMC that had been cultured for 4-5 d. These cells were harvested on the day of assay, labeled with <sup>51</sup>Cr by incubating  $30 \times 10^6$  cells for 60 min with 300 µCi of Na<sub>2</sub><sup>51</sup>CrO<sub>4</sub> (New England Nuclear Corp.), and either TNP-conjugated or left unmodified. All target cell preparations were washed three times before use in the assay. 20,000 target cells were then incubated in 0.2 ml of RPMI 1640 supplemented with 10% fetal calf serum in triplicate cultures with varying numbers of viable effector cells in  $12 \times 75$ -mm plastic tubes (Falcon Labware). After incubation for 6 h at 37°C in a humidified, 5% CO<sub>2</sub> atmosphere, <sup>51</sup>Cr-release was measured as previously described (22). Cytolytic activity is expressed as: % specific release = ([experimental release - spontaneous release]/ [freeze-thaw - spontaneous release]) × 100. Spontaneous release values were consistently <25% of total counts per minute.

Identification of lymphoid cell populations. The percentage of human T cells in various fractions of PBMC was assessed by a standard sheep erythrocyte E-rosetting technique (23). C3 receptor-bearing cells were detected by rosette formation with sheep erythrocytes coated with 19S anti-sheep antibodies (Cordis Labs. Inc., Miami, Fla.) and mouse complement (23). Numbers of monocytes and macrophages were estimated with nonspecific esterase staining (24). Surface immunoglobulin-positive cells were assayed with a fluorescein-conjugated goat anti-human polyvalent antisera (Behring Diagnostics, Somerville, N. J.) according to method of Pernis et al. (25).

Cell fractionation procedures. Unfractionated PBMC contained  $\approx 65\%$  E rosette-positive cells (T cells), 35% C3 receptor-bearing cells (macrophages and B cells) and 25% esterase-positive cells (macrophage-monocytes). To obtain Tcell and non-T-cell lymphoid populations a rapid sheep erythrocyte-rosette procedure was used (26). Briefly, PBMC were rosetted with neuraminidase (20 U/ml Receptor Destroying Enzyme-Vibrio comma extract, Microbiological Associates) treated sheep erythrocytes for 30 min at 4°C and E rosette-positive and non-E-rosette fractions isolated by Ficoll-Isopaque (Pharmacia Fine Chemicals Inc., Piscataway, N. J.) gradient centrifugation. Sheep erythrocytes were lysed with Tris/NH<sub>4</sub>Cl, and cells were washed twice with media. Unfractionated controls were treated in the same manner. The non-E-rosette population contained <8% E rosette-positive cells, >40% esterase-positive cells, and >60% C3 receptorbearing cells. To obtain a T cell-enriched population the Erosette fraction was rosetted a second time. The double E rosette-purified cells contained  $\approx$ 85% E rosette-positive cells, <2% esterase-positive cells, <5% C3 receptor-bearing cells, and <2% surface immunoglobulin-positive cells. All populations were >95% viable as assessed by trypan blue.

Macrophage-monocyte populations were isolated by a plastic dish adherence procedure. 50 million PBMC were incubated in  $100 \times 20$ -mm plastic Petri dishes (Falcon Labware) for 45 min at 37°C in a 5% CO<sub>2</sub> incubator in 10 ml of RPMI 1640 with 10% autologous serum that had not been heat inactivated. Nonadherent cells were removed from Petri dishes by vigorously washing four times at 37°C with 7 ml of RPMI 1640 with 20% fetal calf serum. To recover adherent cells, Petri dishes were then incubated at 37°C for 30 min in the presence of 3 mM of EDTA and then washed twice with Hanks' balanced salt solution at 4°C. The recovered cells were >95% esterase positive and >95% viable. To control for possible adverse effects of EDTA treatment, both adherent and nonadherent cell populations were treated with EDTA in identical fashion in experiments in which adherent cells were isolated.

To obtain non-T lymphocytes, cells that did not adhere to plastic Petri dishes were first fractionated into E rosettepositive and -negative populations. E rosette-negative cells were then depleted of monocytes by passage thru Sephadex G-10 (Pharmacia Fine Chemicals Inc.) columns using a modification of the method of Ly and Mishell (27, 28). The resultant population contained <10% E rosette-positive cells, <5% esterase-positive cells, and >40% surface immuno-globulin-positive cells. HLA typing. Typing for HLA-A- and -B-locus antigens was performed in the Methodist Hospital Tissue Typing Laboratory with standard serologic methods (29). HLA-D-region specificities were similarly detected with B-cell typing procedures (DRw) (30).

#### RESULTS

Characterization of human TNP-conjugate responder cells. The ability of human T cells to form rosettes with sheep erythrocytes was used to determine the responder cell in proliferative responses to hapten-modified cells. Unfractionated PBMC were sensitized to autologous hapten-modified cells and 3 wk later separated into E-rosette and non-E-rosette cell populations. As shown in Table I, the unfractionated, E rosette-positive and E rosette-negative primed responder cells were then restimulated with autologous PBMC which were either TNP-conjugated or unconjugated. The E rosette-positive cells were clearly better responders than the unfractionated cells. Conversely, the responses of the non-E-rosette cells were substantially reduced. The small response noted by the latter population may be explained by the residual contamination,  $\cong 8\%$ , of E rosette-positive cells, although modest intrinsic responsiveness of the non-Erosette population was not ruled out. These experiments thus demonstrate that the cells which proliferated in response to hapten-conjugated PBMC were largely or exclusively T lymphocytes.

Responder cells*	Unconjugated‡	TNP-conjugated	Δcpm§	Percent control
Experiment 1				
Unfractionated	$1588 \pm 147$ ¶	$17,799 \pm 721$	$16,211 \pm 735$	100
E-rosette	$1542 \pm 206$	$27,695 \pm 1218$	$26,153 \pm 1235$	161
Non-E-rosette	$3460 \pm 150$	$7,739 \pm 385$	4,279±413	26
Experiment 2				
Unfractionated	$802 \pm 41$	$9,214\pm50$	$8,412\pm64$	100
E-rosette	$2522 \pm 130$	$15,633\pm872$	$13,111 \pm 882$	156
Non-E-rosette	$894 \pm 124$	$3,304 \pm 223$	$2,410\pm255$	29

 TABLE I

 Characterization of the Responder Cell Population in Secondary

 Hapten-Conjugate Proliferative Responses

\* Primed responder cells were generated by co-culture with autologous TNP-conjugated stimulators for 3-4 wk. These sensitized cells were either left unfractionated or separated into E rosette-positive or non-E-rosette responder populations. Responder cells were cultured at  $1.5 \times 10^5$  cells/microculture well.

‡ TNP-conjugated or unconjugated stimulators were fresh PBMC inactivated with 2,000 rads and cultured at  $1.5 \times 10^{5}$  cells/microculture well together with responder cells. Cultures were harvested 70 h after secondary stimulation.

§ Δcpm were calculated by subtracting counts per minute of responses to unconjugated stimulators from counts per minute of cultures with TNP-conjugated stimulators. <sup>II</sup> Calculated by formula given in Methods.

 $\P$  Data expressed as mean counts per minute of three to four replicate cultures with the SEM.

Characterization of human TNP-conjugate stimulator cells. Previous data from our laboratory suggested that human lymphocytes, at least in part, recognize haptens in the context of autologous MHC determinants. Lymphocytes primed to autologous TNPmodified cells are preferentially restimulated by TNPmodified cells that share HLA-D-region determinants. However, a substantial portion of the secondary TNPconjugate proliferative response did not appear to be genetically restricted. To examine whether a component of secondary proliferative responses was not dependent upon autologous histocompatibility antigens on lymphoid cells, primed responder cells were restimulated with hapten-modified autologous erythrocytes. Erythrocytes express only low levels of HLA-A- and -B-locus determinants and do not express detectable D-region antigens (31). As shown in Table II, TNP-conjugated erythrocytes did not restimulate proliferative responses. In experiments not shown, erythrocyte stimulator concentrations up to six times that of mononuclear cells were still ineffective in inducing responses. These results are consistent with the concept that expression of HLA determinants is necessary for the induction of proliferative responses to haptens. Furthermore, as shown in Table II, viable stimulator cells are required. Thus, neither heat-killed cells nor glutaraldehyde-fixed cells restimulated TNPprimed responder cells. Stimulator cells treated with as little as 0.1% glutaraldehyde for 10 s were almost devoid of immunogenic activity.

Different subpopulations of lymphoid cells express varying amounts of histocompatibility antigens. T cells, B cells, and monocytes all express substantial quantities of HLA-A- and -B-locus antigens. However, only monocytes-macrophages, B cells, and "null" lymphocytes, in contrast with resting T cells, express readily detectable *HLA-D*-region-encoded Ia-like antigens on their cell surface (32, 33). It was, therefore, of particular interest to investigate the ability of PBMC subpopulations to stimulate hapten-conjugate proliferative responses.

Lymphocytes were primed to autologous TNPmodified PBMC and restimulated with different populations of TNP-modified or unmodified stimulator cells as shown in Table III. Hapten-modified T cells (E-rosette fraction) were not effective stimulators. Moreover, hapten-modified non-T cells (non-E-rosette fraction), which contained both monocytes and non-T lymphocytes, restimulated proliferative responses more effectively than modified unfractionated cells.

The ineffectiveness of modified T cells to function as stimulators might be explained by a culture requirement for fresh macrophages. On the basis of nonspecific esterase staining, the E-rosette stimulator cell fraction contained <2% monocytes-macrophages whereas the non-E-rosette stimulator cell fraction possessed  $\approx 50\%$  monocytes-macrophages. Although the primed responder cells contained 25% esterasepositive cells, it is possible that accessory cell function in long-term cultures is defective. It was also possible that the inability of T cells to stimulate haptenconjugate proliferative responses could be explained by a suppressive mechanism. To address these questions, cell-mixing experiments were performed (Table IV). In these experiments, stimulator populations were comprised of equal concentrations of E-rosette and non-E-rosette cells that were either TNP-conjugated or unconjugated. The addition of non-E-rosette cells did not enhance the stimulatory capacity of TNP-modified E-rosette cells. Furthermore, the addition of E rosette-

Secondary Proliferative Responses*					
Percent m control					
1,546 100					
90 2					
137 3					
144 100					
26 4					
87 100					
530 16					
164 10					
371 4					

 TABLE II

 Neither Hapten-Conjugated Erythrocytes nor Dead PBMC Stimulate

 Secondary Proliferative Responses\*

\* Basic experimental protocol and data analysis as in Table I. Unfractionated responder cells were primed by coculture with autologous TNP-conjugated stimulators.

‡ Stimulators were either autologous PBMC or autologous erythrocytes. PBMC were killed after TNP-modification by either a 30-min incubation at 56°C or glutaraldehyde fixation for 10 s.

TABLE III
Stimulatory Capacity of Hapten-Modified E-Rosette and Non-E-Rosette Cells
in Secondary Proliferative Responses*

Stimulators‡	Unconjugated	TNP-conjugated	Δcpm	Percent control
Experiment 1				
Unfractionated	$802 \pm 41$	$9,214\pm508$	$8,412\pm510$	100
E-rosette	$620 \pm 60$	$793 \pm 173$	$173 \pm 183$	2
Experiment 2				
Unfractionated	$520 \pm 14$	$8,087 \pm 402$	$7,567 \pm 402$	100
E-rosette	$669 \pm 63$	$1,981\pm204$	$1,312\pm214$	17
Non-E-rosette	$2,842 \pm 372$	13,846±876	$11,004 \pm 952$	145
Experiment 3				
Unfractionated	$1,304 \pm 124$	$9,780\pm740$	$8,476\pm750$	100
E-rosette	$1,076 \pm 368$	$3,235\pm320$	$2,159 \pm 488$	25
Non-E-rosette	$3,877 \pm 163$	$15,349\pm525$	$11,472\pm550$	135

\* Basic experimental protocol and data analysis as in Table 1. Unfractionated responder cells were primed by coculture with autologous TNP-conjugated stimulators.

‡ Stimulators were either fresh PBMC (unfractionated), or fresh PBMC which were fractionated into E-rosette-positive and non-E-rosette populations.

positive cells to modified non-E-rosette cells did not inhibit responses to the latter population. The data therefore suggested that hapten-conjugated T cells are intrinsically ineffectual as stimulators of secondary proliferative responses.

Additional fractionation procedures were then used to further characterize the cell populations capable of effectively stimulating TNP-conjugate proliferative responses. Unfractionated peripheral blood was first separated into plastic-adherent and nonadherent populations. The adherent fraction was consistently >95% esterase positive, whereas the nonadherent fraction contained 10-15% esterase-positive cells. Nonadherent cells were further separated into E rosette-positive and non-E-rosette fractions, and the non-E-rosette population (35–40% esterase positive) was passed through Sephadex G-10 columns to remove monocytes-macrophages.

Purified plastic-adherent peripheral blood monocytes functioned effectively as hapten-conjugate stimulators (Table V). This adherent monocyte population was, however, less stimulatory than unfractionated cells. Interestingly, the non-E-rosette fraction, which had been depleted of the adherent population, was more effective than the adherent monocytes in stimulating hapten-conjugate proliferative responses. Removal of most of the esterase-positive cells from the non-E-rosette fraction by passage over Sephadex G-10 columns did not affect the stimulatory capacity. That is, a non-T-lymphocyte fraction, which contained <4%

	Stim	Stimulators		
	Unconjugated	TNP-conjugated	cpm	Δcpm
Experiment 1	NER + ER		2,438±146	
	ER -	+ TNP-NER	$13,814 \pm 1044$	11,376±1054
	NER -	⊢ TNP-ER	$3,929 \pm 155$	$1,491\pm213$
Experiment 2	NER + ER		$4,119\pm236$	_
	ER -	+ TNP-NER	$14,547 \pm 382$	$10,427 \pm 450$
	NER -	⊢ TNP-ER	$6,945 \pm 456$	$2,825\pm513$

 TABLE IV

 Hapten-Conjugated T Cells do not Stimulate Secondary Proliferative Responses\*

\* Basic experimental protocol and data analysis as in Table I.

**‡** Stimulator population contained equal concentration of E-rosette (ER) and non-E-rosette (NER) cells that were either TNP-conjugated or unconjugated.

Stimulators‡	Unconjugated	TNP-conjugated	Δcpm	Percent control
Experiment 1				
Unfractionated cells	$857 \pm 75$	$4,110\pm192$	$3,253 \pm 206$	100
Monocytes	$299 \pm 24$	$2,562\pm204$	$2,263\pm205$	70
Non-E-rosette cells	$1,097 \pm 131$	$6,974 \pm 663$	$5,877 \pm 676$	181
E-rosette cells	$284 \pm 15$	$987 \pm 52$	$703 \pm 54$	22
Non-T lymphocytes	$1,591\pm89$	$8,824 \pm 222$	$7,233 \pm 239$	222
Experiment 2				
Unfractionated cells	$1,167 \pm 198$	$10,268 \pm 696$	$9,101\pm724$	100
Monocytes	$841 \pm 156$	$6,530 \pm 325$	$5,689 \pm 361$	63
Non-E-rosette cells	$1,649 \pm 222$	$9,318 \pm 347$	$7,669 \pm 412$	84
E-rosette cells	$587 \pm 85$	$1,426 \pm 158$	$839 \pm 179$	9
Non-T lymphocytes	$1,489 \pm 265$	$8,942 \pm 496$	$7,453\pm 562$	82

 TABLE V

 Ability of Monocytes and Non-T Lymphocytes to Stimulate Secondary

 Hapten-Conjugate Proliferative Responses\*

\* Basic experimental protocol and data analysis as in Table I. Unfractionated responder cells were primed by coculture with autologous TNP-conjugated stimulator.

t Stimulators were either fresh PBMC (unfractionated) or lymphoid populations fractionated from fresh PBMC. Monocytes were plastic adherent cells which contained >95% esterase-positive cells. Non-E-rosette and E-rosette fractions were obtained by separation of the plastic nonadherent cell population. Non-T lymphocytes were non-Erosette cells passaged through Sephadex-G10 columns (Pharmacia Fine Chemicals, Inc., Piscataway, N.J.) to remove monocytes and macrophages. The non-E-rosette population contained 35–40% esterase-positive cells, whereas the non-T-lymphocyte population contained <4% esterase-positive cells.

esterase-positive cells, was as stimulatory as the non-E-rosette fraction, which contained over 40% esterase-positive cells, and was more stimulatory than the adherent population which contained over 95% esterase-positive cells. It was still possible, however, that the residual macrophage contamination was responsible for the ability of this fraction to stimulate. A titration analysis of the stimulator cell populations was therefore performed (Fig. 1). A ninefold dilution of the non-T-lymphocyte stimulator cells, containing <400 monocytes, stimulated responses as great as



FIGURE 1 Titration analysis of proliferative responses to different hapten-conjugated subpopulations of PBMC. Primed responder lymphocytes were restimulated with various concentrations of stimulator cells obtained as described in Table V.

>95,000 purified monocytes. Furthermore, reduction of the monocyte stimulator-cell concentration demonstrated that the initial concentration of these cells was not suppressive.

Requirements for proliferative vs. cytotoxic responses to hapten-modified cells. Our previous data demonstrated that optimal secondary proliferative responses of human lymphocytes to hapten-modified cells require HLA-D-region homology between responder and stimulator cells. On the other hand, Shaw et al. (17, 34) found that T cells primed to TNP-modified cells lysed modified allogeneic targets to the same extent as modified autologous targets, although partial HLA-A- and -B-locus restriction of cytotoxic responses was demonstrated by cold target-blocking studies. To determine whether the specificities involved in these two types of T cell-mediated responses were, in fact, different, and not readily explained by uncontrolled technical variables between laboratories, we compared proliferative and cytotoxic responses within the same experiment (Fig. 2). Primed responder cells were restimulated with hapten-modified or unmodified autologous cells, allogeneic human cells that did not share any HLA determinants with the responder, or mouse cells. Although the modified allogeneic human cells restimulated only 30% of the proliferative response of modified autologous stimulators (Fig. 2, left panel),



FIGURE 2 Antigenic requirements of proliferative vs. cytotoxic responses to hapten-modified cells. In the left panel the bars represent the net proliferative responses of primed A responder cells restimulated with autologous cells (A), allogeneic human lymphoid cells which did not share HLA antigens (B), or murine splenic lymphocytes. In the right panel the cytotoxic response of primed A responder cells restimulated with hapten-modified A cells was assayed against A, B, or mouse targets which were either TNP-conjugated (closed symbols) or unconjugated (open symbols). Proliferative responses were measured on day 3 and cytotoxic responses on day 4 after secondary stimulation. The standard errors for cytotoxic responses were always <2% of the total releasable radioactivity.

modified allogeneic cells were lysed as well or better than the modified autologous target cells (Fig. 2, right panel). TNP-modified murine cells did not stimulate either proliferative or cytotoxic responses, consistent with the specificity shown by other investigators (34).

Because human cytotoxic responses to haptenmodified cells did not appear to be related to the expression of HLA-D-region controlled antigens, we next investigated whether T cells could stimulate cytotoxic responses. Fig. 3 shows the proliferative and cytotoxic responses induced when primed responder cells were restimulated with either hapten-modified unfractionated cells or hapten-modified E-rosette cells. Unmodified non-E-rosette cells were added to all cultures to control for possible macrophage requirements. Recent experiments have shown that macrophages are necessary for accessory function but are not required for hapten presentation in murine TNP-conjugate cytotoxic responses (35). Consistent with the data already presented, modified T cells did not effectively restimulate proliferative responses (Fig. 3A). However, both effector cell populations, those stimulated with modified unfractionated and those with modified T cells, could extensively lyse hapten-modified targets (Fig. 3B). Thus, although T cells could not stimulate proliferative responses, they could induce cytotoxic effector cells.

#### DISCUSSION

Our data support the concept that some human T cells recognize antigen only in association with autologous



FIGURE 3 Stimulator-cell requirements for cytotoxic responses are different from those for proliferative responses. This figure shows the pooled results of three separate experiments in which responder cells were restimulated with either TNP-conjugated unfractionated (Unf) PBMC or with TNPconjugated E-rosette (ER) stimulator cells. Non-E-rosette (NER) cells were added to all cultures to control for possible accessory cell requirements. Four days after stimulation, secondary proliferative responses (panel A) and secondary cytotoxic responses (panel B) were assessed. (panel A) The bars represent the mean cpm±SEM; control responses to unmodified cells are shown in the open bars adjacent to the responses to TNP-modified unfractionated (solid bar) or Erosette (stippled bar) stimulators. (panel B) The ability of effector cells primed with either TNP-unfractionated cells (•) or TNP-E-rosette cells (•) to lyse TNP-conjugated targets (closed circles) or unconjugated targets (open circles) is depicted.

non-T-cell lymphoid determinants. Thus, only those cell populations which express easily detectable amounts of D-region-encoded Ia-like antigens effectively restimulated proliferative responses of TNPconjugate-primed T lymphocytes. This finding extends our previous observations of preferential restimulation of autologous TNP-conjugate-primed cells with modified stimulators that share D-region determinants, DRw and Dw (18, 19). A substantial proliferative response, however, was often generated by modified allogeneic stimulators that did not share detected histocompatibility antigens with the primed responder. The stimulatory capacity of modified allogeneic cells which do not share D-region determinants may be a result of extensive cross-reactivity between human histocompatibility antigens or of as yet undetected specificities. Because non-T lymphocytes which function as effective stimulators also express unique antigens encoded outside of HLA the influence of non-MHC antigens in the recognition of hapten must also be considered (36, 37).

The role of macrophages in proliferative responses to TNP-conjugated cells has previously been studied by Thomas et al. (38) in the guinea pig. In accord with the data reported here, these investigators found that TNP-conjugated thymocytes and erythrocytes did not induce proliferative responses; addition of unmodified macrophages to TNP-modified T cells did not restore responses. Similarly, we found that the addition of nonE-rosette stimulator cells, which contain a high proportion of macrophages ( $\approx$ 50%), had no effect on the net proliferative response to modified T cells. Addition of unmodified T cells to TNP-modified non-E-rosette cells also had no effect. These data militate against the likelihood that macrophage-culture effects, amplifier effects, and suppressor-cell mechanisms explain the failure of modified T cells to induce proliferative responses. Because modified T cells can induce cytotoxic effector cells, these results imply that T cells do not possess structures capable of presenting haptenic determinants to those T cells which respond in proliferative assays.

The histocompatibility requirements for cytotoxic effector-cell function differ from those for proliferative responses. In the human, HLA-A- or -B-locus homology requirements between killer and target cells have been reported for cytotoxic responses generated against the Y male antigen (13), influenza-infected cells (14, 15), and hapten-modified cells (16, 17). In contrast, we and others have reported the importance of HLA-D-region determinants in the generation of human proliferative responses to haptens and other conventional antigens (18-20, 39-41). In this study we have found that antigen presentation requirements for induction of cells capable of cytotoxic effector function also differ from those for proliferative responses. Nonactivated T cells, which do not express easily detectable amounts of D-region-encoded determinants, were capable of inducing secondary TNP-conjugate cytotoxic responses. Thus, the dichotomy between histocompatibility requirements for proliferative and cytotoxic responses is present for response induction as well as at the effector level.

The non-E-rosette population of cells that most effectively presents haptenic determinants is the same PBMC population that stimulates allogeneic mixed leukocyte reactions (42, 43) and autologous mixed leukocyte reactions (43). The observation that glutaraldehyde fixation of TNP-modified cells abrogates the capacity to stimulate secondary proliferative responses is also consistent with previous observations in mixed leukocyte reactions (44). Interestingly, cytotoxic responses to both autologous TNP-modified cells and allogeneic cells can be effectively induced with glutaraldehyde-fixed cells in the mouse (21, 44). Thus the properties of antigenic determinants associated with the induction of proliferative responses differ from those for cytotoxic responses. These results provide further evidence that the distinction between "lymphocyte-defined" or class I molecules (HLA-D) and "serologically defined" or class II molecules (HLA-A and -B), initially demonstrated for reactions to allogeneic cells (45, 46), applies to human immune responses to conventional antigens.

Both adherent monocytes and non-T lymphocytes effectively stimulated secondary TNP-conjugate proliferative responses. The non-T-lymphocyte population contains B cells and nonsurface-immunoglobulinbearing, non-T lymphocytes (null cells), both of which, in contrast to T cells, express Ia-like determinants (32, 33). Additional studies will be required to ascertain whether B cells or null cells, or both, are capable of effective hapten-conjugate presentation. We found that adherent macrophages restimulated proliferative responses less well than either unfractionated cells or non-E-rosette cells. This finding suggests that it may be possible to distinguish between presentation of antigens by subpopulations of macrophages and other Ia-positive cells for development of proliferative responses. Experiments designed to test this hypothesis are in progress.

Lawrence (47) theorized almost two decades ago that delayed hypersensitivity reactions and autoimmune diseases may be expressions of homograft reactions undertaken by the host against certain of its tissues. For delayed hypersensitivity reactions it now appears likely that antigens are indeed recognized in association with autologous determinants. Proliferative responses of lymphocytes from human subjects sensitized in vivo by dinitrochlorobenzene skin painting are preferentially restimulated in vitro with autologous dinitrophenyl-modified cells (48). Available data suggest that the proliferative responses in vitro are related in magnitude and specificity to delayed hypersensitivity reactions (49-51). Investigations of both proliferative responses to hapten-conjugated cells (18-20) and to purified protein derivative (39-41)suggest that the autologous determinants involved in delayed hypersensitivity reactions are encoded in the HLA-D region. Haptenic determinants on soluble proteins or membrane fragments (52) might be presented to T cells by means of adherence to macrophages, transconjugation of haptenic determinants, or macrophage processing, both in vivo and in vitro. Thus it is now reasonable to suggest that in the pathogenesis of delayed hypersensitivity reactions, antigens are presented to host T cells in the context of HLA-Dregion determinants. Epidermal Langerhans cells as well as tissue macrophages express Ia-like antigens (53) and thus might also subserve this function. In addition, it seems possible that the association of HLA determinants with a predisposition to certain diseases is intimately related to the antigen presentation role of these determinants in immue responses. Autoimmune disease, cancer, and ageing may all in part be determined by immune responses to subtle alterations in autologous cell surfaces. Further elucidation of Tcell recognitive units and the cellular requirements involved in responses to modified autologous cells may prove important in understanding the immunopathology of such diseases.

#### ACKNOWLEDGMENTS

The authors wish to thank Dr. John Stobo and Dr. Barton Haynes for their helpful suggestions and Dr. Daniel L. Kastner, Dr. Roger D. Rossen, and Dr. Tom Douglas for their thoughtful discussions of this work. We would also like to acknowledge the technical assistance of Ruth Rickaway and Mai Van and the secretarial assistance of Christina Garibay.

This work was supported in part by grants AI 13810, AI 15394, and RR 00350 from the National Institutes of Health.

#### REFERENCES

- Svejgaard, A., and L. P. Ryder. 1977. Associations between HLA and disease. In HLA and Disease. J. Dausset and A. Svejgaard, editors. Williams & Wilkins Co., Baltimore, Md. 46-71.
- Batchelor, J. R., and P. J. Morris. 1977. HLA and Disease. In Histocompatibility Testing. W. F. Bodmer, J. R. Batchelor, J. G. Bodmer, H. Festenstein, and P. J. Morris, editors. Munksgaard, Copenhagen. 205-258.
- Doherty, P. C., R. V. Blanden, and R. M. Zinkernagel. 1976. Specificity of virus-immune effector T cells for H-2K or H-2D compatible interactions: implications for H-antigen diversity. *Transplant. Rev.* 29: 89-124.
   Bevan, M. J. 1975. The major histocompatibility com-
- Bevan, M. J. 1975. The major histocompatibility complex determines susceptibility to cytotoxic T cells directed against minor histocompatibility antigens. J. Exp. Med. 142: 1349-1364.
- Shearer, G. M., T. G. Rehn, and A-M. Schmitt-Verhulst. 1976. Role of the murine major histocompatibility complex in the specificity of *in vitro* T cell-mediated lympholysis against chemically-modified autologous lymphocytes. *Transplant Rev.* 29: 222-248.
- Schmitt-Verhulst, A-M., G. A. Garbino, and G. M. Shearer. 1977. H-2 homology requirements for secondary cellmediated lympholysis and mixed lymphocyte reactions to TNP-modified syngeneic lymphocytes. J. Immunol. 118: 1420-1427.
- Yano, A., R. H. Schwartz, and W. E. Paul. 1977. Antigen presentation in the murine T-lymphocyte proliferative response.I. Requirement for genetic identity at the major histocompatibility complex. J. Exp. Med. 146: 828-843.
- 8. Zinkernagel, R. M., and R. Welsh. 1976. H-2 compatibility requirement for virus-specific T cell-mediated effector functions *in vivo*. I. Specificity of T cells conferring antiviral protection against lymphocytic choriomeningitis virus is associated with H-2K and H-2D. J. Immunol. 117: 1495-1502.
- Zinkernagel, R. M., A. Althage, B. Adler, R. V. Blanden, W. F. Davidson, V. Kees, M. B. C. Dunlop, and D. C. Shreffler. 1977. H-2 restriction of cell-mediated immunity to an intracellular bacterium. Effector T cells are specific for *Listeria* antigen in association with H-21 regioncoded self-markers. J. Exp. Med. 145: 1353-1367.
- Rosenthal, A. S., and E. M. Shevach. 1976. Macrophage-T lymphocyte interaction: the cellular basis for genetic control of antigen recognition. *In* The Role of Products of the Histocompatibility Gene Complex in Immune Responses. D. H. Katz and B. Benacerraf, editors. Academic Press, Inc., New York. 335–353.
- 11. Yamashita, U., and E. M. Shevach. 1977. The expression

of Ia antigens on immunocompetent cells in the guinea pig. II. Ia antigens on macrophages. J. Immunol. 119: 1584-1588.

- Thomas, D. W., U. Yamashita, and E. M. Shevach. 1977. Nature of the antigenic complex recognized by T lymphocytes. IV. Inhibition of antigen-specific T cell proliferation by antibodies to stimulatory macrophage Ia antigens. J. Immunol. 119: 223-226.
- Goulmy, E., A. Termijtelen, B. A. Bradley, and J. J. van Rood. 1977. Y-antigen killing by T cells of woman is restricted by HLA. *Nature (Lond.)*. 266: 544-545.
- McMichael, A. J., A. Ting, H. J. Zweerink, and B. A. Askonas. 1977. HLA restriction of cell-mediated lysis of influenza virus-infected human cells. *Nature (Lond.).* 270: 524-526.
- Biddison, W. E., S. Shaw, and D. L. Nelson. 1979. Virus specificity of human influenza virus-immune cytotoxic T cells. J. Immunol. 122: 660-664.
- Dickmeiss, E., B. Soeberg, and A. Svejgaard. 1977. Human cell-mediated cytotoxicity against modified targets is restricted by HLA. *Nature (Lond.)*. 270: 526-528.
- Shaw, S., and G. H. Shearer. 1978. Human cytotoxic responses in vitro to trinitrophenyl-modified autologous cells. II. Diversity of self determinants recognized in association with TNP. J. Immunol. 121: 290-299.
- Seldin, M. F., and R. R. Rich. 1978. Human immune responses to hapten-conjugated cells. I. Primary and secondary proliferative responses in vitro. J. Exp. Med. 147: 1671-1683.
- Seldin, M. F., R. R. Rich, and B. Dupont. 1979. Human immune responses to hapten-conjugated cells. II. The roles of autologous and allogeneic histocompatibility determinants in proliferative responses in vitro. J. Immunol. 122: 1828-1833.
- Thorsby, E., and H. Nousiainen. 1979. In vitro sensitization of human T lymphocytes to hapten (TNP) -conjugated and non-treated autologous cells is restricted by self-HLA-D. Scand. J. Immunol. 9: 183-192.
- Forman, J. 1977. T cell-mediated cytotoxicity against trinitrophenyl-modified cells: effects of glutaraldehyde treatment on the immunogenicity and antigenicity of trinitrophenyl-modified cells. J. Immunol. 118: 1755– 1762.
- Kastner, D. L., and R. R. Rich. 1978. Regulation of alloantigen-induced cytotoxic responses by concanavalin A-activated lymphoid cells: suppression by antigen elimination. J. Immunol. 121: 1732-1737.
- Aiuti, F., J-C. Cerottini, R. R. A. Coombs, M. Cooper, H. B. Dickler, S. S. Frøland, H. H. Fudenberg, M. F. Greaves, H. M. Grey, H. G. Kunkel, J. B. Natvig, J-L. Preud'homme, E. Rabellino, R. E. Ritts, D. S. Rowe, M. Seligmann, F. P. Siegal, J. Stjernswärd, W. D. Terry, and J. Wybran. 1974. Identification, enumeration, and isolation of B and T lymphocytes from human peripheral blood. Scand. J. Immunol. 3: 521-532.
- 24. Yam, L. T., C. Y. Li, and W. H. Crosby. 1971. Cytochemical identification of monocytes and granulocytes. *Am. J. Clin. Pathol.* 55: 283-290.
- 25. Pernis, B., L. Forni, and L. Amante. 1970. Immunoglobulin spots on the surface of rabbit lymphocytes. J. Exp. Med. 132: 1001-1018.
- Ettenger, R. B., G. Opelz, J. Walker, P. I. Terasaki, C. Vittenbogaart, A. J. Pennisi, M. H. Malekzadeh, and R. N. Fine. 1978. Antibodies to donor B lymphocytes and mixed lymphocyte culture blocking in cadaveric renal transplantation. *Transplantation (Baltimore).* 25: 169–172.

- Ly, I. A., and R. I. Mishell. 1974. Separation of mouse spleen cells by passage through columns of Sephadex G-10. J. Immunol. Methods. 5: 239-247.
- 28. Schwartz, R. H., A. R. Bianco, B. S. Handwerger, and C. R. Kahn. 1975. Demonstration that monocytes rather than lymphocytes are the insulin-binding cells in preparations of human peripheral blood mononuclear leukocytes: implications for studies of insulin-resistant states in man. Proc. Natl. Acad. Sci. U. S. A. 72: 474-478.
- Terasaki, P. I., and J. D. McClelland. 1964. Microdroplet assay of human serum cytotoxins. *Nature (Lond.)*. 204: 998-1000.
- Bodmer, J. G., P. Pickbourne, and S. Richards. 1977. Ia Serology. *In* Histocompatibility Testing. W. F. Bodmer, J. R. Batchelor, H. Festenstein, and P. J. Morris editors. Munksgaard, Copenhagen. 35-84.
- Doughty, R. W., S. R. Goodier, and K. Gelsthorpe. 1973. Further evidence for HL-A antigens present on adult peripheral red blood cells. *Tissue Antigens*. 3: 189-194.
- Schlossman, S. F., L. Chess, R. E. Humphreys, and J. L. Strominger. 1976. Distribution of Ia-like molecules on the surface of normal and leukemic human cells. *Proc. Natl. Acad. Sci. U. S. A.* 73: 1288-1292.
- 33. Chess, L., R. Evans, R. E. Humphreys, J. L. Strominger, and S. F. Schlossman. 1976. Inhibition of antibodydependent cellular cytotoxicity and immunoglobulin synthesis by an antiserum prepared against a human Bcell Ia-like molecule. J. Exp. Med. 144: 113-122.
- Shaw, S., D. L. Nelson, and G. M. Shearer. 1978. Human cytotoxic response in vitro to trinitrophenyl-modified autologous cells. I. T cell recognition of TNP in association with widely shared antigens. J. Immunol. 121: 281-299.
- Pettinelli, C. B., A-M. Schmitt-Verhulst, and G. M. Shearer. 1979. Cell types required for H-2 restricted cytotoxic response generated by trinitrobenzene sulfonatemodified syngeneic cells on trinitrophenyl-conjugated proteins. J. Immunol. 122: 847-854.
- Legrand, L., and J. Dausset. 1975. A second lymphocyte system (Cy-Li). In Histocompatibility Testing. F. Kissmeyer-Neilsen, editor. Munksgaard, Copenhagen. 665– 670.
- 37. Mann, D. L., W. Strober, S. I. Katz, S. Hsia, and D. B. Amos. Br and Wh, B cell antigens in gluten sensitive enteropathy and dermatitis herpetiformis. *In* The Role of Products of the Histocompatibility Gene Complex in Immune Responses. D. H. Katz and B. Benacerraf, editors. Academic Press, Inc., New York. 59-69.
- Thomas, D. W., G. Forni, E. M. Shevach, and I. Green. 1977. The role of the macrophage as the stimulator cell in contact sensitivity. J. Immunol. 118: 1677-1681.
- Bergholtz, B. O., and E. Thorsby. 1977. Macrophagedependent response of immune human T lymphocytes to PPD in vitro. Scand. J. Immunol. 6: 779-786.

- Bergholtz, B. O., and E. Thorsby. 1978. HLA-D restriction of macrophage-dependent response of human T lymphocytes to PPD in vitro. Inhibition by anti-HLA-DR antisera. Scand. J. Immunol. 8: 63-73.
- Sonderstrup Hansen, G., B. Rubin, S. F. Sorensen, and A. Svejgaard. 1978. Importance of HLA-D antigens for the cooperation between human monocytes and T lymphocytes. *Eur. J. Immunol.* 8: 520-575.
- 42. Lohrmann, H. P., L. Novikous, and R. G. Graw. 1974. Stimulatory capacity of human T and B lymphocyte in human mixed leukocyte culture. *Nature (Lond.).* 250: 144-146.
- Kuntz, M. M., J. B. Innes, and M. E. Weksler. 1976. Lymphocyte transformation induced by autologous cells. IV. Human T-lymphocyte proliferation induced by autologous or allogeneic non-T lymphocytes. J. Exp. Med. 143: 1042-1054.
- Rollinghoff, M., and H. Wagner. 1975. Secondary cytotoxic allograft responses in vitro. I. Antigenic requirements. Eur. J. Immunol. 5: 875-879.
- 45. Bach, F. H., M. B. Widmer, J. L. Bach, and J. Klein. 1972. Serologically defined and lymphocyte-defined components of the major histocompatibility complex in the mouse. J. Exp. Med. 136: 1430-1444.
- 46. Klein, J. 1979. The major histocompatibility complex of the mouse. Science (Wash. D. C.). 203: 516-521.
- 47. Lawrence, H. S. 1960. Delayed sensitivity and homograft sensitivity. Annu. Rev. Med. 11: 207-230.
- Levis, W. R., and A. M. Dattner. 1977. Clonal priming of lymphocytes can measure human immune response gene function. N. Engl. J. Med. 297: 842.
- Mills, J. A. 1966. The immunologic significance of antigen induced lymphocyte transformation in vitro. J. Immunol. 97: 239–247.
- Stulbarg, M., and S. F. Schlossman. 1968. The specificity of antigen-induced thymidine-2-14C incorporation into lymph node cells from sensitized animals. J. Immunol. 101: 764-769.
- 51. Paul, W. E., G. W. Siskind, and B. Benacerraf. 1968. Specificity of cellular immune responses. Antigen concentration dependence of stimulation of DNA synthesis in vitro by specifically sensitized cells, as an expression of the binding characteristics of cellular antibody. J. Exp. Med. 127: 25-42.
- 52. Miller, A. E., and W. R. Levis. 1973. Studies on contact sensitization of man with simple chemicals. I. Specific lymphocyte transformation in response to dinitrochlorobenzene sensitization. J. Invest. Dermatol. 61: 261-269.
- 53. Stingl, G., S. I. Katz, L. D. Abelson, and D. L. Mann. 1978. Immunofluorescent detection of human B cell alloantigens on S-Ig-positive lymphocytes and epidermal Langerhans' cells. J. Immunol. 120: 661–664.