

Effect of Calcium on Superoxide Production by Phagocytic Vesicles from Rabbit Alveolar Macrophages

P. Daniel Lew, Thomas P. Stossel

J Clin Invest. 1981;67(1):1-9. <https://doi.org/10.1172/JCI110000>.

Research Article

Phagocytic vesicles from rabbit lung macrophages produced superoxide in the presence of NADH or NADPH. At 37°C, these vesicles generated 51 ± 7.8 nmol O_2^- /min per mg protein in the presence of 0.5 mM NADPH. The apparent K_m for NADPH and NADH (66 and 266 μ M, respectively), the pH optimum for the reaction (6.9), and the cyanide insensitivity were similar to properties of plasma membrane-rich fractions of stimulated polymorphonuclear leukocytes studied by others. The activity of the phagocytic vesicles was trypsin sensitive. The specific superoxide-generating activity of macrophage phagocytic vesicles isolated from cells incubated up to 90 min with phagocytic particles remained constant.

Calcium in micromolar concentrations inhibited the NADPH-dependent O_2^- -generating activity of phagocytic vesicles. In a physiological ionic medium (100 mM KCl, 2.5 mM $MgCl_2$, 30 mM imidazole-HCl, pH 6.9), a maximal inhibition of O_2^- generation by phagocytic vesicles of 80% was observed at 40 μ M free Ca^{2+} . The half maximum inhibitory effect was at 0.7 μ M Ca^{2+} . Variations of the calcium concentration resulted in rapid and reversible alterations in O_2^- -forming activity. Preincubation of phagocytic vesicles in the presence of EGTA rendered their O_2^- generation rate in the presence of NADPH insensitive to alterations in the free calcium concentration. This desensitization by low EGTA concentrations (≤ 100 μ M) was reversible by the addition of excess calcium, but desensitization by high EGTA concentrations [...]

Find the latest version:

<https://jci.me/110000/pdf>



Effect of Calcium on Superoxide Production by Phagocytic Vesicles from Rabbit Alveolar Macrophages

P. DANIEL LEW and THOMAS P. STOSSEL, *Hematology-Oncology Unit, Massachusetts General Hospital, Department of Medicine, Harvard Medical School, Boston, Massachusetts 02114*

ABSTRACT Phagocytic vesicles from rabbit lung macrophages produced superoxide in the presence of NADH or NADPH. At 37°C, these vesicles generated 51 ± 7.8 nmol O_2^- /min per mg protein in the presence of 0.5 mM NADPH. The apparent K_m for NADPH and NADH (66 and 266 μ M, respectively), the pH optimum for the reaction (6.9), and the cyanide insensitivity were similar to properties of plasma membrane-rich fractions of stimulated polymorphonuclear leukocytes studied by others. The activity of the phagocytic vesicles was trypsin sensitive. The specific superoxide-generating activity of macrophage phagocytic vesicles isolated from cells incubated up to 90 min with phagocytic particles remained constant.

Calcium in micromolar concentrations inhibited the NADPH-dependent O_2^- -generating activity of phagocytic vesicles. In a physiological ionic medium (100 mM KCl, 2.5 mM $MgCl_2$, 30 mM imidazole-HCl, pH 6.9), a maximal inhibition of O_2^- generation by phagocytic vesicles of 80% was observed at 40 μ M free Ca^{2+} . The half maximum inhibitory effect was at 0.7 μ M Ca^{2+} . Variations of the calcium concentration resulted in rapid and reversible alterations in O_2^- -forming activity. Preincubation of phagocytic vesicles in the presence of EGTA rendered their O_2^- generation rate in the presence of NADPH insensitive to alterations in the free calcium concentration. This desensitization by low EGTA concentrations (≤ 100 μ M) was reversible by the addition of excess calcium, but desensitization by high EGTA concentrations (> 1 mM) was not reversible by the addition of calcium either in the presence or absence of purified rabbit lung macrophage or bovine brain calmodulins. Furthermore, trifluoperazine, a drug that inhibits calmodulin-stimulated reactions, did not alter the activity or the calcium sensitivity of the superoxide-generating system of sensitive phagocytic vesicles.

Peripheral plasma membrane vesicles (podosomes) prepared by gentle sonication of macrophages possessed an O_2^- -generating system with similar properties to those of phagocytic vesicles.

We conclude that the activated O_2^- -generating system of rabbit lung macrophages has its initial localization in the plasmalemma and undergoes subsequent internalization into phagocytic vesicles, where it can function for prolonged periods of time. Calcium at concentrations likely to exist in macrophage cytoplasm exerts a regulatory effect on the activated system.

INTRODUCTION

Mammalian phagocytes encountering objects that they recognize convert oxygen molecules to superoxide anions (O_2^-), hydrogen peroxide, and other substances important for microbicidal activity. Reduced pyridine nucleotide oxidase enzymes are the mediators of this respiratory reaction (1). The activation of oxygen metabolism in phagocytes appears to be a process distinct from that maintaining its activity, because experimental manipulations that inhibit activation of the respiratory activity of phagocytes do not necessarily inhibit the activity of the respiratory rate after activation (2, 3). In view of recent evidence that calcium concentrations in the micromolar range can control the activity of peripheral cytoplasmic proteins involved in phagocytosis (4), it is logical to inquire whether this ion also influences the oxygen metabolism of the cells.

In the present investigation, we report that phagocytic vesicles from rabbit alveolar macrophages contain a reduced pyridine nucleotide-linked O_2^- -generating system that is inhibited by concentrations of free calcium expected to occur at times in the cytoplasm.

METHODS

Isolation of macrophages. Cells were obtained from the lungs of New Zealand white rabbits by the procedure of Myrvik et al. (5), as described previously (6), 14–21 d after

Received for publication 7 July 1980 and in revised form 25 August 1980.

the animals received an intravenous injection of 1 ml of Freund's complete adjuvant to increase cell yields. The cells, suspended in 0.15 M NaCl solution, were washed three times at 4°C by centrifugation (250 g, 10 min), suspended in a modified Krebs-Ringer phosphate medium (130 mM NaCl, 4 mM KCl, 1.3 mM MgCl₂, 1 mM CaCl₂, and 10 mM sodium phosphate buffer, pH 7.4, hereafter designated as "medium"), and kept at 0°C until use. Yields ranged from 3 × 10⁸ to 2.4 × 10⁹ cells/rabbit, of which >90% were macrophages.

Serum preparation. Rabbit serum was obtained from freshly clotted blood derived from anesthetized animals by cardiac puncture. Sera were used on the same day or stored at -70°C until use.

Purification of phagocytic vesicles. This technique, described in detail elsewhere (6), is outlined briefly. Diisodecylphthalate (1 ml) (practical grade, Matheson, Coleman and Bell, E. Rutherford, N. J.) was added to 3 ml of medium containing 10 mg/ml of *Escherichia coli* lipopolysaccharide 026:B6 (Difco Laboratories, Detroit, Mich.) and emulsified by sonication. The lipopolysaccharide-coated diisodecylphthalate particles were incubated with fresh or freshly thawed rabbit serum for 20 min at 37°C, during which time an opsonic fragment of the third component of complement was deposited on them (7) (opsonized particles). Cells were suspended in medium and warmed to 37°C. Suspensions of opsonized particles (20%, vol/vol) and cells (5%, vol/vol) were mixed and agitated gently in a shaking bath at 37°C for various amounts of time, during which the cells ingested the opsonized particles. The cells were washed once in cold 0.15 M NaCl, suspended in ice-cold deionized water, and immediately centrifuged at 4°C. During this treatment, which made subsequent homogenization easier, the cells swelled but did not break. The cell pellets were suspended in equal volume of ice-cold 0.34 M sucrose solution containing 5 mM EGTA and 30 mM imidazole-HCl, pH 7.4. The cells were homogenized in this medium in a 40 ml Dounce homogenizer with a tightly fitting pestle. The progress of cell rupture was monitored as the percentage of free nuclei visible by phase-contrast microscopy. About 50 strokes were required to break 90% of the cells. In most experiments, Ca²⁺ (CaCl₂) was then added to a final concentration of 4–5 mM. As explained in Results, the concentration of added calcium was important in the properties of the subsequently purified phagocytic vesicles. EGTA was used at this step because it increases considerably the efficiency of homogenization. The homogenate was transferred to a 30-ml Sorvall centrifuge tube (DuPont Instruments-Sorvall Biomedical Div., DuPont Co., Newtown, Conn.), carefully overlaid by means of a peristaltic pump (Pharmacia Fine Chemicals, Piscataway, N. J.) with 2 vol of a solution containing 0.25 M sucrose, 30 mM imidazole-HCl, pH 6.9, and concentrations of calcium or EGTA, and centrifuged for 60 min at 150,000 g. The floating white layer, representing isolated phagocytic vesicles, was then carefully removed and kept at 0°C until it was further diluted for use. The protein concentration of phagocytic vesicles was determined by the Folin procedure (8).

Preparation of peripheral hyaline vesicles of macrophages (podosomes). This preparation, containing a mixture of right-side-out and inside-out blebs of plasma membrane and virtually free of organelles, was prepared by gentle sonication of a warm suspension of cells according to previously published techniques (9). In brief, macrophages (10⁸ cells) were incubated for 4 min at 37°C in glass conical centrifuge tubes with or without 50 μg of digitonin in 5 ml of medium with 5 mM glucose. After incubation, the suspension was placed in a 37°C temperature sonication bath (Branson Ultrasonic Cleaner, model B 12, purchased from Heat Systems-Ultrasonics, Inc., Plainview, N. Y.) for 8–10 s. The vesicles were separated from

residual cell bodies by sedimenting the latter (250 g, 10 min) and pelleting the resultant supernate. The pellet was washed in a solution containing 0.25 M sucrose, 30 mM imidazole-HCl, pH 6.9, resuspended in the same solution, and kept at 0°C until used.

Assay of O₂⁻ production by phagocytic vesicles and podosomes. O₂⁻ production by phagocytic vesicles or podosomes was assayed by examining the rate of O₂⁻-dependent cytochrome *c* reduction in the presence of reduced pyridine nucleotides in a Gilford spectrophotometer 250 (Gilford Instrument Laboratories, Inc., Oberlin, Ohio) with a controlled temperature regulator. This assay is based on methods used by previous investigators (2).

The standard assay system contained the following, in a total volume of 0.8 ml: cytochrome *c* type VI (50 nmol), NADPH (0.5 μmol) or NADH (2 μmol) (all from Sigma Chemical Co., St. Louis, Mo.), in a medium containing KCl (0.1 mmol), MgCl₂ (2.5 μmol), and imidazole-HCl (24 μmol) pH 6.9. In some experiments, sucrose (0.25 mmol) replaced KCl. Various concentrations of CaCl₂ and EGTA were also added. In addition, superoxide dismutase (30 μg) was added to the reference compartments of the spectrophotometer. The reactions were initiated by the addition of 0.2 ml of phagocytic vesicles or podosomes to both the sample and reference cuvettes, and the absorbance changes at 550 nm were monitored by double-beam spectrophotometry. In addition, podosomes were tested in the presence of KCN (0.5 μmol) (10). An extinction coefficient of 19,500 cm⁻¹ was used for (reduced minus oxidized) cytochrome *c* (11). The sensitivity of this assay is such that it will detect rates of O₂⁻ production as low as 0.5 nmol/min (2).

For experiments in which enzyme activity was measured as a function of free calcium concentration, phagocytic vesicles were washed a second time in a medium containing 0.5 mg/ml bovine serum albumin, 0.25 M sucrose, 30 mM imidazole-HCl, pH 6.9, 2.5 mM MgCl₂, and 100 μM CaCl₂. Samples of the washed vesicles were tested in the medium described above, in solutions containing various concentrations of calcium and EGTA. These experiments were performed at 25°C because the equilibrium constants for EGTA have been established at 20–25°C (12). The free Ca²⁺ concentration of solutions was calculated by means of the computer program of Perrin and Sayce as described by Potter and Gergeley (13). Trifluoperazine dihydrochloride (provided by Dr. Eileen Gallagher, SmithKline and French, Philadelphia, Pa.) was preincubated with vesicles in the presence of 100 μM calcium at 0°C for 1 h.

In some experiments, phagocytic vesicles and podosomes were treated with trypsin. Bovine pancreatic trypsin, type IX, treated with diphenyl carbonyl chloride to inactivate chymotrypsin (Sigma Chemical Co.), was dissolved at a concentration of 10 mg/ml in 100 mM KCl and 30 mM imidazole-HCl, pH 6.9. Vesicles were warmed to room temperature, and the trypsin solution was added to make a final concentration of 500 μg/ml. After 5 min of incubation, soybean trypsin inhibitor (Sigma Chemical Co.) in the same buffer solution was added to a final concentration of 500 μg/ml. The suspension was immediately placed on ice after mixing. Controls contained soybean trypsin inhibitor added simultaneously with trypsin or no additions at all.

Assay of O₂⁻ production of intact macrophages. Spectrophotometric measurements of O₂⁻-dependent cytochrome *c* reduction were performed with a continuous assay, as described above, at 25°C. The sample cuvette contained cytochrome *c* (50 nmol) and macrophages (4 × 10⁶) with or without digitonin (10 μg; Fisher Scientific Co., Pittsburgh, Pa.) in 1 ml of medium with 5 mM glucose. The lag time for activation, as well as the rate of activity of O₂⁻ generation, was calculated as previously described by Cohen et al. (2, 3).

Whole macrophages were treated with trypsin by the same procedure described for phagocytic vesicles. Analysis of basal oxidation of [1-¹⁴C]glucose by intact macrophages was performed as previously described (9).

Purification of macrophage calmodulin. Macrophages were suspended in an equal volume of ice-cold 0.34 M sucrose solution containing 5 mM EGTA, 5 mM dithiothreitol, 1 mM phenylmethylsulfonyl fluoride, 0.25 mg/ml of α -1-antitrypsin, 0.25 mg/ml of soybean trypsin inhibitor, and 20 mM imidazole-HCl buffer, pH 7.4, and homogenized as described above. The homogenate was centrifuged at 100,000 g for 1 h and the supernatant fluid was dialyzed overnight against 10 mM Hepes, 1 mM CaCl₂, pH 7, and fractionated by affinity chromatography in a fluphenazine-agarose column. Fluphenazine (provided by Dr. Salvador Luciano, E. R. Squibb and Sons, Princeton, N. J.) was coupled to Sepharose activated by the bisoxarine procedure as described by Charbonneau and Cormier (14). The activity of activator-deficient phosphodiesterase (provided by Dr. Robert Wallace and Dr. Wai Yui Cheung, St. Jude Children's Research Hospital, Memphis, Tenn.) in the presence of column fractions was determined as described by Strewler et al. (15). The bulk of the protein passed directly through the column and had no phosphodiesterase-stimulating activity. A buffer solution containing 10 mM Pipes, pH 7, 1 mM CaCl₂, and 0.5 M NaCl eluted a small peak of protein that also had no phosphodiesterase activator activity. A 10 mM Pipes solution, pH 7, containing 10 mM EGTA and 0.5 M NaCl, eluted a small protein peak that activated phosphodiesterase in a calcium-dependent manner. This activity was resistant to boiling. Polyacrylamide gel electrophoresis of this fraction in sodium dodecyl sulfate revealed a homogenous polypeptide that comigrated with bovine brain calmodulin (provided by Dr. Wallace and Dr. Cheung). The purified calmodulins were incubated with phagocytic vesicles that had been homogenized and suspended in various concentrations of EGTA at a concentration of 20 μ g/ml in the presence of 100 μ M free Ca²⁺ for 1 h at 0°C.

RESULTS

O₂⁻ production by intact macrophages, macrophage podosomes, and macrophage phagocytic vesicles. As previously reported, rabbit alveolar macrophages stimulated with digitonin released O₂⁻ in the absence of reduced pyridine nucleotides (3). After addition of digitonin to a cuvette containing macrophages and cytochrome *c*, there was an initial period with no change in A₅₅₀, then a gradual rise until a linear rate was achieved. At 25°C, the lag time was 72 ± 6 s and the rate of activity was 0.51 ± 0.06 nmol O₂⁻/10⁶ cells per min (mean ± SEM for three different cell preparations).

Trypsinization of the macrophages (1,000 μ g/2 × 10⁷ cells per ml for 5 min at room temperature) did not detectably alter the O₂⁻ production rate of the cells. The addition of trifluoperazine, an agent that inhibits calmodulin-dependent reactions in the presence of calcium (16), inhibited the rate as well as the lag time of O₂⁻ production (Fig. 1), whether added before or after the activation period, and the effect of the drug was complete <1 min after the addition of the drug. Similar results were obtained using opsonized zymosan to stimulate O₂⁻ release (not shown). Removal of




FIGURE 1 The effect of trifluoperazine on the rate (●) and lag period (○) of O₂⁻ generation by digitonin-stimulated rabbit alveolar macrophages. The indicated amounts of trifluoperazine were added at zero time and O₂⁻ production measured as described in the text.

trifluoperazine by washing the cells did not restore the O₂⁻ release rate to control values over a wide range of concentrations. Trifluoperazine up to 76 μ M did not inhibit the oxidation rates of [1-¹⁴C]glucose of resting macrophages but impaired ¹⁴CO₂ release at higher concentrations (50% decrease at 100 μ M).

Phagocytic vesicles purified from macrophages that had ingested opsonized lipopolysaccharide-coated diisododecylphthalate particles were examined for their ability to generate O₂⁻ in the presence of reduced pyridine nucleotides. At 25°C, O₂⁻ production was constant with time for about 4 min, and under these conditions continuous production was detectable for >30 min. Both NADH and NADPH promoted O₂⁻ generation by the vesicles. The effect of the reduced pyridine nucleotides on the rate of O₂⁻ production was not additive (not shown). O₂⁻ generation by vesicles was not detectable in the absence of NADPH or NADH. The apparent K_m for NADPH was 64 μ M, with a maximum oxidation rate (V_{max}) of 29 nmol O₂⁻/min per mg protein, whereas the apparent K_m for NADH was 266 μ M with a V_{max} of 19 nmol O₂⁻/min per mg protein (Fig. 2a). In contrast to NADPH, NADH elicited substantial cytochrome *c* reduction in the presence of superoxide dismutase (up to 65% of the reduction rate in the absence of superoxide dismutase in NADH compared with up to 14% in NADPH). Therefore, most of our experiments were done in the presence of NADPH. The O₂⁻-generating activity in 0.5 mM NADPH was directly proportional to the




FIGURE 2 Rates of O_2^- production by phagocytic vesicles from rabbit alveolar macrophages incubated at 25°C in a medium containing 0.25 M sucrose, 30 mM imidazole-HCl, pH 6.9, 100 μ M EGTA, 2.5 mM $MgCl_2$, 50 nmol cytochrome *c*, and 0.5 μ mol NADPH, unless otherwise specified. (A) NADH and NADPH oxidase kinetics: O_2^- production was measured at varying concentrations of NADH (○) (100–2,000 μ M) and NADPH (●) (16–1,500 μ M). $1/V$ equals (nanomoles O_2^- generated per milligram vesicle protein per minute)⁻¹; $1/S$ equals (millimoles NADPH or NADH per liter)⁻¹. (B) pH optimum of O_2^- production. The reaction was carried out in a medium containing 100 mM KCl and 30 mM imidazole-HCl at various pH values indicated.

concentrations of added vesicle protein between 10 and 132 μ g/ml. The pH optimum of this reaction was 6.9 (Fig. 2b).

Table I summarizes the effects of various conditions and additions on the initial rate of superoxide forma-

TABLE I
Effect of Temperature and Various Compounds on O_2^- Production by Phagocytic Vesicles from Rabbit Alveolar Macrophages

Treatment	O_2^- production nmol/min/mg protein
37°C	51±7.8 (5)
25°C	25±3.08 (6)
KCN, 2 mM*	25±2 (3)
Azide, 5 mM	24±3 (3)
ATP, 2.5 mM	22.75±1 (2)
N-Ethyl maleimide, 1 mM	17.8±1.5 (2)
Trypsin†	0 (3)

The activity was tested in a medium containing 0.25 M sucrose, 2.5 mM $MgCl_2$, 30 mM imidazole-HCl, pH 6.9, and 100 μ M EGTA at 25°C, unless otherwise specified. All values are mean±SEM. Number of preparations is in parentheses.

* For this experiment, the reference cuvette contained 80 μ g of superoxide dismutase to insure complete detection of the O_2^- formed.

† 500 μ g for 5 min at room temperature.

tion by vesicles in the presence of NADPH. The activity was diminished by 50% at 25°C when compared with 37°C. The superoxide-generating activity was not significantly altered by addition of azide, cyanide, or Mg^{2+} -ATP. N-Ethyl maleimide decreased the activity by 30%. Trypsinization rapidly and totally abolished this activity. If the vesicles were kept at ice temperature for several hours, a decrease in their activity was observed (about 30% after 3 h). It was not possible to compare the specific activity of vesicles with that of homogenates from which they were derived, because of the presence of a powerful superoxide dismutase-resistant, cytochrome *c*-reducing activity, as well as the presence of large concentrations of superoxide dismutase (whether tested in the presence or absence of 0.5 mM KCN) (17).

As shown in Fig. 3, the specific activity of O_2^- -generating activity in 0.5 mM NADPH of phagocytic vesicles isolated from cells incubated with phagocytizable particles for up to 90 min was relatively constant.

Podosomes from resting and digitonin-stimulated macrophages were examined for their ability to generate O_2^- in the presence of NADPH. In the presence of 0.5 mM NADPH, podosomes obtained from resting cells generated 0.4 nmol O_2^- /mg protein per min, whereas podosomes made from macrophages stimulated with digitonin for 4 min generated 3.2 nmol O_2^- /mg protein per min. O_2^- generation by podosomes was not detectable in the absence of NADPH. Trypsinization inactivated the NADPH-dependent superoxide generation of podosomes.




FIGURE 3 Specific activity of the NADPH-dependent O_2^- production by phagocytic vesicles isolated from cells incubated with particles for varying lengths of time. Rabbit alveolar macrophages were incubated with opsonized lipopolysaccharide-coated oil particles for the times noted, and phagocytic vesicles were made as described in the text. NADPH-dependent O_2^- production, (●); total protein, (○).

Effect of calcium on the O_2^- -forming activity of phagocytic vesicles and podosomes. The O_2^- -generating activity of vesicles was very sensitive to the concentration of free calcium in the reaction media, provided that calcium was present in all the steps of the purification of the phagocytic vesicles (see below). Under physiologic ionic conditions (100 mM KCl, 2.5 mM $MgCl_2$, 30 mM imidazole HCl, pH 6.9), 0.1 mM free Ca^{2+} inhibited the initial rate of superoxide formation in the presence of NADPH by $77 \pm 2\%$ (mean \pm SEM for six separate preparations) when compared with the activity measured at $0.016 \mu M$ free calcium.

Fig. 4 shows the effect of the free Ca^{2+} concentration on the rate of O_2^- production. Inhibition of activity was directly proportional to the logarithm of the free Ca^{2+} concentration ($r=0.94$ by linear regression analysis), and the maximum inhibition was observed at a concentration of $40 \mu M$ free Ca^{2+} . The half maximum inhibitory effect was at a concentration of $0.7 \mu M$ free Ca^{2+} .

If the vesicles were suspended in 0.25 M sucrose instead of 0.1 M KCl, the specific activity of the enzyme was not altered in the presence of low calcium concentrations. However, the inhibitory effect of calcium was less marked than in 0.1 M KCl solution. The half maximum inhibitory effect was at $3.5 \mu M$ free Ca^{2+} , and the inhibition of the O_2^- formation rate in 0.1 mM free Ca^{2+} was $48.44 \pm 4.89\%$ (mean \pm SEM for six separate vesicle preparations). Phagocytic vesicles obtained from cells broken in the absence of added EGTA had calcium sensitivity indistinguishable from

that of vesicles obtained from cells homogenized in EGTA, provided that the EGTA was neutralized after cell rupture by addition of $CaCl_2$. The importance of the calcium concentration for maintaining calcium sensitivity is described further below.

As shown in Fig. 5, the effect of calcium on superoxide production by vesicles in a 0.1 M KCl solution and 0.5 mM NADPH was rapid and reversible. In this experiment, the enzyme activity was continuously monitored in the presence of $40 \mu M$ calcium. $400 \mu M$ EGTA was then added to both compartments of the spectrophotometer, making the final free Ca^{2+} concentration $0.035 \mu M$. After a lag period of <1 min, a three-fold stimulation of the superoxide generation rate was observed. Several cycles of stimulation and inhibition under continuous monitoring could be observed, but with a progressive decrease in the measured activities. Similar effects were obtained using other calcium-chelating agents such as EDTA or ATP. The onset of the stimulatory effect of decreasing the calcium concentration of superoxide production was identical after prolonged incubation of the vesicles either on ice (up to 2 h) or at room temperature (up to 30 min), provided that the gradual decrease in the specific activity previously mentioned was taken into account. Furthermore, if the enzyme activity in the presence of calcium was continuously monitored until no further O_2^- was produced, the addition of EGTA stimulated new production of O_2^- . Thus, the inhibition of activity by calcium is not due to proteolytic destruction or to irreversible denaturation of the enzyme system. By con-




FIGURE 4 O_2^- production by phagocytic vesicles from rabbit alveolar macrophages as a function of free Ca^{2+} concentration. Phagocytic vesicles were washed in a medium containing 0.5 mg/ml albumin, 0.25 M sucrose, 30 mM imidazole-HCl, pH 6.9, 2.5 mM $MgCl_2$, and 100 μM $CaCl_2$, and assayed for O_2^- production in the medium described in Fig. 1 (except that 100 mM KCl replaced the sucrose) at various concentrations of Ca and EGTA at 25°C. The results are expressed as percent maximal activity that was obtained at the lowest Ca^{2+} tested. A maximal inhibition of 80% was detected at 40 μM free Ca^{2+} , and half maximal inhibition was documented at 0.7 μM free Ca^{2+} . The line was plotted by linear regression analysis ($r = 0.94$). Free Ca^{2+} concentrations were calculated as described in the test.




FIGURE 5 Time-course and reversibility of the effect of calcium on O_2^- production by phagocytic vesicles. O_2^- production by phagocytic vesicles was measured in a medium containing 100 mM KCl, 30 mM imidazole-HCl, pH 6.9, 2.5 mM $MgCl_2$, 40 μM free Ca^{2+} , 0.5 mM NADPH, 50 μM cytochrome *c*, and 30 μg superoxide dismutase in the reference compartment at 25°C. After 4 min, 400 μM EGTA was added to both compartments of the spectrophotometer.

trast, variations of Mg^{+2} concentration did not result in variations of enzyme activity.

Table II describes experiments that showed that calcium was required to maintain the calcium sensitivity of phagocytic vesicles. If vesicles were first incubated in buffers containing increasing concentrations of EGTA relative to calcium, there was a decrease in the subsequent calcium sensitivity, and the activity tested at either high or low calcium concentrations approached similar values. Reincubation of these desensitized phagocytic vesicles in the presence of higher calcium concentration for 10–20 min at 0°C restored calcium sensitivity to the enzyme activity (Table II, lines 1 and 3), although calcium sensitivity could not be restored to vesicles originally incubated in very high EGTA concentrations (Table II, line 5). These vesicles initially exposed to very high EGTA concentrations did not regain calcium sensitivity even if they were subsequently incubated with calcium for long periods of time (up to 600 μM free Ca^{2+} for 3 h at 0°C). If calcium was not added to the homogenate after cell rupture, permanent desensitization also occurred.

The effects of EGTA on desensitization could be

TABLE II
Effect of EGTA, Trifluoperazine, and Calmodulin on the Specific Activity and Calcium Sensitivity of NADPH-dependent O₂⁻ Production by Purified Phagocytic Vesicles

First incubation		Second incubation			Conditions of assay for O ₂ ⁻ generation	
CaCl ₂	EGTA	Ca ²⁺ (100 μM)	Trifluoperazine (250 μM)	Calmodulin (20 μM)	Ca ²⁺ * (100 μM)	Ca ²⁺ * (0.016 μM)
μM	μM					
20	0	+	-	-	4.7±1.5	26.0±3.0
20	0	+	+	-	4.2±1.3	24.0±2.0
20	100	+	-	-	7.0±0.8	29.0±2
20	100	+	-	+	8.0±1.0	31.0±2.0
20	1,000	+	-	-	14.2±1.5	20.3±1.8
20	1,000	+	-	+	13.37±2.0	19.1±2.2

Purified phagocytic vesicles were suspended in the presence of calcium/EGTA mixtures to give high or low free calcium concentrations as indicated in the left column for 5 min at 0°C (first incubation). Calcium was then added in excess (final free {Ca²⁺} 100 μM) in the presence or absence of calmodulin or trifluoperazine as indicated in the middle column (second incubation). After 1 h at 0°C, samples were tested for O₂⁻ production in the medium described in Fig. 5 at the final calcium concentrations indicated in the right column. The samples incubated in the presence of trifluoperazine were also tested in the presence of trifluoperazine (250 μM).

* Free calcium concentration calculated as described in the text.

compatible with the dissociation of a calcium-binding protein from the phagocytic vesicles at low calcium concentrations. Therefore, as shown also in Table II, several experiments were performed to see whether the observed effect was mediated by the ubiquitous calcium-binding protein, calmodulin. First, vesicles prepared in the presence of calcium were incubated in the presence of trifluoperazine and calcium (Table II, line 2). In contrast to the O₂⁻ release by intact cells, incubation of vesicles in the presence of 250 μM trifluoperazine did not decrease the specific activity or the calcium sensitivity of the O₂⁻-generating activity of sensitive vesicles. Addition of rabbit alveolar macrophage or bovine brain calmodulin did not restore the calcium sensitivity to desensitized vesicles and did not alter their O₂⁻ production rates (Table II, line 6).

As found with phagocytic vesicles, the O₂⁻-generating activity of podosomes obtained from digitonin-stimulated macrophages was inhibited by micromolar concentrations of calcium. In a medium containing 0.25 M sucrose, 2.5 mM MgCl₂, and 30 mM imidazole-HCl, pH 6.9, the activity in the presence of 0.1 mM free Ca²⁺ was 52% of that measured in the presence of 0.016 μM Ca²⁺. We were unable to activate podosomes of unstimulated macrophages by extensive washing in EGTA or by addition of Mg-ATP and extracts of macrophages in the presence of calcium.

DISCUSSION

Alveolar macrophages exposed to a variety of stimuli, including opsonized zymosan, phorbol myristate acetate, *N*-formylmethionyl peptides, concanavalin A, and digitonin, generate superoxide after a lag period (3, 18–20). We have confirmed these previous observations and extended them to show that the O₂⁻-generating system of intact macrophages is insensitive to trypsin and inhibited by trifluoperazine.

Phagocytic vesicles purified from macrophages that have ingested opsonized lipopolysaccharide-coated diisododecylphthalate particles possess an NAD(P)H-dependent, O₂⁻-generating system. It shares several common features with the extensively characterized activity described in several preparations of polymorphonuclear leukocytes, including similar ranges of reduced pyridine nucleotide requirements, pH optimum, and insensitivity to CN⁻ (10, 21–23).

It is possible that phagocytic vesicles might not be representative of the composition of plasma membrane as a whole either because of fusion of the vesicle membranes with lysosomal membranes or because of lateral segregation of membrane molecules out of the vacuoles during phagocytosis. However, we found an NADPH-dependent, O₂⁻-generating system with the features of the phagocyte vesicle system, although of lower specific activity, in peripheral plasma

membrane vesicles prepared by gentle sonication of macrophages (podosomes). These vesicles are peripheral blebs of macrophage plasma membrane, which are representative of the total cell plasma membrane with respect to the marker adenylate cyclase and externally labeled proteins (9). Electron micrographs of these vesicles contain some empty sacs that have filaments on their outer surfaces (9). Although the bulk of these blebs appears to be rightside-out vesicles, the O_2^- generation presumably takes place in the small subpopulation of inverted vesicles, because in contrast to intact macrophages, the O_2^- -generating activity by podosomes as well as phagocytic vesicles was trypsin sensitive and dependent upon the addition of NAD(P)H, which does not permeate intact membranes. The present findings are further evidence that the activated oxidase enzyme is present in the plasmalemma (24–26) and internalized into the phagocytic vacuole during phagocytosis.

When neutrophils are incubated in the presence of opsonized particles and the particulate fraction or phagocytic vesicles isolated at various times after initiation of phagocytosis, a rapid decrease in specific activity occurs after incubations >10 min (23, 27). By contrast, the phagocytic vesicles isolated from macrophages maintained the same specific activity even 90 min after initiation of phagocytosis. Thus, once internalized from the plasma membrane into the phagocytic vesicle, the active enzyme can apparently function for prolonged periods of time. This ability to sustain oxidative activity is important for macrophages that reside for long times in inflammatory sites (28).

The most novel finding of this study was the discovery of an inhibitory effect of calcium on the O_2^- -generating activity of macrophage phagocytic vesicles and podosomes. Although the free Ca^{2+} concentration in the cytoplasm of macrophages has not yet been measured, in other cells, particularly muscle and nerve, the free Ca^{2+} concentrations can fluctuate between 10 nM and 10 μ M, according to the state of stimulation (29, 30). This range of calcium concentrations influenced O_2^- -generating activity by macrophage phagocytic vesicles in NADPH under physiological ionic conditions (100 mM KCl, 2.5 mM $MgCl_2$, pH 6.9). The effect was rapid and reversible, conditions necessary for an effective control of enzyme activity.

The inhibitory effect of calcium that we have observed is exerted upon the activated enzyme that is present in phagocytic vesicles. The role of calcium, if any, in the activation process of the O_2^- -generating system remains to be determined. Podosomes obtained from digitonin-stimulated macrophages had much higher rates of O_2^- production than podosomes obtained from unstimulated macrophages. However, we were unable to activate podosomes of unstimulated macrophages by extensive washing in EGTA or by

addition of Mg-ATP and extracts of macrophages in the presence of calcium.

It is not clear from the present experiments if the effect of calcium is directly on the membrane-bound enzyme or mediated by a calcium-binding component that is washed away in the presence of EGTA. Calmodulin, a ubiquitous calcium-binding protein that acts as a calcium regulatory subunit of many intracellular enzymes (31), is appealing to be this component. However, the results of several experiments argue against a direct involvement of calmodulin in our system. First, purified calmodulin did not restore calcium sensitivity to vesicles that had lost it after exposure to EGTA. Second, trifluoperazine, a drug that binds to calmodulin with high affinity in the presence of calcium, did not change the specific activity nor the calcium sensitivity of O_2^- formation by phagocytic vesicles in the presence of 0.5 mM NADPH. By contrast, trifluoperazine at lower concentrations increased the activation period and inhibited the O_2^- -generating system of intact stimulated macrophages whether added before or after establishment of the maximal rates of O_2^- production.

We recently reported that the plasma membrane of macrophages contains a Mg-ATP-dependent calcium pump that transports calcium from the cytoplasmic to the external membrane surface against an electrochemical gradient. The activity of the pump is stimulated by calmodulin and inhibited by trifluoperazine (32). A slight decrease in the activity of this calcium pump activity could possibly cause an increase of the cytoplasmic free Ca^{2+} concentration. Therefore, some of the inhibitory effect of trifluoperazine on the O_2^- -generating system of intact cells could be due to increased calcium at the inner side of the plasma membrane. However, the trifluoperazine effect on whole cells could be due to other calmodulin-mediated reactions, to competition with NADPH for the oxidase, or to nonspecific or toxic effects (31, 33). In any case, we speculate that, in order to obtain maximal O_2^- -generating rates, the calcium concentration has to decrease in the membrane region containing the activated enzyme.

ACKNOWLEDGMENTS

We are indebted to Mrs. Esther Campos and Ms. Kitty G. Brown for their valuable secretarial assistance.

This work was supported by U. S. Public Health Service training grant CA 09321 and grants from the Swiss Foundation for Scientific Research, Swiss Foundation for Scholarships in Medicine and Biology, and the Swiss Society of Internal Medicine.

REFERENCES

1. Babior, B. M. 1978. Oxygen-dependent microbial killing by phagocytes. *N. Engl. J. Med.* **298**: 659–668; 721–725.
2. Cohen, H. J., and M. E. Chovaniec. 1978. Superoxide

- generation by digitonin-stimulated guinea pig granulocytes. A basis for a continuous assay for monitoring superoxide production and for the study of the activation of the generating system. *J. Clin. Invest.* **61**: 1081-1087.
3. Cohen, H. J., and M. E. Chovaniec. 1978. Superoxide production by digitonin-stimulated guinea pig granulocytes. The effects of *N*-ethylmaleimide, divalent cations, and glycolytic and mitochondrial inhibitors on the activation of the superoxide generating system. *J. Clin. Invest.* **61**: 1088-1096.
 4. Yin, H. L., and T. P. Stossel. 1979. Control of cytoplasmic actin gel-sol transformation by gelsolin, a calcium-dependent regulatory protein. *Nature (Lond.)* **281**: 583-586.
 5. Myrvik, Q. N., E. S. Leake, and B. Fariss. 1961. Studies on pulmonary alveolar macrophages from the normal rabbit: a technique to procure them in a high state of purity. *J. Immunol.* **86**: 128-136.
 6. Stossel, T. P., R. J. Mason, T. D. Pollard, and M. Vaughan. 1972. Isolation and properties of phagocytic vesicles. II. Alveolar macrophages. *J. Clin. Invest.* **51**: 604-614.
 7. Stossel, T. P., R. J. Field, J. D. Gitlin, C. A. Alper, and F. S. Rosen. 1975. The opsonic fragment of the third component of human complement (C3). *J. Exp. Med.* **141**: 1329-1347.
 8. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**: 265-275.
 9. Davies, W. A., and T. P. Stossel. 1977. Peripheral hyaline blebs (podosomes) of macrophages. *J. Cell Biol.* **75**: 941-955.
 10. Cohen, H. J., M. E. Chovaniec, and W. A. Davies. 1980. Activation of the guinea pig granulocyte NAD(P)H-dependent superoxide generating enzyme: localization in a plasma membrane enriched particle and kinetics of activation. *Blood.* **55**: 355-363.
 11. Chance, B., and G. R. Williams. 1956. The respiratory chain and oxidative phosphorylation. In *Advances in Enzymology*. F. F. Nord, editor. Interscience Publishers, Inc., New York.
 12. Sillen, L. G., and A. E. Martell. 1964. *Stability Constants of Metal Ion Complexes*. The Chemical Society, Burlington House, London. 2nd edition.
 13. Potter, J. D., and J. Gergely. 1975. The calcium and magnesium binding sites on troponin and their role in the regulation of myofibrillar adenosine triphosphatase. *J. Biol. Chem.* **250**: 4628-4633.
 14. Charbonneau, H., and M. J. Cormier. 1979. Purification of plant calmodulin by fluphenazine-sepharose affinity chromatography. *Biochem. Biophys. Res. Commun.* **90**: 1039-1047.
 15. Stewler, G. J., V. C. Manganiello, and M. Vaughan. 1978. Phosphodiesterase activator from rat kidney cortex. *J. Biol. Chem.* **253**: 390-394.
 16. Levin, R. M., and B. Weiss. 1977. Binding of trifluoroperazine to the calcium-dependent activator of cyclic nucleotide phosphodiesterase. *Mol. Pharmacol.* **13**: 695-697.
 17. Rister, M., and R. O. Baehner. 1976. A comparative study of superoxide dismutase activity in polymorphonuclear leukocytes, monocytes, and alveolar macrophages of the guinea pig. *J. Cell. Physiol.* **87**: 345-356.
 18. Drath, D. B., and M. L. Karnovsky. 1975. Superoxide production by phagocytic leukocytes. *J. Exp. Med.* **141**: 257-262.
 19. Boxer, L. A., G. Ismail, J. M. Allen, and R. L. Baehner. 1979. Oxidative metabolic responses of rabbit pulmonary alveolar macrophages. *Blood.* **53**: 486-491.
 20. Holian, A., and R. P. Daniele. 1979. Stimulation of oxygen consumption and superoxide anion production in pulmonary macrophages by *N*-formyl methionyl peptides. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* **108**: 47-50.
 21. Babior, B. M., J. T. Curnutte, and B. J. McCurrich. 1976. The particulate superoxide-forming enzyme from human neutrophils. *J. Clin. Invest.* **58**: 989-996.
 22. Badwey, J. A., and M. L. Karnovsky. 1979. Production of superoxide and hydrogen peroxide by an NADH oxidase in guinea pig polymorphonuclear leukocytes. Modulation by nucleotides and divalent cations. *J. Biol. Chem.* **254**: 11530-11537.
 23. Cohen, H. J., P. E. Newburger, and M. E. Chovaniec. 1980. NAD(P)H-dependent superoxide production by phagocytic vesicles from guinea pig and human granulocytes. *J. Biol. Chem.* **255**: 6584-6588.
 24. Briggs, R. T., D. B. Drath, M. L. Karnovsky and M. J. Karnovsky. 1975. Localization of NADH oxidase on the surface of human polymorphonuclear leukocytes by a new cytochemical method. *J. Cell Biol.* **67**: 566-586.
 25. Goldstein, I. M., M. Cerqueira, S. Lind, and M. B. Kaplan. 1977. Evidence that the superoxide-generating system of human leukocytes is associated with the cell surface. *J. Clin. Invest.* **59**: 249-254.
 26. Dewald, B., M. Baggiolini, J. T. Curnutte, and B. M. Babior. 1979. Subcellular localization of the superoxide-forming enzyme in human neutrophils. *J. Clin. Invest.* **63**: 21-29.
 27. Jandl, R. C., J. A. Schwartz, L. Borges-DuBois, R. S. Kipnes, B. J. McMurrich, and B. M. Babior. 1978. Termination of the respiratory burst in human neutrophils. *J. Clin. Invest.* **61**: 1176-1185.
 28. Nathan, C., N. Nogueira, C. Jaunqbhanich, J. Ellus, and Z. Cohn. 1979. Activation of macrophages *in vivo* and *in vitro*. Correlation between hydrogen peroxide release and killing of trypanosoma cruzi. *J. Exp. Med.* **149**: 1056-1068.
 29. Ashley, C. C. 1978. Calcium ion regulation in barnacle muscle fiber and its relation to muscle development. *Ann. N. Y. Acad. Sci.* **307**: 308-329.
 30. Rasmussen, H., and M. C. Gustin. 1978. Some aspects of the hormonal control of cellular calcium metabolism. *Ann. N. Y. Acad. Sci.* **307**: 391-401.
 31. Cheung, W. Y. 1980. Calmodulin plays a pivotal role in cellular regulation. *Science (Wash. D.C.)* **207**: 19-27.
 32. Lew, D. P. and T. P. Stossel. 1980. Calcium transport by macrophage plasma membranes. *J. Biol. Chem.* **255**: 5841-5846.
 33. Cohen, H. J., M. E. Chovaniec, and S. E. Ellis. 1980. Chlorpromazine inhibition of granulocyte superoxide production. *Blood.* **56**: 23-29.