

Mono- and dihydroxyeicosatetraenoic acids alter calcium homeostasis in rabbit neutrophils.

P H Naccache, ... , P Borgeat, E J Goetzl

J Clin Invest. 1981;**67**(5):1584-1587. <https://doi.org/10.1172/JCI110191>.

Research Article

Products of the lipoxygenation of arachidonic acid that express neutrophil chemotactic activity were examined in vitro for effects on the uptake of ^{45}Ca by rabbit peritoneal neutrophils. At optimally chemotactic concentrations, 5- and 11-hydroperoxyeicosatetraenoic acid, 11- and 12L-hydroxyeicosatetraenoic acid, and leukotriene B₄ enhanced ^{45}Ca uptake within 1 min by a mean of 212-694% of the values for control neutrophils incubated in buffer alone, as compared with 75% for 5(S)-hydroxyeicosatetraenoic acid and no effect for 15-hydroperoxyeicosatetraenoic acid and 15-hydroxyeicosatetraenoic acid. The approximate rank of potency of the factors stimulating ^{45}Ca uptake was similar to that observed for chemotaxis. Leukotriene B₄, in addition to stimulating the rate of ^{45}Ca uptake into rabbit neutrophils, also displaced intracellular calcium. The net result of the leukotriene B₄-induced changes in calcium homeostasis was to elevate transiently the intracellular level of exchangeable calcium. That neutrophil lipoxygenase metabolites of endogenous arachidonic acid rapidly enhance the influx of ^{45}Ca supports a possible role for such metabolites in general, and leukotriene B₄ in particular, in the regulation of the intracellular levels of free calcium in the neutrophils and possibly in other hormonally sensitive cells in which this cation is a second messenger.

Find the latest version:

<https://jci.me/110191/pdf>



Mono- and Dihydroxyeicosatetraenoic Acids Alter Calcium Homeostasis in Rabbit Neutrophils

P. H. NACCACHE and R. I. SHA'AFI, *Departments of Pathology and Physiology, University of Connecticut Health Center, Farmington, Connecticut 06032*
P. BORGEAT, *Centre de Recherches en Endocrinologie Moléculaire, Le Centre Hospitalier de l'Université Laval, Sainte-Foy, Quebec, Canada*
E. J. GOETZL, *Howard Hughes Medical Institute of Laboratory at Harvard Medical School, and Department of Medicine of the Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts 02115*

ABSTRACT Products of the lipoxygenation of arachidonic acid that express neutrophil chemotactic activity were examined *in vitro* for effects on the uptake of ^{45}Ca by rabbit peritoneal neutrophils. At optimally chemotactic concentrations, 5- and 11-hydroperoxyeicosatetraenoic acid, 11- and 12L-hydroxyeicosatetraenoic acid, and leukotriene B_4 enhanced ^{45}Ca uptake within 1 min by a mean of 212–694% of the values for control neutrophils incubated in buffer alone, as compared with 75% for 5(S)-hydroxyeicosatetraenoic acid and no effect for 15-hydroperoxyeicosatetraenoic acid and 15-hydroxyeicosatetraenoic acid. The approximate rank of potency of the factors stimulating ^{45}Ca uptake was similar to that observed for chemotaxis. Leukotriene B_4 , in addition to stimulating the rate of ^{45}Ca uptake into rabbit neutrophils, also displaced intracellular calcium. The net result of the leukotriene B_4 -induced changes in calcium homeostasis was to elevate transiently the intracellular level of exchangeable calcium. That neutrophil lipoxygenase metabolites of endogenous arachidonic acid rapidly enhance the influx of ^{45}Ca supports a possible role for such metabolites in general, and leukotriene B_4 in particular, in the regulation of the intracellular levels of free calcium in the neutrophils and possibly in other hormonally sensitive cells in which this cation is a second messenger.

INTRODUCTION

Polymorphonuclear leukocytes (neutrophils) have been shown to metabolize arachidonic acid mostly through the lipoxygenase pathway (1). Several of the lipoxy-

genase metabolites of arachidonic acids (hydroxyeicosatetraenoic acids [HETE])¹ influence diverse functions of the neutrophils. Human and rat neutrophils can be induced to move (2–7), aggregate (6), and degranulate to a limited extent (8) upon the addition of HETE *in vitro*, and guinea pig neutrophils accumulate in the peritoneal cavity in response to the introduction of HETE *in vivo* (9).

We have recently proposed that the neutrophil-directed activities of arachidonic acid and its lipoxygenase metabolites may be mediated by the ability of these fatty acids to alter the homeostatic regulation of the intracellular level of free calcium. This hypothesis was based on the results of experiments showing that calcium fluxes in rabbit neutrophils are specifically altered by exogenous arachidonic acid (10, 11) and that the capacity of rabbit and human neutrophils to migrate and to release lysosomal enzymes is dependent on the integrity of the lipoxygenase pathway (12–16).

We have therefore examined the effects of several monohydroxyperoxyeicosatetraenoic acids (HPETE) isomers, the corresponding mono-HETE, and the predominant natural di-HETE product of neutrophils, 5(S), 12(R)-dihydroxyeicosa-6,8,10 (two-*trans*, one-*cis*)-14-*cis*-tetraenoic acid (or leukotriene B_4), all of which are neutrophil chemotactic factors of differing potency, on the rate of ^{45}Ca uptake and on the steady-state level of exchangeable calcium in rabbit neutrophils. The results of these experiments are presented below.

METHODS

The experiments to be described were performed with rabbit peritoneal neutrophils collected, handled, and suspended in

Address correspondence to Dr. Sha'afi, Department of Physiology, University of Connecticut Health Center, Farmington, Conn. 06032.

Received for publication 22 January 1981 and in revised form 24 February 1981.

¹ Abbreviations used in this paper: HETE, hydroxyeicosatetraenoic acid; HPETE, hydroxyperoxyeicosatetraenoic acid; HPLC, high-pressure liquid chromatography.

modified Hanks' balanced salt solution, as previously reported (13). All the experiments were performed on cells thermally equilibrated at 37°C for 20 min and in the absence of added magnesium and extracellular protein. The rapid-sampling silicone oil method previously described in detail (17) was used for the calcium flux studies.

5-HPETE, 11-HPETE, and 15-HPETE were generated by incubating arachidonic acid (Supelco, Inc., Bellefonte, Pa.) with hydrogen peroxide and cupric chloride in methanol buffered with Tris-HCl (pH 7.8) as described (18). The HPETE were extracted into ethyl ether, resolved, and purified by reverse-phase high-performance liquid chromatography (HPLC), identified by mass spectrometry, and quantified by optical density at 235 nm and by the conversion of triphenylphosphine to triphenylphosphine oxide, as assessed by gas-liquid chromatography (18). 12-L-HETE was produced by incubating arachidonic acid with a partially purified preparation of lipoxygenase from human platelets (4, 9), whereas 5-HETE, 11-HETE, 15-HETE, and leukotriene B₄ were generated from arachidonic acid by neutrophils that had been stimulated with 10 μM calcium ionophore A23187 (5, 9, 14). The mono- and di-HETE were purified by sequential silicic acid column chromatography and reverse-phase HPLC, identified by mass spectrometry, and quantified by optical density at 235 and 280 nm, respectively (5, 14). 15-HPETE and 15-HETE were prepared from the soya bean lipoxygenase, extracted with ether, and fractionated by silicic acid chromatography. The fractions containing the compounds of interest were purified in two steps by HPLC on silica gel and octadecyl silica (reverse-phase). These compounds, whose identity was checked by mass spectrometry, were >95% pure as determined by gas chromatography.

RESULTS AND DISCUSSION

Inhibitors of the lipoxygenation of arachidonic acid have been shown to inhibit the functional responsiveness of rabbit and human neutrophils (12–16) and to concomitantly block both the chemotactic factor and arachidonic acid-induced stimulation of ⁴⁵Ca uptake in rabbit neutrophils (10, 12) and the metabolism of exogenously added arachidonic acid (14, 15). These results suggested that the lipoxygenation of endogenous arachidonic acid may represent one of the biochemical prerequisites for calcium mobilization and the presumably subsequent neutrophil activation by chemotactic factors and other agonists.

Intracellular calcium metabolism in neutrophils and its perturbation by chemotactic factors can be monitored by a variety of indirect radioisotopic tests. One of these, the rate of ⁴⁵Ca uptake, is a composite parameter reflecting both the plasma membrane permeability and the intracellular level of exchangeable calcium. As such it is an adequate initial test for a postulated alteration in calcium homeostasis.

As shown in Fig. 1, leukotriene B₄ significantly and rapidly increases the initial rate of ⁴⁵Ca uptake into rabbit neutrophils. The effects of arachidonic acid on this parameter of calcium metabolism are also illustrated in Fig. 1. Concentrations of the two fatty acids which produce about equivalent increases in ⁴⁵Ca uptake were chosen. Leukotriene B₄ can thus be




FIGURE 1 Effect of leukotriene B₄, 0.02 μg/ml (□), arachidonic acid, 1.5 μg/ml (▽), and f-Met-Leu-Phe, 1 nM (▲) on the time-course of ⁴⁵Ca uptake in rabbit neutrophils. ⁴⁵Ca and the various stimuli were added at zero time. The results shown in the figure depict those of a representative experiment (three different experiments). ●, control.

seen to be close to 100 times more effective on a molar basis than its metabolic precursor, a result in agreement with known biological activities of these two compounds (5–7). Fig. 1, in addition, shows that neither of these two fatty acids is as effective as an optimum concentration of the synthetic chemotactic peptide formyl-methionyl-leucyl-phenylalanine (f-Met-Leu-Phe).

At concentrations which elicit a maximal optimal chemotactic response, 5-HPETE, 11-HPETE, 12-L-HETE leukotriene B₄ and to a lesser extent 5-HETE, but not 15-HPETE or 15-HETE, increased significantly the rate of uptake of ⁴⁵Ca by rabbit neutrophils (Table I). The maximal changes were observed within the 1st min of the additions, following which the rates of uptake paralleled that of unstimulated neutrophils (results not shown). The time-course of the effects of the HETE on ⁴⁵Ca uptake is similar to that of the formyl-methionyl peptides (17).

Leukotriene B₄ (1) exhibits a maximal effect on ⁴⁵Ca uptake at concentrations that are more than 20 times lower than those required for somewhat lesser effects by the HPETE and mono-HETE. In the 5- and 11-series, the HPETE is more active than the corresponding HETE. 11-HPETE and HETE showed no detectable activity on the rate of ⁴⁵Ca uptake at concentrations at high as 3.3 μg/ml (Table I). To the extent that such a correlation can be drawn, the approximate rank order of potencies of the HETE corresponds to their known chemokinetic activity (leukotriene B₄ > 5-HETE > 11-HETE > 12-HETE > 15-HETE; HPETE > HETE) (16, 19).

The effects of leukotriene B₄ on the net movement of ⁴⁵Ca in rabbit neutrophils were then examined. In these experiments, the cells were allowed to reach

TABLE I
Effect of HETE on the Initial Rate of ^{45}Ca Uptake
in Rabbit Peritoneal Neutrophils

Compound	Concentration $\mu\text{g/ml}$	Percent change from control
11-HPETE	1	435 ± 51 (4)
11-HETE	2	250 ± 90 (4)
5-HPETE	0.5	212 ± 67 (5)
5-HETE	1	75 ± 30 (3)
12-HETE	2	247 ± 78 (3)
15-HPETE	3.3	32 ± 70 (3)
15-HETE	3.3	-24 ± 127 (2)
Leukotriene B_4	0.02	694 ± 144 (4)

Calculated as the difference in counts between experimental and control conditions after 1 min of incubation with ^{45}Ca and the stimulus. The results are expressed as means \pm standard error of the mean. The numbers in parentheses refer to the number of experiments, each carried out in duplicate.

isotopic equilibration before being exposed to leukotriene B_4 (Fig. 2). In the presence of 0.5 mM extracellular calcium (intracellularly directed electrochemical gradient for calcium), leukotriene B_4 causes a rapid and significant increase in the steady-state level of cell-associated ^{45}Ca . Leukotriene B_4 therefore unambiguously causes a net increase in the intracellular concentration of exchangeable calcium. In the absence of extracellular calcium, on the other hand, leukotriene B_4 induces a rapid and transient drop in the steady-state level of ^{45}Ca . As detailed elsewhere (23), this last result indicates that leukotriene B_4 causes an intracellular displacement of previously unexchangeable calcium.

Leukotriene B_4 therefore behaves as f-Met-Leu-Phe in causing both an intracellular redistribution of calcium and an increase in the plasma membrane permeability to calcium, the net result of which is to increase, at least transiently, the intracellular concentration of exchangeable calcium (17, 20). As the latter experiments lend themselves more readily to dose-response studies than any of the other parameters of ^{45}Ca fluxes, and because of the unique information that can be derived from these experiments concerning the intracellular status of calcium, we have examined the effects of three different concentrations of leukotriene B_4 on the steady-state level of cell-associated ^{45}Ca in the absence of extracellular calcium. The average drop in the steady-state level of ^{45}Ca 1 min after the addition of 0.0002, 0.002, and 0.02 $\mu\text{g/ml}$ of leukotriene B_4 was found to be 10, 19, and 25%, respectively (means of two experiments, each carried out in duplicate).

The level of stimulation of the rate of ^{45}Ca uptake that was produced by the various HETE with the exception




FIGURE 2 Effect of leukotriene B_4 , 0.02 $\mu\text{g/ml}$, on the steady-state level of cell-associated ^{45}Ca in rabbit neutrophils. In these experiments the cells were incubated with or without "cold" CaCl_2 for 50 min before adding leukotriene B_4 . Each point, taken in duplicate, was then normalized to its respective control. The results shown in the figure depict those of a representative experiment (three different experiments). ●, 0.05 mM Ca^{2+} ; ▲, $\leq 5 \mu\text{M}$.

of leukotriene B_4 (see Fig. 1 and Table I) was significantly lower than that evoked either by f-Met-Leu-Phe or by arachidonic acid. In addition, the high concentrations required for the mono-HETE to exhibit biological activity and to alter ^{45}Ca uptake make it unlikely that these compounds are, under physiological conditions, involved in the alterations of the intracellular level of free calcium. Leukotriene B_4 , on the other hand, affects ^{45}Ca metabolism at concentrations characteristic of the formyl-methionyl peptides and are likely to be attained *in vivo*. It is unknown at present whether the effective concentrations of these metabolites are overestimated because of the method of exogenous presentation to the neutrophils, as contrasted with the intracellular generation by physiological stimuli.

The capacity of exogenously added lipoxygenase products of arachidonic acid to enhance the rate of uptake of ^{45}Ca and to alter its steady-state level supports the hypothesis that such endogenous metabolites are at least in part responsible for the previously described perturbations of the intracellular level of calcium, the postulated second messenger in neutrophils. The recent findings that lipoxygenase inhibitors markedly diminish the anaphylactic contractile response of lung parenchymal strips from ovalbumin-sensitized guinea pigs (21) and the arachidonic acid- and C5_a -induced

contraction of guinea pig trachea (22), make it possible for this hypothesis to have validity in the study of other hormonally sensitive cells.

ACKNOWLEDGMENTS

We would like to thank Mr. T. F. P. Molski for his expert help in these experiments.

This study was supported in part by National Institutes of Health grant AI-13734.

REFERENCES

1. Borgeat, P., and B. Samuelsson. 1979. Arachidonic acid metabolism in polymorphonuclear leukocytes: effects of ionophore A23187. *Proc. Natl. Acad. Sci. U. S. A.* **76**: 2148-2152.
2. Turner, S. R., J. A. Campbell, and W. A. Lynn. 1975. Polymorphonuclear leukocyte chemotaxis towards oxidized lipid components of cell membranes. *J. Exp. Med.* **141**: 1437-1441.
3. Turner, S. R., J. A. Tainer and W. S. Lynn. 1977. Biogenesis of chemotactic molecules by the arachidonate lipoxygenase system of platelets. *Nature (Lond.)*. **257**: 680-681.
4. Goetzel, E. J., and P. R. Gorman. 1978. Chemotactic and chemokinetic stimulation of human eosinophil and neutrophil polymorphonuclear leukocytes by 12-L-hydroxy-5,8,10-heptadecatrienoic acid (HHT). *J. Immunol.* **120**: 526-531.
5. Goetzel, E. J., and W. C. Pickett. 1980. The human PMN leukocyte chemotactic activity of complex hydroxy-eicosatetraenoic acids (HETE's). *J. Immunol.* **125**: 1789-1791.
6. Ford-Hutchinson, A. W., M. A. Bray, M. V. Doig, M. E. Shipley, and M. J. H. Smith. 1980. Leukotriene B, a potent chemokinetic and aggregating substance released from polymorphonuclear leukocytes. 1980 *Nature (Lond.)*. **286**: 264-265.
7. Palmer, R. M. J., R. J. Stepney, G. A. Higgs, and K. E. Eakins. 1980. Chemokinetic activity of arachidonic acid lipoxygenase products on leukocytes of different species. *Prostaglandins*. **20**: 411-418.
8. Stenson, W. F., and C. W. Parker. 1980. Monohydroxy-eicosatetraenoic acids (HETEs) induce degranulation of human neutrophils. *J. Immunol.* **124**: 2100-2109.
9. Goetzel, E. J., F. H. Valone, V. N. Reinhold, and R. R. Gorman. 1979. Specific inhibition of the polymorphonuclear leukocyte chemotactic response to hydroxy-fatty acid metabolites of arachidonic acid by methyl ester derivatives. *J. Clin. Invest.* **63**: 1181-1186.
10. Volpi, M., P. H. Naccache, and R. I. Sha'afi. 1980. Arachidonic metabolite(s) increase the permeability of the plasma membrane of the neutrophils to calcium. *Biochem. Biophys. Res. Commun.* **92**: 1231-1237.
11. Sha'afi, R. I., P. H. Naccache, T. Alobaidi, T. F. P. Molski, and M. Volpi. 1980. Effect of arachidonic acid and the chemotactic factor f-Met-Leu-Phe on cation transport in rabbit neutrophils. *J. Cell. Physiol.* **106**: 215-223.
12. Naccache, P. H., H. J. Showell, E. L. Becker, and R. I. Sha'afi. 1979. Pharmacological differentiation between the chemotactic factor induced intracellular calcium redistribution and transmembrane calcium influx in rabbit neutrophils. *Biochem. Biophys. Res. Commun.* **89**: 1224-1230.
13. Showell, H. J., P. H. Naccache, R. I. Sha'afi, and E. L. Becker. 1980. Inhibition of rabbit neutrophil lysosomal enzyme secretion, nonstimulated and chemotactic factor stimulated locomotion by nordihydroguaiaretic acid. *Life Sci.* **27**: 421-426.
14. Goetzel, E. J. 1980. A role for endogenous mono-hydroxy-eicosatetraenoic acids (HETE's) in the regulation of human neutrophil migration. *Immunology*. **40**: 709-719.
15. Walenga, R., H. J. Showell, M. B. Feinstein, and E. L. Becker. 1980. Parallel inhibition of neutrophil arachidonic acid metabolism and lysosomal enzyme secretion by nordihydroguaiaretic acid. *Life Sci.* **27**: 1047-1053.
16. Goetzel, E. J., A. R. Brash, A. J. Tauber, J. A. Oates, and W. C. Hubbard. 1980. Modulation of human neutrophil function by monohydroxy-eicosatetraenoic acids. *Immunology* **39**: 491-501.
17. Naccache, P. H., H. J. Showell, E. L. Becker and R. I. Sha'afi. 1977. Transport of sodium, potassium and calcium across rabbit polymorphonuclear leukocyte membranes. Effect of chemotactic factor. *J. Cell Biol.* **73**: 428-444.
18. Boeynaems, J. M., J. A. Oates, and W. C. Hubbard. 1980. Preparation and characterization of hydroperoxy-eicosatetraenoic acids (HPETE). *Prostaglandins*. **19**: 81-97.
19. Goetzel, E. J., H. R. Hill, and R. R. Gorman. 1980. Unique aspects of the modulation of human neutrophil function by 12-L-hydroperoxy-5,8,10-14-eicosatetraenoic acids. *Prostaglandins*. **19**: 71-85.
20. Petroski, R. J., P. H. Naccache, E. L. Becker, and R. I. Sha'afi. 1979. Effect of chemotactic factors on calcium levels of rabbit neutrophils. *Am. J. Physiol.* **237**: C43-C49.
21. Yen, S. S., and W. Kreutner. 1980. Effect of inhibition of arachidonic acid metabolism on anaphylaxis of pig lung strips. *Agents Actions*. **10**: 284-287.
22. Regal, J. R., and R. J. Pickering. 1980. Inhibition of C5_a induced contraction of guinea pig trachea. *Fed. Proc.* **39**: 701.
23. Sha'afi, R. I., and P. H. Naccache. 1981. Ionic events in neutrophil chemotaxis and secretion. In *Advances in Inflammation Research*. G. Weissmann, editor. Raven Press, Inc., New York. In press.