

Characterization of resident glomerular cells in the rat expressing Ia determinants and manifesting genetically restricted interactions with lymphocytes.

G F Schreiner, ... , R S Cotran, E R Unanue

J Clin Invest. 1981;68(4):920-931. <https://doi.org/10.1172/JCI110347>.

Research Article

The existence of a subpopulation of rat glomerular cells bearing Ia determinants has been demonstrated with the aid of techniques for the enzymatic isolation and culture of glomerular cells. The Ia-positive cell is normally resident in the uninfamed glomerulus. It resembles a mononuclear phagocyte and consists of a functionally heterogeneous cell population with the capacity of Fc receptor display and phagocytosis, both in vivo and in vitro. A new technique for labeling these cells in situ in intact glomeruli has indicated that Ia-positive cells make up approximately 2% of the total glomerular cell population. The isolated glomerular cells can take up antigen and stimulate immune lymphocytes in an I-region-restricted interaction. They are strongly stimulatory in an allogeneic primary mixed lymphocyte culture. Characterization of this cell type suggests potential new insights into the pathogenesis of renal allograft rejection and immunologically mediated glomerulonephritis.

Find the latest version:

<https://jci.me/110347/pdf>



Characterization of Resident Glomerular Cells in the Rat Expressing Ia Determinants and Manifesting Genetically Restricted Interactions with Lymphocytes

GEORGE F. SCHREINER, JEANNE-MARIE KIELY, RAMZI S. COTRAN, and
EMIL R. UNANUE, *Departments of Pathology, Harvard Medical School and
Brigham and Women's Hospital, Boston, Massachusetts 02115*

ABSTRACT The existence of a subpopulation of rat glomerular cells bearing Ia determinants has been demonstrated with the aid of techniques for the enzymatic isolation and culture of glomerular cells. The Ia-positive cell is normally resident in the uninfamed glomerulus. It resembles a mononuclear phagocyte and consists of a functionally heterogeneous cell population with the capacity of Fc receptor display and phagocytosis, both *in vivo* and *in vitro*. A new technique for labeling these cells *in situ* in intact glomeruli has indicated that Ia-positive cells make up ~2% of the total glomerular cell population. The isolated glomerular cells can take up antigen and stimulate immune lymphocytes in an I-region-restricted interaction. They are strongly stimulatory in an allogeneic primary mixed lymphocyte culture. Characterization of this cell type suggests potential new insights into the pathogenesis of renal allograft rejection and immunologically mediated glomerulonephritis.

INTRODUCTION

Many studies over the last three decades established the pathogenesis of glomerulonephritis as arising from the presence of antibody in the kidney in either of two forms: either deposited as an immune complex formed in a systemic immune reaction or as an antibody directed against glomerular antigenic determinants themselves. Both situations initiate humoral events including the activation of complement that result in functional impairment of glomerular filtration (1). The role of circulating leukocytes became established with the demonstration that neutrophils contributed to the early phase of glomerular damage in experimental glomerulonephritis (2). Recently, we have shown that

Received for publication 13 April 1981 and in revised form 8 June 1981.

some forms of experimental glomerulonephritis are associated with an influx of monocytes that correlates with sustained histological and functional abnormalities in the glomerulus (3). Such investigations have visualized the glomerulus as a somewhat passive participant in its own impairment, providing a vulnerable milieu for a stereotyped inflammatory response whose only reference to the kidney was as an antibody depot. We have now made observations that suggest that the kidney may be a more active contributor to glomerular immune reactions than previously realized.

Using techniques for the isolation and short-term culture of glomerular cells, and their *in situ* labeling, we have found a new cell intrinsic to the uninfamed renal glomerulus. It is phagocytic and bears I-region-associated antigens (Ia), the membrane proteins coded for by the major histocompatibility locus that permit specific interactions between phagocytes and lymphocytes. This Ia-bearing glomerular cell can present antigen in a genetically restricted fashion to syngeneic lymphocytes and can selectively stimulate allogeneic lymphocytes. The presence of this cell indicates that the glomerulus has the potential to process antigen actively and to initiate a cellular immune response *in situ*. These observations not only bear significant implications for defining a new pathogenesis of glomerulonephritis, they also suggest a better way of understanding the mechanisms that underlie the initiation of host rejection of kidney transplants.

METHODS

Animals. Female rats, either of the Lewis or the Buffalo strains, weighing 100–175 g, and at least 5 wk old, were obtained from Microbiological Associates (Walkersville, Md.).

Glomerular isolation. Before harvesting glomeruli, both kidneys were thoroughly perfused to remove circulating erythrocytes and leukocytes with a technique modified from Griffith et al. (4), with the additional step of opening the left renal vein to permit the perfusate to drain. 50–100 ml of a 0.9% NaCl solution injected via the aorta into both kidneys

produced their complete blanching and eventually a clear perfusate. Mottled kidneys, indicative of incomplete perfusion, were discarded. Possible blood contamination was judged minimal because erythrocytes or neutrophils were rare in isolated glomeruli as well as in cultures of enzymatically isolated glomerular cells. Glomeruli were obtained by pressing slices of renal cortex through graded sieves (Tyler sieves, Curtin Matheson Scientific, Inc., Houston, Tex.) of 250, 150, and 75 μg with a rubber stopper (5). Preparations consisted of 90–95% glomeruli with occasional contaminating tubular fragments.

The protocol for the enzymatic release of glomerular cells was extensively modified from previous reports (6). The purified glomeruli were washed twice in Hanks' balanced salt solution (HBSS)¹ at 50 g and then placed, in step 1, in a solution consisting of trypsin (Type III, 10,200 BAEE U/mg) (Sigma Chemical Co., St. Louis, Mo.), 1 mg/ml; collagenase (Type CLS IV, 126 U/mg) (Millipore Corp., Freehold, N. J.), 1 mg/ml; and DNAase (Type III, 1,825 Kunitz U/mg), 0.01 mg/ml, pH 7.2–7.4, for 45 min at 37°C on a rocking platform. The partially digested glomeruli were then washed in HBSS lacking calcium or magnesium (Microbiological Associates) and placed, in step 2, in a solution of 2 mM EDTA (Sigma Chemical Co.) in HBSS without calcium or magnesium for 10 min at 37°C. This preparation was then centrifuged at 50 g for 10 min and divided into a sediment and a suspension of single cells. In step 3, the sediment, consisting of partially digested glomerular fragments, was incubated for an additional 20 min at 37°C in collagenase, 2 mg/ml. The preparation was allowed to settle for 90 s at 1 g, which removed most of the residual debris. The supernate after the collagenase incubation was then centrifuged at 400 g, washed, and recombined with the cells removed after step 2. Cells were spun at 400 g for 10 min and resuspended in HBSS with 1% bovine albumin (Sigma Chemical Co.) and 1 mM Hepes buffer (Microbiological Associates). This process resulted in a suspension containing between 2 and 5 $\times 10^6$ single cells per rat.

Cell adherence and labeling of Ia antigens. 1 ml of cells at 2 $\times 10^6$ /ml was placed in flat-bottom 16-mm wells of multiwell trays (Falcon Labware, Div. of Becton-Dickinson & Co., Oxnard, Calif.) containing glass round coverslips (Bellco Glass, Inc., Vineland, N. J.) previously coated with heat-decomplemented fetal calf serum (Microbiological Associates). The cells were allowed to settle onto the plate by incubation at 37°C for 2 h in a tissue culture incubation chamber containing 5% CO₂ and then used in different experiments. The medium during adherence was the same used for the antigen presentation assay. In some experiments, cells were cultured for longer periods, as specified in the text.

Labeling of rat Ia antigens was done with pooled batches of monoclonal antibody (Accurate Chemical & Scientific Corp., Hicksville, N. Y.). Two preparations previously studied in detail (7, 8) were used: (a) MASO29B, a mouse IgG from clone OX3HL that binds to Ia antigens from Lewis (RT1 haplotype l) and Wistar (haplotype u) strains, that cross-reacts with mouse Ia specificity 9; and (b) MASO28B, a mouse IgG from clone OX4HL that recognizes a common Ia determinant of all rats and cross-reacts with mouse Ia specificity 18. Using our staining protocol, we confirmed that MASO28B recognizes the I region of mouse haplotypes b and s by positively labeling a percentage of peritoneal macrophages from C57BL/6 and

A.TH strains; staining of A.TL and B10.D2 cells was negative. Similarly, MASO29B recognizes mouse haplotypes s and k in that it labels macrophages from A.TH and A.TL strains but not C57BL/6 or B10.D2 strains. We also confirmed McMaster and Williams' (8) findings that, in the rat, both antisera label 20–40% of adherent splenic macrophages, all B lymphocytes, and no peripheral T cells. We also found that each preparation labeled 5–10% of peritoneal exudate cells similar to the level of Ia expression noted in the mouse (9). Labeling was carried out with a 1:4 dilution of decomplemented, pooled (1:1) batches of MASO28B and MASO29B, diluted in HBSS. Coverslips containing the glomerular cells were placed face-side up in new wells to which were added 300 μl of the anti-Ia monoclonal antibodies. They were incubated for 20 min at either 4° or 20°C, depending upon the protocol; the coverslips were then dipped in three washes of cold phosphate-buffered saline (PBS) and placed into fresh wells to which was added a solution of fluorescein-labeled F(ab)₂ rabbit anti-mouse IgG (FITC-anti-Ig) at 100 $\mu\text{g}/\text{ml}$ (10). After incubation at 4°C or room temperature for an additional 20 min, the coverslips were rinsed in two washes of saline and then fixed in 2% paraformaldehyde. The cells were then read for presence of fluorescence on a Leitz microscope with appropriate filters (E. Leitz, Inc., Rockleigh, N. J.). Unless otherwise detailed, results were expressed as percentage of positive labeled cells, for Ia, of total adherent cell population. Controls of labeling with normal mouse IgG or with electrophoretically purified mouse IgG from the plasmacytoma line MOPC195 (Litton Bionetics, Div. Litton Industries, Inc., Kensington, Md.) instead of anti-Ia antibody did not disclose any staining. In some experiments, total cell suspensions, before adherence, were labeled for Ia antigens. 2 $\times 10^6$ cells were suspended in 300 μl of anti-Ig (1:4, as above) for 20 min at 4°C, washed twice, and then placed in FITC-anti-Ig, as above, for 20 min at 4°C. The cells were then washed and fixed in 2% paraformaldehyde. Cells were scored for fluorescent labeling and reported as a percentage of total single cells in suspension. Cells in clumps were not scored. In some experiments, monocytes were obtained by enrichment on Ficoll-Hypaque gradients after subjecting peripheral blood to dextran sedimentation. After adherence for 2 h on coverslips, they were labeled as above. Control experiments indicated that treatment of cells that express Ia, such as monocytes and macrophages, with trypsin and collagenase, as was done to the glomerular cells, neither induced the expression of Ia nor affected the Ia already found on their surfaces.

Labeling intact glomeruli. Whole glomeruli, isolated as noted above, were placed in a solution containing 1 mg/ml trypsin and 0.01 mg/ml DNAase for 30 min at 37°C in HBSS, pH 7.2. The glomeruli were then washed twice and incubated first in a 1:4 dilution of pooled anti-Ia antisera for 30 min at 4°C, washed twice, and then incubated in 100 $\mu\text{g}/\text{ml}$ of FITC-anti-Ig for an additional 30 min at 4°C. After washing two further times, the whole but partially digested glomeruli were then fixed in 2% paraformaldehyde and examined microscopically. Controls included: (a) deleting preincubation with trypsin, (b) deleting incubation with anti-Ia but labeling with FITC-anti-Ig, and (c) substituting for anti-Ia antibody an equal concentration of mouse IgG from plasmacytoma line MOPC195, as detailed above. In none of the controls in any experiment did we observe labeling of any glomerular cells.

Latex bead ingestion. 40 $\times 10^6$ latex beads (1 μm ; Dow Diagnostics, Indianapolis, Ind.) were layered onto the culture wells, which were spun at 150 g for 5 min and then incubated at 37°C for 45 min. The coverslips were washed with warm PBS to remove unbound latex and fixed in 2% paraformaldehyde in PBS. Some coverslips were subsequently studied for

¹Abbreviations used in this paper: EA, erythrocyte antisera; FITC-anti-Ig, fluorescein-labeled F(ab)₂ rabbit anti-mouse IgG; HBSS, Hanks' balanced salt solution; PBS, phosphate-buffered saline.

the presence of Ia-determinants or for Fc receptors. Cells interiorizing three or more particles were scored as positive.

Fc receptor. Receptors for Fc moiety of IgG were assayed by using sheep erythrocytes coated with IgG rabbit anti-erythrocyte antisera (EA), as previously described (11). 1 ml of a 1:100 dilution of the 2% EA solution was layered onto the adherent cells. The tissue culture plates were spun for 5 min at 150 g to sediment the EA onto the adherent cells. After further incubation at room temperature for 5 min, an equal volume of 1.25% glutaraldehyde in PBS was gently added to the wells. Preparations labeled for the presence of both Fc receptors and Ia determinants were labeled with anti-Ia antiserum for 20 min at room temperature before fixation for 10 min in glutaraldehyde, washed four times in PBS, and then labeled with FITC-anti-Ig.

In vivo phagocytosis. Four 150-g Lewis rats were injected with 8 mg of rhodamine-conjugated heat-aggregated human gamma globulin (Miles Laboratories, Inc., Elkhart, Ind.) and the glomeruli harvested 90 min after injection. Individual glomerular cells were prepared and evaluated for the presence of red fluorescent phagocytic vacuoles, which indicate the ingestion of rhodamine-labeled gamma globulin before sacrifice, and green fluorescent membrane staining indicating the presence of labeled Ia determinants, or both.

Antigen presentation assay. The proliferation of immune T lymphocytes in response to antigen presentation by Ia-bearing cells was carried out in a protocol analogous to that previously described for mice (11). Immune lymphocytes were obtained by injecting female Lewis rats, 2 mo old, with 2×10^5 live *Listeria monocytogenes* into the peritoneum 10 d before harvesting. 3 d before harvesting each rat received 5 ml i.p. of 10% proteose-peptone (Difco Laboratories, Detroit, Mich.). T lymphocytes were separated from adherent peritoneal exudate cells by two cycles of 2 h of adherence on plastic tissue culture dishes, followed by a third cycle of adherence in overnight culture. Glomerular cells at 2×10^6 cells/ml were allowed to adhere according to the labeling protocol. Heat-killed, opsonized *Listeria* (2×10^7) were added to the cultures of 2×10^6 adherent glomerular cells and centrifuged at 400 g for 5 min to enhance settling of *Listeria* onto the cells. The antibody-coated *Listeria* were prepared by incubating heat-killed *Listeria*, 5×10^7 /ml, in a 1:10 dilution of whole rabbit anti-*Listeria* serum. The *Listeria*-pulsed cells were incubated at 37°C for 1 h, washed thoroughly, and then mixed with 4×10^5 immune T cells in medium consisting of RPMI 1640 with 10% fetal calf serum, 1% penicillin-streptomycin, 1 mM HEPES buffer, 1 mM pyruvate, and 1 mM L-glutamine (Gibco Laboratories, Grand Island Biological Co., Grand Island, N. Y.). Total length of the co-cultures was 4 d. Cultures were pulsed with 0.4 mCi of tritiated thymidine (sp act 20 μ Ci/ml, New England Nuclear, Boston, Mass.) over the last 16 h of culture. Proliferation was assayed by the incorporation of [3 H]thymidine into trichloroacetic-insoluble material. As has been noted previously, the growth of T cells requires Ia-bearing accessory cells and Ia homology between reacting T cells and antigen-presenting macrophages (11).

Assay of allogeneic reactivity. The responder cells were lymphocytes isolated from the axillary, periaortic, popliteal, and inguinal lymph nodes of adult female Buffalo rats. The stimulatory cells consisted of (a) spleen cells from allogeneic Lewis female rats, (b) adherent glomerular cells from Lewis rats, or (c) syngeneic adherent glomerular cells from Buffalo strain female rats. The medium was the same as for antigen presentation, with the addition of 50 μ M 2-mercaptoethanol. Stimulator cells were placed in culture at a concentration of 2×10^6 /ml (1 ml/well) for spleen cells and 5×10^6 cells/ml for glomerular cells. Stimulator cells were allowed to adhere

for 2 h on flat-bottomed microwells, washed, and then irradiated with 1,500 rad. The responder lymphocytes (2×10^5) were then added and the cultures allowed to proceed for 6 d. Proliferation was assayed exactly as above in the antigen presentation assay by adding tritiated thymidine to the cultures 16 h before harvest on day 6.

RESULTS

Identification of Ia-positive cells in glomeruli. Our interest in the possible existence of a resident, antigen-processing cell in the glomerulus led to the following experiments, in which enzymatically isolated glomerular cells were fluorescently stained for the presence of rat Ia antigens. Cells were examined in suspension and after varying times of adherence on coverslips. As shown in four representative experiments in Table I, we consistently found between 6 and 8% of total glomerular cells, in a single-cell suspension, bearing Ia antigens. Allowing the cells to adhere on coverslips for 2 h enriched the Ia-positive population three- to sixfold to between 21 and 37% of the total adherent glomerular cell population. About 1% of the non-adherent cells were Ia-positive. We found that, after 2 h of culture and after Ia labeling, only ~30–40% of the total cell population remain adherent.

24 h of in vitro culture resulted in a small and variable decrease in the percentage of Ia-bearing cells, as seen in Table I. This appears to be partly a dilutional effect, in that there are always more cells per high-power field after 24 h in culture than after 2 h. It appears that Ia-negative cells are less adherent in short-term culture and more susceptible to being dislodged; after 24 h, they are more adherent. For example, the mean number of cells per high-power field (25 fields) in experiment No. 4 was 29.7 at 2 h, rising to 47.2 at

TABLE I
Ia-positive Glomerular Cells in Culture

Experiment No.	Initial suspension	2-H adherence	24-H adherence
		%	
1	6	37	21
2	7	21	14
3	8	23	18
4	6.5	31	17

In each experiment, glomerular cells were labeled for the presence of Ia determinants at three time periods. "Initial suspension" refers to the percentage of Ia-positive cells in the total glomerular cell population at the beginning of the cultures, before adherence. The figures in the 2-h and 24-h columns refer to the percentage of adherent glomerular cells that were positively labeled for Ia after 2 or 24 h of culture on coverslips. 200–300 cells in each category were counted.

24 h. Nonetheless, there is also an absolute decrease in maintenance of Ia determinants that becomes prominent during more prolonged *in vitro* culture (see below).

Fig. 1A shows a typical Ia-positive glomerular cell. Morphologically, it resembles a mononuclear phagocyte. It ranges between 15 and 25 μm in diameter. It can be amoeboid in appearance but more typically is oval while adherent, with the nucleus asymmetrically placed. The cytoplasm contains few vacuoles. The Ia antigens are diffusely distributed. It should be noted that there is an apparently identical population of cells, with very similar membrane receptors but lacking Ia antigens.

Other cell types are present in the heterogeneous population of adherent cells. The predominant cell type is a smaller (10–15 μm) Ia-negative cell, seen in Fig. 1B, which is an endothelial cell by electron microscopy.² Adherent, large epithelial cells, 40–60 μm in diameter, are prominent in the 2-h cultures, less so after 24 h (Fig. 1C). They are notable for an abundant cytoplasm, ruffled membranes, and numerous vacuoles. Also present in variable numbers are contaminating tubular epithelial cells, as determined by electron microscopy.² Fig. 1D shows an additional cell type present in all cultures that closely resembles the contractile glomerular cell lines described by Ausiello et al. (12). In early culture, it is intermediate in size but quickly elongates with spreading, spindly processes. These cells do not bear Ia determinants and may be of mesangial origin (12).

Functional properties of Ia-positive cells. We assayed three properties of Ia-positive cells: (a) *in vitro* and *in vivo* phagocytosis, (b) *in vitro* display of Fc receptors, and (c) maintenance of the Ia phenotype in culture.

Table II shows representative experiments in which isolated adherent glomerular cells were scored for phagocytosis of latex beads and the presence of Ia antigens. In both experiments, 17 and 18% of adherent glomerular cell population manifested phagocytic capability of which 60 and 54% were Ia-positive. Similarly, glomerular cells were scored for the display of Fc receptors and of Ia antigens. Again, in Table II, ~40% of adherent glomerular cells were Fc receptor positive, of which approximately one-third were Ia-positive.

The above data refer to assays done after 2 h of culture. In evaluating Ia-positive cells both after 2 h and after 24 h, it became clear that there was heterogeneity among Ia-bearing cells with respect to

the expression of one or the other property and that there was a shift in their differential capacities over time in culture. Fig. 2 demonstrates one of two experiments in which Ia-positive cells were scored for phagocytosis and Fc receptors at 2 h and at 24 h. Ia-positive cells were divided into four categories: (a) Fc-receptor negative, latex-phagocytosis negative; (b) Fc-receptor negative, latex-phagocytosis positive; (c) Fc-receptor positive, latex-phagocytosis negative; and (d) positive for both Fc receptor and latex phagocytosis. 30% of the Ia-positive cells at 2 h displayed neither phagocytosis nor Fc receptors; this decreased to <20% over 24 h. 23% of the Ia-positive cells manifested only phagocytosis, with only a modest decrement with time. The largest shifts were seen in Ia-positive cells that also bore Fc receptors. Those cells expressing only Fc receptors decreased from 30 to 10% in longer culture. Ia-positive cells expressing both Fc receptors and phagocytosis increased from 18 to 51% of the Ia-bearing population. In this experiment, the percentage of total adherent cells that were Ia-positive was 34% at 2 h and 13% at 24 h. The mean number of cells per high-power field (25 fields) was 29 at 2 h and 40 at 24 h, which indicates increased adherence of all glomerular cells as discussed above. Thus, within the cells continuing to display Ia antigens there is a shift in functional properties towards a more active state, i.e., with increasing numbers of cells displaying both Fc receptors and phagocytic activity, and fewer cells displaying only one property or neither. We also noted that the range of bead ingestion increased from 2–5 to 5–10 beads/cell after prolonged culture (Fig. 3A). The number of EA bound per cell increased from 3–5 to >10/cell, which indicates an increase in the number or the avidity of Fc receptors or both (Figs. 3B and C).

There is an absolute decrease in the percentage of Ia-positive cells in prolonged culture, whereas the number of cells per high-power field is relatively constant. This is demonstrated in Fig. 4; in this experiment, by 96 h of *in vitro* culture, only 4% of the total glomerular cell population remained Ia-positive. In addition, the intensity of Ia labeling decreases with time among the positive cells, which indicates a generalized progressive loss of Ia antigens. The loss of the Ia phenotype by cells under prolonged culture conditions has been observed in the mouse (13, 14).

Finally, it should be noted that neither the small endothelial cells nor the larger epithelial cells nor the elongated cells of Fig. 1D have been observed to display Fc receptors or to ingest latex particles at any time in culture. Whereas those properties are expressed as a percentage of the total glomerular cell population, only the intermediate-sized cell described above is positive in these assays.

Controls and identification of Ia-positive cells as

²G. F. Schreiner, R. S. Cotran, and E. R. Unanue. Ultrastructural analysis of glomerular Ia-positive cells. Manuscript in preparation.




FIGURE 1 Labeling of Ia determinants on adherent glomerular cells. (A) Phase contrast and fluorescent micrographs of an Ia-positive, adherent glomerular cell after 24 h of culture, $\times 480$. (B) An adherent, Ia-positive glomerular cell with an adjacent, smaller, Ia negative cell (in experiments in progress, electron microcopic analyses have shown that this small Ia-negative cell is the endothelial cell²). In the upper right corner is an intermediate-size Ia-negative cell, $\times 480$. (C) A large epithelial cell in 2 h culture, $\times 550$. (D) An elongated, Ia-negative, adherent glomerular cell with spindle processes, morphologically resembling contractile mesangial cells (12). The other cell in the upper right corner has bound EA and thus displays Fc receptors, $\times 480$.

TABLE II
Functional Properties of Adherent Glomerular Cells

Experiment No.	Phagocytic	Fc receptor-positive	Ia-Positive
		%	
1	18 (54)*	—	37
2	17 (60)	—	21
3	—	42 (28)†	34
4	—	44 (33)	24

"Phagocytic" refers to the percentage of adherent glomerular cells, after 2 h of culture, ingesting three or more polystyrene beads. "Fc receptor positive" refers to the percentage of total adherent glomerular cells displaying Fc receptors under similar conditions. Positive cells were those binding three or more opsonized erythrocytes (EA). "Ia-positive" refers to percentage of total adherent glomerular cells positively labeled for Ia determinants after 2 h in culture. Experiments 1 and 2 studied the uptake of beads and surface Ia; experiments 3 and 4 studied Fc receptors and surface Ia. 200 cells in each category were counted.

* Numbers in parentheses refer to percentage of phagocytic cells that were also Ia-positive.

† Numbers in parentheses refer to percentage of Fc receptor-positive cells that were also Ia-positive.

glomerular cells. Clearly, a major question at this point is whether this cell type is a contaminating blood element or an extraglomerular renal cell. We have developed several approaches to this question and believe, on their basis, that these are specifically glomerular cells.

First, the kidneys were thoroughly perfused with saline with complete blanching before the glomeruli were harvested. This minimized blood contamination, as supported by the fact that erythrocytes were not observed in isolated glomeruli and were quite rare in the dispersed cell preparations, making up much less than 1% of the total cell population before adherence.

The possibility that the Ia-positive cells were interstitial or bound to the tubules that occasionally contaminate the preparation was examined in two ways. We subjected tubular-enriched fragments from sieve sizes other than the ones used to harvest the glomeruli or interstitial fragments to the same enzymatic digestion. This yielded few adherent cells and no Ia-positive cells. Therefore, the Ia-positive cells appeared only in glomerular-enriched fractions. If Ia-positive cells reside in the tubules, these are not detached under our experimental conditions.

More pointedly, we have developed a technique for labeling these cells *in situ* in intact glomeruli. It is based on our observations that trypsin made the glomeruli permeable under appropriate conditions,

permitting antibody to diffuse in and bind to trypsin-resistant antigens, while not grossly disturbing the cytoarchitecture of the intact glomerulus. The results are shown in Figs. 5A and B, in which Ia-positive cells are revealed in the normal glomerulus. They are primarily in an axial distribution with scattered cells located more peripherally. Figs. 5C and D show increased magnification of two glomeruli revealing elongated, serpiginous Ia-bearing cells, as well as the more oval-type cells seen in culture. The average number of cells per glomerulus is 10.2, with considerable heterogeneity between glomeruli, the range being between 5 and 20 cells/glomerulus. Control experiments using a nonspecific myeloma IgG preparation instead of the anti-Ia antisera showed no cellular staining but only a very weak, generalized background fluorescence.

Finally, we have compared the percentage of Ia-positive cells in the adherent glomerular cell population with that in adherent monocyte populations. In three experiments, the percentage of Ia-bearing monocytes was 0.5, 0, and 2%. In one experiment, duplicate coverslips of adherent monocytes were subjected to trypsin and DNAase treatment identical to that experienced by glomerular cells. This resulted in counts of 1 and 2% Ia-positive adherent monocytes. The lack of expression of Ia antigens by circulating monocytes has been observed in the mouse (D. I. Beller, personal communication). Although monocytes may have the capacity ultimately to express Ia, they apparently do so only as a more differentiated or activated cell. In any event, it is clear that even gross contamination by circulating mononuclear cells




FIGURE 2 Division of adherent Ia-positive glomerular cells into functional categories, after 2 or 24 h in culture. 300 Ia-positive cells were scored for demonstration of: (a) Fc receptors (FcR+, defined as binding more than three EA); (b) phagocytosis (defined as ingestion of more than three latex beads), both, or neither. The bars represent percentage of total adherent Ia-positive cells.




FIGURE 3 Phagocytosis and Fc receptor display by Ia-positive glomerular cells. (A) Phase contrast and fluorescent micrographs showing ingestion of a large number of latex beads by an Ia-positive glomerular cell after 24 h of culture. Nearby is a smaller, nonphagocytic, Ia-negative endothelial cell, $\times 480$. (B) An EA-rosette by an adherent Ia-positive glomerular cell after 2 h of culture. The Ia-positive cells after 2 h of culture are somewhat smaller than after 24 h of culture, $\times 480$. (C) Increased binding of EA displayed by Ia-positive glomerular cells after overnight culture, $\times 480$. (D) Phase and fluorescent micrographs of adherent glomerular cells exposed to opsonized rhodamine-conjugated *Listeria*, allowed to phagocytose, and then labeled for Ia determinants with fluorescein-conjugated antibody. The middle panel demonstrates the rhodamine fluorescence of the three cells in the right corner, indicating phagocytosis. The end panel demonstrates the ring fluorescence of membrane Ia labeling on one of the phagocytic cells.

cannot account for the high percentage of Ia-bearing cells isolated from the glomerulus.

In vivo phagocytosis by Ia-positive glomerular cells. To assess in vivo phagocytic function, we injected four Lewis rats with 8 mg i.v. of rhodamine-conjugated heat-aggregated gamma globulin. The rats were sacrificed at 90 min, at which time the labeled gamma globulin could be detected in isolated glomeruli. Harvesting the glomerular cells and then counterlabeling with fluorescein-labeled anti-Ia revealed the following: 30% of the adherent glomerular cells at 2 h demonstrated rhodamine-fluorescent phagosomes, which indicates that they had ingested aggregated gamma globulin. 46% of all adherent Ia-positive cells were similarly labeled, making up 30% of all the phagocytic cells isolated from the glomerulus.

Immune function of glomerular cells. The presence of Ia antigens on phagocytic glomerular cells suggests these should be capable of presenting antigen to immune lymphocytes in a genetically restricted manner. We assessed antigen presentation with an assay based on immunity to *Listeria monocytogenes* (11). In this assay, immunized T lymphocytes are seen to proliferate in response to *Listeria* only when it is presented bound to Ia-positive, syngeneic phagocytes. Our initial attempts to induce adherent glomerular cells to present *Listeria* as an antigen failed. The explanation became clear from experiments elsewhere in our laboratory in which it was observed that trypsin removes from the cell membrane of phagocytes a binding protein for *Listeria*, thus preventing uptake (15). This protein is subsequently regenerated in culture (D. S. Weinberg and E. R. Unanue, experiments to be published). It should be noted that trypsin is used in the isolation procedure that precedes the establishment of glomerular cell cultures. Opsonization of the *Listeria*, however, allowed its uptake by way of the trypsin-resistant Fc receptor. In one experiment, for example, opsonizing *Listeria* increased the percentage of Ia-positive, adherent glomerular cells ingesting *Listeria* from 3 to 37% (Fig. 3D).

Table III, therefore, demonstrates two experiments in which glomerular cells from Lewis rats were pulsed with opsonized heat-killed *Listeria*, washed, and then combined for 3 d in culture with T lymphocytes from Lewis rats immune to *Listeria*. *Listeria*-pulsed adherent glomerular cells were highly effective in presenting antigen to T cells and inducing them to proliferate. The genetic restriction of this antigen presentation is demonstrated in Table IV, in which Lewis glomerular cells were far more effective in presenting antigen to sensitized lymphocytes of the




FIGURE 4 The percentage of total adherent glomerular cells displaying Ia determinants over time in culture. The percentage at time zero represents the total glomerular cell population that is Ia-positive before the adherence step. 200–300 cells were evaluated at each time period. In boxes at the top of the figure are the arithmetic mean number of total cells per high-power field ($n = 25$). The difference noted at 2 and 24 h is discussed in the text.

same haplotype than were comparably treated cells of the allogeneic Buffalo strain. The limited amount of lymphocyte reactivity seen when Lewis T lymphocytes are mixed with Buffalo strain glomerular cells or peritoneal exudate cells, whether or not they have been pulsed with *Listeria*, appears to be due to early lymphocyte alloreactivity.

Finally, the glomerular cells are themselves extremely stimulatory to allogeneic lymphocytes in vitro (Table V). It is interesting to note that syngeneic glomerular cells were capable of inducing some degree of proliferation in lymphocytes, albeit to a much smaller extent than allogeneic stimulation.

DISCUSSION

We have identified in the rat a distinct, new glomerular cell population bearing Ia determinants that can actively function as an antigen-handling cell, capable of presenting antigen to T lymphocytes in a genetically restricted manner. This glomerular cell type encompasses functionally heterogeneous subpopulations that are variably phagocytic and display Fc receptors and that can be enriched by their capacity for adherence. A labeling technique in which entire glomeruli can be made permeable to antibody has shown these cells to be intrinsic to the glomerulus. They are unevenly represented in glomeruli, ranging between 5 and 20 Ia-positive cells/glomerulus.

The other two cells are Ia-negative and somewhat smaller. The intense (red) fluorescence of the *Listeria* conjugate has caused some cross-excitation in the fluorescein (green) range despite the use of appropriate filters and barriers, $\times 480$.




FIGURE 5 Phase and fluorescent micrographs of intact glomeruli permeable to anti-Ia antibody, as described in Methods. (A and B) Entire glomeruli showing the distribution of Ia-positive glomerular cells *in situ*. Glomerular stalks are seen on the right side. Note the varying positions of Ia-positive cells with a tendency toward axial distribution, $\times 240$. (C and D) Higher magnification of axial cells in other glomeruli revealing considerable heterogeneity of morphology of the Ia-positive cells as they are sited in the glomerulus, ranging from ovoid to elongated, almost tortuous shapes, $\times 480$.

Although these cells comprise up to 8% of single-cell suspensions obtained from purified glomeruli, we suspect this is too high a proportion, perhaps because these cells are more easily released by enzymatic digestion than other glomerular cell types.

If one assumes ~ 600 cells in the average glomerulus of a 150- to 200-g rat (16), a range of 5 to 20 Ia-bearing cells per glomerulus yields a more realistic estimate that 1–2% of all glomerulus cells display Ia antigens on their surface.

TABLE III
Proliferative Response of Sensitized Lymphocytes to Antigen Presentation by Glomerular Cells

Lewis T cells	Adherent glomerular cells	<i>Listeria</i> ± pulse	[³ H]Thymidine incorporation*	
			Experiment 1	Experiment 2
+	-	-	130±29	129±39
-	+	-	147±37	135±20
+	-	+	1,071±431	4,926±1,760
-	+	+	102±25	138±50
+	+	-	1,266±312	1,602±420
+	+	+	51,084±2,767	65,930±3,724
	Peritoneal exudate cells	<i>Listeria</i> ± pulse		
-	+	-	212±15	118±9
-	+	+	101±53	498±125
+	+	-	898±338	908±651
+	+	+	28,377±8,160	113,519±6,300

In both experiments, glomerular cells were cultured at 10⁵ cells/well for 2 h and the nonadherent cells removed. Some cultures were exposed to heat-killed, opsonized *Listeria* for 1 h at 37°C to permit phagocytosis; the cultures were subsequently extensively washed to remove unbound *Listeria*. Immune T lymphocytes were then added to the wells at a concentration of 4 × 10⁵ cells/well. Proliferation was assayed by tritiated thymidine incorporation after 4 d of culture. All points were measured in triplicate. As a positive control for antigen presentation, peritoneal exudate cells replaced adherent glomerular cells at a concentration of 10⁴ cells/well. Both lymphocytes and antigen-presenting cells were from Lewis strain rats.

* Arithmetic mean cpm±SE, n = 3.

It should be emphasized that these cells have been identified in well-perfused normal glomeruli and with minimal contamination by circulating cells. Further decreasing the possibility that these cells are simply circulating monocytes is our finding that, as in the mouse, 98–100% of circulating monocytes are Ia-negative; thus, no degree of contamination could account for the high percentage of Ia cells that we have observed in adherent glomerular cell cultures. Both cell-suspension and glomerular-labeling experiments have indicated that this cell type resides to a major extent in the glomerulus. Finally, electron microscopic analysis of intact glomeruli using peroxidase-labeled antibodies has localized the Ia-positive cell to the mesangium in close proximity to Ia-negative typical mesangial cells.² The Ia-positive cell is not, however, the predominant mesangial cell type, which in our cultures appears as an elongated, Ia-negative, non-phagocytic cell with spreading processes, as described by others in established cell lines (12).

Given that the Ia-positive cell is a normal resident of

the glomerulus, there are two possible conjectures concerning its lineage. The first possibility is that it is inherently of renal origin, for which we have no direct proof or disproof. The second, which we favor on morphological grounds, is that it is ultimately derived from the bone marrow, belonging to the heterogeneous grouping of tissue-resident, mononuclear phagocytes that encompass Kupffer cells, splenic and alveolar macrophages, etc. The bone marrow-derived monocytes may infiltrate the mesangium and reside in it for a period of time, acquiring properties of the more differentiated phagocyte. Its variability with respect to Fc receptor display, phagocytic capacity, and density of Ia antigens both at any one time and during sustained in vitro culture indicates that it may exist in the kidney in a variety of differentiated states. Precise definition of these issues is a subject of current investigation.

Why has this cell type resisted definition until this point? Certainly, isolated findings in previous investigations have pointed to the possible existence of such a cell. Two years ago Davies and Butcher (17) demonstrated that alloantisera against Ia antigens of the rat could be absorbed by homogenates prepared from perfused kidneys. Recently, Sakai et al. (18) have shown that unfractionated renal cells can weakly stimulate allogeneic lymphocytes in a mixed lymphocyte response assay. In long-term cultures derived from normal glomerular explants, Holdsworth et al. (19)

TABLE IV
Specificity of Antigen Presentation by Adherent Glomerular Cells to Sensitized Lymphocytes

Lewis T cells	<i>Listeria</i> pulse	[³ H]Thymidine incorporation*		
		Antigen-presenting cells		
		None	Lewis	Buffalo
+	-	311±259		
+	+	108±15		
		Glomerular cells		
-	-		196±18	121±11
-	+		358±103	577±447
+	-		1,226±97	2,724±303
+	+		34,121±2,809	2,564±1,248
		Peritoneal exudate cells		
-	-		105±19	658±272
-	+		220±86	43±3
+	-		131±23	6,989±742
+	+		43,705±4,885	14,100±726

Experimental conditions were the same as in Table III. Antigen-presenting phagocytes were from either the Lewis or Buffalo strain rats. Sensitized T cells were Lewis strain.

* Arithmetic mean cpm±SE, n = 3.

TABLE V
Stimulation of Lymphocytes in Mixed Cultures

Stimulator cells	Responder lymphocytes (Buffalo)	H ³ -Thymidine incorporation*
None	+	289±49
Lewis spleen cells	-	1,176±568
Lewis spleen cells	+	67,561±13,719
Lewis glomerular cells	-	149±11
Lewis glomerular cells	+	33,456±2,433
Buffalo glomerular cells	-	352±251
Buffalo glomerular cells	+	3,226±1,122

Stimulator cells consisted of: (a) adherent Lewis spleen cells at 2×10^5 /well, (b) adherent Lewis glomerular cells at 5×10^5 cells/well, or (c) adherent Buffalo glomerular cells at 5×10^5 cells/well. Stimulator cells were allowed to adhere to culture for 2 h, washed, and then irradiated with 1,500 rad. Allogeneic responder lymphocytes (Buffalo strain) were then added to the cultures at 2×10^5 /well. Proliferation of responder lymphocytes was assessed by measuring tritiated thymidine incorporation after 6 d of co-culture. All points were measured in triplicate.

* Arithmetic mean $\text{cpm} \pm \text{SE}$, $n = 3$.

observed a very rare cell in the outgrowing cell population, 10–20 μm in diameter, that resembled mononuclear cells. Camazine et al. (6) observed a phagocytic cell in short-term cultures of isolated glomerular cells that developed Fc receptors in prolonged culture. In the above studies, functional assays or localization of the cell were not done; nor was the presence of contaminating monocytes ruled out, a contention that has served as the prevalent explanation for the presence of such cells (19, 20).

Indeed, one review (20) has suggested that there is no evidence that mononuclear phagocytes are present in the mesangium in the absence of monocytic infiltrates induced by an inflammatory deposit such as immune complexes. Thus, in evaluating female human kidneys transplanted into males, Schiffer and Michael (21) could not find Y-body-containing cells in the transplanted kidney. After transplanting bone marrow cells with distinctive lysosomal markers into syngeneic mice, Striker et al. (22), using electron microscopy, observed phagocytosis in the glomerulus only by acutely infiltrating monocytes 24 h after injecting immune complexes intravenously. Although further investigation is required to explain the discrepancy between the above observations and our delineation of glomerular phagocytes, we believe a key issue will be the methodologic problems of sampling attendant to histochemical and ultrastructural analysis in the detection of a small phagocytic subpopulation totaling

1–2% of glomerular cells. Our approach has allowed detection of as few as 5 cells/glomerulus. Our conclusions that the phagocytic cell described in the report is normally resident in the glomerulus is consistent with previous reports of *in situ* phagocytosis of non-inflammatory substances including ferritin (23), colloidal carbon (24), and thorium (25).

The display of Ia determinants by a residential glomerular cell bears particular relevance to the subject of renal allograft and rejection. Disparity between donor and host across the I region of the major histocompatibility complex is essential to the induction of host lymphocyte immunity. With the assumption that the kidney does not display intrinsic Ia determinants, previous research devoted to abrogating the induction phase of host immunity has attempted to eliminate from the graft passenger leukocytes, the only hypothesized source of Ia (26). The relative lack of success with this approach, together with our finding an extremely vigorous mixed leukocyte reaction by allogeneic lymphocytes to Ia-positive glomerular cells, indicates that the mononuclear mesangial phagocyte might be the proper focus of future investigation into this issue.

Finally, the observations in this report potentially suggest reconsideration of the contributions of various immune mechanisms to glomerular inflammation. Only recently has the participation of circulating monocytes in experimental glomerulonephritis been delineated (3, 22), and the stimulus causing their infiltration remains obscure, except for the finding that it is not complement dependent (27). The existence of an immanent, mononuclear phagocyte bearing Ia determinants in the glomerulus enlarges the role immune cells may play in initiating or sustaining, via recruitment of additional cells, an immune reaction in the kidney. Considerable evidence has accumulated over the last decade emphasizing the pivotal position occupied by Ia-bearing phagocytes in immunological cellular interactions (28). They degrade and process antigen to selectively initiate cellular and humoral immune responses by lymphocytes. They concomitantly engage in reciprocal, modulatory interactions with lymphocytes that effect proliferation, differentiation, factor secretion, and other functions. We are currently investigating the function of this glomerular cell type in the context of such interchanges, with the additional consideration that the nonimmunological function of the glomerulus may also be affected by such interactions.

ACKNOWLEDGMENTS

We thank Barbara K. Teixeira for her efforts in preparing this manuscript, Anthony Merola and Robert Rubin for photographic assistance, and Melanie McGrory for excellent technical assistance.

REFERENCES

1. Unanue, E. R., and F. J. Dixon. 1967. Experimental glomerulonephritis: immunological events and pathogenetic mechanisms. *Adv. Immunol.* **6**: 1-90.
2. Cochrane, C. G., E. R. Unanue, and F. J. Dixon. 1965. A role of polymorphonuclear leukocytes and complement in nephrotoxic nephritis. *J. Exp. Med.* **122**: 99-119.
3. Schreiner, G. F., R. S. Cotran, V. Pardo, and E. R. Unanue. 1978. A mononuclear cell component in experimental immunological glomerulonephritis. *J. Exp. Med.* **147**: 369-384.
4. Griffith, L. D., R. E. Bulger, and B. F. Trump. 1967. The ultrastructure of the functioning kidney. *Lab. Invest.* **16**: 220-231.
5. Burlington, H., and E. P. Cronkite. 1973. Characteristics of cell cultures derived from renal glomeruli. *Proc. Soc. Exp. Biol. Med.* **142**: 143-149.
6. Camazine, S., G. Ryan, E. R. Unanue, and M. J. Karnovsky. 1976. Isolation of phagocytic cells from the rat renal glomerulus. *Lab. Invest.* **35**: 315-326.
7. Williams, A., G. Gulfre, and C. Milstein. 1977. Analysis of cell surfaces by xenogeneic myeloma-hybrid antibodies: differentiation antigens of rat lymphocytes. *Cell.* **12**: 663-673.
8. McMaster, W. R., and A. F. Williams. 1979. Identification of Ia glycoproteins in rat thymus and purification from rat spleen. *Eur. J. Immunol.* **9**: 426-433.
9. Cowing, C., B. C. Schwartz, and H. Dickler. 1979. Macrophage Ia antigens. I. Macrophage populations differ in their expression of Ia antigens. *J. Immunol.* **120**: 378-384.
10. Beller, D. I., J-M. Kiely, and E. R. Unanue. 1980. Regulation of macrophage populations. I. Preferential induction of Ia-rich peritoneal exudates by immunologic stimuli. *J. Immunol.* **124**: 1426-1432.
11. Farr, A. G., J-M. Kiely, and E. R. Unanue. 1979. Macrophage-T cell interactions involving *Listeria monocytogenes*—role of the H-2 gene complex. *J. Immunol.* **122**: 2395-2404.
12. Ausiello, D., J. Kreisberg, C. Roy, and M. Karnovsky. 1980. Contraction of cultured rat glomerular cells of apparent mesangial origin after stimulation with Angiotensin II and arginine vasopressin. *J. Clin. Invest.* **65**: 754-760.
13. Steinman, R., N. Nogueira, M. Witmer, J. Tydings, and I. Mellman. 1980. Lymphocyte enhances the expression and synthesis of Ia antigens on cultured mouse peritoneal macrophages. *J. Exp. Med.* **152**: 1248-1255.
14. Beller, D. I., and E. R. Unanue. 1981. Regulation of macrophage populations. II. Synthesis and expression of Ia antigens by peritoneal exudate macrophages is a transient event. *J. Immunol.* **126**: 263-269.
15. Weinberg, D. S., and E. R. Unanue. 1981. Antigen-presenting function of alveolar macrophages: uptake and presentation of *Listeria monocytogenes*. *J. Immunol.* **126**: 794-799.
16. Olivetti, G., P. Anversa, M. Melissari, and A. V. Loud. 1980. Morphometry of the renal corpuscle during post-natal growth and compensatory hypertrophy. *Kidney Int.* **17**: 438-444.
17. Davies, H., and G. Butcher. 1978. Kidney alloantigens determined by two regions of the rat major histocompatibility locus. *Immunogenetics.* **6**: 171-181.
18. Sakai, A., K. Yakushiji, and S. Mashimo. 1980. Lymphocyte stimulation by allogeneic tissue cells and in rats: with special reference to differential survival of skin and kidney allografts. *Transplant. Proc.* **12**: 74-81.
19. Holdsworth, S. R., E. F. Glasgow, R. C. Atkins, and N. M. Thomson. 1978. Cell characteristics of cultured glomeruli from different animal species. *Nephron.* **22**: 454-459.
20. Michael A. F., W. F. Keene, L. Raji, R. Vernier, and S. Maur. 1980. The glomerular mesangium. *Kidney Int.* **17**: 141-154.
21. Schiffer, M. S., and A. F. Michael. 1978. Renal cell turnover by Y-chromosome (Y-body) staining of the transplanted human kidney. *J. Lab. Clin. Med.* **92**: 841-848.
22. Striker, G. E., M. Mannik, and M. Tung. 1979. Role of marrow-derived monocytes and mesangial cells in removal of immune complexes from renal glomeruli. *J. Exp. Med.* **149**: 127-136.
23. Farquar, M. G., and G. E. Palade. 1962. Functional evidence for the existence of a third cell type in the renal glomerulus. *J. Cell Biol.* **13**: 55-87.
24. Elema, J. D., J. R. Hoyer, and R. L. Vernier. 1976. The glomerular mesangium: uptake and transport of intravenously injected colloidal carbon in rats. *Kidney Int.* **9**: 394-406.
25. Latta, H., and A. B. Mannsbach. 1962. Relationship of the centrilobular region of the glomerulus to the juxtaglomerular apparatus. *J. Ultrastruct. Res.* **6**: 562-578.
26. Stuart, F. P., E. Bastien, A. Holter, F. W. Fitch, and W. L. Elkins. 1971. The role of passenger leukocytes in the regulation of renal allografts. *Transplant. Proc.* **3**: 461-464.
27. Unanue, E. R., G. F. Schreiner, and R. S. Cotran. 1981. A role of mononuclear phagocytes in immunologically induced glomerulonephritis. In *Immune Mechanisms in Renal Disease*. A. Michael and N. Cummings, editors. In press.
28. Unanue, E. R. 1981. The regulatory role of macrophages in antigenic stimulation. II. Symbiotic relationship between lymphocytes and macrophages. *Adv. Immunol.* **31**: 1-136.