

Hydrogen ion secretion by the collecting duct as a determinant of the urine to blood PCO₂ gradient in alkaline urine.

T D DuBose Jr

J Clin Invest. 1982;69(1):145-156. <https://doi.org/10.1172/JCI110425>.

Research Article

Several theories have been advanced to explain the elevation in urinary PCO₂ during bicarbonate loading and include: (a) H⁺ secretion, (b) countercurrent system for CO₂, (c) the "ampholyte" properties of bicarbonate, and (d) mixing of urine of disparate bicarbonate and buffer concentrations. In this study microelectrodes were used to measure in situ and equilibrium pH (pHis and pHeq) and PCO₂ in control and bicarbonate loaded rats before and after infusion of carbonic anhydrase. The disequilibrium pH method (pH_{dq} = pH_{is} - pH_{eq}) was used to demonstrate H⁺ secretion. Control rats excreting an acid urine (pH = 6.04 +/- 0.06) failed to display a significant disequilibrium pH at the base (BCD), or tip (TCD) of the papillary collecting duct. Urine pH (7.54 +/- 0.12), and urine to blood (U-B) PCO₂ increased significantly during NaHCO₃ loading while PCO₂ at the BCD and TCD also increased (95 +/- 4 and 122 +/- 4). Furthermore, an acid disequilibrium pH was present at both the BCD and TCD (-0.42 +/- 0.04 and -0.36 +/- 0.03) and was obliterated by carbonic anhydrase. Comparison of the PCO₂ in the BCD or TCD with the adjacent vasa recta revealed similar values ($r = 0.97$). It is concluded that H⁺ secretion by the collecting duct into bicarbonate containing fluid with delayed dehydration of H₂CO₃, is the most likely determinant [...]

Find the latest version:

<https://jci.me/110425/pdf>



Hydrogen Ion Secretion by the Collecting Duct as a Determinant of the Urine to Blood PCO_2 Gradient in Alkaline Urine

THOMAS D. DUBOSE, JR., with the technical assistance of LEO R. PUCACCO and
JOHNNIE M. GREEN, *Department of Internal Medicine, University of Texas
Health Science Center at Dallas, Southwestern Medical School,
Dallas, Texas 75235; University of Texas Medical Branch at Galveston,
Texas 77550*

A B S T R A C T Several theories have been advanced to explain the elevation in urinary PCO_2 during bicarbonate loading and include: (a) H^+ secretion, (b) countercurrent system for CO_2 , (c) the "ampholyte" properties of bicarbonate, and (d) mixing of urine of disparate bicarbonate and buffer concentrations. In this study microelectrodes were used to measure *in situ* and equilibrium pH (pH_{is} and pH_{eq}) and PCO_2 in control and bicarbonate loaded rats before and after infusion of carbonic anhydrase. The disequilibrium pH method ($\text{pH}_{\text{dq}} = \text{pH}_{\text{is}} - \text{pH}_{\text{eq}}$) was used to demonstrate H^+ secretion. Control rats excreting an acid urine ($\text{pH} = 6.04 \pm 0.06$) failed to display a significant disequilibrium pH at the base (BCD), or tip (TCD) of the papillary collecting duct. Urine pH (7.54 ± 0.12), and urine to blood (U-B) PCO_2 increased significantly during NaHCO_3 loading while PCO_2 at the BCD and TCD also increased (95 ± 4 and 122 ± 4). Furthermore, an acid disequilibrium pH was present at both the BCD and TCD (-0.42 ± 0.04 and -0.36 ± 0.03) and was obliterated by carbonic anhydrase. Comparison of the PCO_2 in the BCD or TCD with the adjacent vasa recta revealed similar values ($r = 0.97$). It is concluded that H^+ secretion by the collecting duct into bicarbonate containing fluid with delayed dehydration of H_2CO_3 , is the most likely determinant of the U-B PCO_2 in alkaline urine. Similar values for PCO_2 in the collecting

duct and the adjacent vasa recta suggests trapping of CO_2 in the medullary countercurrent system. The rise in PCO_2 occurs both along the collecting duct and after exit from the papilla.

INTRODUCTION

There has been general agreement since the early observations of Mainzer and Bruhn (1), Pitts and Lotspeich (2), and Ryberg (3), that the CO_2 tension of alkaline urine may exceed that of systemic arterial blood by two- to fourfold. Considerable controversy has existed regarding the mechanism by which this elevation in urinary PCO_2 occurs (4-7). Despite this controversy the urine to blood PCO_2 gradient (U-B PCO_2)¹ during bicarbonate administration has been widely used as an index of hydrogen ion secretion by the "distal nephron" in studies involving whole kidney clearance techniques in experimental animals and in man (7-11). In fact, the failure of patients with classical distal renal tubular acidosis to generate an elevated urinary PCO_2 during bicarbonate loading has been proposed as a means of categorizing the type of urinary acidification defect in this disorder ("secretory defect") (9).

Pitts and Lotspeich (2) first proposed that H^+ secretion into bicarbonate-containing fluid in the more distal nephron segments resulted in the formation and subsequent delayed dehydration of H_2CO_3 . Thus, as envisioned by these investigators, CO_2 would be formed in areas of the collecting system where surface-volume relationships would be unfavorable for CO_2 diffusion, resulting in elevated urinary CO_2 tensions.

A preliminary report of this study was presented at the Annual Meeting of the American Society of Nephrology, Washington, D. C., 23-25 November 1980.

Dr. DuBose's present address is Renal-Electrolyte Physiology Laboratory, Department of Internal Medicine, Division of Nephrology, University of Texas Medical Branch, Galveston, Tex.

Received for publication 9 February 1981 and in revised form 18 September 1981.

¹ Abbreviations used in this paper: pH_{dq} , disequilibrium pH; pH_{eq} , equilibrium pH; pH_{is} , *in situ*.

Additional support for this hypothesis was derived from the studies of Ochwaldt and Pitts (12) in which it was observed that systemic administration of carbonic anhydrase, which presumably appeared in the urine, obliterated the U-B PCO_2 gradient. In contrast, Kennedy, Orloff, and Berliner (13) emphasized that the mixing of urine of low bicarbonate and high non-bicarbonate buffer concentration with urine of high bicarbonate, low nonbicarbonate buffer concentration (at different pH) could also serve to elevate the urinary PCO_2 ("mixing hypothesis") (13).

Pak Poy and Wrong (14) noted that high urinary PCO_2 could also be achieved by a medullary countercurrent system but these authors suggested that the CO_2 would be derived primarily from metabolic sources. Uhlich, Baldamus, and Ullrich (15) suggested by indirect techniques, that the CO_2 tension in the collecting tubule exceeded that of the vasa recta by ~ 30 mm Hg. Furthermore, this sizable gradient was abolished by carbonic anhydrase infusion. This study cast doubt on the medullary countercurrent hypothesis and further underscored the importance of delayed dehydration. Although of an indirect nature, this study has served as the foundation of the U-B PCO_2 gradient method used in whole kidney clearance studies. The suggestion of a diffusion barrier for CO_2 across the collecting duct (5, 6, 15) differs markedly from recent findings across the superficial proximal convoluted tubule reported by our laboratory (16, 17).

Recently, the H^+ secretory hypothesis has been called into question by physicochemical considerations. Arruda and associates (5) and Maren (6) have emphasized the "ampholyte" properties of bicarbonate as an explanation for the expected increase in CO_2 tension in highly alkaline aqueous solution or urine when the concentration of carbonate (CO_3^{2-}) and CO_2 are equal. The linear relationship between the U-B PCO_2 gradient and urinary bicarbonate concentration observed by Arruda et al. (8) in a variety of animal models and man have led to the hypothesis that a large component of the U-B PCO_2 gradient is a result of the ampholyte effect initiated by water abstraction in the distal nephron and a subsequent increase in bicarbonate concentration (5). Thus, these investigators suggested that the low U-B PCO_2 gradient observed in distal renal tubular acidosis (9) was critically dependent on urinary concentrating ability, which is often defective in this disease (5). These findings were further supported by the theoretical considerations of Maren (6). In contrast, however, Stinebaugh and associates (7) have recently reevaluated the linear relationship between U-B PCO_2 and urinary bicarbonate concentration in alkaline urine in several species. These studies led these authors to the conclusion that the most plausible explanation for this linear relation-

ship was a H^+ secretory process that increased as a function of the increase in urine bicarbonate concentration (7). It was suggested that a disequilibrium pH could occur as a result of H^+ secretion or alternatively by concentration of bicarbonate in the terminal nephron with simultaneous back diffusion of carbon dioxide (7). These studies were necessarily of an indirect nature, however, since a disequilibrium pH has not been demonstrated previously in the collecting tubule.

The disequilibrium pH method ($\text{pH}_{\text{dq}} = \text{pH}_{\text{is}} - \text{pH}_{\text{eq}}$) has been used by several investigators to demonstrate delayed dehydration, and thus H^+ secretion in other nephron segments (18, 19). Recently, we have used newly developed microelectrode techniques to measure PCO_2 , *in situ* pH (pH_{is}), and equilibrium pH (pH_{eq}).

The purpose of the present study was to examine the role of delayed dehydration in the generation of the U-B PCO_2 gradient by micropuncture of the surgically exposed papilla of the rat in the presence and absence of carbonic anhydrase.

METHODS

Preparation of rats for micropuncture. Studies were performed after 100 mg/kg, i.p. Inactin anesthesia (Promonta, Hamburg, West Germany) on young mutant Munich-Wistar rats weighing 60–150 g. All rats were allowed free access to tap water and standard rat chow until the time of the experiment. The rat was placed on a thermostatically controlled heating table and maintained at 37.5°C. After tracheostomy, polyethylene catheters (PE 50) were inserted into the left jugular vein for infusion and into the left femoral artery for constant blood pressure monitoring and blood collection, and into the bladder for urine collection from the right nonexperimental kidney (under oil). The left kidney was then gently separated from the adrenal gland and peritoneal attachment. The renal papilla was exposed by temporarily displacing the papilla into the renal pelvis and carefully excising the ureter. The kidney was then placed in a lucite cap stabilized by 3% agar and continuously bathed with mineral oil equilibrated with 5% CO_2 –95% O_2 , maintained at 37°C and illuminated with a small fiber optic light source. This technique has been described previously in detail (20). After jugular vein cannulation, rats were infused with Ringer's bicarbonate ($\text{Na}^+ = 140$, $\text{Cl}^- = 110$, $\text{HCO}_3^- = 25$, $\text{K}^+ = 5$ meq/liter) at 1% of body wt/h.

Microelectrode techniques

PCO_2 microelectrode. The *in situ* PCO_2 of tubule fluid at the base of the collecting duct (defined as the earliest accessible portion of the papillary collecting duct), tip of the collecting duct (opening of duct or direct puncture at tip) (distance between tubule puncture sites of 2.1 ± 0.5 mm), and an adjacent vasa recta were obtained by puncture at each site with a PCO_2 microelectrode of 6–9 μm tip Diam. The construction, testing, electrical characteristics, and calibration of these electrodes were exactly as described previously (16). Electrodes having a sensitivity of <57 mV/ $\log_{10} \text{PCO}_2$ were discarded.

***In situ* pH.** The *in situ* pH (pH_{is}) at the base and tip

collecting duct was determined with single or double barreled glass membrane pH microelectrodes of 7–10 μm tip Diam as reported previously (18). Electrodes having a sensitivity of <57 mV/pH unit were not used. Initially all collecting duct punctures were with double-barreled electrodes, however, no difference in pH was noted with single or double-barreled electrodes during any of the physiological conditions examined. Subsequently, in five rats trans-epithelial potential difference was determined with 2–3- μm beveled pipettes filled with 2.5 M KCl and 0.5 M KNO_3 . In 15 tubules the range of potential was –1.0 to +2.0 mV in controls and –3.0 to 0.0 mV during bicarbonate loading. These findings indicate that the transepithelial potential difference in the papillary collecting duct of the exposed papilla (length ~2.0 mm) is not of sufficient magnitude to affect adversely the accuracy of a single-barreled pH electrode having near theoretical slope. Therefore, most of the *in situ* pH data were obtained with single-barreled electrodes because of ease of construction. Calibration before and after *in vivo* use was as described previously (16, 18).

Equilibrium pH. The equilibrium pH (pH_{eq}) was measured *in vivo* with a composite probe consisting of an aspiration pipette into which a single-barreled pH electrode was inserted. This electrode was designed and constructed in our laboratory and has been described in detail (18). This electrode consists of an outer shell (aspiration or collection pipette) of borosilicate glass capillary (2 mm) that was pulled to a taper length of 6–7 mm and beveled at a 50–60° angle to an outer tip Diam of 4–9 μm . A pH electrode was constructed as previously described (18) except that the geometry of the outer shell and pH electrode were taken into account to allow insertion of the pH electrode up to a distance of some 200–400 μm from the tip of the outer shell. After sealing the butt end with epoxy cement and allowing a suitable cure, the pipette was filled with mercury and attached to a hydraulic device via PE-50 tubing. Both the tubing and hydraulics were filled with mercury. Therefore, standard pH buffer or tubule fluid was aspirated behind a mercury column to avoid CO_2 loss. Before use this electrode was calibrated by aspirating standard pH buffer at 37°C ($\text{pH} = 6.84$ and 7.384, respectively) into the pipette under microscopic control to assure that the pH electrode was in contact with the buffer. An attempt was made to withdraw a volume of buffer similar to that expected *in vivo*. Only electrodes with a sensitivity of at least 57 mV/pH unit were accepted. The lucite chamber used for calibration at 37°C was as reported previously (16). In actual use in a micro-puncture experiment, tubule fluid was aspirated and allowed to reach chemical equilibrium while the aspiration pipette tip remained within the tubule lumen. The theoretical advantages of an equilibrium pH electrode constructed in this manner were: (a) it was not necessary to remove tubule fluid to an equilibration chamber *in vitro*, and (b) no assumption regarding the level of papillary PCO_2 was required. This electrode was tested extensively *in vitro* to: (a) assure isolation of the sample with respect to ionic diffusion of H^+ into or out of the tubule sample, (b) assure that CO_2 gas was not lost over the period of time required to make the pH measurement, and (c) to assure that the CO_2 produced from the dehydration of the H_2CO_3 contained in the tubule sample from an “off-equilibrium” source (i.e., where a disequilibrium pH was present) would not erroneously lower the true equilibrium pH. The results of this testing have been reported previously (18). With regard to the third problem however, nonbicarbonate buffers could also contribute as a potential source of CO_2 by this mechanism. For example if the concentration of nonbicarbonate buffers was 5.0 mM

(7), and the disequilibrium pH equal to –0.50 pH U, then 2.5 mM of CO_2 could be produced as the H_2CO_3 was titrated by the nonbicarbonate buffers. This means that if this CO_2 were trapped in the equilibrium pH electrode, the equilibrium pH observed would be falsely acid. Such a problem would be highly unlikely since the tip of the equilibrium electrode through which collection is made is open and remains within the tubule lumen during operation. However, to assure that such a problem did not exist in our system, the determined equilibrium pH was compared with the equilibrium pH calculated from determined PCO_2 and total CO_2 concentration (Henderson-Hasselbach equation, $\alpha = 0.0309$, $\text{pK}_a^1 = 6.13$). In these experiments care was taken to assure that the loss of CO_2 gas from the collected sample was negligible by aspirating tubule fluid directly into a beveled volumetric constriction pipette between oil layers equilibrated with CO_2 . The sample was then transferred immediately to the microcalorimeter (21). The results of the comparison of determined and calculated equilibrium pH (pH_{eq}) from five rats are displayed in Fig. 1. Each point represents a paired determination. Thus, since there is a close correlation between the equilibrium pH determined by these two techniques, the type of consideration noted above could not be adversely affecting the results and, in this system, nonbicarbonate buffers could not contribute significantly to the equilibrium pH measured in the papillary collecting duct.

Physiological conditions

Surgical exposure of the renal papilla in the mutant Mu-nich-Wistar rat would be expected to disrupt the normal anatomical relationship whereby the papilla is bathed in pelvic urine. In addition, exposure would result in the loss of CO_2 gas from the exposed papilla. Therefore, preliminary




FIGURE 1 Comparison of pH_{eq} calculated from $[\text{tCO}_2]$ and PCO_2 determinations (ordinate) vs. pH_{eq} measured by aspiration equilibrium pH microelectrode. Two techniques are not significantly different ($r = 0.956$) ($\text{pH}_{\text{eq}}^{\text{cal}} = 0.74 \text{ pH}_{\text{eq}}^{\text{meas}} + 1.9$).

experiments were designed to allow direct determination of the PCO_2 of urine in the pelvic space. PCO_2 microelectrodes of 15–20 μm tip Diam were assembled. Six hydropenic rats in normal acid-base balance, and excreting an acid urine equal to 6.10 pH U were prepared for micropuncture and the renal pelvis was left intact. The mean PCO_2 of urine in the pelvic space surrounding the renal papilla was 36.5 ± 1.2 mm Hg. When the PCO_2 electrode was then advanced into the papillary interstitium, equal values were obtained. In subsequent experiments (below) in which the renal pelvis was surgically excised, the renal papilla was bathed in heated mineral oil equilibrated with 5% CO_2 . To assure that the mineral oil surrounding the papilla was maintained at 35 mm Hg, the PCO_2 of the oil was frequently measured with a PCO_2 microelectrode of 6–9 μm tip Diam.

Group I (controls) (n = 7)

Seven mutant Munich-Wistar rats served as controls after surgical exposure of the papilla while receiving a maintenance infusion of Ringer's bicarbonate at 1% body wt/h. Care was taken to maintain acid-base balance as described previously (16).

Group II (bicarbonate loading) (n = 25)

25 animals received 300 mM NaHCO_3 at 1.8% body wt/h in an attempt to minimize volume overexpansion. Micropuncture was initiated after a 1.5-h equilibration period.

Group III (bicarbonate load plus carbonic anhydrase) (n = 14)

14 of the same rats in group II then received carbonic anhydrase as an intravenous bolus (10 mg) and maintenance infusion (20 mg/h). Micropuncture was initiated 30 min after the bolus injection. Carbonic anhydrase was prepared from bovine erythrocytes (C-7500, Sigma Chemical Co., St. Louis, Mo.) and assayed for activity *in vitro* before use. The presence of carbonic anhydrase activity in urine from the right kidney was verified by the micromethod of Maren (22).

Arterial blood pH and PCO_2 and urine pH and PCO_2 were determined on a blood gas analyzer (model 165, Corning Medical, Medfield, Mass.). The blood $[\text{HCO}_3^-]$ was calculated with the Henderson-Hasselbalch equation ($\alpha = 0.0301$, $\text{pK}_a = 6.10$), whereas for urine $[\text{HCO}_3^-]$ the α was 0.0309, and the pK_a was corrected for ionic strength by the method of Hastings and Sendroy (23).

The results are expressed as mean \pm SE in each group. Statistical significance was calculated using the Student's *t* test for paired or unpaired data as appropriate.

RESULTS

The systemic arterial and urinary pH, PCO_2 and $[\text{HCO}_3^-]$ values and the U-B PCO_2 for all three groups are displayed in Table I. The control rats (group I) were in normal acid-base balance and excreted an acid urine. These findings are consistent with previous controls in our laboratory (16, 18, 21). The U-B PCO_2 gradient was less than zero in this condition ($P < 0.01$). The animals receiving 300 mM NaHCO_3 (groups II and III) developed an acute metabolic alkalosis and

excreted an alkaline urine (untouched kidney). In group II rats the PCO_2 in urine from the right kidney was 136.3 ± 11.8 mm Hg, so that the U-B PCO_2 gradient was 97.4 ± 10.6 mm Hg. After administration of carbonic anhydrase the urine pH increased significantly (7.54 ± 0.12 to 8.17 ± 0.05), the urine PCO_2 decreased significantly (136.3 ± 11.8 to 59.5 ± 2.3 mm Hg) and the U-B PCO_2 decreased, as expected (97.4 ± 10.6 to 17.7 ± 4.0) ($P < 0.001$). The U-B PCO_2 remained significantly greater than zero after carbonic anhydrase ($P < 0.01$), however.

Micropuncture data

The micropuncture findings are displayed in Tables II, III, IV, and Figs. 2–5.

PCO_2 . Control rats (group I) excreting an acid urine (6.04 ± 0.06) had papillary collecting duct PCO_2 values that were very similar to systemic arterial values (35.2 ± 1.2 at the base, and 36.5 ± 1.5 mm Hg at the tip of the collecting duct) (Table II.). Thus the U-B PCO_2 , and the papillary-blood PCO_2 values were similar in control animals.

Bicarbonate-loaded rats (group II) excreting an alkaline urine displayed markedly elevated values for PCO_2 at the base (95.4 ± 4.1) and tip collecting duct (122.2 ± 4.3) (Table II, Fig. 2). The increase in PCO_2 from base to tip was significant ($P < 0.01$). Thus, the directly measured papillary collecting duct tip minus systemic blood PCO_2 (PCD-B PCO_2) was 54.1 ± 4.1 and 79.6 ± 4.3 , respectively (Tables II and III). As displayed in Table III and Fig. 3 the U-B PCO_2 from the untouched right kidney was slightly, but not significantly, greater than the PCD-B PCO_2 at the tip collecting duct. When 10 rats in this group having a urine pH nearer 8.0 pH units were analyzed separately, similar results were obtained. Specifically, values for arterial blood ($\text{pH}_a = 7.53 \pm 0.05$, $\text{PACO}_2 = 44 \pm 2.1$, $[\text{HCO}_3^-] = 36.8 \pm 1.1$) and urine ($\text{pH} = 7.93 \pm 0.06$ and $\text{PCO}_2 = 92 \pm 4.2$) were more alkaline and the urine PCO_2 slightly lower. The micropuncture findings for PCO_2 in these 10 rats were as follows: BCD = 90 ± 4.7 and TCD = 115 ± 6.8 . These findings do not differ from the results obtained in all 25 rats in group II. Therefore, these results were combined.

In the 14 rats receiving NaHCO_3 and carbonic anhydrase infusion (group III) the PCO_2 at the base and tip collecting duct decreased markedly after infusion of the enzyme (95.4 to 68.1 and 122.2 to 78.3, respectively) (Table II). As shown in Fig. 3, carbonic anhydrase markedly reduced the U-B and PCD-B PCO_2 but did not obliterate this difference so that values of PCO_2 in both collected urine (right kidney) and the microelectrode determined value at the papillary tip (left kidney) remained significantly greater than sys-

TABLE I
Systemic Arterial and Urine Values

Group	Arterial blood			Urine (right-untouched)			U-B P_{CO_2}
	pH	P_{CO_2}	[HCO ₃ ⁻]	pH	P_{CO_2}	[HCO ₃ ⁻]	
I. Controls							
Mean	7.36	37	20.1	6.04	32	0.9	-5
SEM (n)	± 0.01 (7)	± 1	± 0.37	± 0.06 (7)	± 2.5	—	± 1.3 (7)
II. 300 mM NaHCO₃							
Mean	7.50	44	33.0	7.54	136	108.3	97
SEM (n)	± 0.02 (25)	± 1.7	± 0.9	± 0.12 (25)	± 11.8	± 9.3	± 10.6 (25)
III. 300 mM NaHCO₃ + Carbonic anhydrase							
Mean	7.55	41	34.1	8.17	59	162.5	18
SEM (n)	± 0.02 (14)	± 1.9	± 1.1	± 0.05 (14)	± 2.3	± 10.2	± 4.0 (14)
<i>P</i> (II vs. III)	NS	NS	NS	<0.001	<0.001	<0.01	<0.001

temic arterial blood. (17.7 \pm 4.0 and 39.8 \pm 3.0, respectively) (Table III and Fig. 3).

Comparison of collecting duct and vasa recta P_{CO_2} . In five control and five bicarbonate-loaded rats comparison of P_{CO_2} at either the base or tip of the collecting duct with the P_{CO_2} in the immediately adjacent vasa recta was made (Fig. 4). Each point represents a paired tubule-vasa recta determination. Note that a highly significant correlation ($r = 0.97$) was obtained.

Disequilibrium pH. The values of the microelectrode determined *in situ* pH (pH_{is}), equilibrium pH (pH_{eq}) and the difference, or the disequilibrium pH (pH_{dq}) for controls, bicarbonate-loaded, and bicarbonate-loaded and carbonic anhydrase infused rats are displayed in Table IV for each micropuncture site. In control rats values for pH_{is} and pH_{eq} were similar at both the base (6.51 \pm 0.06, 6.44 \pm 0.06) and tip (6.47 \pm 0.07, 6.45 \pm 0.06) of the collecting duct so that a disequilibrium pH was not observed (i.e., pH_{dq} not

TABLE II
Micropuncture Data: P_{CO_2} Papillary Collecting Duct

Condition	P_{CO_2}		PCD-B P_{CO_2} *	
	Base	Tip	Base	Tip
I. Control				
Mean	35	36	-3	-3
SEM (n)	± 1.2 (15)	± 1.5 (20)	± 1.3	± 1.4
II. NaHCO₃				
Mean	95	122	54	80
SEM (n)	± 4.1 (38)	± 4.3 (52)	± 4.1	± 4.3
III. NaHCO₃ + Carbonic anhydrase				
Mean	68	78	29	40
SEM (n)	± 2.4 (44)	± 2.8 (50)	± 3.2	± 3.0
<i>P</i> II vs. III	<0.001	<0.001	<0.001	<0.001

* Papillary collecting duct minus systemic blood P_{CO_2} .

TABLE III
Comparison of Right (Untouched) and Left (Experimental) Kidney

Condition	Urine		Tip PCD	
	P _{CO₂}	U-B P _{CO₂} [*]	P _{CO₂}	P-B P _{CO₂} [†]
II. NaHCO ₃	136 ±11.8 (25)	97 ±10.6 (25)	122 ±4.3 (52)	80 ±4.3 (52)
III. NaHCO ₃ + Carbonic anhydrase	59 ±2.3 (14)	18 ±4.0 (14)	78 ±2.8 (50)	40 ±3.0 (50)
P	<0.001	<0.001	<0.001	<0.001

* Urine minus blood P_{CO₂}.

† Papillary minus blood P_{CO₂}.

different than zero; $+0.07 \pm 0.04$ and $+0.02 \pm 0.04$, respectively). In bicarbonate-loaded rats (group II) *in situ* pH was significantly more acid than equilibrium pH at both sites (7.24 ± 0.08 vs. 7.66 ± 0.06 at the base $P < 0.001$, and 7.35 ± 0.09 vs. 7.71 ± 0.05 at tip $P < 0.01$) (Table IV and Fig. 5). Thus, a significant acid disequilibrium pH was observed at both base (-0.42 ± 0.04) and tip collecting duct (-0.36 ± 0.03). In the 10 rats in group II analyzed separately and having a mean urine pH of 7.93 ± 0.03 U, the values for disequilibrium pH were not different ($P > 0.05$), i.e., base collecting duct $= -0.40 \pm 0.04$ and tip collecting duct $= -0.35 \pm 0.03$ U, (significantly greater than zero, $P < 0.01$). Carbonic anhydrase infusion (group III) completely obliterated the disequilibrium pH, at both sites, and did so by increasing *in situ* pH to values not different from equilibrium pH ($pH_{eq} = +0.06$ and

$+0.03$, respectively) (Fig. 5). Note that equilibrium pH in groups II and III did not differ significantly in either the presence or absence of carbonic anhydrase (Fig. 5).

DISCUSSION

Of the several theories advanced to explain the observed increase in urinary P_{CO₂} above systemic arterial levels during bicarbonate loading, H⁺ secretion into bicarbonate-containing fluid has been the most widely accepted mechanism (2, 4, 7, 10, 12). Recent studies have emphasized the "ampholyte" properties of bicarbonate and the critical importance of urinary concentrating ability (5, 6), thus seriously questioning the contribution of H⁺ secretion to the generation of the elevated U-B P_{CO₂} difference. The present study was

TABLE IV
Disequilibrium pH Papillary Collecting Duct

Condition	Base CD			Tip CD		
	pH _b	pH _{eq}	pH _{de}	pH _b	pH _{eq}	pH _{de}
I. Controls	6.51 ±0.06	6.44 ±0.06	+0.07 ±0.04 (28) NS	6.47 ±0.07	6.45 ±0.06 (33) NS	+0.02 ±0.04 NS
P vs. 0						
II. NaHCO ₃	7.24 ±0.08	7.66 ±0.06	-0.42 ±0.04 (48) <0.001	7.35 ±0.09	7.71 ±0.05 (26) <0.01	-0.36 ±0.03 NS
P vs. 0						
III. NaHCO ₃ + Carbonic anhydrase	7.74 +0.06	7.68 ±0.06	+0.06 ±0.05 (18) NS	7.73 ±0.08 (14) NS	7.70 ±0.05 (14) NS	+0.03 ±0.03 NS
P vs. 0						
II vs. III			<0.001			<0.001




FIGURE 2 Microelectrode PCO₂ values in NaHCO₃ loaded (black bars) and NaHCO₃ loaded and carbonic anhydrase infused rats (stippled bars) at both micropuncture sites, the base and tip of the papillary collecting duct. The dashed line represents mean systemic arterial PCO₂. The reduction in PCO₂ after carbonic anhydrase is highly significant. *P < 0.001.

designed to evaluate PCO₂ and the presence, or absence, of H⁺ secretion by recently developed microelectrode techniques employed in a papillary micropuncture setting. This study represents the first report of direct determination of these parameters in the rat and thus adds insight to the process by which elevated urinary CO₂ tensions are achieved.

Several new findings emerge from these studies: (a) The demonstration of a significant acid disequilibrium pH in conjunction with a significantly elevated CO₂ tension in the papillary collecting duct during bicar-




FIGURE 3 Comparison of right urinary PCO₂ (dots) and left papillary tip PCO₂ (triangles) minus blood PCO₂ (U-B PCO₂) in NaHCO₃ loaded animals before and after carbonic anhydrase (CA).




FIGURE 4 Comparison of PCO₂ in the papillary collecting duct (abscissa) and the immediately adjacent vasa recta (ordinate). Each point represents a paired determination and includes controls and bicarbonate loaded rats. The PCO₂ is similar in both structures ($r = 0.967$, $y = 1.097x - 6.19$).

bionate loading supports the view that hydrogen ion secretion is a determinant of the U-B PCO₂ gradient. (b) The observed increase in PCO₂ above systemic levels occurred both along the papillary collecting duct and after exit from the papilla. (c) Comparison of CO₂




FIGURE 5 Summary of pH determinations in NaHCO₃ loaded rats before and after carbonic anhydrase (CA) infusion (left column) at both micropuncture sites (base and tip). The top (black) bar in both conditions at each site represents *in situ* pH while the hatched bars represent equilibrium pH. The values for the resulting pH_{dq} are displayed in the right column. The starred values for pH_{dq} are significantly greater than zero. (P < 0.001).

tension in both collecting duct fluid and the adjacent vasa recta revealed similar values, suggesting trapping of CO_2 in the medullary countercurrent system.

It has been widely appreciated that the PCO_2 of urine in man and experimental animals is near or below systemic arterial levels during normal or control conditions (4). In the present study it is demonstrated that the PCO_2 measured with a microelectrode by micropuncture of the papillary collecting duct of control rats is similar to the PCO_2 of systemic arterial blood (34.5 vs. 36.6 mm Hg). Furthermore, direct puncture of the pelvic space prior to exposure of the renal papilla revealed similar values for papillary collecting duct and pelvic space PCO_2 . Previous explanations for urinary CO_2 tensions, at or below plasma values during the excretion of an acid urine have included (a) reabsorption of bicarbonate in the terminal nephron decreasing the concentration of HCO_3^- generating a disequilibrium between CO_2 and H_2CO_3 (24), (b) a mixture of urine with differing concentrations of bicarbonate and nonbicarbonate buffers (13), and (c) passive back diffusion of undissociated organic acids (25). As pointed out by Rector (4) none of these explanations serve to support either H^+ secretion or HCO_3^- reabsorption in the collecting duct as a determinate of urinary CO_2 tensions. The failure to demonstrate a significant disequilibrium pH in the papillary collecting duct of control rats in the present study indicates that bicarbonate reabsorption in this segment, during this condition, is exceedingly low. This is not surprising in view of the very low luminal concentrations of bicarbonate and high concentrations of non-bicarbonate buffer in the collecting duct during excretion of an acid urine. Although the value for collecting duct PCO_2 agrees closely with the recent findings of Gruber and associates (26), we have not, in contrast, demonstrated an acid disequilibrium pH in the collecting duct of control rats. Gruber and associates, demonstrated an acid disequilibrium pH in control (-0.26 pH units) and acidotic rats (-0.18 pH U). The microcatheterization technique used by Gruber et al. (27) allows examination of a larger portion of the medullary collecting duct than in the present study. Different electrode techniques were used in the former study, as well, since aspiration into the microcatheter was required for the fluid to make contact with the pH electrode while equilibrium pH was measured *in vitro*. Based on the dehydration rate constant of $49.6/\text{s}$ (28), the equilibrium concentration of H_2CO_3 would be approached with a half-time of 1.4×10^{-2} s. Although nonbicarbonate buffers would reduce the disequilibrium pH by consumption of H_2CO_3 , this effect might be offset by prolongation of the time to reach equilibrium. These differences are not easily resolved, therefore. These investigators have interpreted their

findings as consistent with acidification by a H^+ secretory process along the medullary collecting duct in these conditions (27).

The fact that the PCO_2 of highly alkaline urine exceeds systemic arterial PCO_2 by several fold has been widely appreciated for several decades (1-3). Both the mechanism and clinical relevance of this observation have been a source of debate for many years, however. The major areas of disagreement have been reviewed in detail recently (4, 6, 7). The classical studies of Pitts and Lotspeich (2), and Ochwaldt and Pitts (12) supported a proton secretion mechanism which, in urine containing bicarbonate, would generate H_2CO_3 that would dehydrate forming CO_2 at the uncatalyzed (slow) rate allowing trapping of CO_2 in the collecting system where surface-volume relationships would be unfavorable for diffusion. Thus, these investigators suggested an important role for delayed dehydration of H_2CO_3 , as well as the postpapillary nature of this process in the generation of a high U-B PCO_2 difference (12).

Before the present investigation, no studies have directly determined the disequilibrium pH or PCO_2 in the papillary collecting duct. One micropuncture study by Uhlich, Baldamus, and Ullrich (15) used indirect techniques to estimate PCO_2 . These workers calculated PCO_2 from pH and bicarbonate determinations and demonstrated that collecting duct PCO_2 exceeded vasa recta blood PCO_2 by 40 mm during bicarbonate loading. Furthermore, this difference between tubule lumen and vasa recta was increased by carbonic anhydrase inhibition and obliterated by infusion of excess carbonic anhydrase. Thus, they concluded that the trapping of CO_2 by a medullary countercurrent system could not account for the high PCO_2 of alkaline urine (15).

In contrast, the findings in the present study demonstrate that the PCO_2 in the papillary collecting duct during bicarbonate loading was markedly elevated with respect to systemic arterial blood but was not significantly different than the PCO_2 in the adjacent vasa recta. These findings strongly support capture of CO_2 in the medullary counter-current system. Furthermore, a significant rise in PCO_2 was noted along the length of the papillary collecting duct available for micropuncture. Moreover, a significant acid disequilibrium pH was demonstrated at both the base and tip of the collecting duct in the same condition (-0.42 and -0.36) (Table V). When carbonic anhydrase was administered in amounts which achieved assayable activity in final urine, the disequilibrium pH was completely obliterated while PCO_2 fell dramatically indicating clearly that delayed dehydration of H_2CO_3 was an important factor in the generation of the elevated CO_2 tension in the collecting tubule. The fact that the

TABLE V
Summary of Micropuncture Findings

	BCD		TCD	
	P_{CO_2}	pH_{4q}	P_{CO_2}	pH_{4q}
Controls	35	+0.07	36	+0.02
$NaHCO_3$	95	-0.42	122	-0.36
$NaHCO_3$ + Carbonic anhydrase	68	+0.06	78	+0.03

PCO_2 of urine from the right untouched kidney remained at a level significantly greater than systemic arterial blood (59.5 ± 2.3 vs. 41.0 ± 1.9) after carbonic anhydrase is in agreement with previous studies (6, 29, 30). Very early studies demonstrating a complete return of urinary PCO_2 to systemic arterial levels after carbonic anhydrase infusion were based on calculated, not determined CO_2 tensions (12).

The CO_2 tension measured by the microelectrode at the tip of the collecting duct in the left papilla compared favorably, before and after carbonic anhydrase with that observed in the right final urine (standard macro technique). The observation of a slightly, but significantly higher CO_2 tension in the experimental kidney after carbonic anhydrase deserves further comment. The magnitude of the decrease in CO_2 after carbonic anhydrase depends on the magnitude of the prevailing CO_2 in the medullary interstitium, not systemic blood. When carbonic anhydrase is added to an open beaker of a $NaHCO_3$ solution in vitro, the PCO_2 falls as a result of CO_2 formation throughout the solution and acceleration of CO_2 loss at the liquid-gas interface (5, 6). The papillary interstitium may not be an entirely open system and the dissipation of CO_2 could proceed at a slower rate. Furthermore, the demonstration of similar values for PCO_2 in both collecting duct and vasa recta all along the length of exposed papilla, suggests trapping of CO_2 by a medullary countercurrent system. Since a disequilibrium pH of -0.36 U was observed in the tip of the papillary collecting duct it would be predicted that as chemical equilibrium is achieved, or as the acidic portion of nonbicarbonate buffers react with HCO_3^- at the higher pH (7), CO_2 would be generated and final urine should have a CO_2 tension ~ 25 mm Hg greater than the PCO_2 at the tip of the collecting duct. Comparison of these values in the present study (Table III) indicates that the observed increase in PCO_2 from left papillary tip (122 ± 4.3 mm Hg) to right bladder urine (136 ± 11.8) was not significant. The explanation for this observation is not entirely clear. It may not be appropriate to compare left and right renal findings in the papillary preparation since obvious differences in blood flow, concentrating ability, and electrolyte handing have

been well described (20). Furthermore, it is generally appreciated that urine issuing out of the papillary collecting duct courses back over the papilla, which may allow equilibration of CO_2 between urine and papillary interstitium. Finally, CO_2 could diffuse out of the collecting system (albeit slight) or lost during handling of the urine sample.

In addition to hydrogen ion secretion, other possible mechanisms could result in a relative excess of H_2CO_3 . These possibilities as recently outlined by Stinebaugh et al. (7), and Warnock and Rector (31) include bicarbonate secretion into buffer containing fluid (32), mixture of acid and alkaline tubule fluid from different nephron populations (13), water abstraction from tubule fluid resulting in concentration of H_2CO_3 and HCO_3^- in conjunction with CO_2 loss (24), and finally the dissociation of HCO_3^- to CO_3^- and H_2CO_3 ("ampholyte" effect) (5, 6). Bicarbonate secretion has been observed in various types of acidifying epithelia including the rabbit cortical collecting tubule (32). The contribution of secretion to the final concentration of bicarbonate in urine has not been established, however. Previous micropuncture studies have demonstrated that the concentration of bicarbonate in the superficial distal tubule is quite high during acute metabolic alkalosis (18, 33). The bicarbonate concentration in the distal tubule during bicarbonate loading can be calculated from the equilibrium pH (7.47) and PCO_2 (65 mm Hg) to be 43.9 mM (18). These same studies failed to demonstrate a significant acid disequilibrium pH in the distal nephron during bicarbonate loading (18). Furthermore the anticipated increase in bicarbonate concentration from distal tubule to papillary collecting duct is modest, when water abstraction is considered, and casts doubt on a high capacity secretory process in the collecting tubule. Recently, Lombard and associates (34) have demonstrated that the net secretion of bicarbonate in the cortical collecting tubule of the rabbit is quite low. These same investigators demonstrated bicarbonate reabsorption, not secretion, in the medullary collecting tubule (34). Moreover, the capacity for bicarbonate reabsorption in the medullary collecting tubule exceeded that in the cortical collecting tubule by several fold (34). Compatible with this latter observation are recent micropuncture studies that have demonstrated bicarbonate reabsorption in the papillary collecting duct of the surgically exposed rat papilla (35). It seems highly unlikely, therefore, that bicarbonate secretion could be playing a significant role in the generation of an acid disequilibrium pH in the papillary collecting duct.

Although the admixture of urine of varying bicarbonate and nonbicarbonate buffer concentration could serve to elevate the urinary PCO_2 in certain conditions

(13), the demonstration by Rector and associates of high urinary CO_2 tensions during a combined water and bicarbonate diuresis in phosphate-depleted subjects (36) casts considerable doubt on this possibility as a sole determinant of the U-B PCO_2 . In the present study the excretion of nonbicarbonate buffers would be expected to be low (7) and "mixing" would be minimal from the base to tip of papilla since the majority of branch points occur proximal to this site. Furthermore, a significant acid disequilibrium pH was observed along the length of the papillary collecting duct examined, i.e., did not dissipate. Nonbicarbonate buffers may serve to further increase CO_2 tension in conditions in which the concentration of H_2CO_3 is slightly displaced from equilibrium, as in the present study, since, as pH rises, the acidic portion of the buffer (HA) would react with HCO_3^- to produce H_2CO_3 and thus CO_2 (13, 37). For example, at an equilibrium pH of 7.71 (papillary tip) 2 mM of buffer would produce ~ 0.22 mM of CO_2 and 5 mM of buffer would produce 0.55 mM of CO_2 . Since nonbicarbonate buffer content was not measured in the present study, a correlation between the magnitude of the acid disequilibrium pH, the PCO_2 , and the amount of nonbicarbonate buffer present in the papillary collecting duct is not possible. The extent to which urinary PCO_2 is dependent on the combined effects of proton secretion and nonbicarbonate buffer content cannot be stated with certainty, therefore. A role for nonbicarbonate buffer as an additional determinant of the high CO_2 tension observed in the present study seems likely, however.

Another possible cause for the disequilibrium pH observed in the papillary collecting duct in this study would be CO_2 loss in a segment capable of significant concentration of bicarbonate and carbonic acid as originally proposed by Reid and Hills (24). Loss of CO_2 from collecting duct lumen to vasa recta is suggested by the demonstration of similar values for PCO_2 in these structures (Fig. 4). That there was not adequate concentration in this segment to generate a disequilibrium pH of the magnitude reported in this study is evidenced by the findings in Tables IV and V. The bicarbonate concentration increased from base to tip (104–140 meq/liter) (calculated from the measured pH_{eq} and PCO_2). The tubular fluid to plasma inulin ratio increased slightly (15–22.5) from base to tip in animals similarly prepared for papillary micropuncture in this laboratory. Therefore, $\sim 2\%$ of the filtered load of bicarbonate was reabsorbed between base and tip. To generate a disequilibrium pH of -0.3 U (less than that observed at the base), the concentration of H_2CO_3 would be required to double. For water abstraction to account alone for such an increase in H_2CO_3 concentration, a similar increase in bicarbonate

concentration would be expected (i.e., 104–208 meq/liter). To explain these findings on this basis alone seems untenable. Therefore, proton secretion seems the most likely determinant of the disequilibrium pH and elevated PCO_2 observed in the papillary collecting duct. Finally, in previous studies from this laboratory a disequilibrium pH was not observed in the superficial distal tubule (18). From the calculated concentration of bicarbonate (43.9 mM) in this segment in identically prepared rats, the transit time from distal tubule to base collecting duct of ~ 2 min, and the failure for water abstraction alone to account for the observation of a disequilibrium pH of -0.42 U at the base, it seems unlikely that a mechanism similar to that proposed by Reid and Hills (24), or mixing (13) could account for the observed changes between the distal tubule and base collecting duct as well. Because of the inaccessibility of the nephron segments involved, neither of these mechanisms can be totally eliminated, however.

Recent emphasis has been placed on the physicochemical properties of bicarbonate and the linear relationship between the urinary bicarbonate concentration and the U-B PCO_2 difference (5). The demonstration by Stinebaugh and associates (7) that this relationship can be altered by various physiological maneuvers and pathophysiological conditions as well as the theoretical considerations and calculations of these investigators, make a primary role for the "ampholyte" effect unlikely. The demonstration in the present study of an acid disequilibrium pH in association with elevated papillary CO_2 tensions both of which were markedly reduced by carbonic anhydrase is further evidence against CO_2 generation of this magnitude from a purely physicochemical process.

The recent demonstration that a spontaneous disequilibrium pH did not exist in the superficial distal tubule during bicarbonate loading (18), was interpreted as evidence for a low capacity proton secretory system in this segment. This finding has been strengthened by the recent *in vivo* microperfusion studies of Lucci and associates (38) which demonstrated that significant bicarbonate reabsorption was not present despite the existence of sodium reabsorption and potassium secretion in the superficial distal tubule (38). In this regard, and in view of the demonstration in the present study of a significant acid disequilibrium pH at the base of the papillary collecting duct, it is interesting to speculate that the medullary collecting duct, proximal to the first accessible micropuncture site, participates importantly in the acidification process by proton secretion. It is conceivable therefore, that the disequilibrium pH observed at the base papillary collecting duct (-0.42) pH U, underestimates the maximum disequilibrium pH achievable in the terminal

nephron. Furthermore, since carbonic anhydrase accelerates the conversion of carbonic acid to CO_2 , and CO_2 loss from the collecting duct, the observation of a marked decrease in PCO_2 at the papillary base after carbonic anhydrase administration (Table II) suggests that an effect of the enzyme occurs in nephron segments proximal to the papillary collecting duct prior to accessibility to, and trapping in, the medullary countercurrent system.

The demonstration of an acid disequilibrium pH in the papillary collecting duct, only in association with elevated CO_2 tensions in equilibrium in the collecting duct and vasa recta, suggests that hydrogen ion secretion is a major determinant of the increase in the U-B PCO_2 gradient during an alkaline diuresis. An important role for a medullary countercurrent system, trapping CO_2 in the renal medulla, is also likely. Because of the technical limitations of this study, a role for mixing of dissimilar urines (13), "concentration" with CO_2 loss (24), and the effect of nonbicarbonate buffers cannot be totally eliminated as potential contributors to this process with certainty. In view of these findings, and the recent extensive clearance studies by Stinebaugh and associates (7, 10), Halperin et al. (9), and Arruda et al. (8), it is concluded that when appropriate consideration for urinary concentrating ability, nonbicarbonate buffer content, and urinary bicarbonate concentration is made, the U-B PCO_2 difference can be considered a reliable qualitative index of hydrogen ion secretion by the collecting tubule.

ACKNOWLEDGMENTS

The author wishes to express his sincere appreciation for the advice, wise council, and encouragement of Dr. Norman W. Carter, without whom these studies could not have been completed. Dr. Marjory S. Lucci performed the microcalorimetric measurements, and offered helpful suggestions in the performance of this work. Helpful comments and suggestions were obtained from discussions with Dr. Floyd C. Rector, Jr. and Dr. B. J. Stinebaugh. The manuscript was typed by Mrs. Karen Williams, and Darnell Russo.

This research was supported in part by National Institutes of Arthritis, Metabolism, and Digestive Diseases Research grant 1 RO1 AM 25730, 1 RO1 AM 14677, 1 KO4 AM 01033, and 7 RO1 AM 30603.

REFERENCES

1. Mainzer, F., and M. Bruhn. 1931. Über löslichkeit, Dissoziation und Spannung der kohlen saure im Harn. *Biochem. Ztschr.* **230**: 395-410.
2. Pitts, R. F., and W. D. Lotspeich. 1946. Bicarbonate and the renal regulation of acid-base balance. *Am. J. Physiol.* **147**: 138-154.
3. Ryberg, C. 1948. Some investigations on the carbon dioxide tension of the urine in man. *Acta Physiol. Scand.* **15**: 123-139.
4. Rector, Floyd C., Jr. 1973. Acidification of the urine. In *Handbook of Physiology*. J. Orloff and R. W. Berliner editors. American Physiological Society. The Williams and Wilkins Co., Baltimore, Md. p. 431-454.
5. Arruda, J. A. C., L. Nascimento, P. K. Mehta, D. R. Rademacher, J. T. Sehy, C. Westenfelder, and N. A. Kurtzman. 1977. The critical importance of urinary concentrating ability in the generation of urinary carbon dioxide tension. *J. Clin. Invest.* **60**: 922-935.
6. Maren, T. 1978. Carbon dioxide equilibria in the kidney: the problems of elevated carbon dioxide tension, delayed dehydration, and disequilibrium pH. *Kidney Int.* **14**: 395-405.
7. Stinebaugh, B. J., R. Esquenazi, F. X. Schloeder, W. N. Suki, M. B. Goldstein, and M. L. Halperin. 1980. Control of the urine-blood PCO_2 gradient in alkaline urine. *Kidney Int.* **17**: 31-39.
8. Arruda, J. A. C., L. Nascimento, S. K. Kumar, and N. A. Kurtzman. 1977. Factors influencing the formation of urinary carbon dioxide tension. *Kidney Int.* **11**: 307-317.
9. Halperin, M. C., M. B. Goldstein, A. Haig, M. D. Johnson, and B. J. Stinebaugh. 1974. Studies on the pathogenesis of type I (distal) renal tubular acidosis as revealed by urine PCO_2 tensions. *J. Clin. Invest.* **53**: 669-677.
10. Gougoux, A., P. Vinay, G. Lemieux, R. Richardson, S. Cheung Tam, M. B. Goldstein, B. Stinebaugh, and M. C. Halperin. 1980. Effect of blood pH on distal nephron hydrogen ion secretion. *Kidney Int.* **17**: 615-621.
11. Julka, N. K., J. A. Arruda, and N. A. Kurtzman. 1979. The mechanism of amphotericin-induced distal acidification defect in rats. *Clin. Sci.* **56**: 555-562.
12. Ochwadt, B. K., and R. F. Pitts. 1956. Effect of intravenous infusion of carbonic anhydrase on carbon dioxide tension of alkaline urine. *Am. J. Physiol.* **185**: 426-429.
13. Kennedy, T. J., Jr., J. Orloff, and R. W. Berliner. 1952. Significance of carbon dioxide tension in urine. *Am. J. Physiol.* **169**: 596-608.
14. Pak Poy, R. K., and O. Wrong. 1960. The urinary PCO_2 in renal disease. *Clin. Sci.* **19**: 631-639.
15. Uhlich, E., C. A. Baldamus, and K. J. Ullrich. 1968. Verhalten von CO_2 -Druck und Bicarbonat im Gegenstromsystem des Nierenmarks. *Pflugers Arch. Eur. J. Physiol.* **303**: 31-48.
16. DuBose, T. D., Jr., L. R. Pucacco, D. W. Seldin, N. W. Carter, and J. P. Kokko. 1978. Direct determination of PCO_2 in the rat renal cortex. *J. Clin. Invest.* **62**: 338-348.
17. Lucci, M. S., L. R. Pucacco, N. W. Carter, and T. D. DuBose, Jr. 1979. Direct determination of CO_2 permeability of the rat proximal tubule in vivo. *Clin. Res.* **27**: 765. (Abst.)
18. DuBose, T. D., Jr., L. R. Pucacco, and N. W. Carter. 1981. Determination of disequilibrium pH in the rat kidney *in vivo*: evidence for hydrogen secretion. *Am. J. Physiol.* **240 (Renal, Fluid, Electrolyte Physiol.** **9**): F138-F146.
19. Rector, F. C., Jr., N. W. Carter, and D. W. Seldin. 1965. The mechanism of bicarbonate reabsorption in the proximal and distal tubules of the kidney. *J. Clin. Invest.* **44**: 278-290.
20. Higashihara, E., T. D. DuBose, Jr., and J. P. Kokko. 1978. Direct examination of chloride transport across papillary collecting duct of the rat. *Am. J. Physiol.* **235 (Renal, Fluid, Electrolyte Physiol.** **4**): F219-F226.
21. DuBose, T. D., Jr., L. R. Pucacco, M. Lucci, and N. W. Carter. 1979. Micropuncture determination of pH, PCO_2 , and total CO_2 concentration in accessible structures of the rat renal cortex. *J. Clin. Invest.* **64**: 476-482.

22. Maren, Thomas, H. 1960. A simplified micromethod for the determination of carbonic anhydrase and its inhibitors. *J. Pharmacol. Exp. Ther.* **130**: 26-29.
23. Hastings, A. B., and J. Sendroy. 1925. The effect of variation in ionic strength on the apparent first and second dissociation constants of carbonic acid. *J. Biol. Chem.* **65**: 445-455.
24. Reid, E. L., and A. G. Hills. 1965. Diffusion of carbon dioxide out of the distal nephron in man during anti-diuresis. *Clin. Sci.* **28**: 15-28, 1965.
25. Berliner, R. W. 1957. Some aspects of ion exchange in electrolyte transport by the renal tubules. In *Metabolic Aspects of Transport across Cell Membranes*. Q. R. Murphy, editor. University of Wisconsin Press, Madison 203-220.
26. Gruber, M. C., C. R. Caflisch, H. H. Bengele, E. A. Alexander, 1980. Elevated urinary PCO_2 —An intrarenal event. Abstracts of the Am. Soc. Nephrol. 13th Annual Meeting. 135A. (Abstr.)
27. Gruber, M. C., C. R. Caflisch, H. H. Bengele, J. H. Schwartz, and E. A. Alexander. 1980. pH prolife along the inner medullary collecting duct of the rat. *Clin. Res.* **28**: 446. (Abstr.)
28. Garg, L. C., and T. H. Maren. 1972. The rate of hydration of carbon dioxide and dehydration of carbonic acid at 37°. *Biochim. Biophys. Acta.* **261**: 70-76.
29. Guignard, J. P. 1966. Mecanisme de la reabsorption renale des bicarbonates. *Helv. Physiol. Acta.* **24**: 193-226.
30. Stinebaugh, B. J., E. Ghafary, M. B. Goldstein, M. C. Halperin, F. X. Schloeder, and W. N. Suki. 1978. Dis-equilibrium pH and bicarbonate reabsorption. *Nephron.* **20**: 141-146.
31. Warnock, D. G., and F. C. Rector, Jr. 1981. Renal acidification mechanisms. In *The Kidney*. B. Brenner, and F. C. Rector, Jr., editors. W. B. Saunders Company, Philadelphia. 440-494.
32. McKinney, T. D., and M. B. Burg. 1978. Bicarbonate secretion by rabbit cortical collecting tubules in vitro. *J. Clin. Invest.* **61**: 1421-1427.
33. Malnic, G. M., M. de Mello Aires, and G. Giebisch. 1972. Micropuncture study of renal tubular hydrogen ion transport in the rat. *Am. J. Physiol.* **222**: 147-158.
34. Lombard, W. E., H. R. Jacobson, and J. P. Kokko, 1979. Collecting duct bicarbonate transport: comparison of cortical and medullary segments. *Kidney Int.* **16**: 827A. (Abstr.)
35. Richardson, R. M. A., and R. T. Kunau. 1980. Bicarbonate reabsorption by the rat papillary collecting duct: effect of acetazolamide. Abstracts of the American Society of Nephrology 13th Annual Meeting. 147A. (Abstr.)
36. Rector, F. C., Jr., R. M. Portwood, and D. W. Seldin. 1959. Examination of the mixing hypothesis as an explanation for elevated urinary CO_2 tensions. *Am. J. Physiol.* **197**: 861-864.
37. Kennedy, T. J., M. Eden, and R. W. Berliner. 1957. Interpretation of urine CO_2 tension. *Fed. Proc.* **16**: 72.
38. Lucci, M. S., L. R. Puccio, N. W. Carter, and T. D. DuBose, Jr. 1980. An *in vivo* microperfusion evaluation of bicarbonate transport in the rat distal convoluted tubule. *Clin. Res.* **28**: 454. (Abstr.)