

Restoration of immune responses of aging hamsters by treatment with isoprinosine.

K Y Tsang, H H Fudenberg, M J Gnagy

J Clin Invest. 1983;71(6):1750-1755. <https://doi.org/10.1172/JCI110930>.

Research Article

Immune competence declines with advanced age in hamsters, as in other laboratory mammals and in humans. We found significant alterations in the functional parameters of different populations of immunocytes (natural killer cells, T cells, monocytes, and suppressor cells) in aging hamsters, beginning at approximately 14 mo of age. Natural killer cytotoxicity, phytohemagglutinin-induced lymphocyte stimulation, and monocyte chemotaxis were decreased in aging Lak:LvG(Syr) outbred hamsters. When old hamsters were given a single injection (5 mg/kg body wt) of isoprinosine, a chemical immune potentiator, these three immune parameters increased almost to the levels found in young adult hamsters but returned to pretreatment levels after 7 d. Suppressor cell activity for the lymphocyte response to phytohemagglutinin, which increased with age, was decreased after treatment. In old hamsters treated with weekly injections of isoprinosine, these four immunological parameters remained at or near the levels found in young adults.

Find the latest version:

<https://jci.me/110930/pdf>



Restoration of Immune Responses of Aging Hamsters by Treatment with Isoprinosine

KWONG Y. TSANG, H. HUGH FUDENBERG, and MARK J. GNACY, *Department of Basic and Clinical Immunology and Microbiology, Medical University of South Carolina, Charleston, South Carolina 29425*

ABSTRACT Immune competence declines with advanced age in hamsters, as in other laboratory mammals and in humans. We found significant alterations in the functional parameters of different populations of immunocytes (natural killer cells, T cells, monocytes, and suppressor cells) in aging hamsters, beginning at ~14 mo of age. Natural killer cytotoxicity, phytohemagglutinin-induced lymphocyte stimulation, and monocyte chemotaxis were decreased in aging Lak:LvG(Syr) outbred hamsters. When old hamsters were given a single injection (5 mg/kg body wt) of isoprinosine, a chemical immune potentiator, these three immune parameters increased almost to the levels found in young adult hamsters but returned to pre-treatment levels after 7 d. Suppressor cell activity for the lymphocyte response to phytohemagglutinin, which increased with age, was decreased after treatment. In old hamsters treated with weekly injections of isoprinosine, these four immunological parameters remained at or near the levels found in young adults.

INTRODUCTION

Various immunological functions and responses are affected by the aging process. In humans, responsiveness to thymus-dependent antigens originally contacted in early life decreases with advancing age (1, 2). T cell functions, such as primary delayed skin reaction in response to antigens to which individuals have not been sensitized previously (e.g., dinitrochlorobenzene) and the proliferative capacity of T cells in response to phytohemagglutinin (PHA),¹ also decline with age (3-8). The data of Girard et al. (9)

regarding cell-mediated immunity in 880 hospitalized patients indicated a decrease in the absolute number of circulating T lymphocytes, and also functional impairment of T cells.

In mice, T cell-dependent functions, such as the lymphocyte response to PHA in vitro, decline with age (10-14); resistance to challenge with syngeneic and allogeneic tumor cells in vivo is also dramatically decreased in old animals (15, 16). Decreased absolute numbers of lymphocytes in peripheral blood in aging hamsters have also been reported (17). We have previously found that immune responses such as monocyte chemotaxis, lymphocyte response to PHA stimulation, and natural killer (NK) activity are decreased, and that suppressor cell activity is increased in aging hamsters (18).

Only a few chemical agents possess immunorestorative activity. These include double-stranded polynucleotides (19, 20), thymic hormones (21, 22), certain free radical inhibitors such as vitamin E and coenzyme Q₁₀ (23, 24), levamisole (25, 26), and mercaptoethanol (27, 28). Recently, isoprinosine (ISO), the *p*-acetamido-benzoic acid salt of *N,N*-dimethylamino-2-propanol:inosine complex (3:1 molar ratio) has been shown to be an effective immune potentiator (29, 30). Wybran (31) has demonstrated that ISO can enhance the PHA response of lymphocytes from cancer patients, and Lesourd et al. (32) have reported that it can restore cell-mediated immunity in patients with acute encephalitis. In an animal model of human osteosarcoma, we have shown that ISO can augment proliferation of peripheral blood lymphocytes from tumor-bearing animals in vitro and can increase the functional capacity of their immune system (33). The present investigation was designed to examine the effects of ISO on the immune responses of aging hamsters.

METHODS

Animals. Outbred hamsters, strain Lak:LvG (Syr), were used in this investigation. They were divided into eight

Received for publication 15 June 1982 and in revised form 4 January 1983.

¹ Abbreviations used in this paper: FCI, functional chemotactic index; ISO, isoprinosine; LS, lymphocyte stimulation; MC, monocyte chemotaxis; NK, natural killer; PHA, phytohemagglutinin; SC, suppressor cell; TFS, thymosin.

groups according to age (0.5, 2, 4, 8, 12, 14, 18, and 24 mo). ISO, supplied by Newport Pharmaceuticals International (Newport Beach, CA), was freshly dissolved in saline and injected intraperitoneally at a concentration of 5 mg/kg body wt, the optimal concentration indicated in our previous study (33). Hamsters were bled by cardiac puncture and lymphocytes were isolated on a Ficoll-Hyque gradient as described by Boyum (34).

Lymphocyte stimulation (LS). A modification of the method described by Ron et al. (35) was used for assay of PHA-induced LS. 1×10^5 cells in 0.1 ml RPMI 1640 medium with 10% fetal calf serum were inoculated in the wells of a 96-well flat-bottomed microtest plate and PHA (Wellcome Reagents) was added at 10 μ g/well. Each assay was performed in triplicate. The lymphocyte cultures were incubated for 48 h at 37°C in a 5% CO₂ incubator, 1 μ Ci of [³H]thymidine (Amersham Corp., Arlington Heights, IL, 5 Ci/mmol) was added to each well, and incubation was continued for 24 h. The cultures were harvested in a Mash I cell harvester (Microbiological Associates, Walkersville, MD).

NK cell activity. For measurement of NK activity, Raji cells were used as target cells in an assay as described previously for hamster NK activity (36). Effector cells were prepared from peripheral blood lymphocytes and Ficoll-Hyque-isolated peripheral blood (37). Mononuclear cells were depleted of adherent cells by incubation of the mononuclear cells for 1 h at 37°C on 60-mm plastic culture dishes. The purified mononuclear cells contained <0.5% monocytes based on morphology and nonspecific esterase staining. 100 μ l of nonadherent cells were added with 100 μ l of ⁵¹Cr-labeled target cells (1×10^4) in the wells of a flat-bottomed microtest plate at a ratio of 50:1 or 25:1. Target cells incubated without effector cells were used to determine the level of spontaneous lysis. Maximum lysis was determined by adding 10% Triton X-100 solution (no effector cells) instead of medium to the well. The percentage cytotoxicity was calculated as: Percent cytotoxicity = mean cpm test - mean cpm spontaneous lysis/mean cpm maximum lysis - mean cpm spontaneous lysis \times 100. Each assay was performed in triplicate.

Chemotaxis. The method of Tsang et al. (38) was used to measure monocyte chemotaxis (MC). Chemotactic factor was generated in fresh hamster serum by incubation with 1 mg *Salmonella typhosa* 0901 endotoxin (Difco Laboratories, Inc., Detroit, MI) per ml of serum at 37°C for 1 h followed by 56°C for 30 min. In Boyden chemotaxis chambers, monocytes were separated from the stimulus with a 5- μ m pore size polycarbonate filter. The chambers were incubated at 37°C in a 5% CO₂ incubator for 3 h. The functional chemotactic index (FCI) was calculated by the following formula: FCI = number of migrating cells in experimental chamber/number of migrating cells in control chamber \times 100. Each assay was performed in triplicate.

For the study of the effects of ISO on macrophage chemotaxis in vitro, the following procedures were used. Only adherent cells were used in this assay. Macrophages were incubated with ISO at an optimal concentration (100 μ g/10⁶ cells per ml) for 2 h at room temperature. At the end of the incubation time, ISO-treated macrophages were washed and used in the assay. Macrophages incubated in medium alone were used as controls.

Suppressor cell activity. The method described by Padarathsingh et al. (39) was used to assay suppressor cell activity (SC). Mononuclear cells were isolated on a Ficoll-Hyque gradient. PHA was used as mitogen at a concentration of 10 μ g/well with mononuclear cells cultured at a concentration of 2×10^5 cells/well in microtest plates. For co-cul-

tures 1×10^5 cells collected from old hamsters (18 mo of age) before and after ISO treatment were mixed with 1×10^5 cells from young hamsters (3-4 mo of age). The cultures were incubated at 37°C in a 5% CO₂ incubator for 3 d, and 1 μ Ci of [³H]thymidine was added to each well 24 h before the cultures were harvested. Mononuclear cell cultures from young hamsters or old hamsters alone were used as controls. Percentage suppression was calculated by the following formula: Percent suppression = $[1 - (cpm Y + O)/(cpm Y)] \times 100$. Each experiment was performed in triplicate.

RESULTS

The effects of age on LS, NK, and monocyte chemotaxis (MC) activities are shown in Table I. Each age group contained 15 hamsters (five groups were used for each determination, three hamsters in each group). The highest responses in the three assays were observed in the 8-mo-old group. The 18- and 24-mo-old groups had the lowest responses. The LS response to PHA was significantly decreased in 12-, 14-, 18- and 24-mo-old hamsters as compared with the younger age groups ($P < 0.005$ by Student's *t* test). NK cytotoxicity was significantly decreased in the 14-, 18-, and 24-mo-old groups as compared to 4-, 8-, and 12-mo-old groups ($P < 0.005$). Monocyte chemotaxis was also depressed significantly in the 14-, 18-, and 24-mo-old groups as compared to the younger age groups ($P < 0.01$).

The effects of ISO on macrophage chemotaxis in hamsters of various ages are shown in Table II. Significant increases in macrophage chemotaxis activities were observed in hamsters of 12, 14, and 18 mo of age when treated with ISO in vitro. No significant increases in macrophage chemotaxis activities were seen in ham-

TABLE I
Immune Responses in Hamsters of Various Age Groups

Age mo	PHA-induced lymphocyte proliferation	NK cell activity	MC
	mean cpm/ $10^6 \pm SE$	mean % lysis/ $\pm SE$	FCI
0.5	8,178 \pm 114.6	4.5 \pm 1.8 \ddagger	85.0
2.0	7,972 \pm 156.5	1.2 \pm 4.8 \ddagger	91.8
4.0	8,467 \pm 196.5	47.3 \pm 7.2	86.3
8.0	9,174 \pm 198.7	54.2 \pm 6.5	100.0
12.0	4,354 \pm 106.4 \circ	52.6 \pm 5.9	65.5 \ddollar
14.0	4,106 \pm 117.4 \circ	21.4 \pm 3.8 \ddagger	43.3 \ddollar
18.0	2,316 \pm 98.6 \circ	14.1 \pm 3.1 \ddagger	37.6 \ddollar
24.0	2,410 \pm 79.8 \circ	12.9 \pm 4.8 \ddagger	36.8 \ddollar

Five groups of hamsters were used for each determination, each group consisting of three hamsters. Student's *t* test was used for all statistical analysis.

\circ Statistical significance, $P < 0.005$.

\ddagger $P < 0.01$.

\ddollar $P < 0.01$.

TABLE II
Effects of ISO on MC in Hamsters In Vitro*

Age	Without ISO	With ISO	P
<i>mo</i>			
0.5	82.6±2.51	85.3±4.34	NS
2.0	88.3±2.08	91.3±3.21	NS
4.0	86.3±4.04	91.0±1.00	NS
8.0	100.0±0.57	97.0±3.00	NS
12.0	61.0±2.64	80.6±2.08	P < 0.005
14.0	44.6±4.50	66.0±8.18	P < 0.01
18.0	34.3±4.04	56.3±5.50	P < 0.01

* Results are expressed as FCI. Three groups of hamsters were used for each determination, each group consisting of five hamsters. Student's *t* test was used for all statistical analyses.

sters of 0.5, 2, 4, and 8 mo of age after ISO treatment in vitro.

The effects of ISO injections on the LS responses to PHA are shown in Fig. 1a. Three age groups were used




FIGURE 1 Effects of single injection of ISO on (a) PHA-induced lymphocyte proliferation, (b) NK activity, and (c) MC in aging hamsters: (○) 14 mo, (▲) 18 mo, (×) 24 mo, (○) control (14 mo old).

(14, 18, and 24 mo old) with 15 hamsters in each group. Increases in LS were seen on days 1, 3, and 5 postinjection, returning to control levels on day 7. Fig. 1b shows the effects of ISO on NK cytotoxicity in three groups of hamsters (14, 18, and 24 mo old), each containing 15 hamsters, and Fig. 1c shows the results of MC assays in three groups of 9 hamsters each. Enhancement of both NK and MC activity was seen on days 1, 3, and 5 after ISO injection, and both returned to control levels on day 7 in all three age groups.

The number of macrophages in the mononuclear cell preparations of young hamsters (3-4 mo old) and old hamsters (18 mo old) was determined by the nonspecific esterase staining method. There were no significant differences in the percentage of macrophages between the young (5-7.5%) and old hamsters (6-8.4%). The effects of ISO on SC activity in aging hamsters are shown in Table III. 10 18-mo-old hamsters were used in this study. Lymphocytes from young hamsters were

TABLE III
Effects of ISO on SC Activity in Aging Hamsters

Culture*	Before ISO treatment		After ISO treatment			
	Y	O	cpm†	% suppression§	cpm†	% suppression§
1			6,349±593	—	7,996±449	—
1 + 1			1,097±69	82.8	7,814±560	3.0
1 + 2			1,541±87	75.8	8,062±671	0
1 + 3			1,112±98	82.5	7,094±698	1.2
1 + 4			987±56	84.5	7,126±709	11.0
1 + 5			1,610±107	74.7	7,012±604	12.4
1 + 6			1,812±124	71.5	6,984±587	12.7
1 + 7			1,987±94	69.5	5,837±439	27.1
1 + 8			1,406±119	77.9	6,019±672	24.8
1 + 9			2,013±143	68.3	7,504±704	6.2
1 + 10			2,436±109	61.7	8,126±568	0
2			9,871±746	—	8,964±765	—
2 + 1			1,916±114	80.6	8,166±643	9.0
2 + 2			1,314±134	86.7	8,431±747	6.0
2 + 3			1,659±156	83.2	7,561±564	16.0
2 + 4			1,927±148	80.5	7,844±766	13.0
2 + 5			1,762±114	82.2	7,124±567	20.6
2 + 6			1,845±125	81.6	7,344±671	18.1
2 + 7			1,978±116	80.0	8,126±562	9.4
2 + 8			2,042±187	79.4	6,021±413	32.9
2 + 9			2,137±104	78.4	6,124±409	31.7
2 + 10			1,534±109	84.5	6,849±514	23.6

* Lymphocytes from 18-mo-old hamsters (O₁-O₁₀) were co-cultured with lymphocytes from the pooled blood of two different groups of three 4-mo-old hamsters (Y₁ and Y₂).

† Mean±SE for three determinations.

§ % suppression = [1 - (cpm Y + O)/(cpm Y)] × 100.

Mean percent suppression to Y₁ were 74.9 (before ISO treatment) and 9.84 (after ISO treatment). Mean percent suppression to Y₂ were 81.71 (before ISO treatment) and 17.2 (after ISO treatment).

obtained from the pooled blood of two different groups of three 4-mo-old hamsters. Assays were performed before and on the 4th d after intraperitoneal injection of ISO in the old hamsters. SC activity for PHA-induced lymphocyte proliferation was significantly reduced in all the old hamsters treated with ISO ($P < 0.005$).

The effects of continuous treatment with ISO on the LS, NK, and MC activities of aging hamsters are shown in Fig. 2, and SC activity is shown in Fig. 3. ISO was injected intraperitoneally once per week, beginning at the 14th mo for the LS, NK, and MC studies and at the 18th mo for the SC study, and blood was drawn for assays once per month on the day before the next injection of ISO. All four immune responses, which were significantly decreased in the untreated group, were normal or near normal (as compared to young adult hamsters) in the ISO-treated group of aging hamsters.




FIGURE 2 Effects of weekly injections of ISO on (a) PHA-induced lymphocyte proliferation, (b) NK activity, and (c) MC in aging hamsters. Immune assays were performed before and after the beginning of ISO injections. (●) ISO-treated hamsters, 14 mo old at the beginning of treatment; (▲) age-matched controls.




FIGURE 3 Effects of weekly injections of ISO on suppressor cell activity in aging hamsters. SC assays were performed before and after the beginning of ISO injections. (●) ISO-treated hamsters, 18 mo old at the beginning of treatment; (○) age-matched controls.

DISCUSSION

The results of our investigation in hamsters confirm many findings that cell-mediated immune activity declines with age in mammals (3, 4, 5, 18). LS, NK, and MC activities were significantly decreased and SC activity was significantly elevated in hamsters in the age group of 12–14 mo or older. The present investigation also suggests that ISO can be used to restore or maintain normal immune function in aging animals.

Prostaglandins of the E series have been shown to have inhibitory effects on NK cell activity (40, 41). To eliminate the possibility of inhibition of NK activity by prostaglandins, we used nonadherent cells as effector cells in our assay. Our effector cell preparation contained <0.5% monocytes.

In vitro effects of ISO on macrophage chemotaxis were studied in hamsters of various ages. Our data show that ISO increased MC activity in hamsters 12, 14, and 18 mo old. These hamsters had low MC activities without ISO treatment. However, treatment of macrophages with ISO in vitro did not increase the MC activity in hamsters 0.5, 2, 4, and 8 mo old that had normal or close to normal activity. These results were similar to the results of our study in humans (37).

There were no significant differences in the percent of macrophage contents between the young and old hamsters of the strain we studied. For this reason, we believe that the suppression in PHA-stimulated lymphocyte proliferation in old hamsters was not caused by the difference in the number of macrophages in the cell preparation.

The decline in immune activity in aging animals could result from changes in the immune cells, changes in their milieu, or both. Studies by Makinodan and co-workers (42-46) have suggested that although both types of changes affect the immune response in mice, much of the normal age-related decline in immunity can be attributed to changes in the immune cells. These changes may be in the number of T lymphocytes (49), the capacity of lymphocytes to function (48, 49), and/or alterations in subpopulations of lymphocytes (50).

We attempted to restore the declining immune functions of aging hamsters by administering ISO. We found that a single injection of ISO elevated their LS, NK, and MC activities, with the peak activity on day 5 postinjection as found previously (33), and that continuous injections of ISO restored the LS, NK, MC, and SC activities of aging hamsters to the levels of young adult hamsters. Several studies of other immune potentiators, such as thymic hormones and levamisole, in aging have also been reported. For example, Cowan et al. (51) found that mixed lymphocyte reactivity was significantly increased when thymosin (TF5) was incubated with lymphocytes from elderly individuals. Levamisole, which does not increase immune functions *in vitro*, is effective only when administered *in vivo* over an extended period (52). This would suggest that levamisole itself may not exert its enhancing effect directly on the immune cells, but rather through either another cell type or a metabolite. In contrast, ISO can restore immune function both *in vivo* and *in vitro*. The mechanism of action of ISO in restoring the immune functions of aging hamsters is not known, but it may be mediated partly by factors such as interleukin-2 (53) or interleukin-1 (54) and our data suggest that it acts on both T cells and monocytes. Further investigations have been undertaken to study the effects of ISO on life span, susceptibility to disease, and tumor development in aging hamsters. The information obtained from these studies may be valuable in developing new methods for delaying the onset, decreasing the rate, or reversing the effects of declining immune function in aging animals.

ACKNOWLEDGMENT

We thank Charles L. Smith for editorial assistance.

Research was supported in part by U. S. Public Health Service grant CA-25746.

REFERENCES

1. Walford, R. L. 1968. *Immunologic Theories of Aging*. Munksgaard, Copenhagen.
2. Nossal, G. J. V. 1962. Cellular genetics of immune responses. *Adv. Immunol.* **2**: 163-204.
3. Baer, H., and R. T. Bowser. 1963. Antibody production and development of contact skin sensitivity in guinea pigs of various ages. *Science (Wash. DC)*. **140**: 1211-1212.
4. Gross, L. 1965. Immunological defect in aged population and its relationship to cancer. *Cancer (Phila.)*. **18**: 201-204.
5. Grossman, J., J. Baum, J. Fusner, and J. Condemi. 1975. The effect of aging and acute illness on delayed hypersensitivity. *J. Allergy Clin. Immunol.* **55**: 268-275.
6. Pisciotta, A. V., D. W. Westing, C. Deprey, and B. Walsh. 1967. Mitogenic effect of phytohemagglutinin at different ages. *Nature (Lond.)*. **215**: 193-194.
7. Heine, K. M. 1971. Die reaktionsfähigkeit der lymphozyten im alter. *Folia Haematol. (Leipz.)*. **96**: 29-33.
8. Hallgren, H. M., E. C. Buckley III, V. A. Gilbertsen, and E. J. Yunis. 1973. Lymphocyte phytohemagglutinin responsiveness, immunoglobulins and autoantibodies in aging humans. *J. Immunol.* **111**: 1101-1107.
9. Girard, J. P., M. Paychere, M. Cuevas, and B. Fernandes. 1977. Cell-mediated immunity in an aging population. *Clin. Exp. Immunol.* **27**: 85-91.
10. Meron, M., B. N. Jaroslav, and R. Koesterer. 1974. The decline of cell-mediated immunity in aging mice. *J. Gerontol.* **29**: 499-505.
11. Nielson, H. E. 1974. The effect of age on the response of rat lymphocytes in mixed leukocyte culture, to PHA, and in the graft-vs.-host reaction. *J. Immunol.* **112**: 1194-1200.
12. Goodman, S. A., and T. Makinodan. 1975. Effect of age on cell-mediated immunity in long-lived mice. *Clin. Exp. Immunol.* **19**: 533-542.
13. Hori, Y., E. H. Perkins, and M. K. Halsall. 1973. Decline in phytohemagglutinin responsiveness of spleen cells from aging mice. *Proc. Soc. Exp. Biol. Med.* **144**: 48-55.
14. Becker, M. J., R. Farkas, M. Schneider, I. Druck, and A. Klagman. 1979. Cell-mediated cytotoxicity in humans: age-related declined as measured by a xenogeneic assay. *Clin. Immunol. Immunopathol.* **14**: 204-210.
15. Perkins, E. H., and L. H. Cacheiro. 1977. A multiple-parameter comparison of immunocompetence and tumor resistance in aged BALB/c mice. *Mech. Ageing Dev.* **6**: 15-24.
16. Stjernsward, J. 1966. Age-dependent tumor-host barrier and effect of carcinogen-induced immunodepression on rejection of isografted methylcholanthrene-induced sarcoma cells. *J. Natl. Cancer Inst.* **37**: 505-512.
17. House, E. L., B. Pansky, and M. S. Jacobs. 1961. Age changes in the blood of the Golden hamster. *Am. J. Physiol.* **200**: 1018-1022.
18. Tsang, K. Y., and H. H. Fudenberg. 1981. Effects of inosiplex (ISO) on the immune response of aging hamsters. Proceedings of the American Aging Association, 11th Annual National Meeting. 21.
19. Braun, W., Y. Yajima, and M. Ishizuka. 1970. Synthetic polynucleotides as restorers of normal antibody-forming capacities in aged mice. *J. Reticuloendothel. Soc.* **7**: 418-424.
20. Han, I. H., and A. G. Johnson. 1976. Regulation of the immune system by synthetic polynucleotides. *J. Immunol.* **117**: 423-427.
21. Friedman, D., V. Keiser, and A. Globerson. 1974. Reactivation of immunocompetence of spleen T cells of aged mice. *Nature (Lond.)*. **251**: 545-547.
22. Goldstein, A. L., G. B. Thurman, T. L. Low, G. E. Triv-

- ers, and J. L. Rossio. 1979. *In Physiology and Cell Biology of Aging*. A. Cherkin, C. E. Finch, N. Kharasch, T. Makinodan, F. L. Scott, and B. L. Strehler, editors. Raven Press, New York. 51.
23. Strehler, B. L. 1977. *Time, Cells and Aging*. Academic Press, Inc., New York.
24. Biro, J., and E. Beregi. 1979. Electron microscopical and immunoelectronmicroscopical examination of young and old people after influenza vaccination. *Acta Gerontol.* **9**: 235-239.
25. Renoux, G., and M. Renoux. 1972. Restauration par de phenylimidothiazole de la response immunologique des souris agees. *C.R. Acad. Sci. Ser. D.* **274**: 3034-3035.
26. Morimoto, C., T. Abe, and M. Homma. 1979. Restoration of T-cell function in aged mice with long-term administration of levamisole. *Clin. Immunol. Immunopathol.* **12**: 316-322.
27. Makinodan, T., and J. W. Albright. 1979. Restoration of impaired immune functions in aging animals. II. Effect of mercaptoethanol in enhancing the reduced primary antibody responsiveness *in vitro*. *Mech. Ageing Dev.* **10**: 325-340.
28. Makinodan, T., and J. W. Albright. 1979. Restoration of impaired immune functions in aging animals. III. Effect of mercaptoethanol in enhancing the reduced primary antibody responsiveness *in vivo*. *Mech. Ageing Dev.* **11**: 1-8.
29. Fridman, H., R. Calle, and A. Morin. 1980. Double-blind study of isoprinosine influence on immune parameter in solid tumor-bearing patients treated by radiotherapy. *Int. J. Immunopharmacol.* **2**: 194.
30. Ginsberg, T., and A. J. Glasky. 1977. Inosiplex: an immunomodulation model for the treatment of viral disease. *Ann. NY Acad. Sci.* **284**: 128-138.
31. Wybran, J. 1980. Immunomodulatory properties of isoprinosine in man: *in vitro* and *in vivo* data. *Int. J. Immunopharmacol.* **2**: 197.
32. Lesourd, B., G. Rancurel, J. M. Huraux, A. Pompidou, C. Jacque, D. Denvil, A. Buge, and R. Mauliac. 1980. Immunological restoration *in vivo* and *in vitro*, isoprinosine therapy and prognosis of acute encephalitis. *Int. J. Immunopharmacol.* **2**: 195.
33. Tsang, K. Y., and H. H. Fudenberg. 1982. Isoprinosine as an immunopotentiator in an animal model of human osteosarcoma. *Int. J. Immunopharmacol.* **3**: 383-389.
34. Boyum, A. 1968. Isolation of leucocytes from human blood: further observations. *Scand. J. Clin. Lab. Invest.* **92**(Suppl.): 31-50.
35. Ron, N., A. Laufer, and I. Gery. 1973. An unusual response of hamster lymphocytes to PHA. *Immunology* **25**: 433-439.
36. Tsang, K. Y., M. J. Gnagy, Y. Yamamura, and H. H. Fudenberg. 1981. Detection of natural killer cell-mediated cytotoxicity in hamsters using Raji target cells. *Clin. Immunol. Immunopathol.* **21**: 332-340.
37. Tsang, K. Y., H. H. Fudenberg, J. F. Pan, M. J. Gnagy, and C. B. Bristow. 1982. An *in vitro* study on the effects of isoprinosine on immune responses in cancer patients. *Int. J. Immunopharmacol.* In press.
38. Tsang, K. Y., I. Singh, M. J. Gnagy, and D. Tanski. 1979. Abnormal monocyte chemotaxis in hamsters with human osteosarcoma. *Micron.* **10**: 227-228.
39. Padarathsingh, M. L., J. H. Dean, T. R. Jerrells, J. L. McCoy, D. D. Lewis, and J. W. Northrop. 1979. Evidence for characterization of suppressor cells in BALB/c mice bearing ADJ-PC5 plasmacytomas. *J. Natl. Cancer Inst.* **62**: 1235-1241.
40. Bankhurst, A. D. 1982. The modulation of human natural killer cell activity by prostaglandins. *J. Clin. Lab. Immunol.* **7**: 85-91.
41. Leung, K., and H. S. Koren. 1982. Regulation of human natural killing. II. Protective effect of interferon on NK cells from suppression by PGE₂. *J. Immunol.* **129**: 1742-1747.
42. Singh, I., K. Y. Tsang, and W. S. Blakemore. 1979. A model for human osteosarcoma in hamsters. *Clin. Orthop. Rel. Res.* **144**: 305-310.
43. Albright, J. F., and T. Makinodan. 1976. Decline in the growth potential of spleen-colonizing bone marrow stem cells of long-lived aging mice. *J. Exp. Med.* **144**: 1204-1212.
44. Albright, J. F., and T. Makinodan. 1966. Growth and senescence of antibody-forming cells. *J. Cell. Physiol.* **67**(Suppl. 3): 185-206.
45. Price, G. B., and T. Makinodan. 1972. Immunologic deficiencies in senescence. *J. Immunol.* **108**: 403-412.
46. Price, G. B., and T. Makinodan. Immunologic deficiencies in senescence. *J. Immunol.* **108**: 413-417.
47. Diaz-Jouanen, E., R. C. Williams, Jr., and R. G. Strickland. 1975. Age-related changes in T and B cells. *Lancet.* **I**: 688-689.
48. Phair, J. P., C. A. Kauffman, and A. Bjornson. 1978. Host defenses in the aged: Evaluation of components of the inflammatory responses. *J. Infect. Dis.* **138**: 67-73.
49. Hallgren, H. M., J. H. Kersey, D. P. Dubey, and E. J. Yunis. 1978. Lymphocyte subsets and integrated immune function in aging humans. *Clin. Immunol. Immunopathol.* **10**: 65-78.
50. Gupta, S., and R. A. Good. 1979. Subpopulations of human T lymphocytes. *J. Immunol.* **122**: 1214-1219.
51. Cowan, M. J., P. Fujiwara, D. W. Wara, and A. J. Ammann. 1981. Effect of thymosin on cellular immunity in old age. *Mech. Ageing Dev.* **15**: 29-39.
52. Makinodan T., and M. M. B. Kay. 1980. Age influence on the immune system. *Adv. Immunol.* **29**: 287-330.
53. Morin, A., D. Tello, and J. J. Ballett. 1982. Activation mechanisms of human lymphocytes by the inosine derivative, isoprinosine (Ip). *Int. J. Immunopharmacol.* **4**: 286. (Abstr.)
54. Joseph, M., A. Capron, A. B. Tonel, P. G. Gosset, and C. Meanier. 1982. Immunomodulating properties of isoprinosine *in vivo* and *in vitro*. *Int. J. Immunopharmacol.* **4**: 286. (Abstr.)