## Amendment history:

Correction (April 1985)

# Antineutrophil autoantibodies in Graves' disease. Implications of thyrotropin binding to neutrophils.

S A Weitzman, ..., F Maloof, E C Ridgway

J Clin Invest. 1985;75(1):119-123. https://doi.org/10.1172/JCI111663.

# Research Article

The hyperthyroidism of Graves' disease may be caused by autoantibodies to thyrotropin (TSH) receptors. We have found that patients with this disease have autoantibodies to neutrophils as well, which can be displaced by TSH. Using a radiochemical opsonic assay, we found serum antibodies against homologous neutrophils in 6 of 11 Graves' patients. With a staphylococcal protein A-binding assay, we detected circulating antibodies to homologous neutrophils in 10 of 20 patients, while finding cell-bound antibody on autologous neutrophils in 7 of 8 (including 2 with negative serum tests). Use of human 125I-TSH in a radioligand binding assay revealed that TSH bound to neutrophils rapidly (maximum binding within 10 min at 22 degrees C, pH 7.4), specifically (less than 20% nonspecific binding), and reversibly. Adding TSH to the radiochemical assay resulted in a dose-dependent inhibition of opsonic antibody activity in serum from patients with Graves' disease. In contrast, TSH did not inhibit antibody activity of serum from patients with immune neutropenia not associated with thyroid disease. Our findings suggest a basis for the association of Graves' disease with neutropenia. Furthermore, the discovery of such antineutrophil antibodies in Graves' disease permits detection of [...]



Find the latest version:

https://jci.me/111663/pdf

# Antineutrophil Autoantibodies in Graves' Disease

Implications of Thyrotropin Binding to Neutrophils

Sigmund A. Weitzman, Thomas P. Stossel, David C. Harmon, Gilbert Daniels, Farahe Maloof, and E. Chester Ridgway Hematology-Oncology and Thyroid Units, Massachusetts General Hospital and Department of Medicine, Harvard Medical School, Boston, Massachusetts 02114

## Abstract

The hyperthyroidism of Graves' disease may be caused by autoantibodies to thyrotropin (TSH) receptors. We have found that patients with this disease have autoantibodies to neutrophils as well, which can be displaced by TSH. Using a radiochemical opsonic assay, we found serum antibodies against homologous neutrophils in 6 of 11 Graves' patients. With a staphylococcal protein A-binding assay, we detected circulating antibodies to homologous neutrophils in 10 of 20 patients, while finding cellbound antibody on autologous neutrophils in 7 of 8 (including 2 with negative serum tests). Use of human <sup>125</sup>I-TSH in a radioligand binding assay revealed that TSH bound to neutrophils rapidly (maximum binding within 10 min at 22°C, pH 7.4), specifically (<20% nonspecific binding), and reversibly. Adding TSH to the radiochemical assay resulted in a dosedependent inhibition of opsonic antibody activity in serum from patients with Graves' disease. In contrast, TSH did not inhibit antibody activity of serum from patients with immune neutropenia not associated with thyroid disease.

Our findings suggest a basis for the association of Graves' disease with neutropenia. Furthermore, the discovery of such antineutrophil antibodies in Graves' disease permits detection of cell-bound antibody when free antibody is not present.

## Introduction

The association of Graves' disease with neutropenia was first described by Kocher (1) in 1908, and subsequently confirmed (2). Other groups, studying patients with autoimmune neutropenia, have observed a number of individuals with concurrent or prior thyroid disease. The coexistence of both neutropenia and Graves' disease in some patients led us to study systematically a group of patients with Graves' disease for the presence of antineutrophil autoantibodies. Furthermore, we questioned whether the autoantibodies directed against thyroid thyrotropin (TSH)<sup>1</sup> receptors may cross-react with thyrotropin-binding moieties on neutrophils. To investigate this hypothesis, we studied the binding of thyrotropin to human neutrophils

J. Clin. Invest. © The American Society for Clinical Investigation, Inc. 0021-9738/85/01/0119/05 \$1.00 Volume 75, January 1985, 119–123 and the interactions in vitro of thyrotropin, antineutrophil autoantibodies, and neutrophils.

#### **Methods**

Patients. Sera and citrated whole blood were obtained from patients with Graves' disease and normal volunteers after informed consent. 22 patients were studied. There were 18 females and 4 males, with an age range of 23-74 yr. Their therapy and clinical status is presented in Table I. Patients were classified as having Graves' disease on the basis of the presence of diffuse goiter, exophthalmos, and a current or documented history of hyperthyroidism. Hyperthyroidism was defined on the basis of classical clinical symptoms, elevated serum thyroxine and triiodothyronine concentrations, and undetectable serum TSH levels. Patients were considered euthyroid by an absence of clinical symptoms of hyperthyroidism or hypothyroidism with normal serum thyroxine and triiodothyronine levels.

Neutrophils were purified by sedimentation of citrated blood with dextran (3). The leukocyte-rich supernatants were then centrifuged on Ficoll-sodium diatrizoate gradients, and erythrocyte contamination of the pellets was removed by hypotonic lysis.

IgG was purified by ammonium sulfate precipitation followed by ion exchange chromatography on diethylaminoethyl cellulose (4).

Antineutrophil antibodies were assessed by two methods. The cell:cell recognition assay was used to detect antineutrophil antibody opsonic activity in serum as previously described (3, 5). This assay is based on the observation that normal neutrophils recognize other neutrophils opsonized with antibody and respond with an increase in glucose oxidation.  $2.4 \times 10^6$  normal neutrophils were suspended in 0.2 ml of Krebs-Ringer-phosphate medium (KRP), pH 7.4, and incubated (sensitized) for 30 min at 22°C in 0.4 ml test serum. After being washed, these cells (designated "test cells") were resuspended in 0.2 ml KRP. An equal number of normal neutrophils (designated "indicator cells") was not exposed to serum. Instead, they were incubated for 60 min in 0.8 ml KRP containing 100,000 cpm of [1-14C]glucose. Indicator and washed test cells were then brought together in a total volume of 1 ml KRP in sealed polypropylene vials with a strip of hyamine-impregnated filter paper inserted through the top. The amount of radioactive CO<sub>2</sub> evolved after 30-min incubation at 37°C was determined by counting the filter paper in a scintillation spectrometer. The mean of control incubations (test cells sensitized with normal IgG or serum) was defined as 100% of control activity. In some experiments, varying amounts of bovine TSH (bTSH, Thytropar; Armour Pharmaceutical Co., Tarrytown, NY) were added to test cell suspensions before adding serum.

A second method, the staphylococcal slide assay (6), enabled us to look for IgG antineutrophil antibodies directly bound to autologous neutrophils as well as for free IgG in the serum capable of binding to homologous neutrophils. Dextran-sedimented leukocyte suspensions (autologous or homologous) were allowed to adhere to glass slides, fixed in paraformaldehyde, and washed. The slides were then covered with 50  $\mu$ l of test or control serum and incubated for 30 min at 22°C. Excess serum was then rinsed off with calcium-free KRP buffer, pH 7.4, containing 40 mg/ml bovine serum albumin (Sigma Chemical Co., St. Louis, MO). In each assay, two known positive and two known negative control sera were evaluated in triplicate in parallel with the test sera. In a direct assay, this serum sensitization step was omitted. After sensitization, 50  $\mu$ l of protein A containing *Staphylococcus* 

Address correspondence to Dr. Weitzman, Hematology-Oncology Unit, Cox 6, Massachusetts General Hospital, Boston, MA 02114.

Received for publication 18 January 1984 and in revised form 19 July 1984.

<sup>1.</sup> Abbreviations used in this paper: bTSH, bovine thyrotropin; hTSH, human thyrotropin; KRP, Krebs-Ringer-phosphate medium; TSH, thyrotropin; TSI, thyroid-stimulating immunoglobulins.

Table I	I.	Summary	of	Patients an	d Anti	body	v Test	Results	ŝ
---------	----	---------	----	-------------	--------	------	--------	---------	---

							Antineutrophil antibodies			
					Total	Total	Indirect test (homologous cells)		Direct test (autologous cells)	
Patient no. Age	Sex	Treatment*	Clinical status‡	leukocyte count	neutrophil count	Opsonic test	Slide test	Slide test		
					mm³	mm <sup>3</sup>				
1	36	F	I <sup>131</sup>	Euthyroid		_	Positive	Positive	_	
2	23	F	I <sup>131</sup>	Hyperthyroid	5,100	1,300	_	Positive	_	
3	36	F	Propylthiouracil	Euthyroid	2,650	100 <sup>II</sup>	Negative	Positive		
4	30	F	Methimazole	Euthyroid	7,700	_	Positive	Negative	_	
5	37	F	I <sup>131</sup>	Euthyroid	4,200	2,300	_	Negative	_	
6	53	F	I <sup>131</sup>	Hypothyroid	6,600	3,600	Positive	Positive	_	
7	50	М	I <sup>131</sup>	Euthyroid	7,700	5,200	Positive	Positive	_	
8§	49	М	None	Euthyroid	10,600	5,800	Positive	Positive		
ຶ	37	F	I <sup>131</sup>	Euthyroid	6,800	_	Negative	Negative	_	
10	68	F	I <sup>131</sup>	Euthyroid	6,500	4,000	Negative		_	
11	43	М	I <sup>131</sup>	Euthyroid	_	_	Negative	Negative		
12	26	F	None	Hyperthyroid	2,900	1,400	_	Negative	Positive	
13	34	F	I <sup>131</sup>	Euthyroid	8,900	4,900	_	Negative		
			Propylthiouracil	-				-		
14	74	F	I <sup>131</sup>	Hyperthyroid	8,400	6,700	_	Positive	Positive	
15	34	F	I <sup>131</sup>	Euthyroid	4,600	2,800	—	Positive	Positive	
16§	57	F	None	Euthyroid	_		_	Positive	Positive	
17	36	F	I <sup>131</sup>	Euthyroid	5,500	4,900			Positive	
18	20	F	Methimazole	Hyperthyroid	800	<b>30</b> <sup>  </sup>	Negative	Negative	Positive	
19	32	F	None	Hyperthyroid	3,000	1,900	_	Negative	Negative	
20	53	М	None	Hyperthyroid	6,400	4,000	_	Negative		
21	36	F	None	Hyperthyroid	4,600	2,200	Positive	Positive	Positive	
22	33	F	None	Hyperthyroid	6,000	3,400	_	Negative	<u> </u>	

\* Treatment: I<sup>131</sup>, therapeutic radioiodine at any previous time. Propylthiouracil or methimazole, under treatment with these agents at time of study. \$ Euthyroid (ophthalmic) Graves' disease. 
Patients with acute granulocytopenia thought to be secondary to antithyroid drugs.

aureus Cowan 1 suspensions was lavered onto the cells and incubated for 30 min at room temperature. These organisms, cultured as previously described (7), had been killed by incubation in 0.5% formaldehyde for 3 h at 20°C, and heat-treated at 80°C for 3 min. Bacteria were prepared by suspending 100 ml of the stock in 2 ml of suspension solution (0.05% Triton X-100, 0.15 mM NaCl, 5 mM Na<sub>2</sub> EDTA, 0.2 mg/ml sodium azide, pH 7.4), and centrifuged at 12,000 g for 30 s. To the resultant pellet of 50 µl of packed staphylococci, 1 ml of suspension solution containing 200 mg/ml bovine serum albumin was added and swirled with a vortex mixer to form a suspension. Slides were then immersed sequentially in bovine albumin solutions of decreasing concentrations in KRP buffer. The slides were then stained, inspected under light microscopy, and the percentage of neutrophils that bound three or more staphylococci was determined. Each test was done in triplicate, at least 100 cells were observed, and the average percentage of labeled neutrophils was recorded. A test was considered positive for antineutrophil antibodies if >20% of neutrophils was labeled. Validation of criteria for positive assays has been described (5). The rate of false positive tests is <4%.

Hormone preparation. Highly purified human thyrotropin (hTSH) was provided by the National Pituitary Agency, Baltimore, MD, with an immunopotency of 3-5 U/mg of Mill Hill Reference Standard B. This material was iodinated with <sup>125</sup>I by the lactoperoxidase method (8) to a specific activity of 70-100  $\mu$ Ci/ $\mu$ g and then separated from free iodine by filtration over a 90 × 1.5 cm Sephadex G100 chromato-graphic column. This material was 85-100% immunoprecipitable after

incubation with excess rabbit anti-hTSH serum at final dilution of 1:1,000 (9).

Highly purified bTSH was generously supplied by Dr. John Pierce. Cruder preparations of bTSH were obtained from Armour Pharmaceutical Co. Purified human luteinizing hormone (LER-960) and folliclestimulating hormone (NIH-FSH-HS-1) were obtained from the National Pituitary Agency.

hTSH binding to neutrophils was assessed by a radioligand binding assay.  $5 \times 10^5$  gradient-purified neutrophils were incubated with varying amounts of <sup>125</sup>I-hTSH and test material in isotonic KRP buffer, pH 7.4, with total incubation volumes of 700  $\mu$ l. To terminate incubations,  $500-\mu$ l aliquots of the suspensions were removed and placed immediately into plastic tubes containing 3 ml of iced buffer. These tubes were then immediately inverted over glass fiber filters (GF/A, Whatman Laboratory Products, Inc., Clifton, NJ) that had been mounted in a vacuum suction manifold and presoaked with excess TSH ( $>10^{-5}$  M). This presoaking resulted in negligible counts being bound to the filters in the absence of cells. Under vacuum suction, suspension media were removed and the cells were washed with 12 ml of iced buffer. The filters were then counted for <sup>125</sup>I in an autogamma spectrometer. "Specific binding" was defined as the amount of the total bound radioactivity that was displaced by an excess of unlabeled bTSH (>10<sup>-5</sup> M) that had been added to the incubations concurrently with the radiolabeled TSH.

To assess whether deiodination or degradation of hormone occurred,  $5\times10^5$  neutrophils were incubated with  $^{125}I\text{-}hTSH$  for various times

at both 22° and 37°C. At the completion of incubation, the cells were pelleted by centrifugation at 12,000 g for 1 min, and the supernatant was assayed for immunoprecipitability of radioactivity, using rabbit anti-hTSH serum as noted above. This was compared with immuno-precipitability of preincubation samples of labeled hTSH.

Reversibility of binding was investigated by incubating cells with <sup>125</sup>I-hTSH as described above, subsequently adding excess unlabeled hormone to the incubation, and then determining residual counts bound to cells.

## Results

Antineutrophil antibodies. Using the opsonic method to assay serum against homologous neutrophils, 6 of 11 Graves' patients had antibodies. Using the staphylococcal assay for the same purpose, the sera of 10 of 20 patients were positive. When we studied the autologous neutrophils of eight patients by the staphylococcal slide test, seven of them were found to have cell-bound antibodies, including two in whom the sera were negative for binding of antineutrophil IgG to homologous cells. Antibodies to lymphocytes were not found. As shown in Table I, antibodies were detected even in patients with normal neutrophil counts. Neutropenia ( $<1,800/mm^3$ ) was observed in two patients off treatment as well as in two patients who were possibly neutropenic because of reaction to propylthiouracil or methimazole.

Inhibition of antineutrophil opsonic antibody activity by TSH. When bTSH was added to test cell incubations, the opsonic activity of these antibodies was inhibited in a dosedependent fashion (Fig. 1). bTSH did not, however, inhibit the opsonizing activity of serum from two patients with



Figure 1. Effects of TSH on opsonic activity of antineutrophil antibodies. Test cells were exposed to antisera in the presence of varying amounts of crude bTSH for 30 min at 22°C, washed, and incubated with indicator cells for 30 min. Ordinate shows recognition response ([1-<sup>14</sup>C]glucose oxidation to <sup>14</sup>CO<sub>2</sub> by indicator cells) as percentage of control (recognition response in presence of cells sensitized with normal serum). Solid lines, serum from two different patients with Graves' disease. Dashed lines, serum from patients with autoimmune neutropenia (lower) and drug-induced immune neutropenia (upper). Each point represents mean of 2 or 4 determinations. • and  $\circ$ , Graves'. • and  $\Box$ , other immune neutropenia.

immune neutropenia who had no history of, or evidence of, thyroid disease (Fig. 1).

Binding of hTSH to neutrophils. hTSH was noted to bind rapidly, stably, and reversibly (Fig. 2) to human neutrophils. Specific binding (as defined above) varied between 2.5 and 16% of the added counts. Nonspecific binding was always <20% of the specific binding and generally much lower (in many experiments, there was no nonspecific binding in excess of the cell-free control filter blanks). The effects of pH on this system are also illustrated in Fig. 2. As noted by others (10, 11), specific binding to TSH is greatest at low pH, intermediate in the physiologic range, and lowest at alkaline pH. Effects of temperature are shown in Table II. While there was little difference in binding at 22° and 37°C, parallel incubations at 0°C resulted in smaller amounts of hormone bound. Reversibility of binding was demonstrated by incubating cells with <sup>125</sup>I-hTSH for 15 min, and then adding excess (> $10^{-5}$ ) bTSH. Within 5 min, all labeled TSH was displaced from the neutrophils (Fig. 2, top, dashed line). The displaced <sup>125</sup>I-hTSH was 85-95% immunoprecipitable (the same as control <sup>125</sup>I-hTSH), suggesting that degradation or deiodination did not occur during the incubation.

To investigate further whether the interaction of TSH with neutrophils suggested characteristics of a physiologic receptor, we subjected the radioligand binding data to Scatchard analysis (12, 13). Under conditions used in these assays, we were not



Figure 2. Upper graph: time course of specific hTSH binding to neutrophils. 9,400 cpm human <sup>125</sup>I-TSH was added to  $5 \times 10^5$ neutrophils at 22°C, pH 7.4, and incubated for the times indicated. Parallel incubations with excess unlabeled bTSH (10<sup>5</sup> M) were performed to define nonspecific binding. Ordinate shows TSH specifically bound (total binding minus nonspecific binding) as percent of total labeled TSH added. Dashed line, cells incubated with labeled hormone for 15 min, then excess unlabeled TSH added. Each point represents mean and range of two experiments, each with duplicate determinations. Lower graph: effects of pH on TSH binding. 19,000 cpm <sup>125</sup>I-TSH was added to  $5 \times 10^5$  neutrophils at 22°C for 15 min at the indicated pH. Specific binding was determined as above. Each point represents mean and range of duplicate determinations.

Table II. Effect of Temperature on Specific Binding of Human<sup>125</sup>I TSH to Neutrophils

	% TSH specifically bour	nd
Temperature	10-min incubation	20-min incubation
0°	10.1±0.9	10.8±2.3
22°	14.6±2.6	15.3±2.7
37°	15.7±2.4	14.6±1.7

19,000 cpm of <sup>125</sup>I-TSH were added to  $5 \times 10^5$  neutrophils, pH 7.4, and incubated for the times indicated. Results are expressed as mean±SE (4–6 determinations) percent specific binding.

able to demonstrate consistently the high affinity binding characteristic of peptide hormone receptors.

Additional information about the specificity of human TSH binding to neutrophils is shown in Fig. 3. Unlabeled human TSH was much more efficient at competing with labeled human TSH for binding to neutrophils than was unlabeled purified bTSH. Neither follicle-stimulating hormone nor luteinizing hormone had competing activity.

#### Discussion

The major points in this study are (a) patients with Graves' disease frequently have antineutrophil autoantibodies, (b) TSH inhibits the opsonic activity of these antibodies, and (c) TSH binds to neutrophils rapidly and reversibly.

TSH binding has been described with a variety of tissues (10, 11, 14–16). Published data on the binding characteristics vary widely, depending on both the methods used and the species source of TSH and cell (or membrane) preparations. Reports from other laboratories agree with our observations of certain important features, such as the rapid time course of binding (14) and pH dependence (10, 11). Also, receptors for porcine TSH have been described on human neutrophils (16). Whether these neutrophil binding sites represent biologically important "physiologic receptors" is not certain, but some



*Figure 3.* Specificity of TSH binding. Neutrophils were incubated at 22°C, pH 7.4, for 15 min with human <sup>125</sup>I-TSH and unlabeled ligands in the amounts indicated. Results are expressed as percent of control specific binding (specific binding in the absence of unlabeled ligands). Each point represents mean of duplicate determinations.

studies show that TSH may initiate neutrophil cyclic AMP production and may be relevant in neutrophil function (17).

Hyperthyroidism in Graves' disease is thought to be due to autoantibodies directed against TSH receptors on thyroid cells (18-24). In most cases, immunoglobulins that inhibit binding of radiolabeled TSH to thyroid receptors (thyrotropin binding inhibiting immunoglobulins) are demonstrable in sera from these patients. Other assays, which do not necessarily correlate with the radioligand binding methods, demonstrate that the immunoglobulins from these patients behave as TSH agonists, i.e., activate adenylate cyclase or stimulate colloid droplet formation in thyroid preparations (thyroid-stimulating immunoglobulins or TSI). Recent studies with monoclonal antibodies argue that the antibodies that inhibit TSH binding to thyroid membranes are not the same as those that stimulate the gland (21, 22). Since the antineutrophil antibodies described here behave in a manner similar to the thyroid thyrotropin displacing immunoglobulins, it is possible that these two types of autoantibodies are actually the same, i.e., directed against the same or similar antigens expressed on more than one tissue.

The finding of circulating antineutrophil antibodies in 50-55% of our patients raises several questions, including (a) Why were so few of these individuals neutropenic? and (b) If these antibodies are related to those presumed pathogenic in Graves' disease, why were they not found in a higher fraction of patients? At least two possible explanations exist for the first question. First, it is possible that some of the antibodies may not, in vivo, alter the survival of the neutrophils against which they are directed, or, second, in cases where cell lifespan is shortened, increased marrow production might compensate (similar to the compensated hemolytic state of some erythrocvte disorders). The second question is more complex, but some of our data may help approach its answer. While serum antibodies against homologous cells were not found in almost half the patients, cell-bound IgG antibody was observed on autologous cells of seven of eight patients so studied. These findings are similar to the results of Coombs' antiglobulin test in autoimmune hemolytic anemia. While cell-bound antibody may be found in almost all cases (i.e., the direct antiglobulin test), the indirect test for serum antibody can be negative from 20 to 65% of the time, depending on the method of erythrocyte preparation (25). This analogy might also explain some of the false negative tests for TSI or thyrotropin displacing activity that can be observed in Graves' patients (26). Until recently (21, 22), these negative results occurred often enough to have cast doubt on the model of Graves' disease as an autoimmune process with antibodies directed against TSH receptors. However, it seems reasonable to postulate, on the basis of what is known about autoimmune hemolytic anemia and the neutrophil data shown here, that in some cases of Graves' disease, most all of the TSI or thyrotropin displacing activity could be cell-bound (either on thyroid or other cells), and that serum levels might be below limits of detection.

#### Acknowledgments

We would like to acknowledge the assistance of Mary Desmond, Linda Lorenz, Dr. Lewis T. Williams, and Dr. Laurie Glimcher.

This work was supported by gifts from Edwin H. Hiam and the Edwin S. Webster Foundation.

#### References

1. Kocher, T. 1908. Blutuntersuchungen bei morbus Basedowii mit Beitragen ur Frudiagnose und Theorie der Krankheit. Arch. Klin. Chir. 87:131.

2. Hertz, S., and J. Lerman. 1932. The blood picture in exophthalmic goitre and its changes resulting from iodine and operation. J. Clin. Invest. 11:1179-1196.

3. Weitzman, S. A., M. C. Desmond, and T. P. Stossel. 1979. Antigenic modulation and turnover in human neutrophils. J. Clin. Invest. 64:321-325.

4. Williams, C. A., and M. W. Chase. 1976. Methods Immunol. Immunochem. 5:424-444.

5. Weitzman, S. A., and T. P. Stossel. 1978. Drug-induced immunological neutropenia. *Lancet.* I:1068-1072.

6. Harmon, D. C., S. A. Weitzman, and T. P. Stossel. 1980. A staphylococcal slide test for detection of anti-neutrophil antibodies. *Blood.* 56:64–69.

7. Arvidson, S., T. Holme, and T. Wadstrom. 1970. Formation of bacteriolytic enzymes in batch and continuous culture of *Staphylococcus aureus*. J. Bacteriol. 104:227-235.

8. Wood, P. J., M. M. Shawan, and V. Marks. 1975. Radioimmunoassay of plasma prolactin. *Lancet.* II:1040-1041.

9. Ridgway, E. C., B. D. Weintraub, and F. Maloof. 1974. Metabolic clearance and production rates of human thyrotropin. J. Clin. Invest. 53:895-903.

10. Pekanen, F., and B. D. Weintraub. 1978. Thyrotropin binding to cultured lymphocytes and thyroid cells. *Endocrinology*. 103:1668-1677.

11. Lee, G., E. F. Grollman, S. Dyer, F. Beguinot, L. D. Kohn, W. H. Habig, and M. C. Hardegree. 1979. Tetanus toxin and thyrotropin interactions with rat brain membrane preparations. *J. Biol. Chem.* 254:3826-3832.

12. Rosenthal, H. E. 1967. A graphic method for the determination and presentation of binding parameters in complex systems. *Anal. Biochem.* 20:25-532.

13. Scatchard, G. 1949. The attraction of proteins for small molecules and ions. *Ann. NY Acad. Sci.* 51:660–672.

14. Kishihara, M., Y. Nakao, Y. Baba, S. Matsukura, K. Kuma, and T. Fujita. 1979. Interaction between thyroid-stimulating immunoglobulins and thyrotropin receptors in fat cell membranes. J. Clin. Endocrinol. Metab. 49:706-711.

15. Smith, B. R. 1977. Membrane receptors for polypeptide hormones. Adv. Clin. Chem. 19:91-124.

16. Chabaud, O., and S. Lissitsky. 1977. Thyrotropin specific binding to human peripheral blood monocytes and polymorphonuclear leukocytes. *Mol. Cell Endocrinol.* 7:79-87.

17. Stolc, V. 1972. Regulation of iodine metabolism in human leukocytes by adenosine 3',5'-monophosphate. *Biochim. Biophys. Acta.* 264:285-288.

18. Volpe, R. 1978. The pathogenesis of Graves' disease: an overview. Clin. Endocrinol. Metab. 7:3-29.

19. McKenzie, J. M., M. Zakarija, and A. Sato. 1978. Humoral immunity in Graves' disease. *Clin. Endocrinol. Metab.* 7:31-45.

20. O'Donnell, J., K. Trokoudes, J. Silverberg, V. Row, and R. Volpe. 1978. Thyrotropin displacement activity of serum immunoglobulins from patients with Graves' disease. J. Clin. Endocrinol. Metab. 46:770-777.

21. Valente, W. A., P. Vitti, Z. Yavin, C. M. Rotella, E. F. Grollman, R. S. Toccafondi, and L. D. Kohn. 1982. Monoclonal antibodies to the thyrotropin receptor: stimulating and blocking antibodies derived from the lymphocytes of patients with Graves' disease. *Proc. Natl. Acad. Sci. USA.* 79:6680–6684.

22. Kohn, L. D., W. A. Valente, P. Laccetti, J. L. Cohen, S. M. Aloj, and E. F. Grollman. 1983. Multicomponent structure of the thyrotropin receptor: receptor to Graves' disease. *Life Sci.* 32:15-20.

23. Zakarija, M., and J. M. McKenzie. 1983. Thyroid-stimulating antibody (TSAb) of Graves' disease. *Life Sci.* 32:31-44.

24. Bech, K. 1983. Immunological aspects of Graves' disease and importance of thyroid stimulating immunoglobulins. *Acta Endocrinol.* 254(Suppl):8-38.

25. Petz, L. D., and G. Garraty. 1975. Laboratory correlations in immune hemolytic anemias. *In* Laboratory Diagnostic of Immunologic Disorders. G. N. Vyas, D. P. Stites, and G. Brecher, editors. Grune & Stratton Inc., Orlando, FL. 139–153.

26. Volpe, R. 1979. Immunopathology of Graves' disease. Fed. Proc. 38:2611-2612.