

Prostacyclin production by the deendothelialized rabbit aorta.

J M Boeynaems, ... , N Galand, P Ketelbant

J Clin Invest. 1985;76(1):7-14. <https://doi.org/10.1172/JCI111979>.

Research Article

The acute effect of in vitro deendothelialization on the production of prostacyclin (PGI₂) by the rabbit aorta has been investigated. The effectiveness of removing endothelium by rubbing it against filter paper or scraping it with a scalpel was demonstrated by scanning electron microscopy and en face examination after silver staining. Endothelium removal produced an immediate stimulation of PGI₂ release, resulting in 408% of the control after rubbing and 367% of the control after scraping, during the first 30-min period of incubation. This increased production of PGI₂ gradually declined over time to reach values similar to the control after 2h. At that time, the deendothelialized aorta was totally unresponsive to the stimuli that increase PGI₂ release in the intact aorta (acetylcholine, ADP, ionophore A23187, and arachidonic acid). The enhanced production of PGI₂ in the deendothelialized aorta was associated with an increased release of free arachidonic acid (353% of the control): in contrast with PGI₂, this stimulation was maintained for at least 150 min. A transient exposure of the deendothelialized aorta to ibuprofen (250 microM) was followed by a rebound of PGI₂ production, which was also prolonged by BW-755C (3-10 microM). In conclusion, removal of the endothelium triggered an immediate and sustained mobilization of free arachidonic acid in the rabbit aorta: the resulting increase of PGI₂ production was short-lived, probably as a [...]

Find the latest version:

<https://jci.me/111979/pdf>



Prostacyclin Production by the Deendothelialized Rabbit Aorta

J. M. Boeynaems, N. Galand, and P. Ketelbant

Institute of Interdisciplinary Research and Department of Pathology of Erasme Hospital, Université Libre de Bruxelles, Brussels, Belgium

Abstract

The acute effect of in vitro deendothelialization on the production of prostacyclin (PGI₂) by the rabbit aorta has been investigated. The effectiveness of removing endothelium by rubbing it against filter paper or scraping it with a scalpel was demonstrated by scanning electron microscopy and *en face* examination after silver staining. Endothelium removal produced an immediate stimulation of PGI₂ release, resulting in 408% of the control after rubbing and 367% of the control after scraping, during the first 30-min period of incubation. This increased production of PGI₂ gradually declined over time to reach values similar to the control after 2 h. At that time, the deendothelialized aorta was totally unresponsive to the stimuli that increase PGI₂ release in the intact aorta (acetylcholine, ADP, ionophore A23187, and arachidonic acid). The enhanced production of PGI₂ in the deendothelialized aorta was associated with an increased release of free arachidonic acid (353% of the control): in contrast with PGI₂, this stimulation was maintained for at least 150 min. A transient exposure of the deendothelialized aorta to ibuprofen (250 μ M) was followed by a rebound of PGI₂ production, which was also prolonged by BW-755C (3–10 μ M). In conclusion, removal of the endothelium triggered an immediate and sustained mobilization of free arachidonic acid in the rabbit aorta: the resulting increase of PGI₂ production was short-lived, probably as a consequence of cyclooxygenase self-inactivation. Our results indicate that the subendothelium has a significant capacity to produce PGI₂, but that this capacity is expressed only briefly.

Introduction

The relative contribution of the endothelium and the subendothelium to the production of prostacyclin (PGI₂)¹ by arteries remains a controversial matter. PGI₂ is the major eicosanoid produced by cultured smooth muscle cells from rat (1), rabbit (2, 3), and bovine (4–6) aorta, whereas smooth muscle cells from porcine aorta synthesize mainly prostaglandins E₂ (PGE₂) and F_{2 α} (PGF_{2 α}) (7–9). After in vivo deendothelialization, the release of PGI₂ at the luminal surface of the rabbit aorta, induced by arachidonic acid, is abolished (10); removal of the

endothelium inhibits the capacity of dog femoral arteries to convert exogenous arachidonic acid into PGI₂ (11). Immunoradiometric assays and immunofluorescence studies have shown that the concentrations of PGI₂ synthase are equal in endothelial and smooth muscle cells of bovine aorta, but that the concentration of cyclooxygenase is 20-fold higher in the endothelial than in the smooth muscle cells (12, 13). We have performed a detailed kinetic study of the consequences of in vitro deendothelialization on arachidonic acid release and PGI₂ production in the rabbit aorta. Our results indicate that removal of the endothelium induces a sustained mobilization of free arachidonic acid and a transient stimulation of PGI₂ release: the subendothelium has a significant capacity to produce PGI₂, but, due to cyclooxygenase self-inactivation, the expression of this capacity is short-lived.

Methods

Preparation and incubation of aortic strips. White male rabbits weighing ~3 kg were killed by a blow on the neck. The aorta was quickly dissected from the iliac bifurcation up to the arch, trimmed free of fat and connective tissue, and cut into rings (\pm 3 mm large). These rings were opened by a longitudinal incision, and the intimal face of some of the resulting strips was either rubbed for 30 s against filter paper or scraped with a scalpel. Immediately thereafter, the strips were incubated at 37°C, under constant shaking (80 rpm), in a medium of the following composition: 124 mM NaCl; 5 mM KCl; 1.25 mM MgSO₄; 1.45 mM CaCl₂; 1.25 mM KH₂PO₄; 25 mM Hepes buffer, pH 7.4; and 8 mM glucose. The ratio of tissue weight to medium volume was roughly 25 mg/2 ml. In general the strips were incubated for five 30-min periods (60 min in some experiments), and the medium was replaced at the end of each period. For the measurement of arachidonic acid release, the same medium was used, but it contained in addition indomethacin (1 μ g/ml) and bovine serum albumin (BSA; 1 mg/ml). In some experiments, the strips of rabbit aorta were placed between two plexiglass plates held together by two lateral screws. The upper plate, the base of which was constituted by the aortic luminal surface, contained six holes (5 mm) that served as incubation chambers. The chambers were filled with 300- μ l incubation medium and incubated at 37°C for six 30-min periods. At the end of each period, the incubation medium was collected and replaced. Within an experiment (performed with one aorta), each experimental condition was tested in duplicate or triplicate.

Prostaglandin radioimmunoassay (RIA). The production of PGI₂ was measured by the RIA of prostaglandin-6-keto-F_{1 α} (6-K-PGF_{1 α}), performed directly in the incubation medium without extraction and chromatography. A rabbit antiserum was raised against 6-K-PGF_{1 α} coupled to BSA, as described (14): the limit of detection was 16 pg and the cross-reactions were 1.2% with PGF_{2 α} , 0.3% with PGE₂, and <0.1% with thromboxane B₂. 20- or 50- μ l aliquots of incubation media, ³H-6-K-PGF_{1 α} (11,000 dpm), anti-6-K-PGF_{1 α} antiserum (final dilution, 10⁻⁴), and bovine gamma globulins (0.25%) in Tris buffer (50 mM, pH 7.4) were incubated in a total volume of 0.4 ml for 60 min at room temperature. Then 0.4 ml of a cold 25% (wt/wt) solution of polyethylene glycol was added to separate bound and free antigen. PGE₂ was assayed by a similar method.

Metabolism of exogenous arachidonic acid. Intact and rubbed strips of rabbit aorta (\pm 50 mg) were incubated for four 30-min periods, in 2 ml of the usual medium. [³H]Arachidonic acid (135 Ci/mmol),

Address reprint requests to Dr. Boeynaems, Bldg. C, Campus Erasme, 808 Route de Lennik, 1070 Bruxelles, Belgium.

Received for publication 8 March 1984 and in revised form 26 February 1985.

1. *Abbreviations used in this paper:* ECD, electron capture detection; GLC, gas liquid chromatography; 6-K-PGF_{1 α} , prostaglandin-6-keto-F_{1 α} ; PGE₂, prostaglandin E₂; PGF_{2 α} , prostaglandin F_{2 α} ; PGI₂, prostacyclin; RPHPLC, reversed-phase high performance liquid chromatography.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/85/07/0007/08 \$1.00

Volume 76, July 1985, 7–14

purified shortly before use, was added at the beginning of the fourth period (1 μ Ci/ml, 0.25 μ g/ml). A 1.5-ml aliquot of the incubation medium was directly analyzed by reversed-phase high performance liquid chromatography (RPHPLC), using a μ -Bondapak C₁₈ column (3.9 \times 300 min, 10 μ m particles; Waters Associates, Millipore Corp., Milford, MA). The injector (model U6K), the pumps (model 6000 A), and the gradient programmer (model 600) were also from Waters Associates, Millipore Corp. Elution was performed with a mixture of acetonitrile/water/acetic acid (26:74:0.08; vol/vol) at a flow rate of 2 ml/min for 30 min. Then a linear gradient from this initial composition to acetonitrile/acetic acid (100:0.01; vol/vol) was developed in 37 min (2 ml/min). 2-ml fractions were collected and counted by liquid scintillation.

Assay of free arachidonic acid release. Arachidonic acid was measured by gas liquid chromatography (GLC) with electron capture detection (ECD). Aortic strip incubations were performed in a medium containing BSA (1 mg/ml) and indomethacin (1 μ g/ml). After the addition of 1 μ g docosahexaenoic acid as an internal standard, the incubation medium was extracted with 1 vol ethyl acetate. The FFA were converted into pentafluorobenzyl esters by a modification of the method of Wickramasinghe et al. (15). To the dry residue of the extract were added 290 μ g pentafluorobenzylbromide (in 5 μ l acetonitrile) and 44 μ g diisopropylethylamine (in 5 μ l acetonitrile). After 5 min at 40°C and evaporation of the solvent under nitrogen, the samples were redissolved in 50 μ l hexane. The 20-fold reduction in the amounts of reagents used, as compared with the original procedure, made it possible to directly analyze the samples without prior purification. The aorta strips did not release significant amounts of endogenous docosahexaenoic acid. GLC analysis was performed in a Varian instrument (model 3700; Varian Associates, Palo Alto, CA) equipped with a ⁶³Ni ECD. 2-m columns of 3% OV-1 on Gas chrom Q (Applied Science Laboratories, Div. of A. S. Laboratories, Inc., Waltham, MA) were used isothermally at 235°C.

Morphological examinations. The effectiveness of endothelium removal was checked by *en face* microscopic examination after silver staining (16). Aortic strips were immersed for 1-min periods in the dark at room temperature sequentially in 5% glucose, 0.25% AgNO₃, 5% glucose, 3% CoBr₂ + 1% NH₄Br, and 5% glucose (twice). Deendothelialization was also checked by scanning electron microscopy (17). Rabbit aorta strips were fixed in 4% glutaraldehyde in 0.1 M phosphate buffer, pH 7.4, at room temperature, washed in the same buffer, postfixed in 1% osmium tetroxide, dehydrated in acetone, processed through the critical point, and finally coated with gold. They were examined with an electron microscope (80 kV, model EM 400; Philips Electronic Instruments, Inc., Mahwah, NJ).

Statistical analysis. The statistical significance of observed differences was established using the analysis of variance for repeated measurements with two within factors (time period and treatment) and one grouping factor (animal). Computerizations were performed with the use of the P₂V program of the BMDP statistical software (18).

Materials. Arachidonic and docosahexaenoic acids were purchased from Sigma Chemical Co., St. Louis, MO. Ionophore A23187 was obtained from Boehringer Mannheim Diagnostics, Inc., Mannheim, FRG. BSA (fatty acid poor) was purchased from Calbiochem-Behring Corp., La Jolla, CA. Indomethacin, ibuprofen, and BW-755C were gifts from Merck Sharp & Dohme Div., West Point, PA, Upjohn Co., Kalamazoo, MI, and The Wellcome Foundation, Ltd., The Wellcome Research Laboratories, Beckenham, Kent, England, respectively. 6-K-PGF_{1 α} and PGE₂ were purchased from Upjohn Co.; [³H]6-K-PGF_{1 α} , [³H]PGE₂, and [³H]arachidonic acid were from Amersham Corp., Amersham, England. The antiserum against PGE₂ was obtained from Institut Pasteur de Paris, France. PGH₂ synthase, purified to homogeneity from ram seminal vesicles, was purchased from Oxford Biomedical Research, Oxford, MI. Acetonitrile, pentafluorobenzylbromide, and diisopropylethylamine were purchased respectively from Burdick & Jackson Laboratories, Inc., American Hospital Supply Corp., Muskegon, MI, Pierce Chemical Co., Rockford, IL, and Sigma Chemical Co.

Results

The endothelium was completely removed from the rabbit aorta strips after their luminal surfaces were rubbed against filter paper or scraped with a scalpel. The *en face* examination of the complete surfaces of numerous scraped strips, after silver staining, revealed the homogenous loss of the mosaic pattern characteristic of the endothelium (16). Scanning electron microscopy provided similar results. Endothelium removal induced a transient increase of 6-K-PGF_{1 α} release from the rabbit aorta strips (Fig. 1). The amounts of 6-K-PGF_{1 α} (nanograms per 100 milligrams of tissue) released by the strips during the first 30-min incubation period after their preparation were for the intact, rubbed, and scraped strips, respectively, 67 \pm 8 (n = 23), 248 \pm 26 (n = 14), and 219 \pm 19 (n = 23) (mean \pm SE; n , number of experiments). Rubbing and scraping produced an equal stimulation of 6-K-PGF_{1 α} output of 408% of the respective controls in the case of rubbing and 367% in the case of scraping. To confirm that the product formed after endothelium removal was authentic 6-K-PGF_{1 α} , RIA was repeated after RPHPLC purification of several samples. All the immunoreactivity was indeed recovered in the fractions where a standard of 6-K-PGF_{1 α} was eluted (data not shown). Scraping the endothelium of the dog aorta also stimulated the release of 6-K-PGF_{1 α} (270% of the control, mean of three experiments). The increased production of 6-K-PGF_{1 α} in the deendothelialized aorta progressively declined with time, both when the strips were incubated continuously in the same medium (Fig. 1) and when the medium was changed every 30 min (Fig. 4). Indomethacin abolished the release of 6-K-PGF_{1 α} induced by deendothelialization (Fig. 2). Endothelium removal also increased the release of PGE₂ by rabbit aorta strips but in smaller amounts (32 ng/100 mg during the first 30 min, or nine times less than the corresponding amounts of 6-K-PGF_{1 α} ; mean of four experiments). PGE₂ output decreased with time parallel to 6-K-PGF_{1 α} (data not shown).




Figure 1. Kinetics of 6-K-PGF_{1 α} release from intact and deendothelialized rabbit aorta strips. Rings of rabbit aorta were incubated for three 30-min periods, after which all the rings were opened by a longitudinal incision. The intimal surface of half the resulting strips was scraped with a scalpel. Intact (●) and scraped (○) strips were then incubated for 150 min. Aliquots of incubation medium were collected at various times. The results represent the accumulation of 6-K-PGF_{1 α} in the incubation medium (mean \pm SD of quadruplicate determinations in two separate experiments).




Figure 2. Inhibition by indomethacin of 6-K-PGF_{1α} release from the deendothelialized rabbit aorta. The aortic strips were incubated for six 30-min periods with (■) or without (□) indomethacin (10 µg/ml). At the end of each period, the medium was collected and replaced. At the end of the third period, the intimal surface of all the strips was scraped with a scalpel before the incubation was resumed. Results represent the mean±SD of six determinations in two separate experiments.

Previous experiments had shown that rubbing the luminal surface abolished the stimulatory effects of acetylcholine (19) and ADP or ATP (20) on PGI₂ production by the rabbit aorta. As shown on Figs. 3 and 4, the responses to ionophore A23187 and arachidonic acid were also lost in strips of deendothelialized aorta obtained either by rubbing or by scraping. The 30-fold stimulation of 6-K-PGF_{1α} output by ionophore A23187 (5 µM) in the intact rabbit aorta was reduced to 1.3-fold in deendothelialized strips (mean of eight experiments). Continuous exposure of intact strips to arachidonic acid produced an increase of 6-K-PGF_{1α} output, which was maximal at the beginning and progressively declined. The deendothelialized aorta was unresponsive to arachidonic acid from the start to the end of the incubation (Fig. 4). Intact strips of rabbit aorta converted exogenous [³H]arachidonic acid mainly into 6-K-PGF_{1α}, and to a smaller extent into PGE₂, PGF_{2α}, and a compound coeluting with 12-hydroxyeicosatetraenoic acid (Fig. 5). Deendothelialized strips of aorta failed to metabolize exogenous arachidonic acid, although, as expected, they released larger amounts of endogenous 6-K-PGF_{1α} than the intact ones (Fig. 5). When only half of the luminal surface of the aorta strips was scraped, the initial stimulation of 6-K-PGF_{1α} release was equal to that in completely denuded strips, whereas the response to ionophore A23187 was partially maintained, roughly in proportion to the amount of residual endothelium (38% of the response in intact strips, mean of three experiments) (data not shown). The template procedure (10) was used to




Figure 3. Unresponsiveness of 6-K-PGF_{1α} production from the deendothelialized rabbit aorta to ionophore A23187 and arachidonic acid. Strips of rabbit aorta, intact or rubbed against filter paper, were incubated for five 30-min periods. Ionophore A23187 (5 µM; ▨) and arachidonic acid (1 µg/ml; ■) were added at the beginning of the last period. Results express the release of 6-K-PGF_{1α} during this period (mean±SD of triplicate determinations in one representative experiment). (□), Control.

determine the effect of endothelium removal specifically on the release of PGI₂ at the luminal surface of the aorta. As shown on Fig. 6, deendothelialization increased the luminal release of 6-K-PGF_{1α} and abolished the responsiveness to ionophore A23187.

Endothelium removal from the rabbit aorta induced the mobilization of free arachidonic acid, measured by its release into the albumin-containing incubation medium (Fig. 7). Arachidonate release (nanomoles per 100 mg and per 30 min) was maximal during the first incubation period: 6.1±2.1 in scraped aorta strips compared with 1.7±0.5 in the intact ones (mean±SD; five experiments). During the next 90 min, arachidonic acid was mobilized at a lower rate, which was 3.5-fold higher in the scraped strips than in the intact ones (*P* = 0.04). This rate of release did not decrease significantly during this period (*P* = 0.12). Ionophore A23187 increased the release of free arachidonic acid in the intact aorta. This effect was lost after endothelium removal (Fig. 7).

Further experiments were performed to explain the discrepancy between the progressive decline of 6-K-PGF_{1α} output (Figs. 1 and 4) and the continuous release of arachidonic acid (Fig. 7), observed between 30 and 120 min in the deendothelialized aorta. Addition of pure PGH₂ synthase, in the absence of exogenous arachidonic acid, increased dose dependently the




Figure 4. Comparative kinetics of 6-K-PGF_{1α} release by intact and deendothelialized rabbit aorta, continuously exposed to arachidonic acid. Strips of rabbit aorta, intact or rubbed against filter paper, were incubated for five 30-min periods. The medium was collected and replaced at the end of each period. Results are expressed as the mean of duplicate determinations in one representative experiment out of three. (●), Control; (○), arachidonic acid (5 μg/ml) added at the beginning of each period.

release of 6-K-PGF_{1α} from scraped aorta strips but had almost no effect on the intact aorta (Fig. 8). Purification of samples by RPHPLC before RIA demonstrated that the immunoreactivity measured was due to authentic 6-K-PGF_{1α} and not to the cross-reaction of PGE₂ or other products of the nonenzymatic breakdown of the endoperoxides generated by PGH₂ synthase (data not shown). A transient inhibition of cycloo-




Figure 6. Kinetics of 6-K-PGF_{1α} release at the luminal surface of intact (●) and deendothelialized (○) rabbit aorta strips. The luminal surface of aortic strips, intact or deendothelialized by scraping with a scalpel, was exposed at the bottom of incubation chambers (5-mm diam holes in a plexiglass plate) containing 0.3 ml medium. The incubation was performed at 37°C for six 30-min periods. The medium was collected and replaced at the end of each period. Ionophore A23187 was present during the last period. Results represent the mean±SD of six determinations, in two separate experiments.

xygenase by ibuprofen resulted in a rebound of 6-K-PGF_{1α} output after ibuprofen withdrawal (Fig. 9): the 90% inhibition during the 90 min in the presence of ibuprofen was followed




Figure 5. RPHPLC separation of the products formed from [³H]arachidonic acid by the intact and deendothelialized aorta. Strips of rabbit aorta, intact or rubbed against filter paper, were incubated for four 30-min periods. The medium was collected and replaced at the end of each period. [³H]Arachidonic acid (2 μCi, 0.5 μg) was added at the beginning of the fourth period. 6-K-PGF_{1α} was measured by RIA in the media from the initial incubations (*insets*), whereas the medium collected at the end of the last period was analyzed by RPHPLC, as described in Methods. The arrows indicate the retention times of various standards. The figure shows representative results, which have been obtained in duplicate in two separate experiments. 12-HETE, 12-hydroxyicosatetraenoic acid.




Figure 7. Kinetics of arachidonic acid release by the deendothelialized aorta. Strips of rabbit aorta, intact (●) or scraped (○) with a scalpel, were incubated for five 30-min periods in a medium containing BSA (1 mg/ml) and indomethacin (1 μg/ml). The medium was collected and replaced at the end of each period. Arachidonic acid was measured by GLC-ECD, as described in Methods. Results represent the mean±SD of six determinations in three separate experiments.

by a threefold enhancement during the next 120 min ($P = 0.004$). A similar rebound was observed when scraped aorta strips were incubated for 90 min at 0°C and then for 60 min at 37°C: a 61% inhibition followed by a threefold increase, by reference to scraped strips continuously incubated at 37°C (data not shown). BW-755C has a dual effect on prostaglandin synthase in acellular systems, stimulation at low concentrations, and inhibition at high concentrations. The stimulatory effect has been explained by the ability of BW-755C to scavenge radicals that are generated by the hydroperoxidase component




Figure 8. Stimulation of 6-K-PGF_{1α} release from the deendothelialized rabbit aorta by exogenous PGH₂ synthase. Aortic strips, intact (●) or deendothelialized (○) by scraping with a scalpel, were incubated for five 30-min periods. The incubation medium was replaced at the end of each period. At the start of the last period, PGH₂ synthase (12.5 or 37.5 U/ml) and arachidonic acid (20:4, 2 μg/ml) were added as indicated. In the experiments depicted in B, all the strips were deendothelialized, and the concentration of PGH₂ synthase was 12.5 U/ml. Negligible amounts of material reactive with the anti-6-K-PGF_{1α} antibody were generated during an incubation of PGH₂ synthase with arachidonic acid, in the absence of tissue. Results represent the mean±SD of six determinations in three separate experiments (A) and six determinations in two separate experiments (B).




Figure 9. Kinetics of 6-K-PGF_{1α} release by deendothelialized rabbit aorta strips: effect of a transient exposure to ibuprofen. Immediately after its removal, the aorta was divided into two equal portions, one of which was immersed and manipulated in a medium containing ibuprofen (250 μM). After scraping the luminal surface with a scalpel, the strips were incubated for seven 30-min periods. The incubation medium was collected and replaced at the end of each period. Results are the mean±SD of quadruplicate determinations in two separate experiments. (●), Control, without ibuprofen; (○), ibuprofen (250 μM) present during the manipulation of the tissue and during the three initial periods of incubation. The inset represents the cumulated release of 6-K-PGF_{1α} during the three initial and four final periods of incubation, respectively. (□), Control; (▨), ibuprofen (250 μM) present during the three initial periods.

of prostaglandin synthase and that inactivate the cyclooxygenase (21). BW-755C prolonged the release of 6-K-PGF_{1α} by the deendothelialized rabbit aorta (Fig. 10). 3 μM of BW-755C caused a negligible inhibition of 6-K-PGF_{1α} output during its 60-min exposure to the scraped aorta strips, followed by a doubling of 6-K-PGF_{1α} release during the 90 min after the removal of the drug. 10 μM of BW-755C produced a 34% inhibition followed by a 2.7-fold stimulation. This stimulatory effect of BW-755C was highly significant ($P = 0.002$). Other antioxidants reported to protect the synthesis of PGI₂, ascorbic acid (22), and nafazatrom (23) did not modify the kinetics of 6-K-PGF_{1α} release from the deendothelialized aorta (data not shown).

Discussion

Our results indicate that the removal of the aortic endothelium induced a sustained mobilization of free arachidonic acid, a transient stimulation of PGI₂ synthesis, and an increased release of PGI₂ at the luminal surface, accompanied and followed by a complete unresponsiveness to all the stimuli that increase PGI₂ in the intact aorta, including arachidonic acid itself. Since we checked carefully to see that the endothelium was completely removed by the mechanical procedures used, it is most likely that the events measured in this study occurred in the smooth muscle cells. Increased PGI₂ synthesis




Figure 10. Effect of BW-755C on the release of 6-K-PGF_{1α} by the deendothelialized rabbit aorta. Strips of rabbit aorta, deendothelialized by scraping with a scalpel, were incubated for five 30-min periods. The medium was collected and replaced at the end of each period. Results are the mean±SD of eight determinations, in four separate experiments. (●), Control; (○), BW-755C (3 μM) present during the two initial periods; (×), BW-755C (10 μM) present during the two initial periods. The inset represents the cumulated release of 6-K-PGF_{1α} during the two initial and three final periods of incubation, respectively. (□), Control; (▨), 3 μM BW-755C; (■), 10 μM BW-755C.

after endothelium removal is likely to be the consequence of the increased availability of free arachidonic acid. This spontaneous mobilization of endogenous arachidonic acid could also explain the lack of stimulatory effect of exogenous arachidonic acid on PGI₂ production (saturation of the cyclooxygenase) and the inability of ionophore A23187 to release additional free arachidonic acid. The progressive decline of PGI₂ release, despite the continuous availability of free arachidonic acid, is likely to be due to cyclooxygenase inactivation, a contention supported by three arguments. The output of PGE₂ was initially enhanced and decreased progressively, parallel to PGI₂, but at a much lower level, thus excluding PGI₂ synthase inactivation as the (only) explanation of the data. The stimulatory effect of exogenous PGH₂ synthase indicates that, in the deendothelialized aorta, the limiting factor for PGI₂ synthesis was indeed the cyclooxygenase activity, and confirms the availability of free arachidonic acid in this situation. Finally, the prolongation of PGI₂ synthesis by a transient exposure to the reversible cyclooxygenase inhibitor ibuprofen or the radical scavenger BW-755C suggests that inactivation of cyclooxygenase might result from a suicidal oxidation by prostaglandin endoperoxides.

Whether smooth muscle cells contribute significantly to the production of PGI₂ by arteries remains a controversial issue. Studies performed with cultured aortic smooth muscle

cells have not provided conclusive answers. PGI₂ is the major eicosanoid produced by cells from rat, rabbit, and bovine aorta (1–6), whereas PGE₂ and PGF_{2α} are the main products formed in cells cultured from porcine aorta (7–9). Aortic smooth muscle cells dedifferentiate in culture and lose some of their native properties, such as contractility (24). Our observation that, after complete removal of the endothelium, the aorta was able to produce large amounts of PGI₂, but less than the intact aorta stimulated by arachidonic acid and only for a short period (Fig. 4), is perfectly consistent with the recent study of De Witt et al. (13) who have shown that the aortic smooth muscle cells contain as much PGI₂ synthase as, but 20 times less cyclooxygenase than the aortic endothelial cells. Our data are also consistent with the observation by Hornstra et al. (25) that disruption of the endothelium results in the increased release of a heat-labile inhibitor of platelet aggregation by the rat and the rabbit aorta, and with the results of Goldsmith (26) who showed that the release of PGI₂ at the luminal surface of the bovine aorta was initially increased after *in vitro* removal of the endothelium. On the contrary, De Mey et al. (11) have concluded, from the inability of deendothelialized strips of canine femoral arteries to convert exogenous [¹⁴C]arachidonic acid into 6-K-PGF_{1α}, that the endothelium is the key site of PGI₂ synthesis in the arterial wall. We have made the same observation in the deendothelialized rabbit aorta (Fig. 5), but the simultaneous measurement of the endogenous production of 6-K-PGF_{1α} leads us to an opposite conclusion. The deendothelialized aorta released more PGI₂ than the intact one, and the lack of conversion of exogenous radiolabeled arachidonic acid may be the consequence of isotopic dilution by the large amounts of endogenous free arachidonic acid mobilized after endothelium removal. This underscores the danger of using the conversion of exogenous arachidonic acid as the only method to evaluate eicosanoid biosynthetic capacity. This is true also for the work of Eldor et al. (10) who have shown that removal of the endothelium from the rabbit aorta *in vivo* abolishes the conversion of exogenous arachidonic acid into PGI₂ measured at the luminal surface. These authors did not report an increase of the spontaneous release of PGI₂ after endothelium removal. Since the stimulation observed in our study was transient, it might have been missed in the time interval between the deendothelialization performed *in vivo* and the start of the *in vitro* incubation.

The short-lived release of PGI₂, in spite of a continuous mobilization of endogenous free arachidonic acid, that we have observed in the deendothelialized rabbit aorta is similar to the transient production of PGI₂ by the intact rabbit aorta continuously perfused with exogenous arachidonic acid (27) and to the observation that cultured endothelial cells respond to arachidonic acid only once (6, 28). In each case, the rapid decline of PGI₂ production is likely to be due to self-inactivation of the cyclooxygenase, involving the irreversible oxidation of the enzyme by a free radical generated during the reduction of prostaglandin G₂ into prostaglandin H₂ (21, 27, 28). Our results suggest that BW-755C, which, depending on the concentration, stimulates or inhibits cyclooxygenase in a cell-free preparation (21), constitutes the model of a specific drug protecting against this inactivation in vascular smooth muscle

cells. Indeed a concentration of BW-755C as low as 3 μ M prolongs the release of PGI₂ by the deendothelialized aorta.

Mechanical removal of the endothelium seems to represent a straightforward method to evaluate its contribution to phenomena occurring in arteries. In particular, this procedure has been used to demonstrate that acetylcholine and other agents relax the rabbit aorta and other arteries by acting on endothelial cells (29, 30). Our study indicates that experiments involving endothelium removal should be interpreted with great caution, since this procedure dramatically alters the metabolism of arachidonic acid in the subendothelial layer. Other acute biochemical modifications after endothelium removal have been noticed in the aorta, including increased cyclic GMP content (31) and phosphorylation of one specific protein (32).

One question raised by our observations is whether the mobilization of arachidonic acid and the resulting PGI₂ synthesis after endothelium removal is a mere consequence of the trauma imposed to the aortic smooth muscle or is due to the disruption of endothelial-smooth muscle cell interactions. Control and deendothelialized samples were submitted to the same trauma when the aorta was cut into rings. This trauma, followed by recovery, might be responsible for the initial and transient phase of free arachidonic acid release. It could be speculated that the sustained release observed in the deendothelialized aorta results from the permanent loss of some endothelium-smooth muscle interaction, of which several examples have been described (33–36), but additional studies are required to support this hypothesis. Whatever the mechanism involved, our study suggests that the local release of PGI₂ induced by endothelial lesions might contribute to the increased systemic production of PGI₂ observed in patients with atherosclerosis (37).

Acknowledgments

We thank Dr. R. Pochet for his help in the microscopic examinations, Dr. V. de Maertelaer for performing the statistical analysis, Mr. D. Martins for designing the template device, and Mrs. D. Leemans for typing the manuscript. We are grateful to Dr. J. E. Dumont for his continuous interest and support.

This work was performed under contract of the Ministère de la Politique Scientifique (Action Concertée).

References

- Whiting, J., K. Salata, and J. M. Bailey. 1980. Aspirin: an unexpected side effect on prostacyclin synthesis in cultured vascular smooth muscle cells. *Science (Wash. DC)*. 210:663–665.
- Larrue, J., M. Rigaud, D. Daret, J. Demond, J. Durand, and H. Bricaud. 1980. Prostacyclin production by cultured smooth muscle cells from atherosclerotic rabbit aorta. *Nature (Lond.)*. 285:480–483.
- Larrue, J., B. Dorian, J. Demond-Henri, and H. Bricaud. 1981. Endogenous arachidonic acid metabolism by cultured arterial smooth muscle cells. *Biochem. Biophys. Res. Commun.* 101:861–867.
- Ingerman-Wojenski, C., M. J. Silver, J. B. Smith, and E. Macarak. 1981. Bovine endothelial cells in culture produce thromboxane as well as prostacyclin. *J. Clin. Invest.* 67:1292–1296.
- Coughlin, S. R., M. A. Moskowitz, H. N. Antoniades, and L. Levine. 1981. Serotonin receptor-mediated stimulation of bovine smooth muscle cell prostacyclin synthesis and its modulation by platelet-derived growth factor. *Proc. Natl. Acad. Sci. USA*. 78:7134–7138.
- Dejana, E., G. Balconi, C. De Castellarnau, B. Barbieri, M. Vergara-Dauden, and G. De Gaetano. 1983. Prostacyclin production by human endothelial and bovine smooth muscle cells in culture. Effect of repeated stimulation with arachidonic acid, thrombin and ionophore A23187. *Biochim. Biophys. Acta*. 750:261–267.
- MacIntyre, D. E., J. D. Pearson, and J. L. Gordon. 1978. Localisation and stimulation of prostacyclin production in vascular wall. *Nature (Lond.)*. 271:549–551.
- Siess, W., F. Dray, C. C. Seillan, C. Ody, and F. Russo-Marie. 1981. Prostanoid synthesis by vascular slices and cultured vascular cells of piglet aorta. *Biochem. Biophys. Res. Commun.* 99:608–616.
- Ody, C., C. Seillan, and F. Russo-Marie. 1982. 6-K-PGF_{1 α} , PGE₂, PGF₂ and TxB₂ production by endothelial cells, smooth muscle cells and fibroblasts cultured from piglet aorta. *Biochem. Biophys. Acta*. 712:103–110.
- Eldor, A., D. J. Falcone, D. P. Hajjar, C. R. Minick, and B. B. Weksler. 1981. Recovery of prostacyclin production by de-endothelialized rabbit aorta. *J. Clin. Invest.* 67:735–741.
- De Mey, J. G., M. Claeys, and P. M. Van Houtte. 1982. Endothelium-dependent inhibitory effects of acetylcholine, ATP, thrombin and arachidonic acid in the canine femoral artery. *J. Pharmacol. Exp. Ther.* 222:166–173.
- Smith, W. L., D. L. De Witt, and M. L. Allen. 1983. Bimodal distribution of the PGI₂ synthase antigen in smooth muscle cells. *J. Biol. Chem.* 258:5922–5926.
- De Witt, D. L., J. S. Day, W. K. Sonnenburg, and W. L. Smith. 1983. Concentrations of prostaglandin endoperoxide synthase and prostaglandin I₂ synthase in the endothelium and smooth muscle of bovine aorta. *J. Clin. Invest.* 72:1882–1888.
- Maclof, J. 1981. A radioimmunoassay for 6-keto-PGF_{1 α} . *Methods Enzymol.* 86:273–286.
- Wickramasinghe, J. A. F., W. Morozowich, W. E. Hamlin, and S. R. Shaw. 1973. Detection of PGF_{2 α} as pentafluorobenzyl ester by electron-capture GLC. *J. Pharm. Sci.* 62:1428–1431.
- Poole, J. C. F., A. G. Sanders, H. W. Florey. 1958. The regeneration of aortic endothelium. *J. Pathol. Bacteriol.* 75:133–142.
- Morrison, A. D., L. Orci, L. Berwick, A. Perrelet, and A. I. Winegrad. 1977. The effects of anoxia on the morphology and composite metabolism of the intact aortic intima-media preparation. *J. Clin. Invest.* 59:1027–1037.
- Dixon, W. J., M. B. Brown, L. Engleman, J. W. Frane, M. A. Hill, R. I. Jennrich, and J. D. Toporek. 1981. BMDP Statistical Software. University of California Press, Berkeley, CA.
- Boeynaems, J. M., and N. Galand. 1983. Cholinergic stimulation of vascular prostacyclin synthesis. *Prostaglandins*. 26:531–545.
- Boeynaems, J. M., and N. Galand. 1983. Stimulation of vascular prostacyclin synthesis by extracellular ADP and ATP. *Biochem. Biophys. Res. Commun.* 112:290–296.
- Marnett, L. J., P. H. Siedlik, and L. W. M. Fung. 1982. Oxidation of phenidone and BW-755C by prostaglandin endoperoxide synthetase. *J. Biol. Chem.* 257:6957–6964.
- Beetens, J. R., M. Claeys, and A. G. Herman. 1981. Antioxidants increase the formation of 6-oxo-PGF_{1 α} by ram seminal vesicle microsomes. *Biochem. Pharmacol.* 30:2811–2815.
- Sevilla, M. D., P. Neta, and L. J. Marnett. 1983. Reaction of the anti-thrombotic and antimetastatic agent nafazatrom with oxidizing radicals. *Biochem. Biophys. Res. Commun.* 115:800–806.
- Chamley, J. H., G. R. Campbell, J. D. McConnell, and U. Groschel-Stewart. 1977. Comparison of vascular smooth muscle cells from adult human monkey and rabbit in primary culture and subculture. *Cell Tissue Res.* 177:503–522.
- Hornstra, G., E. Haddeman, and J. A. Don. 1978. Some investigations in the role of prostacyclin in thromboregulation. *Thromb. Res.* 12:367–374.

26. Goldsmith, J. C. 1982. Contribution of the subendothelium to prostacyclin release after vascular injury. *J. Lab. Clin. Med.* 100:574-584.
27. Kent, R. S., S. L. Diedrich, and A. R. Whorton. 1983. Regulation of vascular prostaglandin synthesis by metabolites of arachidonic acid in perfused rabbit aorta. *J. Clin. Invest.* 72:455-465.
28. Brotherton, A. F. A., and J. C. Hoak. 1983. Prostacyclin biosynthesis in cultured vascular endothelium is limited by deactivation of cyclooxygenase. *J. Clin. Invest.* 72:1255-1261.
29. Furchgott, R. F., and J. V. Zawadzki. 1980. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature (Lond.)*. 288:373-376.
30. Chand, N., and B. M. Altura. 1981. Acetylcholine and bradykinin relax intrapulmonary arteries by acting on endothelial cells: role in lung vascular diseases. *Science (Wash. DC)*. 213:1376-1379.
31. Rapoport, R. M., and F. Murad. 1983. Agonist-induced endothelium-dependent relaxation in rat thoracic aorta may be mediated through cGMP. *Circ. Res.* 52:352-357.
32. Rapoport, R. M., M. B. Draznin, and F. Murad. 1983. Endothelium-dependent relaxation in rat aorta may be mediated through cGMP-dependent protein phosphorylation. *Nature (Lond.)*. 306:174-176.
33. Stet, E. M., G. Mayno, and I. Joris. 1979. Cellular pathology of the rat aorta. Pseudo-vacuoles and myo-endothelial herniae. *Virchows Arch. A Pathol. Anat. Histol.* 383:135-148.
34. Castellot, J. J., M. L. Addonizio, R. Rosenberg, and M. J. Karnovsky. 1981. Cultured endothelial cells produce a heparin-like inhibitor of smooth muscle cell growth. *J. Cell. Biol.* 90:372-379.
35. Hajjar, D. P., D. J. Falcone, S. Fowler, and C. R. Minick. 1981. Endothelium modifies the altered metabolism of the injured aortic wall. *Am. J. Pathol.* 102:28-39.
36. Wight, T. N., K. D. Curwen, M. M. Litrenta, D. R. Alonso, and C. R. Minick. 1983. Effect of endothelium on glycosaminoglycan accumulation in injured rabbit aorta. *Am. J. Pathol.* 113:156-164.
37. Fitzgerald, G. A., B. Smith, A. K. Pedersen, and A. R. Brash. 1984. Increased prostacyclin biosynthesis in patients with severe atherosclerosis and platelet activation. *N. Engl. J. Med.* 310:1065-1068.