

## Human Recombinant Interleukin 1 Stimulates Collagenase and Prostaglandin E<sub>2</sub> Production by Human Synovial Cells

Jean-Michel Dayer, Beatrice de Rochemonteix, Beatrice Burrus, Stephen Demczuk, and Charles A. Dinarello

Division of Immunology and Allergy, Hans Wilsdorf Laboratory, Department of Medicine, University Hospital Cantonal, 1211 Geneva 4, Switzerland; and Division of Experimental Medicine, Department of Medicine, Tufts University School of Medicine, Boston, Massachusetts 02111

### Abstract

The pathogenesis and progression of rheumatoid arthritis involves the production of biologically active lymphokines and monokines. Of these, interleukin 1 (IL-1) has been somewhat of a controversial molecule because it seems to evoke various biological responses in several different tissues. In these studies we demonstrate that three biological properties of human monocyte-derived IL-1 (T-lymphocyte activation and human synovial cell prostaglandin E<sub>2</sub> and collagenase production) co-purify. The complementary DNA for the prominent pI 7 form of human IL-1 was expressed, purified, and tested. Any controversy now appears resolved since homogeneous recombinant human IL-1 stimulates prostaglandin E<sub>2</sub> and collagenase from human synovial cells as well as activates T cells in vitro.

### Introduction

Mononuclear cell factor (MCF)<sup>1</sup> is a human blood monocyte-derived cytokine that stimulates adherent synovial cells, dermal fibroblasts, chondrocytes, and bone-derived cells to produce increasing amounts of collagenase and prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) (1-3). Because of these activities, MCF plays a major role in the pathogenic progression of various arthritides, particularly rheumatoid arthritis (4). In addition to its inflammatory and tissue-destructive activities, MCF shares biochemical and biological properties with lymphocyte-activating factor (LAF) (5), a polypeptide that augments lymphocyte responses to mitogens and antigens. MCF also possesses properties similar to endogenous pyrogen (EP), which produces fever by elevating hypothalamic PGE<sub>2</sub> (6). The current hypothesis is that MCF, LAF, and EP represent the biological activities of either a single peptide or

related but distinct proteins grouped together under the name IL-1 (7). IL-1 production is often associated with disease processes. Thus messenger RNA (mRNA) from endotoxin-stimulated human blood monocytes directs oocyte translation of MCF and LAF activity, whereas mRNA from unstimulated blood monocytes lack this property (8-10). We report here that MCF activity coelutes with LAF activity during gel-filtration and elutes from an affinity column of immobilized anti-human EP. Furthermore, homogeneous recombinant human IL-1 stimulates human synovial cells to produce collagenase and PGE<sub>2</sub>. Thus, these studies settle any previous dispute since these multiple biological properties do reside in a single polypeptide.

### Methods

**Human MCF bioassay.** Target cells used for the measurement of MCF activity were obtained from freshly dispersed synovium of rheumatoid arthritis patients and from human infant foreskin tissue. Details concerning the culture conditions of these cells have been described previously (8, 9). Various test specimens were diluted in Dulbecco's modified Eagle's medium (DME; Gibco, Grand Island, NY) supplemented with 10% fetal calf serum (FCS) (Gibco) and 5 µg/ml of polymyxin B (Sigma Chemical Co., St. Louis, MO) and then added to synovial cells or dermal fibroblasts (8). After 72 h of incubation, media were removed and assayed for collagenase and PGE<sub>2</sub>. Supernatants were treated with trypsin (TRTPCK, Worthington Biochemical Corp., Freehold, NJ), followed by soybean trypsin inhibitor (Worthington Biochemical Corp.). Collagenase activity was measured by incubation with <sup>14</sup>C-acetylated salt-soluble rat tail tendon collagen (8). PGE<sub>2</sub> level was measured by a double-antibody radioimmunoassay using an antibody provided by L. Levine (Brandeis University, Waltham, MA) (8). Standard PGE<sub>2</sub> was obtained from Upjohn Corp., Kalamazoo, MI.

**Murine LAF assay.** Samples were assayed for LAF using the incorporation of [<sup>3</sup>H]thymidine into thymocytes of 6-8-wk-old C3H/HeJ mice (11) or the murine clonal T cell line, D10.G4.1 (12) stimulated with phytohemagglutinin (PHA, Burroughs Wellcome Co., Triangle Park, NC), 1 µg/ml and in the presence of 5 µg/ml polymyxin B.

**MCF preparation.** Human peripheral blood mononuclear cells from freshly isolated buffy coats were separated on Ficoll-Hypaque gradients. After 4 h of incubation at 4 × 10<sup>6</sup> cells/ml in RPMI 1640 supplemented with 2 mM glutamine, 25 mM Hepes, 5% heat-inactivated FCS, 100 µg/ml penicillin, and 100 µg/ml streptomycin, floating cells were removed and adherent cells were washed three times and incubated for 48 h with 1 µg/ml PHA and 10 µg/ml concanavalin A (Pharmacia Fine Chemicals, Piscataway, NJ). Supernatants were dialyzed for 48 h against water at 4°C and lyophilized. 50 mg of the lyophilized material was resuspended in 10 mM Tris HCl, pH 7.5, 165 mM NaCl, 5 mM CaCl<sub>2</sub>, applied to a 2.6 × 90 cm of Ultrogel AcA54 (LKB Produkter; Bromma, Sweden) and eluted with the same buffer at 4°C. Fractions containing MCF activity were pooled and lyophilized.

**Preparation of human monocyte interleukin-1 (IL-1).** Adherent

Address all correspondence to Dr. Dinarello, Tufts University School of Medicine, 136 Harrison Avenue, Boston, MA 02111.

Received for publication 11 September 1985.

1. *Abbreviations used in this paper:* DME, Dulbecco's modified Eagle's medium; EP, endogenous pyrogen; FCS, fetal calf serum; hrIL-1, human recombinant interleukin 1; IL-1, interleukin 1; LAF, lymphocyte-activating factor; MCF, mononuclear cell factor; PAGE, polyacrylamide gel electrophoresis; PG, prostaglandin; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>; PHA, phytohemagglutinin.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/86/02/0645/04 \$1.00

Volume 77, February 1986, 645-648

monocytes were stimulated with heat-killed *Staphylococcus albus* by previously published methods (13). The crude supernate was purified by elution from the anti-human EP affinity gel followed by gel filtration over Sephadex G-50 (170 × 5 cm). Fractions were assayed for EP and LAF activity. Active fractions in the 15–18-kD range were pooled and subjected to chromatofocusing. Using [<sup>35</sup>S]methionine labeling of monocytes, this three-step procedure yielded a homogenous 17.5-kD protein as judged by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and fluorography that possessed LAF and EP activities (13).

**Anti-EP immunoabsorbent.** Anti-EP IgG was purified over immunoabsorbents to remove unwanted antibodies (14). The antibody used in the present study was prepared with high specific activity human EP (10) and adsorbed before immobilization on Sepharose. 1 ml of phosphate-buffered saline/bovine serum albumin (PBS/BSA) (1 mg/ml BSA) containing 100 mg/ml of lyophilized MCF preparation obtained after AcA54 purification was added to 3 ml of the anti-EP affinity gel and rocked for 2 h at room temperature. The gel was poured into a 1-cm diameter column, washed with PBS, and the MCF was eluted with 3 M NaSCN. Fractions were dialyzed against PBS, lyophilized, and resuspended in DME 10% FCS for detecting MCF and LAF activities. Preimmune rabbit IgG bound to Sepharose served as a control column.

**Human recombinant IL-1 (hrIL-1).** The IL-1 cDNA was expressed by isolating the NcoI-XmnI fragment (bp 295–1,407) from the plasmid pCD 12-1-8 (10) and inserting it into an *Escherichia coli* expression vector. The hrIL-1 expressed is 223 amino acids long and contains 199 of the 269 amino acid long IL-1 precursor form. The alanine amino acid 117 is the NH<sub>2</sub>-terminus of the processed IL-1 which is found in supernates of stimulated human blood monocytes (15). The calculated molecular weight of the hrIL-1 is 24,556. hrIL-1 was extracted from the insoluble cellular fraction and purified by sequential ion-exchange and either gel-filtration chromatography or high-performance liquid chromatography. The hrIL-1 was stored at –70°C in physiological saline. The identity of the purified hrIL-1 was confirmed by amino acid composition, the sequence of the amino acid terminal, and by SDS-PAGE (Fig. 1). The endotoxin concentration of the homogeneous hrIL-1 was <20 pg/μg of IL-1 protein using the Limulus ameocyte lysate test (Associates of Cape Cod, Woods Hole, MA). *E. coli* endotoxin as high as 5 μg/ml has no effect in the MCF assay.

## Results

**Coelution of MCF and LAF activity.** Crude MCF elutes from the sizing chromatography column in two peaks; the major peak contains the three biological activities eluted between 14 and 25 kD (Fig. 2 A). Another peak of activity eluted between the oval-




Figure 1. SDS-PAGE (12%, reduced) of hrIL-1 stained with Coomassie blue. Lane 1, standards; lane 2, crude extract of IL-1-containing bacterial lysate; lane 3, lysate after DEAE cellulose chromatography; lane 4, hrIL-1 (lot 042285) after purification.

bumin and the BSA but possessed only collagenase-stimulating and LAF activity but no significant PGE<sub>2</sub>-stimulating activity. Active fractions (molecular weight range of 14,000–25,000) were pooled and applied to the anti-EP affinity column. As shown in Fig. 2 B, most of the activity was retained on the column and eluted with the addition of 3 M NaSCN. In contrast, when the same material was applied to a control affinity column, only small amounts of PGE<sub>2</sub> and collagenase-inducing activities were eluted from the column after addition of NaSCN, and these are likely due to nonspecific binding to the Sepharose.

**Effects of hrIL-1 on production of collagenase and PGE<sub>2</sub> by synovial cells.** As shown in Table I, both purified human monocyte-derived IL-1 and hrIL-1 markedly stimulated collagenase and PGE<sub>2</sub> production by human adherent synovial cells. Similar results were obtained using PGE<sub>2</sub> production from dermal fibroblasts (data not shown). The crude *E. coli* extract possessed significantly greater MCF activity when compared with identical dilutions of extract from the same recombinant *E. coli* strain without the IL-1 insert. Two lots of homogeneous hrIL-1 were active in all assays at subnanogram per milliliter concentrations.




Figure 2. (A) Ultrogel AcA54 chromatography of PHA/Con A-stimulated monocyte-macrophage supernatants with MCF and LAF activities as indicated. The thymocyte assay was used for LAF activity. Calibration markers are BSA (bovine serum albumin), ovalb. (ovalbumin), carb. anh. (carbonic anhydrase), SBTI (soybean trypsin inhibitor), lyso. (lysozyme). (B) LAF and MCF activity elution profiles from anti-EP affinity column (see Methods). Control column used immobilized preimmune serum.

Table I. Effect of Human Recombinant IL-1 on Collagenase and PGE<sub>2</sub> Production by Human Synovial Cells

Preparations	Concentrations*	Human adherent synovial cells		T cells‡ (index of stimulation)
		PGE <sub>2</sub> ng/ml	Collagenase U/ml	
Monocyte-derived IL-1	1:10	667±30	0.22	10.0
	1:100	603±11	0.18	13.5
	1:1,000	187±3	0.08	8.9
	1:10,000	43±10	<0.01	4.7
<i>E. coli</i> extract (with IL-1 insert)	1:1,000	324±17	1.11	30.6
	1:10,000	101±2	0.99	15.3
	1:100,000	28±5	0.40	3.8
<i>E. coli</i> extract (without IL-1 insert)	1:1,000	118±13	0.86	8.3
	1:10,000	9±2	0.19	1.7
	1:100,000	2±1	0.05	1.2
Homogeneous hrIL-1 (lot 69)	500 ng/ml	664±63	0.60	9.2
	50 ng/ml	268±55	0.24	6.0
	5 ng/ml	48±11	0.20	4.0
	0.5 ng/ml	41±8	0.09	3.0
Homogeneous hrIL-1 (lot 042285)	1 µg/ml	628±42	0.24	16.2
	100 ng/ml	352±37	0.19	8.0
	10 ng/ml	330±33	0.16	4.7
	1 ng/ml	223±14	0.15	2.4
Medium alone		2±1	<0.01	1

\* The purified monocyte-derived IL-1 is indicated as dilution of rabbit pyrogenic doses where one dose equals peak fever of 0.6–0.9°C (13). The *E. coli* extracts were diluted after the extraction procedure and the undiluted hrIL-1 protein was estimated on SDS-PAGE to be ~50 µg/ml; the concentration in the assays of the two homogeneous hrIL-1 lots are indicated. ‡ Stimulation index was calculated by setting the PHA response at 1 (11). D10.G4.1 cells were used.

## Discussion

To support the hypothesis that MCF and LAF activities are likely due to the same molecule, two methods have been frequently employed: first, the coelution of biological activities during protein purification, and second, the use of apparently homogeneous preparations that possess these biological properties. We initiated our experiments by showing that MCF and LAF activities are found in the same fractions after either gel filtration or antibody affinity chromatography. EP and LAF activities have been followed during protein purifications; apparently homogeneous preparations show a single band on SDS-PAGE (13, 16), and this material also possesses MCF activities. Although both methods have drawbacks, our data support previous observations that MCF and LAF activities are not easily separable. It always seemed likely that MCF and EP were related since MCF production of PGE<sub>2</sub> from synovial cells was biologically similar to EP stimulation of hypothalamic PGE<sub>2</sub> leading to fever (6).

During the course of these studies, human IL-1, as measured by the LAF assay, was cloned (10) and the cDNA expressed in recombinant *E. coli*. As soon as the crude extracts became available, we began testing for MCF activity to help resolve the issue that MCF and LAF reside in the same molecule. Our results now establish that homogeneous hrIL-1 possesses MCF and LAF activities. The ability of a single substance to manifest inflam-

matory, catabolic, and immunoregulatory activities clarifies an important aspect of the multiple biological functions attributed to IL-1. The hrIL-1 used in the present studies stimulates significant PGE<sub>2</sub> and collagenase activity at <1 ng/ml. To reduce species differences which may influence specific activity, we employed human target cells for MCF activity. The hrIL-1 used in these studies corresponds to the pI 7 form which is the predominant IL-1 isolated from human pathological fluids. It is distinct from the recombinant murine IL-1 which possesses a pI of 5 (17). Although these two forms exhibit <26% amino acid homology, they are related, particularly at the carboxyl terminus. However, the specific activity of the human pI 5 form remains to be established in the human MCF assay.

The clinical and pharmacologic implications of these studies may help explain the chronicity of some arthritides, especially rheumatoid arthritis, since IL-1 results in destruction of connective tissue but is also a source of immunologic stimulation for continued production of antibodies and lymphokines (1–4, 7). Thus therapies designed to alleviate or halt the progression of the disease would best be designed at the level of reduced IL-1 production.

## Acknowledgments

We are grateful to Scott D. Putney of Repligen Corp., Cambridge, MA; Gail LoPreste and Sheldon M. Wolff of Tufts University School of Med-

icine and the New England Medical Center, Boston, MA; Walter O. Fredericks of Cistron Technology, Inc.; and M.-C. Seydoux for editorial assistance.

This work was supported in part by grant no. 3.449.0.83 (to Dr. Dayer) from the Swiss National Science Foundation, by a grant from the Fondation Centre de Recherches Medicales Carlos et Elsie de Reuter (to Dr. Dayer), by National Institutes of Health grant AI 15614 (to Dr. Dinarello), and by Cistron Technology Inc., Pine Brook, NJ.

## References

1. Dayer, J.-M., J. Breard, L. Chess, and S. M. Krane. 1979. Participation of monocytes-macrophages and lymphocytes in the production of a factor that stimulates collagenase and prostaglandin release by rheumatoid synovial cells. *J. Clin. Invest.* 64:1386-1392.
2. Dayer, J.-M., M. L. Stephenson, E. Schmidt, W. Karge, and S. M. Krane. 1981. Purification of a factor from human blood monocyte-macrophages which stimulates the production of collagenase and prostaglandin E<sub>2</sub> by cells cultured from rheumatoid synovial tissues. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 124:253-256.
3. Dayer, J.-M., and S. Demczuk. 1984. Cytokines and other mediators in rheumatoid arthritis. *Springer Sem. Immunopathol.* 7:387-413.
4. Krane, S. M., S. R. Goldring, and J.-M. Dayer. 1982. Interactions among lymphocytes, monocytes and other synovial cells in the rheumatoid synovium. In *Lymphokines*. E. Pick and M. Landy, editors. Academic Press, Inc., New York. 75-95.
5. Mizel, S. B., J.-M. Dayer, S. M. Krane, and S. E. Mergenhagen. 1981. Stimulation of rheumatoid synovial cell collagenase and prostaglandin by partially purified lymphocyte-activating factor (interleukin-1). *Proc. Natl. Acad. Sci. USA.* 78:2474-2478.
6. Dinarello, C. A., and S. M. Wolff. 1982. Molecular basis of fever in humans. *Am. J. Med.* 72:799-819.
7. Dinarello, C. A. 1984. Interleukin-1. *Rev. Inf. Dis.* 6:51-95.
8. Dayer, J.-M., C. Zavadil-Grob, C. Ucla, and B. Mach. 1984. Translation of mRNA into biologically active mononuclear cell factor (interleukin-1) which stimulates prostaglandin E<sub>2</sub> and collagenase production by synovial cells. *Eur. J. Immunol.* 14:898-901.
9. Demczuk, S., B. Mach, and J.-M. Dayer. 1985. Differential expression of messenger RNA's for mononuclear cell factor from stimulated human monocyte-macrophage cultures. *Br. J. Rheumatol.* 24(Suppl):77-81.
10. Auron, P. E., A. C. Webb, L. J. Rosenwasser, S. F. Mucci, A. Rich, S. M. Wolff, and C. A. Dinarello. 1984. Nucleotide sequence of human monocyte interleukin-1 precursor cDNA. *Proc. Natl. Acad. Sci. USA.* 81:7909-7911.
11. Fontana, A., E. Weber, and J.-M. Dayer. 1984. Synthesis of interleukin-1/endogenous pyrogen in the brain of endotoxin-treated mice: a step in fever induction? *J. Immunol.* 133:1696-1698.
12. Kaye, J., S. Gillis, S. B. Mizel, E. M. Shevach, T. R. Maleck, C. A. Dinarello, L. B. Lachman, and C. A. Janeway, Jr. 1982. Growth of a cloned helper T-cell line induced by a monoclonal antibody specific for the antigen receptor interleukin-1 is required for the expression of receptors for interleukin-2. *J. Immunol.* 133:1339-1345.
13. Dinarello, C. A., H. A. Bernheim, J. C. Cannon, G. LoPreste, S. J. C. Warner, A. C. Webb, and P. E. Auron. 1985. Purified <sup>35</sup>S-met, <sup>3</sup>H-leu-labelled human monocyte interleukin-1 with endogenous pyrogen activity. *Br. J. Rheumatol.* 24(Suppl):59-64.
14. Dinarello, C. A., L. Renfer, and S. M. Wolff. 1977. Human leukocytic pyrogen: purification and development of a radioimmunoassay. *Proc. Natl. Acad. Sci. USA.* 74:4624-4629.
15. Van Damme, J., M. DeLey, G. Opdenakker, and A. Billiau. 1985. Homogeneous interferon-inducing 22k factor is related to endogenous pyrogen and interleukin-1. *Nature (Lond.)* 314:266-268.
16. Hanson, D. F., and P. A. Murphy. 1984. Demonstration of interleukin-1 activity in apparently homogeneous specimens of the pl 5 form of rabbit endogenous pyrogen. *Infect. Immun.* 45:483-490.
17. Lomedico, P. T., U. Gubler, C. P. Hellman, M. Dukovich, J. G. Giri, Y. E. Pan, K. Collier, R. Semionow, A. O. Chua, and S. B. Mizel. 1984. Cloning and expression of murine interleukin-1. *Nature (Lond.)* 312:458-462.