

## Stimulation of inositol trisphosphate and diacylglycerol production in renal tubular cells by parathyroid hormone.

K A Hruska, ... , S Westbrook, M Huskey

*J Clin Invest.* 1987;**79**(1):230-239. <https://doi.org/10.1172/JCI112788>.

### Research Article

Parathyroid hormone (PTH) produced a dose-dependent immediate stimulation of inositol triphosphate and diacylglycerol production in the opossum kidney cell line, primary culture proximal tubular cells, and basolateral membranes from canine proximal tubular segments. The increase in inositol triphosphate production was accompanied by a minor increase in inositol phosphate and no significant increase in inositol bisphosphate production. Associated with the changes in inositol triphosphate and diacylglycerol, there was an immediate hydrolysis of phosphatidylinositol 4'5-bisphosphate. The effect on phospholipid hydrolysis was followed by stimulation of phosphorylation of phosphatidylinositol 4' monophosphate and phosphatidylinositol. PTH produced a sudden increase in cytoplasmic  $\text{Ca}^{2+}$  in opossum kidney cells that persisted for approximately 1 min. Inositol triphosphate transiently increased cytoplasmic  $\text{Ca}^{2+}$  in saponin-treated opossum kidney and primary culture proximal tubule cells. The effects of PTH were not mimicked by cyclic nucleotides. In fact, cyclic AMP appeared to diminish inositol triphosphate production. These results demonstrate that PTH may activate renal tubular epithelial cells by the production of inositol triphosphate and diacylglycerol.

**Find the latest version:**

<https://jci.me/112788/pdf>



# Stimulation of Inositol Triphosphate and Diacylglycerol Production in Renal Tubular Cells by Parathyroid Hormone

Keith A. Hruska, David Moskowitz, Pedro Esbrit, Roberto Civitelli, Scott Westbrook, and Margaret Huskey  
Renal Division, Department of Medicine, Jewish Hospital and Washington University, St. Louis, Missouri 63110

## Abstract

Parathyroid hormone (PTH) produced a dose-dependent immediate stimulation of inositol triphosphate and diacylglycerol production in the opossum kidney cell line, primary culture proximal tubular cells, and basolateral membranes from canine proximal tubular segments. The increase in inositol triphosphate production was accompanied by a minor increase in inositol phosphate and no significant increase in inositol bisphosphate production. Associated with the changes in inositol triphosphate and diacylglycerol, there was an immediate hydrolysis of phosphatidylinositol 4/5-bisphosphate. The effect on phospholipid hydrolysis was followed by stimulation of phosphorylation of phosphatidylinositol 4' monophosphate and phosphatidylinositol. PTH produced a sudden increase in cytoplasmic  $\text{Ca}^{2+}$  in opossum kidney cells that persisted for  $\sim 1$  min. Inositol triphosphate transiently increased cytoplasmic  $\text{Ca}^{2+}$  in saponin-treated opossum kidney and primary culture proximal tubule cells. The effects of PTH were not mimicked by cyclic nucleotides. In fact, cyclic AMP appeared to diminish inositol triphosphate production. These results demonstrate that PTH may activate renal tubular epithelial cells by the production of inositol triphosphate and diacylglycerol.

## Introduction

Phosphatidylinositol 4,5-bisphosphate ( $\text{PtdIns}(4,5)\text{-P}_2$ )<sup>1</sup> is a negatively charged phospholipid which, quantitatively, is a minor component of the plasma membrane phospholipid of renal tubular cells. Phosphorylation of  $\text{PtdIns}$  and  $\text{PtdIns}(4,5)\text{-P}_2$  hydrolysis by phospholipase C produces a high membrane turnover rate of  $\text{PtdIns}(4,5)\text{-P}_2$  making it an effective source of second messages for cell activation. In recent years, a variety of hormones and neurotransmitters have been shown to produce cell activation by stimulating phosphodiesteratic hydrolysis of  $\text{PtdIns}(4,5)\text{-P}_2$  (1, 2). The products of phospholipase C-stimu-

lated hydrolysis, diacylglycerol (DAG), and inositol 1,4,5-triphosphate ( $\text{IP}_3$ )<sup>2</sup>, are messages which in turn activate phospholipid-dependent protein kinases (C-kinase) and produce transient rises in cytoplasmic calcium ( $[\text{Ca}^{2+}]_i$ ). Thus, DAG and  $\text{IP}_3$  control two branches of the calcium messenger system, the C-kinase branch and the calmodulin branch (4). The action of  $\text{IP}_3$  to increase  $[\text{Ca}^{2+}]_i$  results from calcium release from the endoplasmic reticulum (5–7). However, there are additional mechanisms of cell signalling through  $\text{Ca}^{2+}$ , because several hormones increase calcium entry from the extracellular space to produce their biologic effect (4,8–11). The increase in  $[\text{Ca}^{2+}]_i$  produced by agonist stimulated  $\text{IP}_3$  production or influx across the plasma membrane activates cellular secretion, contraction, and several other functions (1, 2, 12). Many of the functions stimulated by the rise in  $[\text{Ca}^{2+}]_i$  are mediated by calmodulin or other intracellular  $\text{Ca}^{2+}$  receptor proteins, thus its designation as the calmodulin branch of the  $\text{Ca}^{2+}$  messenger system (4).

Cell activation by parathyroid hormone (PTH) remains enigmatic. Most reports (13, 14) suggest that PTH binds to a single class of noninteractive receptors, although Scatchard analysis of the binding of  $^{125}\text{I}$ -labeled (Nle8, Nle18, Tyr34) b-PTH 1-34 revealed possible heterogeneity of receptors (15). PTH-receptor complexes clearly stimulate cell function through coupling to adenylate cyclase (16). However, recent studies have demonstrated that PTH increases  $[\text{Ca}^{2+}]_i$  in proximal tubular cells (PTC) of the mammalian kidney (17) independent of cyclic nucleotides. This indicates that the PTH-receptor complex also couples to a mechanism producing a  $\text{Ca}^{2+}$  transient. One such mechanism would be receptor-mediated  $\text{PtdIns}(4,5)\text{-P}_2$  hydrolysis and  $\text{Ins}(1,4,5)\text{-P}_3$  production. Several investigators (18–21) have shown an effect of PTH on renal tubular phosphoinositide metabolism, and one report shows the effect was present in the apical membrane of the proximal renal tubule (18). Some have reported that the stimulation of phosphoinositide metabolism by PTH was cyclic AMP dependent (19). This would remove it from a mechanism of cell activation, and place it in the realm of a biologic response. However, other investigators (20, 21) have not found a stimulation of phosphoinositide metabolism by cyclic AMP in the renal cortex. To date, no report has demonstrated production of the messages of  $\text{PtdIns}(4,5)\text{-P}_2$  hydrolysis ( $\text{IP}_3$  and DAG) by PTH. Thus, the question of whether cell activation by PTH involves stimulation of a calcium-related signal remains a critical biologic issue.

The studies reported here have been designed to examine the action of PTH on phosphoinositide metabolism in the opossum kidney (OK) cell line (which exhibits proximal tubular characteristics). Primary cultures of canine proximal tubules, and basolateral cell membranes (BLM) prepared from isolated

Address correspondence to Dr. Hruska, Renal Division, Jewish Hospital of St. Louis, 216 S. Kinghighway, St. Louis, MO 63110.

Received for publication 2 June 1986 and in revised form 7 August 1986.

1. Abbreviations used in this paper: BLM, basolateral cell membrane;  $[\text{Ca}^{2+}]_i$ , cytoplasmic calcium; C-kinase, phospholipid-dependent protein kinase; DAG, diacylglycerol; FCCP, (carbonyl) cyanide *p*-trifluoromethoxyphenylhydrazone; GPI, glycerophosphoinositol; GPIIP, GPI phosphate; GPIIP<sub>2</sub>, GPI diphosphate; IP, inositol phosphate; OK, opossum kidney; PA, phosphatidic acid; PTC, proximal tubular cells;  $\text{PtdIns}$ , phosphatidylinositol;  $\text{PtdIns}$  4-P,  $\text{PtdIns}$  4'-monophosphate;  $\text{PtdIns}(4,5)\text{-P}_2$ ,  $\text{PtdIns}$  4,5-bisphosphate; TBAS, tetrabutylammonium sulfate.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/87/01/0230/10 \$1.00

Volume 79, January 1987, 230–239

2. The product of  $\text{PtdIns}(4,5)\text{-P}_2$  hydrolysis may be inositol 1,2(cyclic)-4,5-triphosphate (3). This compound is converted to  $\text{IP}_3$  by acidic extraction procedures, and is not discussed separately from  $\text{IP}_3$  for the purposes of this paper.

segments of the proximal tubule were also studied. The BLM of proximal tubular cells is known to contain specific PTH receptors and adenylate cyclase sensitive to PTH (13–15). The studies reported here demonstrate that PTH stimulates phosphoinositide turnover through activation of phospholipase C, resulting in the production of DAG and IP<sub>3</sub>. The studies using the BLM in vitro indicate that this effect is closely associated with receptor binding. The stimulation of IP<sub>3</sub> may participate in the production of an elevated [Ca<sup>2+</sup>]<sub>i</sub> that follows stimulation of cultured renal tubular cells by PTH.

## Methods

**Sources of chemicals.** Myo-[2-<sup>3</sup>H]inositol (1.25 µCi/ml) was obtained from American Radio Label (St. Louis, MO) and [1-<sup>14</sup>C]arachidonic acid (0.42 µCi/ml) was obtained from Amersham Corp., (Arlington Heights, IL). Other chemicals were the highest purity available.

### Experimental protocol

The OK cell line passages (Nos. 78–83), were grown in Dulbecco's minimal essential medium containing 10% fetal calf serum in a 5% CO<sub>2</sub>/95% air atmosphere at 37°C in 25 cm<sup>2</sup> polystyrene flasks. At confluency, and 72 h before study, the growth medium was changed to an inositol-free culture medium. 24 h before the experiments, the media was changed to one containing myo-[2-<sup>3</sup>H]inositol (3–4 × 10<sup>6</sup> cpm/3 ml) or [1-<sup>14</sup>C]arachidonic acid 9–9.5 × 10<sup>4</sup> cpm/ml). Preliminary experiments (data not shown), revealed that labeling of the various inositol pools was at equilibrium by 24 h. On the day of the experiment, bPTH 1-84 at varying concentrations or its vehicle (0.01 M acetic acid), were added to the culture media, and the flasks were gently shaken and returned to the incubator for variable periods.

The reactions were stopped by removing the media, adding 1 ml of H<sub>2</sub>O at 4°C, and immediate freezing at –50°C. For analysis, cells were thawed, removed from flasks with a rubber policeman, and rinsed with water. After removing appropriate samples for determination of DNA, the tissue was extracted according to Grove (22). To the tissue suspension, 2 ml of chloroform/methanol (1:1) with 10 mM tetrabutylammonium sulfate (TBAS) was added, then 1 ml of 0.1 N HCl, and an additional 1 ml of chloroform were added. The extracts were vortexed and centrifuged. The upper phases were removed and the organic phases washed with 1 ml chloroform/0.1 N HCl (1:1 vol/vol). The aqueous phases of the washes were added to the original aqueous phases.

**Chromatographic techniques.** Aqueous phases of the cell extracts were added to columns containing 1.5 ml of anion exchange resin AG 1-X8 (Bio-Rad Laboratories, Richmond, CA). The IPs were eluted as described by Berridge (23, 24) with 15 ml of water followed by 9 ml of 5 mM tetraborate, 60 mM sodium formate followed by 16 ml of 0.1 M formic acid, 0.15 M ammonium formate followed by 10 ml of 0.1 M formic acid, 0.3 M ammonium formate, followed by 10 ml of 0.1 M formic acid, 0.75 M ammonium formate, and finally 10 ml of 0.1 M formic acid and 1 M ammonium formate.

The lipid phases of the cell extracts were deacylated by the transacylation procedure of Clark and Dawson (25). The deacylated lipids were added to columns containing anion-exchange resins similar to those used for the aqueous phases of the extracts and eluted by 5 mM sodium tetraborate, 0.15 M ammonium formate (for glycerophosphoinositol [GPI]); 0.1 mM formic acid, 0.3 M ammonium formate (for GPI phosphate [GPIP]); and 0.1 M formic acid, 0.75 M ammonium formate (for GPIP<sub>2</sub>). Recovery of radiolabeled phospholipid from the deacylation procedure as GPIP<sub>2</sub> was 99% or greater. Overall, between 25 and 35% of the myo-[2-<sup>3</sup>H]inositol added to the culture media was incorporated into the cells. Recoveries from the extractions and chromatographic procedures were between 80 and 90%, and PTH had no effect on recovery. Between 50 and 60% of the myo-[2-<sup>3</sup>H]inositol in the aqueous phases of the cell extracts eluted as free inositol.

For the determination of [<sup>14</sup>C]DAG, lipid phases of the cell extracts

and authentic standards were spotted on activated silica gel G-25 thin-layer chromatographic plates and developed in the solvent system of hexane/ethylether/acetic acid (80:20:2). The lipids were identified by staining with iodine and scrapped from the thin-layer plate. [<sup>14</sup>C]DAG in the spots was determined by liquid scintillation spectroscopy.

**Determination of [Ca<sup>2+</sup>]<sub>i</sub>.** OK cells were removed from tissue culture flasks by a 20-min exposure to 5 mM EDTA at 37°C. Cell suspensions were allowed to reequilibrate in growth media for 1 h in a shaking water bath at 37°C, and [Ca<sup>2+</sup>]<sub>i</sub> was determined using the fluorescent indicator indo-1 of Grynkiewicz et al. (26) as previously described (27). The acetoxymethyl ester of indo-1 (10 µM) was added to cell suspensions for 40 min. Cells were centrifuged at 200 g for 1 min and resuspended in a 10 mM buffer containing 127 mM NaCl, 3.8 mM KCl, 1.2 mM KH<sub>2</sub>PO<sub>4</sub>, 1.2 mM CaCl<sub>2</sub>, 0.8 mM MgCl<sub>2</sub>, 5 mM glucose, and 10 mM Hepes (pH 7.40). These cell suspensions were placed in a 2-ml quartz cuvette in a spectrofluorometer (model 4800; SLM Instruments, Inc., Urbana, IL). The chamber was constantly stirred and the temperature maintained at 37°C. Excitation was at 355 nm with an 8-nm slit width, and emission was monitored at 400 nm with an 8-nm slit width and at 490 nm using an interference filter (Corion Corp., Holiston, MA). The ratio of emitted light 400/490 nm was continuously recorded and used to calculate [Ca<sup>2+</sup>]<sub>i</sub> as described by Grynkiewicz et al. (26).

**Primary culture of proximal convoluted tubule cells.** Proximal tubular cells (PTC) from the cortex of canine kidneys were isolated and cultured as previously described (17). The experimental protocols used for studies with PTC was the same as those described above for the OK cell line except for the differences in culture techniques.

**Calcium uptake in permeabilized cells.** OK and PTC were grown in 24-well culture plates to confluency in their respective growth media. Cells were permeabilized with 75 µg/ml saponin for 10 min at 37°C. Plates were then washed with a cytoplasm-like buffer containing 100 mM KCl, 20 mM NaCl, 5 mM MgSO<sub>4</sub>, 0.96 mM NaH<sub>2</sub>PO<sub>4</sub>, 25 mM NaHCO<sub>3</sub>, and 1 mM EGTA. The calcium concentration was adjusted to either 20 nM or 3.3 µM. Ca<sup>2+</sup> concentrations were determined using the method of Bers (28). Then 0.5 ml of cytoplasmic medium containing 1 µCi/ml <sup>45</sup>CaCl and an ATP regeneration system (5 mM creatine phosphate, 5 U/ml creatine phosphokinase, and 1.5 mM ATP) was added to the cells. Incubation was carried out at 37°C, and calcium uptake was terminated by aspiration of the medium and washing with ice-cold cytoplasmic buffer ×2. Then 0.5 ml of Trypsin (0.05%)-EDTA (0.02%) was added to the wells and incubated for 30–60 min at 37°C. 250 µl was aspirated from each well and counted for radioactivity in a liquid scintillation spectrometer.

**Isolated proximal tubular segments.** Proximal tubular segments were isolated from the cortex of mongrel dog kidneys as previously described (20). The isolated proximal tubular segments were suspended in a modified Krebs-Hensleit bicarbonate buffer containing 5 mM glucose, 5 mM alanine, and 5 mM sodium lactate. They were gassed with 95% O<sub>2</sub>/5% CO<sub>2</sub>. The tubules were resuspended in 50 ml of Krebs-Ringer bicarbonate containing 7 mM ATP and 100 µCi of myo-[2-<sup>3</sup>H]inositol for 3 h. BLM were prepared from the tubular segments as previously described (29). BLM were suspended in a buffer containing 130 mM sucrose, 26 mM KCl, 5 mM MgCl<sub>2</sub>, and 50 mM Tris-HCl (pH 7.5). Characterization of the BLM preparation was previously reported (29), and the results of the characterization in the present studies were similar. Enrichment of Na<sup>+</sup>/K<sup>+</sup> ATPase activity was 8.2 ± 1.5, *n* = 10, compared with the superficial cortical homogenate. Enrichment of alkaline phosphatase was 0.8 ± 0.3, *n* = 7 and that of 6-NAD hydrogenase was 1.0 ± 0.3, *n* = 7. The activities of the above enzymes were used as markers of the BLM, the apical cell membrane, and the mitochondrial membrane, respectively.

**Phosphorylation of BLM.** Phosphorylation of BLM was carried out as previously described (30, 31) for proximal tubular brush border membranes. Suspensions of BLM containing 200–400 µg of protein were incubated in a water bath at 37°C for 1 min. Then 4 vol of solutions containing 1 mM Mg<sup>2+</sup>-ATP and 5–20 µCi of [<sup>32</sup>P]-ATP, 26 mM KCl, 5 mM MgCl<sub>2</sub>, 1 mM EDTA, and 1 µM free Ca<sup>2+</sup>, 50 mM Tris-HCl (pH 7.6) with or without PTH and/or calmodulin according to the experiment, were added to the membrane suspension. This solution was hypotonic

compared with the membrane suspension to cause swelling of vesicles in the preparation (which is ~50% sheets, reference 29), and to increase the access of reagents to both sides of the membranes. Preliminary experiments revealed the lipid products of phosphorylation to be phosphatidic acid (PA), PtdIns 4-P, and PtdIns(4,5)-P<sub>2</sub>.

The phosphorylation reaction was stopped at various times of incubation by addition of 3 ml ice-cold chloroform/methanol (1:1 vol/vol) containing 10 mM TBAS. Extraction and chromatography were completed as described above.

**Analysis of <sup>32</sup>Pi phospholipids of BLM.** Lipid extracts of BLM were spotted on Silica gel 60 thin-layer chromatographic plates (E. Merck, Darmstadt, Germany) and developed in two different solvent systems. The first system consisted of chloroform/methanol/15 N ammonium hydroxide/water (90:90:5:20, vol/vol), and it was used for separation of the polyphosphoinositides. The second system consisted of chloroform/acetone/methanol/acetic acid/water (10:40:20:20:10, vol/vol) for PA. Phospholipids were localized by exposing to iodine vapors and spots labeled with <sup>32</sup>Pi were detected by autoradiography on Kodak X-Omat RP film. Authentic standards were used for identification of the phosphoinositides and PA. Phospholipid spots were scraped from the plates and their radioactivity counted in a Tri-Carb 460CD liquid scintillation counter (Packard Instrument Co., Inc., United Technologies, Downer's Grove, IL).

**BLM protein phosphorylation.** BLM and 0.2–0.4 mg protein, were suspended in 26 mM KCl, 130 mM sucrose, 5 mM MgCl<sub>2</sub>, and 50 mM Tris-HCl (pH 7.5). They were reacted with similar solutions for phosphorylation without sucrose containing 1 mM Mg-ATP, and 5–20 μC of [<sup>32</sup>Pi]-ATP, and where appropriate: 1 μM Ca<sup>2+</sup>, 1 mM EGTA, 3 × 10<sup>-7</sup> M calmodulin, 10<sup>-7</sup> M bPTH 1-84, 10<sup>-6</sup> M 3'5'-cAMP, and 2 × 10<sup>-4</sup> M W13. Reactions were stopped by addition of an ice-cold, 60 mM Tris-HCl (pH 5.8) containing 10% sodium dodecyl sulfate (wt/vol) and 30% glycerol (wt/vol). Samples were boiled for 2 min, cooled, and frozen at -70°C. BLM samples containing 100–200 μg protein were added to 10% acrylamide slab gels and electrophoresis was performed for 4 h with 6 milliA/gel. Gels were stained with Coomassie Blue, dried, and exposed to radiographic film. Autoradiograms were developed after 3–7 d of exposure.


**Chemical analyses.** DNA content of renal tubular cells was determined by the method of Labarca and Paigen (32). Na<sup>+</sup>/K<sup>+</sup> ATPase, 6 NAD hydrogenase and alkaline phosphatase were measured by methods previously described (29).

**Statistical analysis.** The significance of differences between the results for vehicle-treated and PTH-treated cells and membranes were determined by the use of the unpaired *t* test.

## Results

**Myo-[2-<sup>3</sup>H]inositol labeling of OK cells and separation of phosphoinositols.** The OK cells were grown as described in Methods for 7–10 d to confluency. After incubation in inositol-free media and labeling with myo-[2-<sup>3</sup>H]inositol as described in Methods, the cells were exposed to PTH, cyclic nucleotides, or the respective vehicle for variable periods. After the experimental incubations, the tissues were extracted and the aqueous phases of the extracts were added to columns containing an anion-exchange resin, and eluted as described in Methods. Fig. 1 A shows a representative elution profile. There were six peaks of myo-[2-<sup>3</sup>H]inositol representing inositol (I), GPI, IP, IP<sub>2</sub>, and IP<sub>3</sub> (23, 24, 33)<sup>3</sup>. These were identified by the elution pattern of pure

3. The finding that IP<sub>3</sub> as identified in this paper is a mixture of inositol 1,3,4-triphosphate and inositol 1,4,5-triphosphate is beyond the objective of this paper (34). Ins 1,3,4, P<sub>3</sub> may be a breakdown product of Ins 1,3,4,5 P<sub>4</sub> formed by phosphorylation of Ins 1,4,5 P<sub>3</sub> (35). The generic term IP<sub>3</sub> will be used to refer to the product isolated from the anion-exchange columns which probably represents both Ins 1,4,5 P<sub>3</sub> and Ins 1,3,4 P<sub>3</sub>.



**Figure 1.** Anion-exchange chromatography of aqueous IPs, and deacylated phosphoinositides extracted from OK cells. Confluent cells in an inositol-free media were incubated with myo-[2-<sup>3</sup>H]inositol (3–5 × 10<sup>-6</sup> cpm/3 ml) for 24 h. Experiments were conducted as described in Methods. Aqueous phases of the cell extracts were applied to columns containing AG 1X-8 formate form anion-exchange resin and eluted with: a, water for inositol; b, 5 mM sodium tetraborate/60 mM sodium formate for GPI; c, 0.1 M formic acid/0.15 M ammonium formate for IP; d, 0.1 M formic acid/0.3 M ammonium formate for IP<sub>2</sub>; e, 0.1 M formic acid/0.75 M ammonium formate for IP<sub>3</sub>; and f, 0.1 M formic acid/1 M ammonium formate. The lipid phases of the extract were deacylated as described in Methods and applied to similar anion-exchange columns and eluted with the deletion of solutions b and f and substitution of 5 mM sodium tetraborate/150 mM sodium formate for solution c. Fractions of 1 ml were collected and counted for <sup>3</sup>H by liquid scintillation spectroscopy.

standards. The acidic extraction procedure eliminated inositol cyclic phosphates from the elution profile (23). The small peak eluting after IP<sub>3</sub> has not been identified as yet.

The lipid phases of the extracts were deacylated and eluted from columns containing anion-exchange resins as described in Methods. Myo-[2-<sup>3</sup>H]inositol was present in four peaks: free inositol, GPI, GPIIP, and GPIIP<sub>2</sub>. The latter three peaks represented the original inositol phospholipids, PtdIns, PtdIns 4-P, and PtdIns(4,5)-P<sub>2</sub>, respectively (Fig. 1 B).

Table I shows the distribution of incorporated myo-[2-<sup>3</sup>H]inositol among the various inositides. Most of the label appeared in PtdIns, whereas the levels of PtdIns 4-P and

Table I. Incorporated Myo-[2-<sup>3</sup>H]inositol in OK Cells

Inositide	DNA	Distribution
	cpm/μg	%
GPI	7,250±540	4.1
IP	19,293±1,590	11.0
IP <sub>2</sub>	500±110	0.3
IP <sub>3</sub>	1,417±173	0.8
PtdIns	145,624±2,074	83
PtdIns 4-P	1,084±74	0.6
PtdIns(4,5)-P <sub>2</sub>	370±12	0.2

Data are mean±SE, *n* = 4.

PtdIns(4,5)-P<sub>2</sub> in the cell extracts were low. The largest fraction in the aqueous phases of the cell extracts was IP, as expected from the pattern of phospholipid labeling.

**Effect of PTH on IP<sub>3</sub> release.** Addition of PTH to OK cells labeled with myo-[2-<sup>3</sup>H]inositol incubations produced a prompt and sustained increase in IP<sub>3</sub> production (Table II). Although the levels of IP, which were the largest IP fraction, also tended to increase, this effect was smaller than the effect on IP<sub>3</sub>. IP<sub>2</sub> levels were not affected by PTH. In concert with the stimulation of IP<sub>3</sub> production, DAG levels were also increased by PTH (Fig. 2).

The effect of PTH on the levels of myo-[2-<sup>3</sup>H]inositol-containing lipids were as predicted by the effects on IP production. There was a rapid decrease in the levels of [<sup>3</sup>H]PtdIns(4,5)-P<sub>2</sub> (Table III). Thereafter, [<sup>3</sup>H]PtdIns(4,5)-P<sub>2</sub> levels recovered to values not different from control in 1 min and increased above control at 5 min.

The effect of PTH was dose related (Table IV). Maximal effects were observed at 10<sup>-7</sup> M. At 10<sup>-10</sup> M PTH, the effects on IP<sub>3</sub> production were still detectable. The dose-dependent stimulation of IP<sub>3</sub> production was more sensitive than the reported dose response of cAMP production in these cells (36). However, modulation of cAMP and IP<sub>3</sub> by PTH in OK cells will have to be studied together to adequately address the questions of sensitivity and modulation of one pathway without the other. The IP<sub>3</sub> response Table IV shows still reflected an insensitivity to PTH. The accessibility of the receptor for hormone

Table II. Effects of PTH and cAMP on IP<sub>3</sub> and IP Production by OK Cells

	10 s	30 s	5 min
	cpm/μg DNA	cpm/μg DNA	cpm/μg DNA
IP <sub>3</sub>			
Control <i>n</i> = 7	1,314±137	1,417±173	1,404±195
PTH, 10 <sup>-7</sup> M <i>n</i> = 4	1,717±270*	2,128±270*	2,165±292*
cAMP, 10 <sup>-3</sup> M <i>n</i> = 3	1,313±281	1,277±179	1,685±298
IP			
Control <i>n</i> = 7	19,950±1,000	19,256±810	19,256±1,300
PTH, 10 <sup>-7</sup> M <i>n</i> = 4	21,115±890	20,602±1,460	21,001±2,357
cAMP × 10 <sup>-3</sup> M <i>n</i> = 3	17,900±358	16,711±240	17,825±900

\* *P* < 0.05.




Figure 2. Effects of PTH on DAG production in OK cells. Experiments were conducted as described in Methods, and [<sup>14</sup>C]arachidonic acid was substituted for myoinositol 24 h before the experiment. Analysis of DAG content in the lipid extracts was performed as described in Methods. Results are mean±SE of three experiments, PTH > control, *P* < 0.05. Hatched bars, control; open bars, 10<sup>-7</sup> M PTH.

binding, the expression of PTH receptors in OK cells, and insensitivity of the detection methods must be considered as potential and unexplored factors leading to apparent lack of sensitivity to PTH in this model system of polarized anchorage-dependent cells on a solid support.

**Stimulation of IP release from PTC grown in primary culture by PTH.** As a transformed cell line, OK cells may potentially exhibit a response to PTH not found in nontransformed cells. Thus, the effects of PTH on PTC grown in primary culture were tested. PTC were grown to confluency in a defined media as previously described (17). When the experimental protocols described in Methods for the OK cells were repeated in the primary culture of PTC, there were significant differences in the incorporation of myo-[2-<sup>3</sup>H]inositol. In primary culture of PTC, incorporation of the myo-[2-<sup>3</sup>H]inositol label was 5–10% compared with the 25–35% observed in the OK cells. Despite the lower level of inositol labeling, the aqueous phases of the cell extracts of the primary culture of PTC had IP levels that were readily detectable. The predominance of PtdIns and IP shown in the OK cells was again present in the PTC. PTH produced a prompt increase in the production of the IPs by 30 s Table V shows. The largest effect was on IP<sub>3</sub> production as in the OK cells.

**Effects of PTH on [Ca<sup>2+</sup>]<sub>i</sub> in OK cells.** If PTH indeed increases IP<sub>3</sub> production, an effect of the hormone to transiently increase [Ca<sup>2+</sup>]<sub>i</sub> would be expected. We have previously shown (17) that PTH increases [Ca<sup>2+</sup>]<sub>i</sub> in confluent PTC grown in defined media. As Fig. 3 shows, PTH also produced a prompt calcium transient in suspensions of OK cells. Basal [Ca<sup>2+</sup>]<sub>i</sub> was 145±15 nM, *n* = 15. During multiple experiments similar to that shown in Fig. 5, PTH produced an immediate 40–50% increase in [Ca<sup>2+</sup>]<sub>i</sub> which returned to basal levels in the course of ~1 min.

**Calcium uptake in saponin-treated OK and PTC.** An effect of IP<sub>3</sub> production in other cells, has been to cause a transient release of Ca<sup>2+</sup> from an intracellular nonmitochondrial pool

Table III. Effect of PTH on Phosphatidylinositol 4,5-bisphosphate, PtdIns(4,5)-P<sub>2</sub>, Hydrolysis in OK Cells

	PtdIns(4,5)-P <sub>2</sub>		
	10 s	30 s	5 min
	cpm/μg DNA	cpm/μg DNA	cpm/μg DNA
Control <i>n</i> = 4	390±20	370±21	195±14
PTH <i>n</i> = 4	187±17*	217±22*	233±23

\* *P* < 0.05.

Table IV. Dose-dependent Effect of PTH on IP<sub>3</sub> Production

	cpm/ $\mu$ g DNA	Control %
Control	1354 $\pm$ 117	—
bPTH 1-84 10 <sup>-7</sup> M	2107 $\pm$ 24*	155
bPTH 1-84 10 <sup>-8</sup> M	1714 $\pm$ 112†	127
bPTH 1-84 10 <sup>-9</sup> M	1648 $\pm$ 10†	122
bPTH 1-84 10 <sup>-10</sup> M	1600 $\pm$ 100†	118

Data are mean $\pm$ SE of three experiments performed in duplicate or triplicate.

\* >C, *P* < 0.01.

† >C, *P* < 0.05.

(4–6). To study the homeostasis of cellular calcium in the PTC and OK cells, the cells were grown as described in Methods and exposed to Saponin. Then the media were changed to a buffer made to resemble the cytosol as described in Methods. The uptake of <sup>45</sup>Ca was examined in the presence and absence of an uncoupler and a mitochondrial ATPase inhibitor, (carbonyl) cyanide *p*-bifluoromethoxyphenylhydrazone (FCCP), and oligomycin. At low media calcium concentrations (0.02  $\mu$ M), ATP-stimulated calcium uptake was similar in both conditions (Fig. 4). When the media calcium was raised to 3  $\mu$ M, ATP-stimulated calcium uptake in the FCCP and oligomycin-treated cells was similar to the calcium uptake observed at 0.02  $\mu$ M calcium. However, in the untreated cells there was a dramatic increase in ATP-stimulated calcium uptake. These results indicate that at low calcium concentrations, a nonmitochondrial intracellular organelle is largely responsible for ATP-stimulated calcium uptake. However, at higher calcium concentrations, the mitochondria represents a much larger potential calcium reservoir than the nonmitochondrial pool. These data are similar to those of Bonventre et al. in porcine kidney-derived continuous cell line cells (37). At extracellular calcium concentrations of 0.02  $\mu$ M, ATP-stimulated calcium uptake was rapidly released by adding 5.0  $\mu$ M IP<sub>3</sub>. The release of calcium by IP<sub>3</sub> was short lived and by 5–10 min ATP-stimulated calcium uptake was restored. These results demonstrate that IP<sub>3</sub> produces a prompt release of calcium from the intracellular store accumulating calcium in the presence of ATP. Data from other investigators suggest that the nonmitochondrial store, which is responsible for this ATP-dependent calcium uptake, is the endoplasmic reticulum (4–6, 38).

#### Effects of PTH on IP and PtdIns(4,5)-P<sub>2</sub> turnover in BLM.

Table V. Effect of PTH on Production of Inositol Phosphates in Primary Cultures of Canine Proximal Tubule Cells

	IP	IP <sub>2</sub>	IP <sub>3</sub>
	cpm/ $\mu$ g DNA	cpm/ $\mu$ g DNA	cpm/ $\mu$ g DNA
Control	7500 $\pm$ 400	1075 $\pm$ 100	2625 $\pm$ 250
PTH (10 <sup>-7</sup> )	8725 $\pm$ 550*	1550 $\pm$ 105*	3900 $\pm$ 300*

Data are mean $\pm$ SE, *n* = 4 at 30 s of incubation.

\* *P* < 0.05.




Figure 3. Effect of PTH on [Ca<sup>2+</sup>]<sub>i</sub> in OK cells. OK cells grown to confluency were suspended with indo-1 as described in Methods. The ratio of emitted light at 400/490 nm was continuously recorded. The break in the tracing represents the addition of 10<sup>-7</sup> M PTH.

Because we have previously shown that PTH produces an increase in the phosphoinositides of the brush border membrane of PTC (18), experiments were designed to analyze whether PTH could directly effect PtdIns(4,5)-P<sub>2</sub> turnover in isolated basolateral membranes. This would indicate that the effect of PTH on IP<sub>3</sub> release and PtdIns(4,5)-P<sub>2</sub> turnover was closely linked to receptor binding. To perform these experiments, isolated proximal tubular segments were prepared from the cortex of canine kidneys as described in Methods. The segments were labeled with myo-[2-<sup>3</sup>H]inositol as described in Methods. After labeling with myo-[2-<sup>3</sup>H]inositol, BLM were prepared from the proximal tubular segments. Characterization of the BLM were similar to results previously reported from this laboratory for freshly prepared BLM. Incorporation of the labeled inositol into the proximal tubular segments was only 2–5% of the label. Despite these low rates of incorporation, the levels of IPs in the aqueous phases of the extracts for BLM were detectable although the levels of IP released were much lower than that observed in the whole cell extracts of OK and PTC cultures. This is probably due to the cytoplasmic nature of a PtdIns-specific phospholipase-C (unpublished results, Suzuki, Y., and K. Hruska), whereas the






Figure 4. Effect of IP<sub>3</sub> on ATP-stimulated calcium uptake in saponin-treated OK and PTC. Suspensions of OK and PTC were treated with saponin as described in Methods. The results are shown for OK cells but they were exactly similar in PTC. A, calcium uptake stimulated by ATP in an extracellular calcium buffer of 0.2  $\mu$ M. B, ATP-stimulated calcium uptake in the presence and absence of FCCP and oligomycin at 3.0  $\mu$ M extracellular calcium. C, effects of IP<sub>3</sub> 0.5  $\mu$ M on ATP-stimulated calcium uptake with an extracellular calcium concentration of 0.2  $\mu$ M. Results are the mean of four experiments done in triplicate. Solid circles, control; open circles, FCCP-oligomycin.



**Figure 5.** Effect of PTH on  $\text{IP}_3$  release from BLM. BLM were prepared from isolated proximal tubular segments labeled with myo-[2- $^3\text{H}$ ]inositol as described in Methods. The elution of  $\text{IP}_3$  (steps a-e, C) was described in the legend to Fig. 1. Data are the mean of two experiments.  $10^{-7}$  M PTH produced a prompt and sustained increase in  $\text{IP}_3$  (fractions 39-43) release from the BLM. Similar results were observed in two other experiments, but with lower counts in all of the  $\text{IP}$  fractions. Solid circles, b-PTH 1-84; open circles, control.


membrane-bound phospholipase-C is specific for  $\text{PtdIns}(4,5)\text{-P}_2$ . As Fig. 5 shows, PTH produced a prompt and sustained increase in the production of  $\text{IP}_3$ . The effect of PTH on the production of  $\text{IP}_2$  lagged behind  $\text{IP}_3$  and was transient, whereas the effect on  $\text{IP}$  was less.

To further study the effects of PTH on phosphoinositide turnover in BLM, the effects of PTH on  $^{32}\text{P}$ i incorporation into phospholipids from [ $\gamma\text{-}^{32}\text{P}$ ]ATP was studied as described in



**Figure 6.** Effect of PTH on phospholipid phosphorylation of BLM. BLM were prepared and phosphorylated as described in Methods. The incorporation of  $^{32}\text{P}$ i from  $\gamma\text{-}^{32}\text{P}$ -ATP was increased in BLM treated with PTH at 5 min. Data are shown as  $^{32}\text{P}$ i incorporated into phospholipid as the mean  $\pm$  SE of five experiments. \* PTH > control,  $P < 0.05$ . Solid triangles, b-PTH  $1\text{-}84 \cdot 10^{-6}$  M; solid circles, control.

Methods. As reported previously (18, 30), the phospholipids labeled in these experiments were PA,  $\text{PtdIns 4-P}$ , and  $\text{PtdIns}(4,5)\text{-P}_2$ .  $\text{PtdIns}(4,5)\text{-P}_2$  turnover stimulates  $^{32}\text{P}$ i incorporation from ATP as a result of phosphorylation of  $\text{PtdIns}$  and  $\text{PtdIns 4-P}$  (1, 2, 20, 39). Also, the product of  $\text{PtdIns}(4,5)\text{-P}_2$  turnover, DAG, is phosphorylated to PA. The enzyme DAG kinase has been shown to maintain very low DAG levels in the plasma membrane using substrate in the reaction to near completion in producing PA (39, 40). As Fig. 6 shows, incubation of BLM with PTH produced an increase in  $^{32}\text{P}$ i incorporation into PA,  $\text{PtdIns 4-P}$ , and  $\text{PtdIns}(4,5)\text{-P}_2$  at 5 min. After 5 min, the levels of [ $^{32}\text{P}$ ]PA, [ $^{32}\text{P}$ ]PtdIns 4-P, and [ $^{32}\text{P}$ ]PtdIns(4,5) $\text{-P}_2$  declined in BLM treated with PTH. By 30 min of incubation, the levels of [ $^{32}\text{P}$ ]PtdIns(4,5) $\text{-P}_2$  were significantly lower in the PTH-treated BLM, which indicated a greater rate of  $\text{PtdIns}(4,5)\text{-P}_2$  hydrolysis in PTH-treated BLM as capacity for [ $^{32}\text{P}$ ] phosphorylation was exhausted. These data indicate that PTH-stimulated phosphoinositide turnover in BLM resulted in an increased rate of  $^{32}\text{P}$ i incorporation into the polyphosphoinositides through phosphorylation of  $\text{PtdIns}$  and  $\text{PtdIns 4-P}$ . The hydrolysis of



**Figure 7.** Effect of dibutyryl cAMP on  $\text{IP}_2$  and  $\text{IP}_3$  production in BLM prepared from proximal tubular segments. Shown are the terminal phases of the elution from anion-exchange columns, fractions 30-38,  $\text{IP}_2$ , and 40-45,  $\text{IP}_3$ . Results are mean of an experiment performed in triplicate. A, control incubations. B, incubations with dibutyryl cAMP ( $10^{-3}$  M). At 10 s ( $10''$ )  $\text{IP}_2$  and  $\text{IP}_3$  levels were decreased in the incubations of BLM with dibutyryl cAMP as compared with control. Solid circles,  $10''$ ; open triangles,  $1', 1'+$ ; solid squares,  $5', 5'+$ .


PtdIns(4,5)-P<sub>2</sub> decreased product inhibition of PtdIns kinase and the stimulation of phosphorylation resulted (41).

**Effect of 3'5'-cAMP on PtdIns(4,5)-P<sub>2</sub> turnover.** To study the effects of cyclic nucleotides on the turnover of PtdIns(4,5)-P<sub>2</sub> hydrolysis and IP<sub>3</sub> release, experiments identical to those Figs. 5 and 6 show were repeated using additions of dibutyl cAMP 10<sup>-3</sup> M instead of PTH. BLM were prepared from isolated proximal tubular segments prelabeled with myo-[2-<sup>3</sup>H]inositol as described in the previous section and Methods. Fig. 7 shows production of IP<sub>2</sub> and IP<sub>3</sub> from labeled BLM. The results suggested that IP<sub>3</sub> production was decreased by dibutyl cAMP at 10 s of incubation but not affected at 30 s and 5 min of incubation. In addition, dibutyl 3'5'-cAMP did not mimic the PTH effect on <sup>32</sup>Pi incorporation from [ $\gamma$ -<sup>32</sup>Pi]ATP into PA, PtdIns 4-P, and PtdIns(4,5)-P<sub>2</sub> (Fig. 8).

**Effect of PTH on calmodulin-dependent phosphorylation of BLM.** To study direct stimulation of Ca<sup>2+</sup>-calmodulin-dependent functions in BLM, the effects of PTH on calmodulin-dependent protein phosphorylation was studied. As Fig. 9 shows, PTH added to BLM stimulated the phosphorylation of a protein with a molecular weight of ~38,000 D. The phosphorylation of this protein was stimulated by calmodulin and inhibited by specific calmodulin inhibitors. Thus, PTH was shown to stimulate calmodulin-dependent protein kinases in the BLM in vitro. PTH addition in the absence of Ca<sup>2+</sup>-calmodulin failed to stimulate phosphorylation of the 38,000-D protein (data not shown). Although the mechanism of the PTH effect on the Ca<sup>2+</sup>-calmodulin-dependent protein kinase requires further study, these results indicate the presence of a system in the BLM dependent on Ca<sup>2+</sup>-calmodulin and responsive to PTH. Thus, the Ca<sup>2+</sup> transient produced by PTH in PTC may affect the activity of this kinase.

## Discussion

In three experimental settings, OK cells, PTC, and BLM, the data presented above demonstrate a prompt increase in IP<sub>3</sub> production, DAG production, and PtdIns(4,5)-P<sub>2</sub> hydrolysis after PTH addition. The high rate of incorporation of myo[2-<sup>3</sup>H]inositol by OK cells was used to unequivocally demonstrate the stimulation of IP<sub>3</sub> and DAG production by PTH. The studies



**Figure 8.** Effect of dibutyl cAMP on BLM phospholipid phosphorylation. BLM were prepared and phosphorylated as described in Fig. 8 and Methods. Addition of dibutyl cAMP 10<sup>-3</sup> M had no significant effect on <sup>32</sup>Pi incorporation of  $\gamma$ -<sup>32</sup>P ATP. Similar results were observed with additions of 8-bromo cAMP (10<sup>-4</sup> M). Open circles, 10<sup>-4</sup> M db-cAMP.

in PTC and BLM in vitro indicate that this is an effect of PTH directly associated with receptor binding in nontransformed cells. The effects of PTH on PtdIns(4,5)-P<sub>2</sub> hydrolysis and IP<sub>3</sub> production were not mimicked by cAMP. In OK cells, PTH (10<sup>-10</sup> M) was sufficient to stimulate production of IP<sub>3</sub> and DAG. This dose-dependent stimulation of IP<sub>3</sub> production compared favorably with previous reports of PTH-stimulated cAMP production in these cells (34).

In OK cells shown here, and in PTC (17), PTH produces an increase in [Ca<sup>2+</sup>]<sub>i</sub>. Furthermore, in PTC and OK cells whose plasma membrane has been permeabilized by saponin, ATP produces an accumulation of calcium in a nonmitochondrial cellular pool. IP<sub>3</sub> releases calcium rapidly and transiently from this pool. Taken together, the data presented indicate that PTH-associated receptor binding stimulates the hydrolysis of PtdIns(4,5)-P<sub>2</sub> producing an increase in IP<sub>3</sub> and DAG. These substances have been shown to be important second messages for other neurotransmitters and hormones (1–10). The IP<sub>3</sub> may participate in the production of a calcium transient that would activate calmodulin-dependent processes. The finding that PTH stimulated calmodulin-dependent protein phosphorylation supports the contention that PTH activates the calmodulin-dependent pathway of cell activation through production of a calcium transient.

To relate IP<sub>3</sub> production and PTH-stimulated Ca<sup>2+</sup> transients to biologic effects in PTC, we have studied PTC membrane topography (42) and gluconeogenesis (43). In these studies, we were able to differentiate the effects of Ca<sup>2+</sup> transients from those of increased cAMP levels. For instance, increases in [Ca<sup>2+</sup>]<sub>i</sub> produce a transient shortening and rarefaction of PTC brush border membrane microvilli, whereas cAMP produces elongation (42). The effect of PTH on brush border membrane topography is biphasic, immediate shortening and rarefaction followed by recovery and elongation after 5 min. In contrast, agents such as angiotensin II and  $\alpha_1$ -adrenergic agents produced microvillar shortening and rarefaction with much longer recovery times. These latter agents are thought to produce cell activation through IP<sub>3</sub>-induced Ca<sup>2+</sup> transients and Ca<sup>2+</sup>-dependent protein kinases (C-kinase and calmodulin-dependent kinase) (1, 2, 4) and do not stimulate cAMP production. In addition, calmodulin inhibitors abolished the effects of PTH.

PTH and  $\alpha_1$ -adrenergic agonists stimulated PTC to become gluconeogenic (43, 44). This action of PTH and  $\alpha_1$  agonists was mimicked by Ca<sup>2+</sup> ionophores and prevented by buffering the increase in [Ca<sup>2+</sup>]<sub>i</sub> produced by these agents. Whereas high doses of cAMP also stimulated gluconeogenesis, PTH in the presence of an intracellular Ca<sup>2+</sup> buffering agent was unable to stimulate gluconeogenesis despite stimulation of cAMP production. Thus, the data suggest that the Ca<sup>2+</sup> transient produced by PTH may be sufficient to produce activation of certain metabolic processes independent of cAMP.

Activation of multiple mechanisms of cell signaling by PTH may be important in regulating the cellular response to any one signal. Rasmussen et al. have suggested that interaction of cellular messages stimulated by hormonal activation is an important feature of cell activation (4, 45). Modulation of one message by another affects the response to each message. In the setting of the proximal nephron, activation of adenylate cyclase and cAMP production may serve to inhibit the effect of diacylglycerol production. Diacylglycerol activation of protein kinase C has been shown to be the mechanism stimulating cell alkalization




Figure 9. The effect of PTH on  $\text{Ca}^{2+}$ -calmodulin-dependent BLM protein phosphorylation. BLM were phosphorylated as described in Methods in the presence or absence of  $\text{Ca}^{2+}$ -calmodulin, and PTH. Autoradiograms from two separate experiments are shown (lanes 1–

12). The additions for each lane are listed under the respective lane. W13 is a specific calmodulin antagonist. Similar results were observed in experiments using W7, another calmodulin antagonist.

through the sodium/hydrogen antiport (46). This stimulation of the sodium/hydrogen antiport producing cell alkalization has been shown to be an important step in cell growth and hypertrophy (47, 48). PTH has been shown to inhibit the sodium/hydrogen antiporter of this cell and to produce cellular acidification (49). This effect of PTH has been suggested to be regulated through cAMP-stimulated phosphorylation. Furthermore, stimulation of adenylate cyclase activity and production of cAMP participates in many of the effects of PTH on transport functions of the PTC (31, 44, 49). However, PTH-stimulated changes in topography of the apical membrane and its effects on gluconeogenesis suggest important roles of the calcium messenger system in signal transduction induced by PTH. Further studies will be required to fully elucidate the role of  $\text{IP}_3$  and  $\text{PtdIns}(4,5)\text{-P}_2$  turnover in the control of proximal tubular function by PTH.

The mechanism whereby PTH and its receptor couple to both adenylate cyclase and phospholipase C also remains to be elucidated. Receptor coupling to adenylate cyclase is through the guanine nucleotide binding protein ( $G_s$ , reference 50). The  $\alpha$  subunit of  $G_s$  confers the stimulatory effect, but the  $\beta$ - $\gamma$  subunit may induce an inhibitory effect on another guanine nucleotide binding protein ( $G_i$ ). The  $G_i$  protein stimulates the activity of phospholipase-C in some systems (50, 51). For PTH, if, as most of the available experimental data suggests a single class of PTH receptors exist in the BLM of renal tubular cells (13, 14), then signal transduction through a single receptor is responsible for activation of both adenylate cyclase and  $\text{PtdIns}(4,5)\text{-P}_2$  phospholipase C. This unique mechanism of cell

stimulation might occur through association of the PTH-receptor complex with both  $G_s$  and  $G_i$  proteins in the plasma membrane in different degrees. Increased levels of both  $\alpha$  subunits of the two proteins could result in dual pathway activation. However, recent data may indicate that a low affinity receptor for PTH is present in the PTC membrane (52). If this is in fact the case, signal transduction by PTH may occur through two receptors linked to different enzymatic mechanisms. This may also relate to our previous observation that PTH increases turnover of the  $\text{PtdIns}$  cycle in the apical membrane of the PTC (18). Thus, while the present study identifies PTH stimulation of  $\text{IP}_3$  and DAG production as a mechanism of signal transduction, the mechanism of receptor coupling to phospholipase C and the biologic roles of these signals await further study.

### Acknowledgments

We thank Allan Pollock for providing us with the OK cell line. We also thank Helen Odle for her excellent secretarial assistance.

This work was supported by a grant from the National Institutes of Health AM-09976 and AM-32087. Dr. Esbrit was a Fogarty International Fellow. Dr. Hruska was an established investigator of the American Heart Association during a portion of this work.

### References

- Berridge, M. J. 1984. Inositol triphosphate and diacylglycerol as second messengers. *Biochem. J.* 220:345–360.

2. Berridge, M. J., and R. F. Irvine. 1984. Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature (Lond.)*. 312:315–321.
3. Wilson, D. B., T. E. Bross, W. R. Sherman, R. A. Berger, and P. W. Majerus. 1985. Inositol cyclic phosphates are produced by cleavage of phosphatidylinositols (polyphosphoinositides) with purified sheep seminal vesicle phospholipase C enzymes. *Proc. Natl. Acad. Sci. USA*. 82:4013–4017.
4. Rasmussen, H. 1986. The calcium messenger system. *New Engl. J. Med.* 314:1164–1170.
5. Joseph, S. K., A. P. Thomas, R. J. Williams, R. F. Irvine, and J. R. Williamson. 1984. Myo-inositol 1,4,5-trisphosphate. *J. Biol. Chem.* 259:3077–3081.
6. Prentki, M., T. J. Biden, D. Janjic, R. F. Irvine, M. J. Berridge, and C. B. Wollheim. 1984. Rapid mobilization of  $\text{Ca}^{2+}$  from rat insulinoma microsomes by inositol-1,4,5-trisphosphate. *Nature (Lond.)*. 309:562–564.
7. Muallem, S., M. Schoeffield, S. Pandol, and G. Sachs. 1985. Inositol trisphosphate modification of ion transport in rough endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA*. 82:4433–4437.
8. Kojima, I., K. Kojima, and H. Rasmussen. 1985. Characteristics of angiotensin II-,  $\text{K}^{+}$ - and ACTH-induced calcium influx in adrenal glomerulosa cells. *J. Biol. Chem.* 260:9171–9176.
9. Putney, J. W., Jr. 1986. A model for receptor-regulated calcium entry. *Cell Calcium*. 7:1–12.
10. Albert, P. R., and A. H. Tashjian, Jr. 1984. Thyrotropin-releasing hormone-induced spike and plateau in cytosolic free  $\text{Ca}^{2+}$  concentrations in pituitary cells. *J. Biol. Chem.* 259:5827–5832.
11. Bates, M. D., and P. M. Conn. 1984. Calcium mobilization in the pituitary gonadotrope: relative roles of intra- and extracellular sources. *Endocrinology*. 115:1380–1385.
12. Michell, R. H. 1975. Inositol phospholipids and cell surface receptor functions. *Biochim. Biophys. Acta*. 415:81–147.
13. Bellorin-Font, E., and K. J. Martin. 1981. Regulation of the PTH-receptor-cyclase system of canine kidney: effects of calcium, magnesium, and guanine nucleotides. *Am. J. Physiol.* 241:F364–F373.
14. Nissenson, R. A., and C. D. Arnaud. 1979. Properties of the parathyroid hormone receptor-adenylate cyclase system in chicken renal plasma membranes. *J. Biol. Chem.* 254:1469–1475.
15. Segre, G. V., M. Rosenblatt, B. L. Reiner, J. E. Mahaffey, and J. T. Potts, Jr. 1979. Characterization of parathyroid hormone receptors in canine renal cortical plasma membranes using a radioiodinated sulfur-free hormone analogue. *J. Biol. Chem.* 254:6980–6986.
16. Chase, L. R., and G. D. Aurbach. 1967. Parathyroid function and the renal excretion of 3'5' adenylic acid. *Proc. Natl. Acad. Sci. USA*. 58:518–525.
17. Hruska, K. A., M. Goligorsky, J. Scoble, M. Tsutsumi, S. Westbrook, and D. Moskowitz. 1986. Effects of parathyroid hormone on cytosolic calcium in renal proximal tubular primary cultures. *Am. J. Physiol.* 251:F188–F198.
18. Khalifa, S., S. Mills, and K. A. Hruska. 1983. Stimulation of calcium uptake by parathyroid hormone in renal brush-border membrane vesicles. *J. Biol. Chem.* 258:14400–14406.
19. Bidot-Lopez, P., R. V. Farese, and M. A. Sabir. 1981. Parathyroid hormone and adenosine-3',5'-monophosphate acutely increases phospholipids of the phosphatidate-polyphosphoinositide pathway in rabbit kidney cortex tubules in vitro by a cycloheximide-sensitive process. *Endocrinology*. 108:2078–2081.
20. Meltzer, V., S. Weinreb, E. Bellorin-Font, and K. A. Hruska. Parathyroid hormone stimulation of renal phosphoinositide metabolism is a cyclic nucleotide-independent effect. 1982. *Biochim. Biophys. Acta*. 712:258–267.
21. Lo, H., D. C. Lehotay, D. Katz, and G. S. Levey. 1976. Parathyroid hormone-mediated incorporation of  $^{32}\text{P}$ -orthophosphate into phosphatidic acid and phosphatidylinositol in renal cortical slices. *Endocr. Res. Commun.* 3(Suppl. 6):377–385.
22. Grove, R. I., D. Fitzpatrick, and S. D. Schimmel. 1981. Effect of  $\text{Ca}^{++}$  on triphosphoinositide extraction in fusing myoblasts. *Lipids*. 16:691–693.
23. Berridge, M. J., C. P. Downes, and M. R. Hanley. 1982. Lithium amplifies agonist-dependent phosphatidylinositol responses in brain and salivary glands. *Biochem. J.* 206:587–595.
24. Berridge, J. J. 1983. Rapid accumulation of inositol trisphosphate reveals that agonists hydrolyse polyphosphoinositides instead of phosphatidylinositol. *Biochem. J.* 212:849–858.
25. Clarke, N. G., and R. M. C. Dawson. 1981. Alkaline O leads to N-transacylation. A new method for the quantitative deacylation of phospholipids. *Biochem. J.* 195:301–306.
26. Grynkiewicz, G., M. Poenie, and R. Y. Tsien. 1985. A new generation of  $\text{Ca}^{2+}$  indicators with greatly improved fluorescence properties. *J. Biol. Chem.* 260:3440–3450.
27. Reid, I. R., R. Civitelli, L. R. Halstead, L. V. Avioli, and K. A. Hruska. 1986. Effect of PTH on  $\text{Ca}^{2+}$  and  $\text{pH}_i$  of osteoblast-like cells. *Clin. Res.* 34:432A. (Abstr.)
28. Bers, D. M. 1982. A simple method for the accurate determination of free  $[\text{Ca}]$  in Ca-EGTA solutions. *Am. J. Physiol.* 242:C404–C408.
29. Scoble, J. E., S. Mills, and K. A. Hruska. 1985. Calcium transport in canine renal basolateral membrane vesicles: Effects of parathyroid hormone. *J. Clin. Invest.* 75:1096–1105.
30. Hruska, K. A., S. C. Mills, S. Khalifa, and M. R. Hammerman. 1983. Phosphorylation of renal brush-border membrane vesicles. *J. Biol. Chem.* 258:2501–2507.
31. Hammerman, M. R., and K. A. Hruska. 1982. Cyclic AMP dependent protein phosphorylation in canine renal brush border membrane vesicles is associated with decreased phosphate transport. *J. Biol. Chem.* 257:992–999.
32. Labarca, C., and K. Paigen. 1980. A simple, rapid, and sensitive DNA assay procedure. *Anal. Biochem.* 102:344–352.
33. Shayman, J. A., and A. R. Morrison. Bradykinin-induced changes in phosphatidyl inositol turnover in cultured rabbit papillary collecting tubule cells. *J. Clin. Invest.* 76:978–984.
34. Irvine, R. F., A. J. Letcher, D. J. Lander, and C. P. Downes. 1984. Inositol trisphosphates in carbachol-stimulated rat parotid glands. *Biochem. J.* 223:237–243.
35. Irvine, R. F., A. J. Letcher, J. P. Heslop, and M. J. Berridge. 1986. The inositol tris tetrakisphosphate pathway-demonstration of  $\text{Ins}(1,4,5)\text{P}_3$  3-kinase activity in animal tissues. *Nature (Lond.)*. 320:631–634.
36. Teitelbaum, A. P., and G. J. Stewler. 1984. PTH receptors coupled to cyclic adenosine monophosphate formation in an established renal cell line. *Endocrinology*. 114:980–985.
37. Bonventre, J. V., and J. Y. Cheung. 1986. Cytosolic free calcium concentration in cultured renal epithelial cells. *Am. J. Physiol.* 250:F329–F338.
38. Somlyo, A. P. 1984. Cellular site of calcium regulation. *Nature (Lond.)*. 309:516–517.
39. Fain, J. N., S.-H. Lin, I. Litosch, and M. Wallace. 1985. Relationship between vasopressin activation of rat hepatocyte glycogen phosphorylase, inhibition of  $\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ -ATPase and phosphoinositide breakdown. In *Inositol and Phosphoinositides*. J. E. Bleasdale, J. Eichberg, and G. Hauser, editors. Humana Press, Clifton, NJ. 255–274.
40. Nishizuka, Y., Y. Takai, A. Kishimoto, U. Kikkawa, and K. Kaibuchi. 1984. Phospholipid turnover in hormone action. *Recent Prog. Horm. Res.* 40:301–345.
41. Palmer, F. B. St. C., and S. E. Mack. 1985. Hydrolysis of phosphatidylinositol 4-phosphate and phosphatidylinositol(4,5)-bisphosphate by specific phosphomonoesterases in human erythrocytes. In *Inositol and Phosphoinositides*. J. E. Bleasdale, J. Eichberg, and G. Hauser, editors. Humana Press, Clifton, NJ. 199–212.
42. Goligorsky, M. S., D. N. Menton, and K. A. Hruska. 1986. Parathyroid hormone-induced changes of the brush border topography and cytoskeleton in cultured renal proximal tubular cells. *J. Membr. Biol.* 92:151–162.

43. Goligorsky, M. S., K. A. Hruska, and I. Karl. 1986. Hormonal control of gluconeogenesis in primary cultures of renal proximal tubular cells: role of calcium. *Clin. Res.* 24:596A. (Abstr.)
44. Klahr, S., and K. A. Hruska. 1983. Effects of parathyroid hormone on the renal reabsorption of phosphorus and divalent cations. In *Advances in Bone and Mineral Research*. W. A. Peck, editor. Elsevier/North Holland, Amsterdam. 65-124.
45. Rasmussen, H., and D. B. P. Goodman. 1977. Relationships between calcium and cyclic nucleotides in cell activation. *Physiol. Rev.* 57:421-509.
46. May, W. S., E. G. Lapetina, and P. Cuatrecasas. 1986. Intracellular activation of protein kinase C and regulation of the surface transferrin receptor by diacylglycerol is a spontaneously reversible process that is associated with rapid formation of phosphatidic acid. *Proc. Natl. Acad. Sci. USA.* 83:1281-1284.
47. Allemain, G. L., A. Franchi, E. Cragoe, Jr., and J. Pouyssegur. 1984. Blockade of the  $\text{Na}^+/\text{H}^+$  antiport abolishes growth factor-induced DNA synthesis in fibroblasts. *J. Biol. Chem.* 259:4313-4319.
48. Allemain, G. L., S. Paris, and J. Pouyssegur. 1984. Growth factor action and intracellular pH regulation in fibroblasts. *J. Biol. Chem.* 259:5809-5815.
49. Pollock, A. S., D. G. Warnock, and G. J. Stewler. 1986. Parathyroid hormone inhibition of  $\text{Na}^+/\text{H}^+$  antiporter activity in a cultured renal cell line. *Am. J. Physiol.* 250:F217-F225.
50. Litosch, I., and J. N. Fain. 1986. *Minireview*: regulation of phosphoinositide breakdown by guanine nucleotides. *Life Sci.* 39:187-194.
51. Blackmore, P. R., S. B. Bocchino, L. E. Waynick, and J. H. Exton. 1985. Role of a guanine nucleotide-binding regulatory protein in the hydrolysis of hepatocyte phosphatidylinositol 4,5-bisphosphate by calcium-mobilizing hormones and the control of cell calcium. *J. Biol. Chem.* 260:14477-14483.
52. Rouleau, M. F., J. Mitchell, A. Tenenhouse, and D. Goltzman. 1986. In vivo and in vitro demonstration of parathyroid hormone receptor down-regulation in vitamin D deficient rat. *J. Bone & Min. Res.* 1(Suppl. 1):322. (Abstr.)