

Effect of formate on volume reabsorption in the rabbit proximal tubule.

L Schild, ... , L P Karniski, P S Aronson

J Clin Invest. 1987;79(1):32-38. <https://doi.org/10.1172/JCI112803>.

Research Article

Studies on microvillus membrane from rabbit kidney cortex suggest that chloride absorption may occur by chloride/formate exchange with recycling of formic acid by nonionic diffusion. We tested whether this transport mechanism participates in active NaCl reabsorption in the rabbit proximal tubule. In proximal tubule S2 segments perfused with low HCO₃ solutions, the addition of formate (0.25-0.5 mM) to the lumen and the bath increased volume reabsorption (JV) by 60%; the transepithelial potential difference remained unchanged. The effect of formate on JV was completely reversible and was inhibited both by ouabain and by luminal 4,4'-diisothiocyanostilbene-2,2'-disulfonate. Formate (0.5 mM) failed to stimulate JV in early proximal convoluted tubules perfused with high HCO₃ solutions. As measured by miniature glass pH microelectrodes, this lack of formate effect on JV was related to a less extensive acidification of the tubule fluid when high HCO₃ solutions were used as perfusate. These data suggest that chloride/formate exchange with recycling of formic acid by nonionic diffusion represents a mechanism for active, electroneutral NaCl reabsorption in the proximal tubule.

Find the latest version:

<https://jci.me/112803/pdf>



Effect of Formate on Volume Reabsorption in the Rabbit Proximal Tubule

Laurent Schild, Gerhard Giebisch, Lawrence P. Karniski, and Peter S. Aronson

Departments of Physiology and Internal Medicine, Yale University School of Medicine, New Haven, Connecticut 06510

Abstract

Studies on microvillus membrane from rabbit kidney cortex suggest that chloride absorption may occur by chloride/formate exchange with recycling of formic acid by nonionic diffusion. We tested whether this transport mechanism participates in active NaCl reabsorption in the rabbit proximal tubule.

In proximal tubule S2 segments perfused with low HCO_3^- solutions, the addition of formate (0.25–0.5 mM) to the lumen and the bath increased volume reabsorption (J_V) by 60%; the transepithelial potential difference remained unchanged. The effect of formate on J_V was completely reversible and was inhibited both by ouabain and by luminal 4,4'-diisothiocyanostilbene-2,2'-disulfonate.

Formate (0.5 mM) failed to stimulate J_V in early proximal convoluted tubules perfused with high HCO_3^- solutions. As measured by miniature glass pH microelectrodes, this lack of formate effect on J_V was related to a less extensive acidification of the tubule fluid when high HCO_3^- solutions were used as perfusate. These data suggest that chloride/formate exchange with recycling of formic acid by nonionic diffusion represents a mechanism for active, electroneutral NaCl reabsorption in the proximal tubule.

Introduction

In the early portion of the proximal tubule of the mammalian kidney, isotonic volume reabsorption (J_V)¹ is associated with preferential reabsorption of bicarbonate and organic solutes (1, 2). This results in increased chloride concentration in the later portion of the proximal tubule and a shift of the transepithelial potential from lumen negative to lumen positive. The generation of a chloride diffusion potential by the transepithelial chloride gradient reflects a relatively large conductance for Cl^- , most probably across the intercellular pathway (3, 4). Given the favorable passive driving force for Cl^- in the late proximal tubule, it is thought that a significant fraction of Cl^- reabsorption occurs passively.

On the other hand, chloride reabsorption across the proximal tubule has also been shown to continue in the absence of any passive driving force for chloride (5, 6). This suggests an active, transcellular mechanism of chloride reabsorption. It has been

proposed that transcellular, electroneutral NaCl transport is involved in this component of chloride reabsorption, but little is known about its specific mechanism (7).

On the basis of studies on rabbit renal cortical microvillus membrane vesicles, Karniski and Aronson have suggested that chloride/formate exchange with recycling of formic acid by nonionic diffusion could be a potential mechanism for active Cl^- transport across the luminal membrane (8). If this anion exchanger were to operate in parallel with Na^+/H^+ antiport, coupled electroneutral NaCl reabsorption would take place.

Our experiments were designed to test whether electroneutral formate/chloride exchange plays a significant role in proximal chloride reabsorption. The present study was performed to test the following four predictions.

Physiological concentrations of formate should reversibly stimulate J_V without changes in the transepithelial potential.

The formate-stimulated J_V should be subject to inhibition both by 4,4'-diisothiocyanostilbene-2,2'-disulfonate (DIDS), an inhibitor of formate/chloride exchange in renal microvillus membrane vesicles, and by ouabain, the inhibitor of active Na transport.

Stimulation of J_V by formate should be demonstrable in the absence of any passive driving force for Cl^- .

Formate-induced water reabsorption should be affected by luminal pH, to the extent that recycling of formic acid by nonionic diffusion is rate limiting for formate/chloride exchange and thus also for NaCl reabsorption.

Our study demonstrates that these four predictions were accurate.

Methods

Preparation and perfusion of isolated tubules in vitro. Segments of rabbit proximal tubules were isolated and perfused in vitro with techniques first described by Burg et al. (9). Briefly, New Zealand White rabbits weighing 1.5–2 kg were used. The right kidney was rapidly removed and cut in coronal slices ~ 0.5 mm thick. The dissection was done at 4°C in an ultrafiltratelike solution containing 1 g/100 ml albumin, buffered at pH 7.4 with Hepes and gassed with 100% O_2 .

Two different segments of the proximal tubule were used in this study. S2 segments, which include the last convolutions and the beginning of the pars recta of the proximal tubule, were dissected using bundles radiating from the cortico-medullary junction to the surface of the cortex. Only tubules reaching the kidney surface, S2 superficial tubules, were used. Early proximal convoluted tubules, defined as S1 segments, were selected on the basis of their attachment to the glomerulus and their localization in the superficial cortex. The dissected tubules were transferred into a temperature-regulated Lucite chamber containing 2 ml of bathing solution at 37°C (see Table I) and mounted between appropriate concentric pipettes. The flow rate of the perfusion was regulated by gravity (hydrostatic pressure, 15–20 cm H_2O) to maintain a tubular perfusion rate between 10 and 20 nl/min. Net J_V (nanoliter · minute⁻¹ · millimeter⁻¹) was determined as the difference between the calculated perfusion rate and the measured collection rate. The collection rate was measured by collecting samples of the perfusate over timed intervals in calibrated constriction pipettes with a volume > 50 nl. Exhaustively dialyzed meth-

Part of this material has been presented in abstract form. (1986. *Kidney Int.*, 29:407a.)

Dr. Karniski's present address is Department of Internal Medicine, University of Iowa, Iowa City, IA 52242.

Received for publication 22 May 1986.

1. *Abbreviations used in this paper:* DIDS, 4,4'-diisothiocyanostilbene-2,2'-disulfonate; J_V , volume reabsorption; V_{te} , transepithelial potential difference.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/87/01/0032/07 \$1.00

Volume 79, January 1987, 32–38

Table I. Composition of Solutions

	Bath		Perfusate	
	Control	Low HCO ₃ ⁻ , high Cl ⁻	Low HCO ₃ ⁻ , high Cl ⁻	High HCO ₃ ⁻ , low Cl ⁻
NaCl	115	135	135	115
NaHCO ₃	20	4	4	20
NaH ₂ PO ₄	4	4	4	4
KCl	5	5	5	5
CaCl ₂	1.8	1.8	1.8	1.8
MgSO ₄	1	1	1	1
Acetate	5	5	—	—
Glucose	5	5	—	—
Alanine	5	5	—	—
Mannitol	—	—	15	15
Albumin	6 g/100 ml	6 g/100 ml	—	—
CO ₂	5%	1%	5% or 1%	5

oxy-³H]inulin (New England Nuclear, Boston, MA) was used as volume marker and added to the perfusate at a concentration of 20 μCi/ml. The perfusion rate was calculated from the collection rate and from the difference of inulin concentration between the perfusate and the collectate.

The compositions of the perfusion solutions and the bath solutions are listed in Table I. S2 segments of proximal tubules were perfused with a high-chloride, low-bicarbonate solution similar in composition to late proximal tubule fluid. Solutions simulating serum and containing 6 g/100 ml albumin were used as bath solution. S1 segments were perfused and bathed with symmetrical solutions containing either 20 or 4 mM HCO₃⁻. To obtain the same pH (7.30 to 7.35), the serumlike bath solutions were bubbled with either 95% O₂-5% CO₂ or 99% O₂-1% CO₂ gas mixture. Dialyzed albumin (BSA Fraction V; Sigma Chemical Co., St. Louis, MO) was added to the bath solution at a concentration of 6 g/100 ml. No organic substrates were present in the perfusion solutions. All solutions were adjusted to a total osmolality of 300 mosmol/kg H₂O by adding H₂O or NaCl salt.

During the experiments, bathing solutions continuously flowed through the chamber at a rate of ~0.3 ml · min⁻¹. The solutions were preheated through tubing at a temperature of 40°C, and the bath temperature was maintained in the chamber at 38°C with a specially designed temperature-regulating system. The bath was continuously bubbled with the appropriate gas mixtures, and its pH was continuously monitored with a miniature pH electrode (Beetode; World Precision Instruments, Inc., New Haven, CT) placed in the vicinity of the tubule.

The protocols were designed to study the effects of formate and different transport inhibitors on water reabsorption and the transepithelial potential difference in the proximal tubule. The first collection began after an equilibration time of 30 min, to be followed by several experimental periods. Before starting collections, the holding pipette was carefully rinsed and the first collected sample was discarded. The tubular collections were performed for at least 10 min to obtain a minimum of two samples per period. Tubules were discarded if irreversible morphological changes such as cell swelling or cell granulation occurred during the experiment.

Transepithelial potential difference measurements. The transepithelial potential difference (V_{te}) was measured as in previous studies performed in this laboratory (10). The perfusion pipette was used as a bridge to the tubular lumen; the perfusion solution and the bath were connected to calomel electrodes via 150 mM NaCl-agarose bridges. The potential difference of this circuit was measured with a high-impedance electrometer (model 750; World Precision Instruments, Inc.). The output of the electrometer was displayed on a digital voltmeter (Newport Electronics, Inc., Santa Ana, CA) and a strip chart recorder (model 220; Gould, Inc., Cleveland, OH). With no tubule in place this system was stable for several hours. The reference baseline potential, measured at 38°C in the absence

of a tubule, never exceeded 0.5 mV. Correction of the liquid junction potential was made in the following way. Before mounting a tubule between the pipettes, the measured small circuit potential was compensated and the potential was set to zero when the chamber was filled with the bath at 37°C. The perfusion pipette contained the perfusate. The tubule was then mounted and cannulated, and the change in circuit potential was recorded as the transepithelial potential. Tubules were perfused uniformly throughout the experiments with solutions of identical ionic composition, or with such additions as submillimolar concentrations of transport inhibitors and formate or acetate. Accordingly, corrections for changes in liquid junction potentials were minimized in our experimental settings.

pH measurements of the tubular fluid. pH-sensitive glass electrodes were used to measure the pH of the tubular fluid at the distal end of the perfused tubule. The pH-sensitive glass electrodes were constructed by fusing pH-sensitive glass (World Precision Instruments, Inc.) to the tip of a non-pH-sensitive glass capillary (Drummond Scientific Co., Broomall, PA) to form a bulb with a diameter of ~40–50 μm. This miniature pH electrode was filled with a citrate pH buffer (pH 6) and connected via an electrode holder (World Precision Instruments, Inc.) to a high impedance amplifier (model 750; World Precision Instruments, Inc.). The electrodes were calibrated before and after the experiments in a standard buffer solution. The slope of the pH electrode was 59.09 ± 1.8 mV/pH U (mean ± SD). The pH electrode assembly was inserted into the collecting pipette. Its shape was modified by a constriction at the tip of the pipette to accommodate the pH electrode. This made it possible to measure the pH of the emerging tubular fluid in a small compartment (100 nl). This minimized loss of CO₂ from the collected fluid (11). The pH of the initial perfusion fluid was measured by two methods. The first was to measure pH in the collection pipette assembly when tubules were perfused under pressure at very high flow rate (in excess of 600 nl/min), because pH changes by tubule activity are virtually abolished under these conditions. This was demonstrated directly, as the pH values so obtained were not different from values measured with the miniature glass pH electrode in the initial perfusion fluid, bubbled with the appropriate gas mixture.

Statistics. The results are expressed as the mean ± SE of individual measurements. In general, paired experiments were performed. Thus, when appropriate, the mean paired *t* test was used (12).

Results

A summary of the effect of formate on transepithelial water reabsorption and on V_{te} across the S2 segment of the proximal tubule is presented in Fig. 1. Significant net fluid absorption is apparent during perfusion of S2 segments with high-chloride-low-bicarbonate solutions and a bath solution containing 20 mM HCO₃⁻. Addition of formate in submillimolar concentrations, 0.5 and 0.25 mM to the bath and the lumen, increased J_v significantly, from 0.58 ± 0.08 nl · min⁻¹ · mm⁻¹ (mean ± SE) to 0.94 ± 0.08 nl · min⁻¹ · mm⁻¹ (mean ± SE) and from 0.38 ± 0.03 nl · min⁻¹ · mm⁻¹ (mean ± SE) to 0.63 ± 0.04 nl · min⁻¹ · mm⁻¹ (mean ± SE), respectively.² Since the perfusion solution contains

2. The mean value of J_v obtained in the S2 segment was 0.47 nl/min · mm ± 0.05 nl/min · mm (mean ± SE). This value is between 0.56 nl/min · mm, reported by Berry in the proximal convoluted tubule (1983, *J. Clin. Invest.* 71:268–281.) and 0.38 nl/min · mm, reported by Schafer et al. in the pars recta of rabbit kidney (1981, *Kidney Int.* 20:588–597.) It is likely that our mean value of J_v obtained in the S2 segment reflects the transition between the proximal convoluted and straight tubule. We also note the difference in J_v between the two control groups using low and high formate concentration. We have no explanation for this difference, but the design of our experiment includes the use of paired samples, each tubule serving as its own control.




Figure 1. Effect of formate, 0.5 mM ($n = 10$) and 0.25 mM ($n = 13$), added to the bath and the lumen on J_v and V_{te} . S2 segments of the proximal tubule were perfused with low HCO_3^- solutions simulating late proximal tubule fluid. The bath contained 20 mM HCO_3^- . In the control conditions and after the addition of formate the mean perfusion rates were 14.27 ± 4 and 14.56 ± 2.2 nl/min (mean \pm SE), respectively.

mainly NaCl, we conclude that the increase in J_v reflects stimulation of NaCl transport.

The V_{te} was positive during the control period and remained unchanged after the addition of formate, 0.5 and 0.25 mM, to the bath and lumen. The lumen positive potential under these experimental conditions reflects the diffusion potential due to the passive, intercellular movement of Cl^- across the epithelium (4, 5, 7, 13). The failure of V_{te} to change in parallel with the increase in J_v suggests that formate stimulates an active, electroneutral component of NaCl transport, and that formate does not affect the passive paracellular component of Cl^- movement across the epithelium.

As shown in Fig. 2, the increase in water reabsorption induced by formate was fully reversible. Thus, formate at the concentrations used in these experiments did not cause irreversible alteration of the tubular epithelium. To rule out a nonspecific metabolic effect as responsible for stimulation of J_v by formate, we measured J_v after addition of 0.5 mM acetate to the bath and the lumen. As illustrated in Fig. 3, 0.5 mM acetate had no significant effect on transepithelial water reabsorption. These data suggest that formate increases the reabsorption of water in the proximal tubule by stimulating a specific electroneutral transport mechanism.

The effects of two transport inhibitors, ouabain and DIDS, were also investigated. As shown in Figs. 4 and 5, these agents significantly modify the effects of formate on proximal fluid transport. The results of experiments performed on S2 segments perfused with solutions simulating late proximal tubule fluid are shown. Ouabain, at a concentration of 0.05 mM, suppressed the increase in J_v due to addition of formate to the bath and the lumen. Transepithelial fluid reabsorption was not completely abolished by ouabain because a favorable passive driving force for NaCl absorption was present (4, 14). However, J_v in the presence of ouabain was significantly lower ($P = 0.018$) than that observed during the control period in which active transport was present. The transepithelial potential difference remained




Figure 2. Reversibility of the effect of formate (0.5 and 0.25 mM) in the bath and lumen on J_v and V_{te} . S2 segments of the proximal tubule were perfused with low HCO_3^- solutions and bathed with a solution containing 20 mM HCO_3^- .

positive and unchanged after inhibition of active transport by ouabain. Again this confirms that an important component of NaCl transport is active and electroneutral and can be stimulated by formate. The addition of the anion-exchange inhibitor DIDS to the tubular lumen at a concentration of 1 mM, also suppressed the stimulation of J_v induced by formate. This inhibitory effect of DIDS is consistent with participation of a specific anion-exchange mechanism in the stimulation of NaCl transport by formate. Interestingly, water reabsorption during perfusion with luminal DIDS was significantly lower than that in the control period ($P = 0.003$). The observation that the lumen positive transepithelial potential difference dropped to 0 after addition of DIDS suggests that DIDS, at a concentration of 1 mM, may decrease the passive movement of chloride via the paracellular shunt pathway in addition to inhibiting the luminal membrane anion exchanger. This interpretation is consistent with the suppression of J_v below control values by DIDS, because a reduction of the transepithelial chloride permeability would interfere with that component of transepithelial fluid movement driven by the chloride gradient.

The experiments described so far were performed in the S2 proximal tubules perfused with a solution simulating late prox-




Figure 3. Effect of addition of 0.5 mM acetate to the perfusate and the bath on J_v . S2 segments of the proximal tubule were perfused with low HCO_3^- solutions and bathed in a solution containing 20 mM HCO_3^- .




Figure 4. Effect of ouabain (0.05 mM in the bath) on J_v and V_{te} in S2 segments of the proximal tubule in the presence of formate (0.5 and 0.25 mM), in the perfusion fluid and the bath. The perfusion fluid and the bath contained 4 and 20 mM HCO_3^- , respectively.




Figure 6. Effect of formate (0.5 and 0.25 mM) on J_v and V_{te} in the proximal convoluted tubule (S1 segment), perfused and bathed with symmetrical low HCO_3^- solutions (4 mM).

imal fluid. In a second protocol we examined the ability of formate to stimulate NaCl reabsorption in the early proximal convoluted tubule (segment S1), in which transepithelial gradients of Cl and bicarbonate were initially absent.

Early proximal convoluted tubules were perfused and bathed with symmetrical solutions high in NaCl and low in HCO_3^- (Table I). Under these conditions, in the presence of a low HCO_3^- concentration in the lumen and the bath (4 mM), it is unlikely that a significant chloride gradient could develop across the proximal tubule due to the preferential reabsorption of HCO_3^- (14). As shown in Fig. 6, formate in concentrations of 0.5 and 0.25 mM in the bath and the lumen increased reversibly the reabsorption of fluid. V_{te} was not significantly different from zero (-0.36 ± 0.13 mV, mean \pm SE). The absence of a significant lumen negative potential indicates that under these conditions the passive driving force for transepithelial chloride movement was minimal. Changes in V_{te} were not observed after addition of formate.

We also tested the effect of formate on fluid movement across the proximal convoluted tubule under conditions in which the HCO_3^- concentration in the bathing and in the perfusion solutions was 20 mM. As shown in Fig. 7, in the presence of such high HCO_3^- concentrations in the tubule fluid, addition of for-

mate, 0.5 mM in bath and lumen, failed to stimulate significantly the reabsorption of fluid.

It has been pointed out elsewhere (8) that in order for chloride/formate exchange to drive significant amounts of Cl^- into the cell, recycling of formate from the lumen to cell is required. One possible mechanism for such formate recycling into the cell is nonionic back diffusion of formic acid across the apical membrane (8). Hence inefficient recycling of formic acid at a high luminal pH could be responsible for the lack of effect of formate observed in tubules perfused with high HCO_3^- solutions.

In order to test this hypothesis we performed pH measurements in the tubular fluid. As summarized in Table II, different pH levels were obtained in the perfusion experiments with different HCO_3^- solutions. In proximal convoluted tubules perfused and bathed with symmetrical solutions containing 20 mM HCO_3^- , the pH of the perfusate decreased from 7.32 to 7.12 along the tubule. In this experimental setting, formate failed to stimulate water reabsorption. On the other hand, in experiments in which tubules were perfused and bathed with symmetrical low HCO_3^- solutions (4 mM HCO_3^-), the fall in luminal pH was greater, from 7.34 to 6.85. Under this condition formate stimulated fluid reabsorption. These observations support the suggestion that pH-sensitive recycling of formic acid across the apical membrane may be rate limiting for the stimulation of NaCl transport by formate in the proximal tubule.




Figure 5. Effect of DIDS (1 mM in the perfusate) on J_v and V_{te} in the presence of formate (0.5 and 0.25 mM) in the perfusion fluid and the bath. S2 segments of the proximal tubule were perfused with a low HCO_3^- solution, and bathed in a 20 mM HCO_3^- solution.




Figure 7. Effect of formate (0.5 mM) on J_v and V_{te} in the proximal convoluted tubule (S1 segment), perfused and bathed with symmetrical high HCO_3^- solutions (20 mM).

Table II. pH Data

	pH in perfusion fluid	pH in collected fluid	pH in bath	Volume perfusion nl/min	pH difference (perfusion, collected fluid)
4 mM HCO ₃ 1% CO ₂	7.34±0.02	6.85±0.05	7.34±0.02	13.18±1.40	0.49
20 mM HCO ₃ 5% CO ₂	7.32±0.02	7.12±0.03	7.35±0.01	12.97±1.9	0.20
4 mM HCO ₃ 1% CO ₂	7.32±0.02	6.82±0.02	7.32±0.02	13.12±0.9	0.50

Direct pH measurements were performed in seven paired convoluted proximal tubules, first perfused and bathed with symmetrical low HCO₃⁻ solutions, then with solutions containing 20 mM HCO₃⁻ during the second period. A third period of recovery was performed in five tubules.

Discussion

The major question addressed by this study is whether formate-coupled transport of chloride contributes significantly to transepithelial chloride reabsorption in the proximal tubule. The chloride/formate exchange system and the nonionic diffusion of formic acid have recently been proposed by Karniski and Aronson to provide an electroneutral mechanism for secondary active NaCl reabsorption across the luminal membrane of the rabbit proximal tubule as depicted in Fig. 8 (8). Proton secretion via the Na⁺/H⁺ exchanger allows formic acid to enter the cell by nonionic diffusion and maintains the intracellular formate concentration above electrochemical equilibrium across the luminal membrane. The outwardly directed electrochemical formate gradient represents the driving force for an active Cl⁻ absorption via the chloride/formate exchanger. An interesting feature of this model concerns the recycling of formate from the lumen into the cell by nonionic diffusion. This mechanism potentially allows low concentrations of formate to drive significant amounts of Cl⁻ from the tubular lumen into the cell despite the fact that formate is normally present in the plasma at levels of 0.1 to 1.4 mM (15). Our experiments show that low concentrations of formate in the bath and the lumen stimulate isotonic volume reabsorption, and as predicted by the model, the increase in NaCl reabsorption is sensitive to anion-exchange inhibitors, ouabain, and luminal pH.

It is generally agreed that both passive and active transport contribute to NaCl reabsorption in the proximal tubule. Micro-puncture experiments in the rat proximal tubule concluded that a major fraction of NaCl reabsorption is dependent on active transport processes because metabolic inhibitors reduce water

reabsorption by one half to two thirds in the presence of a normal chloride gradient (6, 14, 16).

There are basically two ways that active Na⁺ transport could affect chloride reabsorption. First, simple electrogenic Na⁺ transport could result in a lumen negative transepithelial potential difference that would drive passive chloride flux through the paracellular pathway. For instance, Cardinal et al. observed in the superficial proximal tubule that in the presence of a chloride gradient, ouabain shifts the positive transtubular potential in the positive direction, thus suggesting a component of simple electrogenic Na⁺ transport (17). Similar results suggesting the presence of an electrogenic Na⁺ transport have been reported in the rabbit pars recta (4). Second, active Na⁺ transport may be coupled to secondary active transcellular Cl⁻ transport. Recently, Baum and Berry, studying the electrical nature of NaCl transport in the rabbit proximal convoluted tubule, showed that ouabain decreased the active transport of NaCl without affecting the transepithelial potential (7). They concluded that active NaCl is electroneutral and transcellular.

In agreement with the later observations, our experiments presented in Fig. 4 indicate that reabsorption of NaCl in the S2 segment of the proximal tubule from a high-chloride solution is stimulated by formate and inhibited by ouabain. Both of these events occur without changes in the transepithelial potential. The active component of NaCl reabsorption inhibited by ouabain in these experiments is clearly electroneutral. Failure to find any difference in the V_{te} suggests that simple electrogenic Na⁺ transport providing the electrochemical driving force for passive Cl⁻ reabsorption cannot account for the active, ouabain-sensitive component of NaCl reabsorption. These results differ from those reported by Schafer et al. for the pars recta (18) and suggest that intrinsic differences regarding the mechanism of NaCl reabsorption may exist along the proximal tubule. Instead this finding, in agreement with the result reported by Baum and Berry in the proximal convoluted tubule, supports the notion that the active component of NaCl transport is an electroneutral and transcellular process stimulated by formate. It should be pointed out that ouabain did not completely abolish the transepithelial water flux (Fig. 4). This ouabain-insensitive component of J_v probably reflects the passive diffusion of Cl⁻ due to the transepithelial Cl⁻ gradient present in these experiments.

The experiments performed on the early proximal convoluted tubule (S1 segment) perfused and bathed with symmetrical high-chloride solutions are noteworthy in two respects. First they indicate that the increase in J_v due to formate accounts only for a stimulation of the reabsorption of NaCl. For instance, if formate would stimulate HCO₃⁻ reabsorption, then the transepi-




Figure 8. Model proposed by Karniski and Aronson (8) for the role of chloride/formate exchange in mediating Na-coupled active reabsorption of Cl across the luminal membrane of the proximal tubule cell.

thelial HCO_3^- flux, calculated from the difference in J_V before and after the addition of formate, would exceed the quantity of HCO_3^- perfused into the tubular lumen.³ Second, these experiments constitute additional evidence for an active transcellular NaCl reabsorption stimulated by formate. Under these conditions where no passive driving forces are present for Na^+ and Cl^- reabsorption, formate reversibly stimulated an electrically silent NaCl reabsorption.

A possible mode of sodium-coupled electroneutral Cl^- transport involves parallel ion exchangers such as Na^+/H^+ and Cl^-/OH^- or $\text{Cl}^-/\text{HCO}_3^-$ as identified in the Necturus proximal tubule (19). The presence of a $\text{Cl}^-/\text{HCO}_3^-$ or Cl^-/OH^- exchange has not been reproducibly demonstrated in microvillus membrane vesicles isolated from rabbit proximal tubule cells (20, 21), and Schwartz was unable to demonstrate the presence of this parallel anion exchange system in the intact rabbit proximal tubule (22). Chloride/formate exchanger in parallel with the Na^+/H^+ exchanger represents an alternative mechanism for transcellular sodium-coupled Cl^- transport.

There are several possible ways for formate to stimulate water reabsorption in the proximal tubule. An increase in NaCl transport related to stimulation of cell metabolism can be excluded on the basis of the lack of effect of acetate on water reabsorption. Acetate is involved in generating substrates for cellular metabolism but does not share the chloride/formate exchanger (8). Moreover, luminal DIDS abolished the NaCl reabsorption stimulated by formate. This suggests that a specific anion exchange mechanism located at the luminal membrane is involved in the active NaCl reabsorption induced by formate.

Luminal DIDS also resulted in a drop in the transepithelial diffusion potential for Cl^- and in a lower J_V observed in the presence of luminal DIDS ($0.35 \text{ nl} \cdot \text{min}^{-1} \cdot \text{mm}^{-1}$) compared with the control condition ($0.54 \text{ nl} \cdot \text{min}^{-1} \cdot \text{mm}^{-1}$, $P = 0.003$) in Fig. 5. The reduction of J_V below the control level could reflect the inhibition of another anion exchange like $\text{HCO}_3^-/\text{Cl}^-$ involved in part of the volume reabsorption observed during the control period. An effect of 4-acetamido-4'-isothiocyanostilbene-2,2'-disulfonate on the paracellular permeability pathway similar to the effect of DIDS on V_{te} has been reported previously (13). The persistent volume reabsorption after the addition of DIDS in the lumen is likely due either to an incomplete inhibition of the luminal chloride/formate exchanger, as Karniski and Aronson reported a 60% inhibition of the chloride/formate exchanger by 0.75 mM DIDS in rabbit microvillus membrane vesicles (8), or to incomplete inhibition of passive intercellular NaCl reabsorption.

Since the quantity of Cl^- reabsorbed greatly exceeds any possible net secretion of formate, formate must be recycled from the lumen into the cell for the formate/chloride exchanger to be operative in the active reabsorption of Cl^- . The quantity of Cl^- reabsorbed will therefore depend on formate's ability to be recycled back into the cell. One possible mechanism for the

recycling of formate is the nonionic back diffusion of formic acid, as proposed by Karniski and Aronson. Accounting for the pK_a of formate (3.9), the permeability coefficient for diffusion of formic acid across lipid bilayers ($10^{-2} \text{ cm} \cdot \text{s}^{-1}$) (23), and assuming a luminal pH of 6.8, as measured in the tubular fluid, and an intracellular pH of 7.2, the calculated lumen-to-cell unidirectional flux of formic acid is $\sim 30 \text{ pmol} \cdot \text{min}^{-1} \cdot \text{mm}^{-1}$ when the concentration of formate in the lumen was 0.5 mM. The magnitude of this flux is sufficient to sustain significant volume reabsorption in the proximal tubule.

On the other hand, we observed in the early proximal convoluted segment that formate failed to stimulate NaCl reabsorption when these tubular segments were perfused with a high HCO_3^- solution (Fig. 5), whereas in those perfused with low HCO_3^- solution formate did result in significant stimulation of J_V (Fig. 6). One possible explanation is that the tubular fluid is less effectively acidified during perfusion with a solution of a high buffer content (20 mM bicarbonate) than with a solution with a low buffer content (4 mM bicarbonate), so that the nonionic back diffusion of formic acid is compromised enough to render the chloride/formate mechanism ineffective.

The pH measurements of the tubular fluid (Table II) indicate that the absence of formate-induced NaCl reabsorption when tubules were perfused with a high bicarbonate perfusate can be correlated with a smaller drop in luminal pH. This observation supports the hypothesis that nonionic back diffusion of formic acid into the cell is essential for maintaining an intracellular formate concentration high enough to drive active Cl^- uptake into the cell. However, we cannot exclude the possibility that HCO_3^- directly inhibited the chloride/formate exchanger when tubules were perfused with 20 mM HCO_3^- .

Even if the formate/chloride exchanger represents an important step in the active electroneutral NaCl reabsorption in the proximal tubule, it is still not clear how Cl^- is reabsorbed in the absence of added formate and any electrochemical driving force. For example, in the experiments illustrated in Fig. 6, the small negative transepithelium potential (-0.3 mV) can passively drive at most 10% of the Cl^- reabsorption observed during the control condition, assuming a permeability coefficient (P_{Cl}) for chloride of $0.73 \cdot 10^{-4} \text{ cm} \cdot \text{s}^{-1}$, as reported in the rabbit proximal tubule by Schafer et al (4). One possibility is that the intracellular production of formate is sufficient to sustain a physiologically relevant transcellular Cl^- transport. The concentration of formate produced within proximal tubule cells is unknown.

As discussed previously, another possibility is that $\text{Cl}^-/\text{HCO}_3^-$ exchange takes place. Certainly it is possible that transport systems functional in the intact cell could be inactive in isolated membrane vesicles. Finally, the low apical conductance to chloride in the rabbit proximal tubule (24), and the probability that intracellular Cl^- activity is above its electrochemical equilibrium, as in the rat proximal tubule (25), makes passive diffusion of Cl^- across the luminal membrane very unlikely. Accordingly the mechanism of the baseline electroneutral Cl^- absorption in the absence of added formate requires further investigation.

In summary, we have shown that formate in submillimolar concentrations stimulates active electroneutral NaCl transport in the S1 and S2 segments of the proximal tubules of the rabbit kidney. A specific anion exchange located at the luminal membrane is involved in this transport mechanism. Our results are compatible with the model already proposed where chloride/formate exchange in parallel with Na^+/H^+ antiport allows an

3. Considering a luminal pH of 6.85, as measured in the collected fluid of proximal convoluted tubules perfused and bathed with low HCO_3^- solution under control conditions (Tables I and II), this drop in luminal pH represents a reabsorptive flux of HCO_3^- of $37 \text{ pmol} \cdot \text{min}^{-1} \cdot \text{mm}^{-1}$. If formate stimulates HCO_3^- reabsorption, the increase in J_V would mean an additional $60 \text{ pmol} \cdot \text{min}^{-1} \cdot \text{mm}^{-1}$ of HCO_3^- reabsorbed. Under this condition, HCO_3^- reabsorption would be twofold greater than the quantity of HCO_3^- perfused through the tubule.

electroneutral sodium-coupled, chloride-active reabsorption. This mechanism of active NaCl transport contributes significantly to the transepithelial reabsorption of NaCl in the proximal tubule.

Acknowledgments

We are extremely grateful to Dr. S. K. Agulian for generously supplying us with the manufacture of the pH-sensitive electrodes.

This work was supported by National Institutes of Health grants AM-17433 and AM-33793.

References

1. Burg, M. B. 1986. Renal handling of sodium, chloride, water, amino acids and glucose. In *The Kidney*. B. M. Brenner and F. C. Rector, editors. W.B. Saunders Company, Philadelphia. Third ed. 145-176.
2. Giebisch, G., and P. S. Aronson. 1986. The proximal nephron. In *Physiology of Membrane Disorders*. T. E. Andreoli, J. F. Hoffman, D. D. Fanestil, and S. G. Schultz, editors. Plenum Medical Book Company, New York and London. Second ed. 669-700.
3. Boulpaep, E. L. 1971. Electrophysiological properties of the proximal tubule: importance of cellular and intercellular transport pathways. In *Electrophysiology of Epithelial Cells*. G. Giebisch, editor. F. K. Schattauer Verlag, Stuttgart. 199-246.
4. Schafer, J. A., S. L. Troutman, and T. E. Andreoli. 1974. Volume reabsorption, transepithelial potential differences, and ionic permeability properties in mammalian superficial proximal straight tubules. *J. Gen. Physiol.* 64:582-607.
5. Alpern, R. J., K. J. Howlin, and P. Preisig. 1985. Active and passive components of chloride transport in the rat proximal convoluted tubule. *J. Clin. Invest.* 76:1360-1366.
6. Green, R., J. H. V. Bishop, and G. Giebisch. 1979. Ionic requirements of proximal tubular sodium transport. III. Selective luminal ion substitution. *Am. J. Physiol.* 236:F268-F277.
7. Baum, M., and C. A. Berry. 1984. Evidence for neutral transcellular NaCl transport and neutral basolateral chloride exit in the rabbit proximal convoluted tubule. *J. Clin. Invest.* 14:205-211.
8. Karniski, L. P., and P. S. Aronson. 1985. Chloride/formate exchange with formic acid recycling: a mechanism for active chloride transport across epithelial membranes. *Proc. Natl. Acad. Sci. USA.* 82: 6362-6365.
9. Burg, M. B., J. Grantham, M. Abramow, and J. Orloff. 1966. Preparation and study of fragments of single rabbit nephrons. *Am. J. Physiol.* 210:1293-1298.
10. Biagi, B. A., and G. Giebisch. 1979. Temperature dependence of transepithelial potential in isolated and perfused proximal tubules. *Am. J. Physiol.* 236:F302-F310.
11. Koeppe, B. M., and S. I. Helman. 1982. Acidification of luminal fluid by the rabbit cortical collecting tubule perfused in vitro. *Am. J. Physiol.* 242:F521-F531.
12. Snedecor, G. W., and W. G. Cochran. 1980. *Statistical Methods*. Iowa State University Press, Ames, IA. Seventh ed. 1-106.
13. Weinstein, S. W., and A. Williams. 1983. Effect of amiloride and SITS on NaCl permeability in isolated perfused proximal tubule. *Kidney Int.* 23:269. (Abstr.)
14. Green, R., R. J. Moriarty, and G. Giebisch. 1981. Ionic requirements for proximal tubule fluid reabsorption: flow dependence of fluid transport. *Kidney Int.* 20:580-587.
15. Annison, E. F. 1954. Studies on the volatile fatty acids of sheep blood with special reference to formic acid. *Biochem. J.* 58:670-680.
16. Chantrelle, B. M., M. G. Cogan, and F. C. Rector, Jr. 1985. Active and passive components of NaCl absorption in the proximal tubule of the rat kidney. *Mineral. Electrolyte Metab.* 11:209-214.
17. Cardinal, J., M. D. Lutz, M. B. Burg, and J. Orloff. 1975. Lack of relationship of potential difference to fluid reabsorption in the proximal tubule. *Kidney Int.* 7:94-102.
18. Schafer, J. A., C. S. Patlak, and T. E. Andreoli. 1975. A component of fluid absorption linked to passive ion flows in the superficial pars recta. *J. Gen. Physiol.* 66:445-471.
19. Seifter, J. L., and P. S. Aronson. 1984. Cl transport via anion exchange in Necturus renal microvillus membranes. *Am. J. Physiol.* 247: F888-F895.
20. Seifter, J. L., R. Knickelbein, and P. S. Aronson. 1984. Absence of Cl-OH exchange and NaCl cotransport in rabbit microvillus membranes vesicles. *Am. J. Physiol.* 247:F753-F759.
21. Warnock, D. G., and V. J. Yee. 1981. Chloride uptake in brush border membrane vesicles isolated from rabbit renal cortex. Coupling to proton gradients and K diffusion potentials. *J. Clin. Invest.* 67:103-115.
22. Schwartz, G. J. 1983. Absence of Cl⁻/OH⁻ or Cl⁻/HCO₃⁻ exchange in the rabbit proximal tubule. *Am. J. Physiol.* 245:F462-F469.
23. Walter, A., and J. Gutknecht. 1984. Monocarboxylic acid permeation through lipid bilayer membranes. *J. Membr. Biol.* 77:255-264.
24. Lapointe, J. Y., R. Laprade, and J. Cardinal. 1986. Characterization of the apical membrane ionic permeability of the rabbit proximal convoluted tubule. *Am. J. Physiol.* 250:F339-F347.
25. Cassola, A. C., M. Mollenhauer, and E. Fromter. 1983. The intracellular chloride activity of rat kidney proximal tubular cells. *Pfluegers Arch. Eur. J. Physiol.* 399:259-265.