Vitamin D₃ and Cardiovascular Function in Rats

Ronald E. Weishaar and Robert U. Simpson

Department of Pharmacology, University of Michigan, Ann Arbor, Michigan 48109

Abstract

We have previously identified a receptor for 1,25-dihydroxyvitamin D₃ in myocardial cells (Simpson, R. U. 1983. Circulation. 68:239.). To establish the relevance of this observation, we evaluated the role of the prohormone vitamin D₃ in regulating cardiovascular function. In rats maintained on a vitamin D₃-deficient diet for nine weeks, increases in systolic blood pressure (BP) and serum creatine phosphokinase (CPK) were observed. These increases coincided with a reduction of serum calcium from 10.3 to 5.6 mg/dl. However, while serum calcium remained depressed throughout the study, increases in BP and serum CPK were transient. After nine weeks of vitamin D₃-depletion, but not after six weeks, ventricular and vascular muscle contractile function were also markedly enhanced. The increase in ventricular contractile function could not be prevented by maintaining serum calcium at 9.0 mg/dl during the period of D_3 -depletion. These observations suggest a primary role for the vitamin D₃-endocrine system in regulating cardiovascular function.

Introduction

The importance of vitamin D_3 for maintaining plasma levels of calcium is well known and is due to the effect that the vitamin D_3 metabolite 1,25-dihydroxyvitamin D_3 (1,25[OH]₂ D_3)¹ exerts on calcium metabolism in the bone, kidney, and intestine (1–3). A number of recent studies have indicated that 1,25(OH)₂ D_3 may also play a direct role in regulating metabolic events in other tissues and cells, including the pancreas (4), brain (5), pituitary gland (4, 6), and cancer cells (7). This direct role has been suggested by the identification of a specific receptor for 1,25(OH)₂ D_3 in these cells, as well as the presence of 1,25(OH)₂ D_3 -dependent calcium-binding proteins (8–10).

In previous reports we demonstrated that a specific receptor for $1,25(OH)_2D_3$ is present in cultured heart and skeletal muscle (11, 12). Subsequently, Walters et al. identified a receptor for $1,25(OH)_2D_3$ in a low-salt chromatin preparation from normal rat hearts (13), and Thomasett and co-workers have shown that

J. Clin. Invest. © The American Society for Clinical Investigation, Inc. 0021-9738/87/06/1706/07 \$1.00 Volume 79, June 1987, 1706-1712 myocardial tissue contains a vitamin D_3 -dependent calciumbinding protein (10). These observations suggest a role for $1,25(OH)_2D_3$ in regulating cardiac metabolism.

Although administration of toxic quantities of vitamin D_3 have been shown to produce heart failure (14), presumably through an indirect elevation of plasma calcium and subsequent myocardial calcinosis, a direct role for $1,25(OH)_2D_3$ in contributing to the regulation of myocardial metabolism has not been described. There is considerable evidence, however, to indicate that changes in myocardial calcium homeostasis are important in mediating certain aspects of cardiovascular dysfunction (15–17).

In the present study, the role of vitamin D_3 in regulating cardiovascular function was evaluated by examining the response of rats maintained on a vitamin D_3 -deficient diet for either 6 or 9 wk. The results of this study clearly demonstrate that vitamin D deficiency produces significant changes in normal cardiovascular function.

Methods

Male weanling Sprague-Dawley rats (Holtzman Co., Madison, WI) were obtained from mothers maintained on a low vitamin D₃ diet. Immediately upon arrival the rats were housed in an area removed from ultraviolet light, including fluorescent light. The rats were maintained on a vitamin D3-deficient diet that contained 0.4% calcium and 0.4% phosphorus (diet TD84475, Teklad Laboratories, Madison, WI). In a subsequent study rats were also maintained for 9 wk on a vitamin D₃-deficient diet containing 2.5% calcium and 1.5% phosphate (diet TD86029, Teklad Laboratories). After 1 wk the rats were weighed and measurements of systolic blood pressure (BP) and heart rate (HR) were made. Blood samples were also taken from an incision in the tail for measurements of serum Ca, PO₄, and creatine phosphokinase (CPK). Measurements of BP, HR, and serum Ca, PO₄, and CPK were made at weekly or biweekly intervals throughout the remainder of the study. Following these initial measurements the rats were randomly divided into two groups. The D⁺ group was maintained on the 0.4% Ca/0.4% PO4 diet, which was supplemented with 2 IU vitamin D₃/g diet. The vitamin D₃ was dissolved in corn oil and added directly to the diet. Rats in the D⁻ group received the same diet, containing an equal amount of corn oil. Both groups received an equal amount of the diet (15 g/rat/d). Changes in body weight were recorded weekly or biweekly.

After 6 wk half the animals in each group were sacrificed for evaluation of in vitro cardiac and vascular muscle contractility. At 9 wk the remainder of the animals were sacrificed and measurements of in vitro contractility repeated.

Analysis of serum. Blood samples (2-3 ml) were collected under vacuum into side arm test tubes previously rinsed with deionized water. The blood was allowed to clot for 30 min, centrifuged, and the serum was decanted and stored at -20° C until analysis (usually within 1 mo). Serum calcium was measured using atomic absorption spectrophotometry (Atomic Absorption Spectrophotometer, model 2380, Perkin-Elmer Corp., Norwalk, CT). The spectrophotometer was equipped with a single-element hollow cathode lamp (excitation wavelength of 422.7 nm). A nitrous oxide-acetylene flame was employed to reduce interference from serum proteins and increase sensitivity. 0.1% lanthanum chloride was included in all samples to control ionization interference. Serum phosphate was measured by the method of Fiske and SubbaRow (18), using

Portions of this study were published in abstract form at the 1986 Federation of American Societies for Experimental Biology meeting, St. Louis, MO.

Address reprint requests to Robert U. Simpson, Department of Pharmacology, University of Michigan, Ann Arbor, MI 48109.

Received for publication 27 May 1986 and in revised form 5 January 1987.

^{1.} Abbreviations used in this paper: CPK, creatine phosphokinase; HR, heart rate; LVSP, left ventricular systolic pressure; $25(OH)D_3$, 25-hydroxyvitamin D₃; $1,25(OH)_2D_3$, 1,25-dihydroxyvitamin D₃; PSS, physiological salt solution.

a commercial kit (Kit 670-C, Sigma Chemical Co., St. Louis, MO). Serum CPK was measured using the colorimetric method described by Hughes (19), also using a commercial kit (Kit 520, Sigma Chemical Co.). Serum levels of 25-hydroxyvitamin D₃ (25[OH]D₃) were measured using radioimmunoassay after extraction of 25(OH)D₃ and other hydroxylated metabolites with acetonitrile (20), using a commercial kit (Catalogue No. 5800, Immuno Nuclear Corp., Stillwater, MN). Serum 1,25(OH)2D3 was measured by the method of Reinhardt et al. (21), also using a commercial kit (Immuno Nuclear Corp.). For this measurement vitamin D₃ metabolites were extracted using acetonitrile and then partially purified using a C_{18} cartridge. Further purification was achieved using a silica cartridge. Percent recovery averaged 50-55%. Quantification was achieved using a 1,25(OH)₂D₃ receptor protein competitive binding assay. Dextrancoated charcoal suspension was used to separate bound from free hormone. For measurements of 25(OH)D₃ and 1,25(OH)₂D₃, blood from five pentobarbital-anesthetized rats was taken via heart puncture and pooled to obtain sufficient serum for analysis of both 25(OH)D₁ and 1.25(OH),D1.

Measurement of blood pressure and heart rate. Systolic blood pressure and heart rate were measured weekly or biweekly in conscious, restrained animals by tail cuff occlusion using standard equipment and techniques (Narco Bio-Systems, Healthdyne Co., Houston, TX). An average of six recordings was made for each rat.

Measurement of in vitro contractility. Rats were injected with heparin (100 U/kg) 30 min before sacrifice via cervical dislocation. After stunning, the chest cavity was rapidly opened and the heart and thorasic aorta were quickly removed and placed in a beaker containing ice-cold saline until contractions ceased (within 20 s). The heart and aorta were then rinsed in a second beaker of ice-cold saline, after which the aorta was removed and placed in an ice-cold physiological salt solution (PSS) containing the following: 130 mM NaCl, 4.7 mM KCl, 1.18 mM KH₂PO₄, 1.17 mM MgSO₄, 4.9 mM NaHCO₃, 5.5 mM glucose, 1.6 mM CaCl₂, and 0.03 mM CaNa₂EDTA (pH 7.4). The aorta was refrigerated overnight and evaluated the following day.

The heart was then mounted via the aortic root and perfused Langendorf fashion at 37°C. The composition of the perfusate was as follows: 118 mM NaCl, 4.7 mM KCl, 1.2 mM KH₂PO₄, 1.2 mM MgSO₄, 25.0 mM NaHCO₃, 11.0 mM glucose, and 2.0 mM CaCl₂ (2.5 mM CaCl₂ + 0.5 mM Na₂EDTA). The perfusate was bubbled continuously with 95% $O_2/5\%$ CO₂, which maintained the pH at ~ 7.4. The heart was perfused in this manner for 10 min, during which the aortic perfusion pressure was adjusted to 80 mmHg. A Millar pressure transducer was also placed into the lumen of the left ventricle through a small incision in the left atria. Once properly positioned, the pressure transducer was sutured in place. This transducer was used for measuring left ventricular systolic pressure (LVSP), the first derivative of the rise and fall in left ventricular pressure $(\pm dP/dt)$, and HR. After the 10-min equilibration period the concentration of calcium in the perfusate was reduced to 1.0 mM (1.5 mM CaCl₂ + 0.5 mM Na₂EDTA), and once a new steady state of contractility was achieved (within 5 min) measurements of LVSP, $\pm dP/dt$, and HR were made. The concentration of calcium in the perfusate was then incrementally increased to 1.5, 2.0, 2.5, and 3.0 mM at 5-min intervals. Measurements of contractility and HR were made at each new concentration of extracellular calcium.

The following day rings 3–4 mm in length were prepared from the thorasic aortae obtained the day before. These rings were always taken from the same location (~ 3 mm below the aortic arch), because preliminary studies showed that the response to different constricting agents, e.g., KCl, can vary depending upon the site of origin of the aortic ring (personal observation). The rings were mounted vertically between two stainless steel triangles in a 50-ml water-jacketed tissue bath containing PSS maintained at 37°C. The composition of this salt solution was the same as previously described. After mounting, resting tension was gradually increased to 5.0 g. This degree of resting tension resulted in a maximum contractile response to 100 mM KCl or to $1.0 \,\mu$ M norepinephrine (personal observation). The aortic rings were allowed to equilibrate for 90 min, during which PSS was changed once. The rings were then constricted by increasing the concentration of KCl in the PSS to 100 mM. After the maximum response was obtained, the rings were rinsed three times at 10-min intervals with fresh PSS. Resting tension was then readjusted to 5.0 g, and increasing concentrations of norepinephrine were added to the bath, until a maximum response was obtained. The rings were again washed three times at 10-min intervals with fresh PSS, after which resting tension was readjusted, and the rings were again challenged with increasing concentrations of norepinephrine. The response to the second challenge of norepinephrine was used for all calculations, because preliminary results showed that the first response to norepinephrine occasionally varied, whereas the second response to norepinephrine was very reproducible. Following the second challenge with norepinephrine, the rings were removed from the bath, blotted dry, and weighed.

Statistical analyses. Changes in serum calcium, phosphate, CPK, systolic BP, and HR were evaluated using an unpaired Student t test (22). Changes in the relationship between extracellular calcium and cardiac contractility and the relationship between increasing concentrations of norepinephrine and vascular contractile response between the two groups were evaluated using analysis of variance (22).

Results

Body weight and serum changes. Fig. 1 illustrates the rate of growth for the vitamin D₃-deficient rats maintained on a 0.4% calcium diet (D⁻ rats), and for the vitamin D₃-sufficient rats maintained in the same diet (D⁺ rats). As can be seen, the D⁻ rats gained slightly less weight during the 9-wk course of the study than did the D⁺ rats. After 9 wk, however, the difference in body weight between the two groups was < 10%.

Changes in serum Ca, PO₄, and serum CPK, which accompanied vitamin D₃-depletion are shown in Fig. 2. Within 4 wk, serum Ca levels in the D⁻ rats had fallen to roughly 50% of the level observed in the D⁺ rats (10.5 \pm 0.3 mg/dl for the D⁺ rats and 5.6 \pm 0.2 mg/dl for the D⁻ rats). Serum Ca remained at this reduced level in the D⁻ rats throughout the remainder of the study (Fig. 2).

A large increase in serum CPK was observed in the D⁻ rats, which coincided roughly with the development of hypocalcemia (Fig. 2). Within 3-4 wk after the onset of vitamin D₃-depletion, the serum CPK in the D⁻ rats was more than double that observed in the D⁺ rats (188±11 IU/dl for the D⁻ rats and 85±13 IU/dl for the D⁺ rats). However, whereas the level of serum Ca remained low throughout the study, the increase in serum CPK was transient, and after 8 wk the level of serum CPK in the D⁻ rats had returned to normal (Fig. 2). A slight increase in serum PO₄ was also observed in the D⁻ rats (Fig. 2). This increase, however, was never > 15% above the level in the D⁺ rats.

In addition to a profound reduction in serum calcium, changes in the circulating levels of $1,25(OH)_2D_3$ and $25(OH)D_3$ were also used to establish vitamin D₃-deficiency. As Table I



Figure 1. Representative age-dependent changes in body weight (g) for vitamin D₃-deficient rats maintained on a diet containing 0.4% calcium and 0.4% phosphate (\odot), and vitamin D₃sufficient rats maintained on the same diet (\bullet). Each symbol represents the mean±SEM of at least eight rats.



Figure 2. Age-dependent changes in serum calcium (mg/dl), serum phosphorus (mg/dl), and serum CPK (IU/dl) in the vitamin D₃-deficient rats maintained on a diet containing 0.4% calcium and 0.4% phosphate (hatched box), and vitamin D₁-sufficient rats maintained on the same diet (solid box). Bars represent the mean±SEM for 3-6 serum samples. *P < 0.05.

illustrates, serum levels of both $1,25(OH)_2D_3$ and $25(OH)_2D_3$ were below the level of detection in rats maintained on the vitamin D_3 -deficient diet for 6 or 9 wk.

BP and HR changes. The effect of vitamin D₃-depletion on systolic blood pressure and HR is shown in Fig. 3. Removal of vitamin D₃ from the diet was associated with a significant increase in systolic BP in the vitamin D3-deficient rats, as compared with the vitamin D₃-sufficient rats. An early increase in systolic BP was observed in both groups, which may be age related, due to equilibration of the rats to their environment or the change in diet for the weanling rats. This initial increase, however, was much greater in the D^- rats than in the D^+ rats. As was the case with the increase in serum CPK, the increase in systolic BP in the vitamin D₃-deficient rats was transient and by 8 wk there was no difference in BP between the two groups (Fig. 3). Subsequent studies have shown that BP remains the same in both groups for up to 18 wk (personal observation). No difference in HR between the two groups was observed at any time during the study.

Changes in in vitro contractile function. At 6 wk and again at 9 wk half of the rats in both groups were sacrificed for evaluation of cardiac and vascular muscle contractile function in vitro. Changes in cardiac contractility were evaluated by examining the response of Langendorf-perfused hearts from the

Table I. Levels of $1,25(OH)_2D_3$ and $25(OH)D_3$ in Serum From Vitamin D_3 -Deficient and Vitamin D_3 -Sufficient Rats

Serum 1,25-(OH) ₂ D ₃	Serum 25-(OH)D	
pgm/ml	ngm/ml	
*	*	
*	*	
132.5	9.5	
97.0	14.1	
	Serum 1,25-(OH) ₂ D ₃ pgm/ml * * 132.5 97.0	

Each value represent the level of $1,25-(OH)_2D_3$ or $25-(OH)D_3$ in pooled serum from five rats.

* Below the level of detection (10 pgm/ml for $1,25(OH)_2D_3$ and 0.1 ngm/ml for $25(OH)D_3$).



Figure 3. Age-dependent changes in systolic BP (mmHg) and HR (bpm) in the vitamin D₃-deficient rats maintained on a diet containing 0.4% calcium and 0.4% phosphate (\odot), and vitamin D₃-sufficient rats maintained on the same diet (\bullet). Each symbol represents the mean±SEM for 5-8 rats. *P < 0.05.

 D^- and D^+ rats to changes in extracellular Ca. Changes in vascular muscle contractility were evaluated by examining the response of isolated aortic rings from the D^- and D^+ rats to increasing concentrations of exogenous norepinephrine.

Changes in cardiac contractile function. After 6 wk of vitamin D₃ depletion, a slight increase in the rate of pressure development (+dP/dt) in isolated hearts from the D⁻ rats was observed (Fig. 4 A). In addition, a slight increase in the rate of relaxation (-dP/dt) was also noted (Fig. 4 B). These increases, however, were not statistically significant (P > 0.05). After 9 wk on the vitamin D3-deficient diet, however, large and statistically significant increases in +dP/dt and -dP/dt were observed in the hearts from the D⁻ rats compared with hearts from the D⁺ rats (Fig. 5, Aand B). In both groups the percent increase in +dP/dt and -dP/dtdt after each incremental elevation of extracellular calcium was comparable, and the shape of the calcium-contractility concentration-response curves over the same concentration range was identical for both the D^- and D^+ hearts. As shown in Fig. 5, A and B, the major difference between the two groups appears to be the magnitude of the response evoked by extracellular calcium.

Changes in vascular contractile function. Fig. 6, A and B illustrate the sensitivity of aortic rings from the D^+ and D^- rats to exogenous norepinephrine following 6 and 9 wk of vitamin



Figure 4. Changes in +dP/dt(A) and -dP/dt (B) in response to changes in extracellular calcium in isolated, Langendorf perfused hearts from vitamin D₃-deficient rats maintained on a diet containing 0.4% calcium and 0.4% phosphate (0), and vitamin D₃-sufficient rats maintained on the same diet (•) after 6 wk of study. Each symbol represents the mean±SEM of 4-6 hearts from each group. No significant differences were observed between the two groups.



Figure 5. Changes in +dP/dt(A) and -dP/dt (B) in response to changes in extracellular calcium in isolated, Langendorf perfused hearts from vitamin D₃-deficient rats maintained on a diet containing 0.4% calcium and 0.4% phosphate (0), and vitamin D₃-sufficient rats maintained on the same diet (•) after 9 wk of study. Each symbol represents the mean±SEM of 5-7 hearts from each group. A statistically significant difference (P < 0.05) was observed between the two groups.



Figure 7. Changes in the magnitude of the response of isolated aortic rings to exogenous norepinephrine in vitamin D3-deficient rats maintained on a diet containing 0.4% calcium and 0.4% phosphate (0), and vitamin D₃-sufficient rats maintained on the same diet (•) after 6 wk (A) or 9 wk (B) of study. Each symbol represents the mean±SEM of 4-6 rings from each group. Statistically significant differences (P < 0.05) were observed after 9 wk of vitamin D₃ depletion but not after 6 wk of vitamin D_3 depletion.

 D_3 depletion. As can be seen, no difference in norepinephrine sensitivity was observed between the two groups after 6 wk on the vitamin D_3 -deficient diet (Fig. 6 A). In both groups, the EC₅₀ values for norepinephrine were roughly 30 nM. Because serum Ca in the D^- rats was reduced by 50% at this time, these experiments were repeated in PSS containing 0.8 mM Ca instead of the usual 1.6 mM calcium. As before, no differences in norepinephrine sensitivity were observed between the two groups (data not shown).

After 9 wk of vitamin D_3 depletion, a significant increase in the sensitivity of aortic rings from the D^- rats to exogenous norepinephrine was observed (Fig. 6 *B*). The EC₅₀ values for norepinephrine were 25 nM for the aortae from the D^+ rats and 9 nM for aortae from the D^- rats.

Changes in the magnitude of the contractile response of aortae from the vitamin D₃-deficient and vitamin D₃-sufficient rats to norepinephrine are shown in Fig. 7, A and B. As can be seen, norepinephrine evoked comparable increases in tension in aortae from both groups after 6 wk of vitamin D₃ depletion (Fig. 7 A). However, after 9 wk on the vitamin D₃-deficient diet, norepinephrine produced much greater increases in tension in the aortae from the D⁻ rats compared with the D⁺ rats (0.542 \pm 0.038



Figure 6. Changes in the sensitivity of isolated aortic rings to exogenous norepinephrine in vitamin D₃-deficient rats maintained on a diet containing 0.4% calcium and 0.4% phosphate (0), and vitamin D3-sufficient rats maintained on the same diet (•) after 6 wk (A) or 9 wk (B) of study. Each symbol represents the mean±SEM of 4-6 rings from each group. Statistically significant differences (P < 0.05) were observed after 9 wk of vitamin D3 depletion but not after 6 wk of vitamin D₃-depletion.

g/mg wet wt. for the D⁻ rats vs. 0.258 ± 0.027 g/mg wet wt. for the D⁺ rats) (Fig. 7 B).

Direct/indirect effects of vitamin D_3 on cardiac contractile function. To determine whether vitamin D_3 exerts a direct effect on cardiac contractile function or whether its effect represents a secondary response to the hypocalcemia that accompanies vitamin D_3 depletion, it was necessary to develop a vitamin D_3 deficient diet in which serum Ca and PO₄ are maintained at normal or near-normal levels. For these studies diets containing varying amounts of Ca and PO₄ were fed to vitamin D_3 -deficient rats and serum levels of Ca and PO₄ were recorded.

Table II illustrates the steady-state levels of serum Ca and PO₄ obtained after long-term administration of the various diets to vitamin D₃-deficient rats, compared with levels obtained in rats maintained on the vitamin D₃-sufficient diet containing 0.4% Ca and 0.4% PO₄. Although increasing the amount of Ca in the vitamin D₃-deficient diet from 0.4% to 2.5% normalized serum Ca, failure to also increase the amount of PO₄ in the diet results in profound hypophosphatemia (Table II). This reduction is not significantly altered by increasing the level of PO₄ in the diet from 0.4% to 1.0%. Increasing PO₄ content to 1.5%, however, was found to prevent hypophosphatemia in the vitamin D₃-de-

Table II. Levels of Serum Electrolytes after Chronic Administration of Diets Varying in Calcium and Phosphate Composition to Vitamin D₃-Deficient Rats

Vitamin D ₃	Diet composition		Serum electrolytes	
	% Calcium	% Phosphate	Calcium	Phosphate
			mg/dl	mg/dl
+	0.4	0.4	10.3±0.2 (47)	5.8±0.2 (75)
_	0.4	0.4	5.6±0.2 (32)	6.3±0.2 (66)
	2.5	0.4	9.7±0.5 (16)	2.3±0.3 (31)
-	2.5	1.0	9.9±0.4 (15)	2.6±0.2 (27)
-	2.5	1.5	9.0±0.4 (14)	5.6±0.2 (31)

Data represent the mean±SEM of the number of samples in parentheses.

ficient diet containing 2.5% Ca. This latter diet was therefore used for all subsequent studies.

To assess the possible involvement of hypocalcemia with the increases in cardiac contractile function which accompany vitamin D_3 depletion, rats were placed on the vitamin D_3 -deficient diet containing high Ca (2.5%) and high PO₄ (1.5%) for 9 wk to determine whether maintenance of serum Ca during the period of vitamin D_3 depletion could prevent these changes.

After 9 wk of vitamin D₃ depletion, rats maintained on either the 0.4% Ca, 0.4% PO4 diet (D⁻, 0.4% Ca) or the 2.5% Ca, 1.5% PO_4 diet (D⁻, 2.5% Ca) were sacrificed and the contractile response of isolated Langendorf-perfused hearts to increasing concentrations of extracellular Ca was evaluated. For comparative purposes, the response of isolated hearts from rats maintained for 9 wk on the vitamin D₃-sufficient diet (D⁺, 0.4% Ca) was also evaluated. As Fig. 8 illustrates, large and statistically significant increases in +dP/dt were again observed in hearts from the D^- , 0.4% Ca rats compared with hearts from the D^+ , 0.4% Ca rats. Fig. 8 also illustrates that similar increases were also observed in the D⁻, 2.5% Ca rats. Because this latter diet raised serum Ca to near-normal levels in the vitamin D₃-deficient rats, these results suggest that this change represents a direct response to vitamin D₃ depletion, rather than a secondary response to the hypocalcemia that normally accompanies vitamin D_3 depletion.

Discussion

Recently, Simpson et al. identified a specific receptor for $1,25(OH)_2D_3$, the metabolically active form of vitamin D_3 , in heart cell preparations (11, 12). Walters and co-workers subsequently demonstrated a receptor for $1,25(OH)_2D_3$ in normal rat hearts (13), and Stumpf et al., using autoradiographic techniques, demonstrated nuclear receptors for $1,25(OH)_2D_3$ in rat heart cells (23). Thomasset et al. have also shown that myocardial tissue contains a 10,000 K vitamin D_3 -dependent Ca-binding protein that is immunochemically similar to that found in intestine (10). Given the close relationship between changes in Ca homeostasis in cardiac muscle and changes in myocardial con-



Figure 8. Changes in +dP/ dt in response to changes in extracellular calcium in isolated, Langendorf perfused hearts from vitamin D3-deficient rats maintained for 9 wk on a low (0.4%) calcium, low (0.4%) phosphate diet (0), from vitamin D₃deficient rats maintained for 9 wk on a high (2.5%) calcium, high (1.5%) phosphate diet (\triangle), and from vitamin D₃-sufficient rats maintained for 9 wk on the low calcium, low phosphate diet (.). Each symbol represents the mean±SEM of 5-7 hearts from each group.

Statistically significant differences (P < 0.05) were observed between the vitamin D₃-sufficient rats and both groups of vitamin D₃-deficient rats. No statistically significant differences were observed between the two groups of vitamin D₃-deficient groups. tractility, these results indicate that vitamin D_3 may play a direct role in regulating cardiovascular function.

Little information is presently available regarding the direct influence of vitamin D₃ on cardiovascular function under normal conditions or conditions such as ischemia, heart failure, or hypertension, where changes in cellular Ca homeostasis are apparent. In skeletal muscle, vitamin D₃ depletion is associated with a prolongation of muscle contraction kinetics (24). This change is apparently not the result of hypocalcemia or alterations in parathyroid hormone metabolism. Wassner et al. have also found that vitamin D₃ depletion increases skeletal muscle degradation (25). Several investigators have described a specific cardiomyopathy associated with end-stage renal disease (26, 27). Administration of either 25(OH)D₃ (28) or $1-\alpha$ -hydroxyvitamin D_3 (27) has been shown to improve left ventricular function in these patients. Whether this improvement was due to restoration of a 1,25(OH)₂D₃-dependent metabolic process in cardiac muscle or the ability of $1,25(OH)_2D_3$ to suppress the hyperparathyroidism observed in these patients is not known.

In the present study, vitamin D₃ deficiency was associated with significant changes in cardiovascular function, including increases in cardiac and vascular muscle contractile responses and systolic BP. The changes in cardiac contractile function are of interest in that whereas only modest differences in cardiac contractile function were apparent after 6 wk of vitamin D₃ depletion, after 9 wk a profound increase in cardiac contractility was observed in the vitamin D3-deficient rats. This change in contractile function appears to represent the direct response to vitamin D₃ depletion and not a response to hypocalcemia that accompanies depletion, because increasing serum Ca from 5.6 to 9.0 mg/dl during the 9-wk period of vitamin D₃ depletion does not prevent the change in cardiac contractile function. The possibility exists, however, that hypocalcemia could still contribute to the changes in ventricular contractile function observed in the vitamin D₃-deficient rats, because rats placed on the 2.5% Ca vitamin D₃-deficient diet remained slightly hypocalcemic compared to rats on the vitamin D₃-sufficient diet (9.0 vs. 10.3 mg/dl).

In addition to an increase in the rate of myocardial pressure development, hearts from the vitamin D3-deficient rats also displayed an increase in the rate of myocardial relaxation. Katz and others have suggested that the rate of myocardial relaxation is closely associated with the rate of Ca sequestration by the sarcoplasmic reticulum (29-31), possibly indicating that vitamin D_3 deficiency can increase the rate at which such sequestration occurs. Likewise, an increase in the rate of Ca release by the sarcoplasmic reticulum may represent the basis for the increase in contractility observed in the hearts of the vitamin D₃-deficient rats, because Ca release by the sarcoplasmic reticulum represents the principle initiating event in myocardial contraction (30, 31). Fabiato and Fabiato have shown that Ca release by the sarcoplasmic reticulum is triggered by an increase in cytosolic free Ca (30), which occurs during each depolarization when Ca enters the cell through the Ca slow channel.

An alternative explanation for the increase in myocardial contractility in the vitamin D_3 -deficient rats is the absence of $1,25(OH)_2D_3$ -dependent Ca-binding proteins, which under normal conditions might serve as a Ca buffer in cardiac muscle. Thomasset et al. has previously demonstrated that cardiac muscle from rats maintained on a vitamin D_3 -sufficient diet contains a $1,25(OH)_2D_3$ -dependent Ca-binding protein immunochemically similar to that found in intestinal cells (10), and it has been

suggested that this protein may play an important role in calcium handling in a number of organs and cells (32, 33). The loss of such a protein in cardiac muscle following vitamin D_3 depletion might allow the cytosolic free Ca level to rise to a higher than normal level after each depolarization, thereby triggering the release of a greater quantity of Ca from the sarcoplasmic reticulum, resulting in a greater contraction. Both alternative explanations are currently under investigation in our laboratory.

Vitamin D₃ depletion also resulted in profound changes in vascular muscle contractile function. These changes temporally paralleled the changes in cardiac muscle contractility, in that whereas no significant differences in the contractile function of isolated aortic rings were observed after 6 wk of vitamin D₃ depletion, after 9 wk large differences in vascular contractility were observed between the vitamin D₃-deficient and vitamin D₃-sufficient groups. The direct/indirect role of vitamin D₃ in modulating these changes in vascular contractility are not currently known, but preliminary evidence suggests the hypocalcemia may play a more important role in modulating these changes than vitamin D₃ (manuscript in preparation).

Whereas the changes in vascular and cardiac contractility may represent direct alterations in calcium homeostasis in cardiac and vascular muscle, the basis for the change in BP that accompanies vitamin D_3 depletion is not clear. It is noteworthy that the increase in blood pressure in the vitamin D₃-deficient rats coincided with the development of hypocalcemia in these animals. McCarron has previously demonstrated that hypocalcemia is associated with increases in BP, both in patients and experimental animals (34). Administration of Ca to young spontaneously hypertensive rats has also been shown to attenuate the development of hypertension in these animals (35), and will reverse "fixed" hypertension in adult animals (36). However, while serum Ca remained reduced throughout the present study, the increase in BP in the vitamin D_3 -deficient rats was transient. This observation suggests that a compensatory mechanism exists which is capable of overcoming the initial hypertensive response. One such compensatory mechanism might be an increase in serum PO₄ which can accompany vitamin D_3 depletion (24, 25). Lau et al. have previously reported a strong correlation between increases in serum PO_4 and hypotension (37). However, in the present study only slight increases in serum PO₄ were observed in the vitamin D₃-deficient rats, which were never more than 15% above levels in the vitamin D₃-sufficient rats.

A significant increase in serum CPK was also observed in the vitamin D_3 -deficient rats, which is indicative of muscle damage in these animals. Because only total CPK was measured in the present study, it cannot be said for certain whether the increase in serum CPK in the vitamin D_3 -deficient rats is due to damage to myocardial or skeletal muscle or both. As with the increase in BP, the increase in serum CPK appeared to correlate with the onset of hypocalcemia, suggesting that both events may represent a metabolic consequence of the reduction in circulating Ca. The subsequent decline in serum CPK, like the return of BP to normotensive levels, suggests the presence of adaptive mechanisms to compensate for the initial insult.

The present study demonstrates that vitamin D_3 deficiency produces profound changes in cardiovascular function, including heightened contractile responses of isolated cardiac and vascular smooth muscle, hypertension, and elevation of serum CPK. These observations indicate that vitamin D_3 plays an important role in maintaining normal cardiovascular function. This involvement appears to be a direct one in cardiac muscle. However,

in vascular muscle the change in function may be mediated by hypocalcemia secondary to vitamin D₃ deficiency (manuscript in preparation). Whether changes in vitamin D₃ also play an important role in cardiovascular disease states is not known. It is important to note that there is a decline in circulating levels of both 1,25(OH)₂D₃ and Ca with increasing age (38), and a correlation between increased BP and hypocalcemia has been demonstrated in patients and animal models of hypertension (34). In addition, the possibility exists that vitamin D_3 may play a role in the changes in cardiovascular function that can accompany diabetes (39), lengthy bedrest or immobilization (40, 41), or prolonged periods of weightlessness (41), because circulating levels of $1,25(OH)_2D_3$ are reduced by these conditions (42-45). In conclusion, the results of this study strongly suggest that the vitamin D₃-endocrine system contributes to the regulation of cardiovascular function, either directly as is the case with cardiac muscle or indirectly through the effects of 1,25(OH)₂D₃ on Ca or other circulating factors.

Acknowledgments

This research was supported by a grant-in-aid from the American Heart Association, with funds contributed in part by the Michigan Heart Association (85-846) and the National Institutes of Health (CA-36507).

References

1. Martin, D. L., and H. F. DeLuca. 1969. Calcium transport and the role of vitamin D. Arch. Biochem. Biophys. 134:139-148.

2. Norman, A. W., J. Roth, and L. Orci. 1982. The vitamin D endocrine system. *Endocr. Rev.* 3:331-366.

3. DeLuca, H. F., and H. L. Schnoes. 1983. Vitamin D: recent advances. Annu. Rev. Biochem. 52:411-439.

4. Pike, J. W., L. L. Gooze, and M. R. Haussler. 1979. Biochemical evidence for 1,25-dihydroxyvitamin D receptor macromolecules in parathyroid, pancreatic, pituitary and placental tissues. *Life Sci.* 26:407–412.

5. Stumpf, W. E., W. Sar, S. A. Clark, and H. F. DeLuca. 1982. Brain target sites for 1,25-dihydroxyvitamin D_3 . Science (Wash. DC). 215:1403-1404.

6. Stumpf, W. E., M. Sar, F. A. Reid, Y. Tanaka, and H. F. DeLuca. 1979. Target cells for 1,25-dihydroxyvitamin D_3 in intestinal tract, stomach, kidney, skin, pituitary and parathyroid. *Science (Wash. DC).* 206: 1108–1109.

7. Thomas, G. A., and R. U. Simpson. 1986. High performance liquid chromatography analysis of 1,25-dihydroxyvitamin D_3 receptor in malignant cells. Correlation of effects on cell proliferation and receptor concentration. *Cancer Biochem. Biophys.* 45:221–224.

8. Sielvanayagam, C. S., S. Y. Tsai, M.-J. Tsai, P. Selvanayagam, and G. F. Saunders. 1984. Multiple origins of transcription for the human placental lactogen genes. J. Biol. Chem. 259:14642-14646.

9. Bainbridge, K. G., and C. D. Parkes. 1981. Vitamin D-dependent calcium binding protein in the chick brain. *Cell Calcium*. 2:65-76.

10. Thomasset, M., C. D. Parkes, and P. Cuisinier-Eleizes. 1982. Rat calcium-binding proteins: distribution, development, and vitamin D dependence. *Am. J. Physiol.* 243:E483-E488.

11. Simpson, R. U. 1983. Evidence for a specific 1,25-dihydroxyvitamin D₃ receptor in rat heart. *Circulation*. 68:239.

12. Simpson, R. U., G. A. Thomas, and A. J. Arnold. 1985. Identification of 1,25-dihydroxyvitamin D_3 receptors and activities in muscle. *J. Biol. Chem.* 260:8882–8891.

13. Walters, M. R., D. C. Wicker, and P. C. Riggle. 1986. 1,25dihydroxyvitamin D₃ receptors identified in the rat heart. J. Mol. Cell. Cardiol. 18:67-72. 14. Bonucci, E., and R. Sadun. 1973. Experimental calcification of the myocardium. Am. J. Pathol. 71:167-184.

15. Suko, J., J. H. K. Vogel, and C. A. Chidsey. 1970. Intracellular calcium and myocardial contractility. III. Reduced calcium uptake and ATPase of the sarcoplasmic reticular fraction prepared from chronically failing calf hearts. *Circ. Res.* 27:235–247.

16. Dhalla, N. S., D. K. Das, and G. P. Sharma. 1978. Subcellular basis of cardiac contractile failure. J. Mol. Cell. Cardiol. 10:363-385.

17. Sarma, J. S. M., S. Ikeda, R. Fischer, Y. Maruyama, R. Weishaar, and R. J. Bing. 1976. Biochemical and contractile properties of heart muscle after prolonged alcohol administration. *J. Mol. Cell. Cardiol.* 8: 951–972.

18. Fiske, C. H., and V. SubbaRow. 1925. The colorimetric determination of phosphorus. J. Biol. Chem. 66:375-379.

19. Hughes, B. P. 1962. A method for the estimation of serum creatine kinase and its use in comparing creatine kinase and aldolase activity in normal and pathologic sera. *Clin. Chim. Acta.* 7:579–603.

20. Hummer, L., L. Nilas, L. Tjellesen, and C. Christiansen. 1984. A selective and simplified radioimmunoassay of 25-hydroxyvitamin D₃. Scand. J. Clin. Lab. Invest. 44:163–170.

21. Reinhardt, T. A., R. L. Horst, J. W. Orf, and B. W. Hollis. 1984. A microassay for 1,25-dihydroxyvitamin D₃ not requiring high performance liquid chromatography: application to clinical studies. J. Clin. Endocrinol. & Metab. 58:91-97.

22. Winner, B. J. 1971. Statistical Principles in Experimental Design. McGraw-Hill Book Co., New York.

23. Stumpf, W. E., S. Madhabananda, and H. DeLuca. 1985. Sites of action of $1,25(OH)_2$ vitamin D₃ identified by thaw-mount autoradiography. *In* Proceedings of Sixth Workshop on Vitamin D. A. W. Norman, editor. Walter DeGrutyer, Berlin. 416-418.

24. Rodman, J. S., and T. Baker. 1978. Changes in the kinetics of muscle contraction in vitamin D-depleted rats. *Kidney Int.* 13:189-193.

25. Wassner, S. J., J. B. Li, A. Sperduto, and M. E. Norman. 1983. Vitamin D deficiency, hypocalcemia and increased skeletal muscle degradation in rats. J. Clin. Invest. 72:102–112.

26. Drücke, T., K. Le Pailleur, B. Meilhac, C. Koutoudis, J. Zingraff, J. Dimatteo, and J. Crosnier. 1977. Congestive cardiomyopathy in uremic patients on long-term hemodialysis. *Br. Med. J.* i:350–353.

27. McGonigle, R. J. S., M. B. Fowler, A. B. Timmis, M. J. Weston, and V. Parsons. 1984. Uremic cardiomyopathy: potential role of vitamin D and parathyroid hormone. *Nephron.* 36:94–100.

28. Coratelli, P., F. Petratrulo, E. Buongiorno, M. Giannattasio, G. Antonelli, and A. Amerio. 1984. Improvement in left ventricular function during treatment of hemodialysis patients with 25-OHD₃. *Contrib. Nephrol.* 41:433–437.

29. Ebashi, S., and M. Endo. 1960. Calcium and muscle contraction. Prog. Biophys. Mol. Biol. 18:125-183. 30. Fabiato, A., and F. Fabiato. 1977. Calcium release from the sarcoplasmic reticulum. *Circ. Res.* 40:119-129.

31. Katz, A. M. 1979. Role of the contractile proteins and sarcoplasmic reticulum in the response of the heart to catecholamines: an historical review. *Adv. Cyclic Nucleotide Res.* 11:303-344.

32. Wasserman, R. H., R. A. Corradino, and A. N. Taylor. 1968. Vitamin D-dependent calcium-binding protein. J. Biol. Chem. 243:3978–3986.

33. Zerwekh, J. F. 1979. Vitamin D-dependent intestinal calcium absorption. *Gastroenterology*. 76:404–410.

34. McCarron, D. A. 1985. Is calcium more important than sodium in the pathogenesis of essential hypertension? *Hypertension (Dallas)*. 7: 607-627.

35. McCarron, D. A., P. A. Lucas, R. J. Shneidman, B. Lacour, and T. Drüeke. 1985. Blood pressure development of the spontaneously hypertensive rat after concurrent manipulations of dietary Ca²⁺ and Na⁺. J. Clin. Invest. 76:1147-1154.

36. McCarron, D. A., N. N. Yung, B. A. Ugoretz, and S. Krutzik. 1981. Disturbances of calcium metabolism in the spontaneously hypertensive rat. *Hypertension (Dallas).* 3:1162–1167.

37. Lau, K., G. Chen, and B. Eby. 1984. Evidence for the role of PO₄ deficiency in antihypertensive action of a high-Ca diet. *Am. J. Physiol.* 246:H324-H330.

38. Fujisawa, Y., K. Kida, and H. Matsuda. 1984. Role of calcium in vitamin D metabolism with age in calcium and phosphorous metabolism in normal human subjects. J. Clin. Endocrinol. & Metab. 59:719-726.

39. Regan, T. J. 1983. Congestive heart failure in the diabetic. Annu. Rev. Med. 34:161-168.

40. Nixon, J. V., R. G. Murray, and C. Bryant. 1979. Early cardiovascular adaptation to simulated zero gravity. J. Appl. Physiol. Respir. Environ. Exercise Physiol. 46:541-548.

41. Dietlein, L. F. 1983. Spaceflight and the telltale heart. Am. J. Surg. 145:703-706.

42. Nyomba, B. L., R. Bouillon, W. Lissens, H. Van Baelen, and P. de Moor. 1985. 1,25-dihydroxyvitamin D and vitamin D-binding protein are both depressed in streptozotocin-diabetic rats. *Endocrinology*. 116: 2483–2488.

43. Ushakov, A. S., V. B. Spirichev, M. S. Belakovsky, I. N. Sergeev, and Y. I. Kondratyev. 1982. Calcium-phosphorous metabolism and prevention of its disorders in hypokinetic rats. *Physiologist*. 25:S45–S48.

44. Stewart, A. F., M. Adler, C. M. Byers, G. V. Segre, and A. E. Broadus. 1982. Calcium homeostasis in immobilization: an example of resorptive hypercaliuria. *N. Engl. J. Med.* 306:1136-1140.

45. Leach, C. S., S. I. Altchuler, and N. M. Cintron-Trevino. 1983. Circulating 1,25(OH)₂D₃ decreased following prolonged weightlessness. *Med. Sci. Sports Exercise.* 15:432–440.