

Effects of acid-base variables on ion transport in rat colon.

D S Goldfarb, ... , R W Egnor, A N Charney

J Clin Invest. 1988;**81**(6):1903-1910. <https://doi.org/10.1172/JCI113537>.

Research Article

Alterations in arterial acid-base variables have important effects on colonic electrolyte transport in vivo. To confirm the relative effects of these variables and to characterize the transport processes involved, we measured unidirectional ^{22}Na and ^{36}Cl fluxes across short-circuited, distal colonic mucosa of Sprague-Dawley rats. Stripped tissues were studied in Hepes buffer and in Ringer's solutions at HCO_3 concentrations of 11, 21, and 39 mM, and CO_2 tensions between 0 and 69.6 mmHg. Increases in PCO_2 , but not in either pH or HCO_3 concentration, caused similar increases in JNanet and JClnet (net flux of sodium and chloride, respectively) from -0.2 ± 0.3 and $-1.5 \pm 0.4 \mu\text{eq/cm}^2$ per h at $\text{PCO}_2 = 0$ to 6.8 ± 0.6 and $7.6 \pm 0.7 \mu\text{eq/cm}^2$ per h, respectively, at $\text{PCO}_2 = 69.6$ mmHg. These increases were accounted for by changes in Jms and were accompanied by small decreases in Isc . 1 mM acetazolamide decreased both JNanet and JClnet and their responses to increases in CO_2 . 0.75 mM luminal amiloride prevented the increase in sodium absorption, but did not affect the CO_2 -induced increase in chloride absorption. In the presence of amiloride, CO_2 increased JR (residual flux). 0.1 mM luminal furosemide did not affect the CO_2 -induced increases in JNanet in the absence or presence of amiloride. Changes in HCO_3 concentration did not alter JR . [...]

Find the latest version:

<https://jci.me/113537/pdf>



Effects of Acid-Base Variables on Ion Transport in Rat Colon

David S. Goldfarb, Richard W. Egnor, and Alan N. Charney

Nephrology Section, Veterans Administration Medical Center, New York, New York 10010; and Department of Medicine, New York University School of Medicine, New York, New York 10016

Abstract

Alterations in arterial acid-base variables have important effects on colonic electrolyte transport *in vivo*. To confirm the relative effects of these variables and to characterize the transport processes involved, we measured unidirectional ^{22}Na and ^{36}Cl fluxes across short-circuited, distal colonic mucosa of Sprague-Dawley rats. Stripped tissues were studied in Hepes buffer and in Ringer's solutions at HCO_3^- concentrations of 11, 21, and 39 mM, and CO_2 tensions between 0 and 69.6 mmHg. Increases in PCO_2 , but not in either pH or HCO_3^- concentration, caused similar increases in $J_{\text{net}}^{\text{Na}}$ and $J_{\text{net}}^{\text{Cl}}$ (net flux of sodium and chloride, respectively) from -0.2 ± 0.3 and -1.5 ± 0.4 $\mu\text{eq}/\text{cm}^2$ per h at $\text{PCO}_2 = 0$ to 6.8 ± 0.6 and 7.6 ± 0.7 $\mu\text{eq}/\text{cm}^2$ per h, respectively, at $\text{PCO}_2 = 69.6$ mmHg. These increases were accounted for by changes in J_{ms} and were accompanied by small decreases in J_{sc} . 1 mM acetazolamide decreased both $J_{\text{net}}^{\text{Na}}$ and $J_{\text{net}}^{\text{Cl}}$ and their responses to increases in CO_2 . 0.75 mM luminal amiloride prevented the increase in sodium absorption, but did not affect the CO_2 -induced increase in chloride absorption. In the presence of amiloride, CO_2 increased J^{R} (residual flux). 0.1 mM luminal furosemide did not affect the CO_2 -induced increases in $J_{\text{net}}^{\text{Na}}$ in the absence or presence of amiloride. Changes in HCO_3^- concentration did not alter J^{R} . We conclude that ambient CO_2 effects active, electroneutral sodium absorption in the rat distal colon. The process stimulated by CO_2 is dependent on mucosal carbonic anhydrase activity and most likely represents Na/H and Cl/ HCO_3^- ion exchange.

Introduction

The *in vivo* observation that arterial acid-base variables have a major effect on intestinal electrolyte and water transport suggested a mechanism other than neurohumoral by which the body maintains electrolyte homeostasis (1-3). The distal colon of the rat is of particular interest in this regard, because its electrolyte transport systems are well characterized (4-9), and its response to arterial PCO_2 and HCO_3^- concentration is specific and spans the physiological range of values for these variables. As in other transporting epithelia, the effects of CO_2 apparently were mediated through changes in the intracellular

pH (pHi)¹ of the colonic epithelium (10, 11), and were dependent on mucosal carbonic anhydrase activity (12). The data also suggested that the transport processes affected were electroneutral and included both Na/H ion exchange and Cl/ HCO_3^- ion exchange (3, 11, 13).

Several questions were raised by these findings that could not be addressed in studies of intact animals. These included determining whether (a) the action of PCO_2 was entirely a local effect on the colonic epithelium, (b) the electroneutral sodium absorptive process stimulated by CO_2 reflected coupled NaCl cotransport as well as Na/H exchange, (c) the effect of the plasma HCO_3^- concentration on net HCO_3^- secretion reflected transcellular (as opposed to intercellular) HCO_3^- movement, and (d) whether the responsive transport processes were pH-sensitive in the absence of ambient CO_2 .

These issues were addressed in the present study by examining distal colonic tissue in the Ussing (E. W. Wright, Guilford, CT) ion flux chamber. This experimental approach was especially suitable because the response of this tissue to CO_2 is immediate, of significant magnitude, and completely reversible (2, 3, 11-13). As compared with *in vivo* studies, this method allows for the precise control and manipulation of acid-base variables in the absence of secondary cardiovascular and systemic neurohumoral influences. In addition, we could examine the role of various electrolyte transport processes by comparing the effects of PCO_2 in the presence and absence of specific inhibitors of each process.

Methods

Male Sprague-Dawley rats weighing 250-350 g were maintained on a standard diet with free access to water. While the rats were under pentobarbital sodium anesthesia (5 mg/100 g body wt), the distal 10 cm of descending colon was removed, rinsed with 0.9% saline, and placed over a glass rod. The tissue was stripped of the serosa and segments were mounted in modified Ussing half-chambers (E. W. Wright) exposing 1.12 cm² surface area. One tissue pair was obtained from each animal.

The transepithelial potential difference (PD) was measured using calomel electrodes connected to the half-chambers by bridges containing 3% agar constituted in the same Ringer's solution used as bathing solution. The PD was expressed as serosal side positive with respect to the mucosa. Tissues were studied under short-circuited conditions except for 1-s intervals every 100 s, during which bipolar pulses of 0.5 mV yielded electrical current values that were used to calculate tissue conductance (G). The short-circuit current (J_{sc}) was passed across the tissue through 3% agar bridges via a voltage clamp (University of Iowa, Iowa City, IA). J_{sc} , by convention, was given the same polarity as PD. Tissues were paired for ion flux studies on the basis of differences in G no greater than 25%.

1. *Abbreviations used in this paper:* DIDS, 4,4'-diisothiocyanostilbene-2,2'-disulfonate; G, conductance; J_{sc} , short-circuit current; $J_{\text{ms}}^{\text{Cl}}$, mucosal to serosal chloride flux; $J_{\text{net}}^{\text{Cl}}$, net chloride flux; J_{ms} , mucosal to serosal flux; $J_{\text{ms}}^{\text{Na}}$, mucosal to serosal sodium flux; $J_{\text{net}}^{\text{Na}}$, net sodium flux; J_{net} , net flux; J_{sm} , serosal to mucosal flux; PD, potential difference; pHi, intracellular pH.

Portions of this work were previously published as abstracts in 1986. *Clin. Res.* 34:649a (Abstr.), and 1987. *Gastroenterology.* 92:1407 (Abstr.).

Address reprint requests to Dr. Alan N. Charney, Nephrology Section, Veterans Administration Medical Center, 408 First Avenue, New York, NY 10010.

Received for publication 13 April 1987 and in revised form 30 November 1987.

The Journal of Clinical Investigation, Inc.
Volume 81, June 1988, 1903-1910

Unidirectional fluxes of Na and Cl were measured by adding 2 μCi of ^{22}Na and 1 μCi of ^{36}Cl (100 Ci/g sp act; New England Nuclear, Boston, MA) to the mucosal side of one member of each tissue pair and the serosal side of the other. After an initial 15–30-min equilibration period, samples of the hot side were collected at the beginning and end of a 45-min experimental period for determination of each isotope's specific activity. Samples of the cold side were taken every 15 min during the 45-min period for determination of unidirectional ion flux. Samples from both sides were replaced with the appropriate Ringer's solutions. Ready-Solv HP (Beckman Instruments, Inc., Fullerton, CA) was added to each sample and counts were obtained in a liquid scintillation system (LS-3133P) and a gamma counter (both obtained from Beckman Instruments, Inc.). Mucosal to serosal (J_{ms}) and serosal to mucosal (J_{sm}) fluxes were calculated by standard formulae (4), and net flux (J_{net}) was calculated as the difference between them ($J_{ms} - J_{sm}$). The residual flux (J^R) was calculated by $J^R = I_{sc} - (J_{net}^{Na} - J_{net}^{Cl})$.

Acid-base variables

Paired tissues were bathed in 10 ml of identical Ringer's solutions at 37°C containing (in millimoles per liter): NaCl, 96; NaHCO_3 , 21; KCl, 4; Na_2HPO_4 , 2.4; NaH_2PO_4 , 0.4; Na gluconate, 18; CaSO_4 , 1.0; MgSO_4 , 1.2; and glucose, 10. Tissue pairs were studied during two or four consecutive 45-min experimental periods in which bathing solution HCO_3 concentration and pH, PCO_2 and pH, or HCO_3 concentration and PCO_2 were altered. Bicarbonate concentrations were altered by substituting Na gluconate or Hepes buffer for NaHCO_3 over the range 0 to 39 mM. In HCO_3 -free Ringer's solution, Hepes buffer was titrated with 1 M HCl or NaOH, and the solution was gassed with 100% O_2 . In HCO_3 - H_2CO_3 -buffered solutions, tissues were gassed with either room air ($\text{PCO}_2 = 0$ mmHg), 3% $\text{CO}_2/97\%$ O_2 ($\text{PCO}_2 = 21.9$ mmHg), 5% $\text{CO}_2/95\%$ O_2 ($\text{PCO}_2 = 34.5$ mmHg), or 11% $\text{CO}_2/89\%$ O_2 ($\text{PCO}_2 = 69.6$ mmHg). In preliminary time-control experiments, stable levels of sodium and chloride flux were observed for more than 6 h, and changes in transport induced by the acid-base variables were fully reversible.

The pH and PCO_2 of the bathing solutions studied in consecutive periods were measured before isotope addition with a Radiometer BMS 3 MK 2 system; with a (PHM 73 acid-base analyzer; London, Cleveland, OH). Bicarbonate concentration was calculated by the Henderson-Hasselbach equation as previously described (1, 2). In salt solutions, $pK' = 6.33 - 0.52(I)^{1/2} \pm 0.6(I)$, where 6.33 is the pK of the H_2CO_3 - HCO_3 buffer system in water, and I is the sodium plus potassium concentrations in moles per liter (2, 14). To correct pK' for pH, an empiric factor was determined in this laboratory: $pK'' = pK' - \log(1 + \log^{pH-8.7})$. The CO_2 solubility in salt solutions was adjusted for ionic strength: $\alpha = 10^{-1.482-0.085(I)}$ (2, 14).

Inhibitors

We then examined whether inhibitors affected the change in transport caused by the acid-base variables. Electrolyte transport was measured at a bathing solution HCO_3 concentration of 21 mM during consecutive periods of $\text{PCO}_2 = 21.6$ and 69.6 mmHg before and after the addition of the specific inhibitor.

Acetazolamide. Sodium acetazolamide (Lederle Parenterals, Inc., Carolina, PR) was added to both mucosal and serosal bathing solutions at a final concentration of 0.1 or 1.0 mM. Electrolyte transport was measured at a bathing solution HCO_3 concentration of 21 mM during consecutive periods of $\text{PCO}_2 = 21.6$ and 58.6 mmHg, and $\text{PCO}_2 = 21.6$ and 69.6 mmHg.

Amiloride. Amiloride (Merck Sharp & Dohme Research Laboratories, West Point, PA) was added to the mucosal bathing solution at a final concentration of 0.75 mM. The effect of amiloride also was studied in a bathing solution in which choline chloride was substituted for sodium chloride so as to reduce the sodium concentration to 50 mM.

Furosemide. Furosemide (Sigma Chemical Co., St. Louis, MO) was added to the mucosal bathing solution at a final concentration of 0.1 mM.

Amiloride and furosemide. To determine whether amiloride and furosemide were additive in their effects and whether the effect of furosemide was affected by the presence of amiloride, 0.75 mM amiloride and 0.1 mM furosemide were both added to the mucosal bathing solution. The effects of increasing PCO_2 were then compared with the changes caused by CO_2 in the presence of amiloride or furosemide.

4,4'-Diisothiocyanostilbene-2,2' disulfonate (DIDS). DIDS was added to the mucosal side of a 21-mM HCO_3 Ringer's solution at a final concentration of 0.1 mM or 1 mM.

Statistics

Calculated values for the 3 15-min samples for each condition were averaged and the values for each experimental group were expressed as the mean \pm SE. Statistical analyses consisted of paired and unpaired, two-tailed t tests, one-way analysis of variance, and linear regression analysis by least squares (15). Correlation coefficients were calculated from the mean values of the experimental groups. Corrections were made for multiple t tests when appropriate, and a $2P$ value < 0.05 was considered significant.

Results

Acid-base variables

The effects of varying the PCO_2 on electrolyte transport at a bathing solution HCO_3 concentration of 21 mM are shown in Table I. At $\text{PCO}_2 = 0$ mmHg, net Na absorption was zero and net chloride secretion was observed. The residual ion flux, J^R , also was zero. Net Na and Cl absorption were noted at $\text{PCO}_2 = 21.6$ mmHg, and progressively increased so that at $\text{PCO}_2 = 69.6$ mmHg, net Na and Cl absorption had increased $\sim 60\%$. The increase in absorption was accounted for by an increase in J_{ms} rather than by a reduction in J_{sm} . Increases in PCO_2 between 0 and 69.6 mmHg did not alter PD, but did increase J^R ($P < 0.05$) and decrease the I_{sc} ($P < 0.01$) and conductance ($P < 0.02$). These changes in ion flux and G were completely reversible, and occurred regardless of the order in which PCO_2 was altered.

To determine to what extent the observed responses were effects of PCO_2 and not pH, we measured electrolyte fluxes at bathing solution HCO_3 concentrations of 39 and 11 mM at a constant PCO_2 of 34.5 mmHg. As shown in Table II, there were no significant differences in the net absorption of sodium or chloride in these solutions of markedly different HCO_3 concentration and pH. Indeed, the pH range examined in this experiment (7.65–7.14) was similar to that caused by changing PCO_2 from 21.6 to 69.6 mmHg (compare Tables I and II). There also was no effect of pH or HCO_3 concentration on J^R , although J^R tended to be greater in 39 mM HCO_3 than in 11 mM HCO_3 Ringer's solution.

Nevertheless, because these data were obtained (by necessity) in separate groups of tissues, a small effect of pH may have been obscured by intertissue variability. We therefore measured ion flux in a HCO_3 -free Hepes-Ringer solution gassed with 100% O_2 that was titrated during consecutive periods with HCl or NaOH. As shown in Table II, J_{ms}^{Na} , J_{net}^{Na} , J_{ms}^{Cl} , and J_{net}^{Cl} were markedly reduced in the virtual absence of ambient HCO_3 and CO_2 . J^R was zero under these conditions. When bathing solution pH was reduced from 7.59 to 7.09, very small but significant changes were noted in I_{sc} , PD, G , and the mucosal to serosal and net fluxes of Na and Cl. These changes were qualitatively similar to, although much smaller than, those seen when a similar pH change was induced by altering PCO_2 (Table I). In addition, the decreases in I_{sc} and

Table I. Effect of CO₂ on Colonic Electrolyte Transport

PCO ₂	pH	Isc	PD	G	Na flux			Cl flux			J ^R
					J _{ms}	J _{sm}	J _{net}	J _{ms}	J _{sm}	J _{net}	
mmHg		μeq/cm ² per h	mV	mS/ cm ²	μeq/cm ² per h			μeq/cm ² per h			
0	8.42	1.8	4.4	11.1	5.9	6.1	-0.2	7.4	8.9	-1.5	0.4
	±0.03	±0.1	±0.1	±0.4	±0.3	±0.3	±0.3	±0.4	±0.6	±0.4	±0.5
21.6	7.63	1.6	5.0	8.9	9.2	5.0	4.2	12.8	8.0	4.8	2.2
	±0.7	±0.01	±0.2	±0.5	±0.6	±0.5	±0.3	±0.4	±0.5	±0.3	±0.5
34.5	7.41	1.4	4.7	8.5	10.1	4.7	5.5	14.6	8.0	6.6	2.5
	±0.8	±0.01	±0.2	±0.7	±0.6	±0.8	±0.2	±0.8	±0.8	±0.4	±0.8
69.6	7.10	0.9	3.2	7.9	11.7	4.8	6.8	16.5	9.0	7.6	1.7
	±1.2	±0.01	±0.1	±0.5	±0.5	±0.9	±0.3	±0.6	±0.7	±0.4	±0.7
P*	<0.001	<0.001	<0.025			<0.01	<0.001	<0.001	<0.001	<0.001	<0.05

Values are means±SE. Number of tissue pairs studied at a PCO₂ of 0 = 4, at a PCO₂ of 21.6 = 17, at a PCO₂ of 34.5 = 15, and at a PCO₂ 69.6 = 17. Data were obtained in separate groups of tissue pairs bathed in 140 mM Na and 21 mM HCO₃-Ringer's solutions at one or more CO₂ tensions. * Four experimental groups were compared by one-way analysis of variance.

PD in Hepes buffer were entirely accounted for by increases in net Cl transport since as shown in Table II, $I_{sc} = J_{net}^{Na} - J_{net}^{Cl}$ and $J^R = 0$.

We then examined the possibility that the concentration of HCO₃ altered the effect of CO₂ on Na and Cl absorption. We measured ion fluxes in the 39-mM HCO₃ Ringer's solution at CO₂ = 69.6 mmHg (pH 7.35) and in the 11-mM HCO₃ Ringer's solution at PCO₂ = 21.6 mmHg (pH 7.35). The differences in net Na flux (1.3 μeq/cm² per h), net Cl flux (3.6 μeq/cm² per h), I_{sc} (-0.9 μeq/cm² per h), and G (0.1 mS/cm²) at these CO₂ tensions were similar to the differences observed between similar PCO₂ levels in the 21-mM HCO₃ Ringer's solution shown in Table I.

The relative effects of bathing solution PCO₂, pH, and HCO₃ concentration on ion flux were confirmed by correlating these variables in all 75 tissue pairs in all the acid-base groups described above (except the groups exposed to room air or bathed in Hepes-Ringer's solution). As shown in Fig. 1, net sodium absorption correlated only with bathing solution PCO₂ ($r = 0.83$, $P < 0.02$) and did not correlate with either pH or HCO₃ concentration. When unidirectional fluxes were examined, J_{ms}^{Na} correlated with PCO₂ ($r = 0.97$, $P < 0.001$). The slopes of both of these relations were 0.04, indicating that for each 10-mmHg increment in PCO₂, J_{net}^{Na} flux increased 0.4 μeq/cm² per h.

Although the relation between CO₂ and chloride absorp-

Table II. Effect of pH on Colonic Electrolyte Transport in HCO₃ and Hepes-buffered Ringer's Solutions

HCO ₃	PCO ₂	pH	Isc	PD	G	Na flux			Cl flux			J ^R	
						J _{ms}	J _{sm}	J _{net}	J _{ms}	J _{sm}	J _{net}		
mM	mmHg		μeq/cm ² per h	mV	mS/cm ²	μeq/cm ² per h			μeq/cm ² per h				
HCO ₃ Ringer's	39	35.8	7.65	1.3	3.8	9.2	10.2	5.7	4.5	13.2	7.4	5.8	2.6
		±1.0	±0.01	±0.1	±0.4	±0.6	±0.7	±0.7	±0.8	±0.7	±0.7	±0.7	±0.4
	11	34.3	7.14	1.4	5.2	7.8	10.4	5.4	5.1	15.8	10.4	5.4	1.7
		±0.7	±0.01	±0.2	±0.8	±2.1	±0.7	±0.3	±0.8	±0.9	±0.4	±0.8	±0.4
P*		<0.001		<0.001						<0.05	<0.001		
Hepes Ringer's	0	0	7.59	1.7	6.5	7.0	5.5	4.4	1.1	8.1	8.4	-0.2	0.3
			±0.01	±0.1	±0.6	±0.2	±0.2	±0.2	±0.4	±0.7	±0.5	±0.8	±0.5
	0	0	7.09	1.3	5.7	6.2	6.1	4.0	2.2	8.9	7.7	1.2	0.3
			±0.01	±0.2	±0.7	±0.3	±0.4	±0.3	±0.6	±0.7	±0.5	±0.8	±0.3
P†		<0.001	<0.005	<0.05	<0.001	<0.05	<0.05	<0.01	<0.01	<0.025	<0.005	<0.005	

Values are means±SE. Number of tissue pairs studied at 39 mM HCO₃-Ringer's solution = 12, 11 mM HCO₃-Ringer's = 14, and 21 mM Hepes = 9. Data for HCO₃-Ringer's solution were obtained in separate groups of tissue pairs bathed in 39 mM HCO₃-Ringer's solution or 11 mM HCO₃-Ringer's solution. Data for Hepes-Ringer's solution were obtained in one group of tissues bathed in 21 mM Hepes-Ringer's solution titrated during consecutive periods with HCl or NaOH. * Unpaired *t* test, † paired *t* test.




Figure 1. Relation of sodium absorption to bathing solution PCO_2 . J_{ms}^{Na} and J_{net}^{Na} were measured in Ringer's solutions containing 11, 21, or 39 mM HCO_3^- during consecutive periods in which PCO_2 was altered as indicated by the dashed lines. Both J_{ms}^{Na} and J_{net}^{Na} increased with increasing PCO_2 . Values are means \pm SE.

tion appeared to be curvilinear (Fig. 2), when analyzed by linear regression analysis, similar results were obtained for chloride flux. Net chloride absorption correlated with PCO_2 ($r = 0.92$, $P < 0.005$), but not with either pH or HCO_3^- concentration. A correlation also was found for J_{ms}^{Cl} and PCO_2 ($r = 0.72$, $P < 0.05$). The slopes of these relations were similar to the slope relating sodium flux to PCO_2 . There was an $\sim 0.6 \mu eq/cm^2$ per h increase in J_{net}^{Cl} per 10-mmHg increase in PCO_2 .

Inhibitors

In another series of experiments, inhibitors of various electrolyte transport processes were added to 21 mM HCO_3^- Ringer's bathing solution and the effect of increasing CO_2 tension was measured. Alterations in Na and Cl fluxes in the presence and absence of the inhibitor were then compared.

Acetazolamide. To examine the effect of inhibiting mucosal carbonic anhydrase activity, 0.1 mM sodium acetazolamide was added to both mucosal and serosal bathing solutions. At $PCO_2 = 21.6$ mmHg, this addition reduced J_{ms} and J_{net} for chloride $\sim 50\%$ (Table III). As shown in Table IV, however, the increments in net sodium and chloride flux caused by increasing PCO_2 from 21.6 to 69.6 mmHg, although somewhat smaller, were not significantly different in the presence and absence of acetazolamide. The presence of acetazolamide nevertheless was associated with an increase in J^R in response to this increase in PCO_2 .

This partial effect of acetazolamide suggested the possibility that the 48-mmHg increase in CO_2 tension (69.6 minus 21.6 mmHg) had overcome the effects of 0.1 mM acetazolamide. The effect of increasing PCO_2 from 21.6 to 58.6 mmHg on net sodium and chloride fluxes was therefore compared with the increments caused by increasing PCO_2 from 21.6 to 69.6 mmHg. In fact, these increments were similar. When




Figure 2. Relation of chloride absorption to bathing solution PCO_2 . J_{ms}^{Cl} and J_{net}^{Cl} were measured in Ringer's solutions containing 11, 21, or 39 mM HCO_3^- during consecutive periods in which PCO_2 was altered as indicated by the dashed lines. Both J_{ms}^{Cl} and J_{net}^{Cl} increased with increasing PCO_2 . Values are means \pm SE.

PCO_2 was raised from 21.6 to 58.6 mmHg in the presence of 0.1 mM acetazolamide, J_{net}^{Na} increased from 2.7 ± 0.8 to $5.0 \pm 1.2 \mu eq/cm^2$ per h and J_{net}^{Cl} increased from 2.8 ± 0.7 to $6.7 \pm 1.0 \mu eq/cm^2$ per h (compare with Tables III and IV). These findings suggested that a 0.1-mM concentration of acetazolamide was insufficient to completely inhibit mucosal carbonic anhydrase in our system. Consequently, we repeated the experiment in the presence of 1.0 mM acetazolamide. Under these conditions, J_{ms} for both sodium and chloride were markedly reduced, resulting in zero net sodium absorption and net chloride secretion at $PCO_2 = 21.6$ mmHg. In addition, I_{sc} tended to decrease and J^R changed direction (Table III). When PCO_2 was increased from 21.6 to 69.6 mmHg, the increases in J_{ms} for both Na and Cl were significantly smaller than in the absence of acetazolamide (Table IV).

Amiloride. The participation of electrogenic Na transport and Na/H exchange in the colonic response to PCO_2 were examined by adding 0.75 mM amiloride to the mucosal bathing solution. At a CO_2 tension of 21.6 mmHg, amiloride decreased J_{ms}^{Na} 43%, J_{net}^{Na} 69%, and J_{ms}^{Cl} 23% (Table III). As shown in Table IV, the presence of amiloride markedly reduced but did not completely prevent the increments in J_{ms}^{Na} and J_{net}^{Na} produced by increasing the PCO_2 to 69.6 mmHg. In addition, the increments in J_{ms}^{Cl} and J_{net}^{Cl} caused by increasing PCO_2 were unaffected by the presence of amiloride. These increments in chloride flux were associated with increases in J^R that were not observed in the absence of amiloride.

We also studied amiloride in a HCO_3^- -Ringer's bathing solution in which the sodium concentration was reduced to 50 mM (Table V). In the absence of amiloride, increases in PCO_2

Table III. Effect of Various Inhibitors on Colonic Electrolyte Transport at a PCO₂ of 21.6 mmHg

Addition	Isc ^a μeq/cm ² per h	PD mV	G ^b mS/cm ²	Na flux			Cl flux			J ^R
				J _{ms} ^c	J _{sm}	J _{net} ^c	J _{ms} ^c	J _{sm}	J _{net} ^c	
				μeq/cm ² per h			μeq/cm ² per h			
None	1.6 ±0.2	5.0 ±0.5	8.9 ±0.6	9.2 ±0.5	5.0 ±0.3	4.2 ±0.4	12.8 ±0.5	8.0 ±0.3	4.8 ±0.4	2.2 ±0.3
Acetazolamide (0.1 mM)	1.3 ±0.1	4.8 ±0.7	8.4 ±0.7	7.7 ±0.7	5.1 ±0.5	2.6 ±0.6	10.2 [§] ±0.7	8.1 ±0.4	2.1 ±0.5	0.8 ±0.6
Acetazolamide (1.0 mM)	0.9 ±0.2	4.2 ±0.5	5.7 [§] ±0.4	3.8 ±0.1	3.8 ±0.2	0.0 ±0.2	5.3 ±0.4	7.6 ±0.7	-2.3 ±0.5	-1.3 ±0.5
Amiloride (0.75 mM)	1.2 ±0.1	6.2 ±0.4	5.4 ±0.5	5.0 ±0.5	3.7 ±0.4	1.3 [§] ±0.7	10.0 [§] ±0.7	6.9 ±0.7	3.1 ±0.8	3.1 ±0.6
DIDS (0.1 mM)	2.0 ±0.0	7.4 ±1.0	7.7 ±0.4	6.3 [§] ±0.7	4.5 ±0.3	1.8 [§] ±0.8	10.0 [§] ±0.7	7.9 ±0.5	2.1 ±1.2	2.3 ±0.4
Furosemide (0.1 mM)	1.6 ±0.3	6.3 ±1.2	7.3 ±0.4	6.5 [§] ±0.7	4.6 ±0.4	1.9 [§] ±1.1	9.6 [§] ±0.8	7.6 ±0.4	2.0 [§] ±0.6	1.8 ±0.6
Amiloride (0.75 mM) Furosemide (0.1 mM)	1.7 ±0.3	6.4 ±0.9	7.1 ±0.3	4.1 ±0.4	4.5 ±0.3	-0.4 ±0.5	8.8 ±0.5	9.4 ±0.5	-0.6 ±0.6	1.6 ±0.6

Values are means±SE. Number of tissue pairs studied with no addition = 17, 0.1 mM acetazolamide = 15, 1.0 mM acetazolamide = 6, 0.75 mM amiloride = 8, 0.1 mM DIDS = 6, 0.1 mM furosemide = 6, and 0.75 mM amiloride plus 0.1 mM furosemide = 7. Separate groups of tissue pairs were studied in 140 mM Na and 21 mM HCO₃⁻-Ringer's solution at a CO₂ tension of 21.6 mmHg (pH 7.63). * P < 0.002, † P < 0.001 when the mean values for all groups were compared by one-way analysis of variance. § P < 0.05, || P < 0.005 when compared with addition: none by unpaired t test.

Table IV. Effect of Increasing PCO₂ on Colonic Electrolyte Transport in the Presence of Various Inhibitors

Addition	ΔIsc μeq/cm ² per h	ΔPD mV	ΔG mS/cm ²	ΔNa flux			ΔCl flux			ΔJ ^R
				J _{ms}	J _{sm}	J _{net}	J _{ms}	J _{sm}	J _{net}	
				μeq/cm ² per h			μeq/cm ² per h			
None	-0.7 ±0.1	-1.8 ±0.4	-1.0 ±0.2	2.5 ±0.5	-0.2 ±0.1	2.6 ±0.5	3.8 ±0.4	1.0 ±0.3	2.8 ±0.5	-0.6 ±0.3
Acetazolamide (0.1 mM)	-0.6 ±0.2	-2.7 ±0.5	-1.1 ±0.2	1.5 ±0.3	-0.5 ±0.2	2.0 ±0.4	3.0 ±0.7	-0.6* ±0.4	3.6 ±0.7	1.0* ±0.6
Acetazolamide (1.0 mM)	-0.4 ±0.1	-1.5 ±0.4	-0.5 ±0.2	0.9* ±0.2	-0.4 ±0.3	1.3* ±0.4	1.1* ±0.3	-1.5* ±0.5	2.6 ±0.4	0.9* ±0.4
Amiloride (0.75 mM)	-0.7 ±0.9	-2.9 ±0.5	-0.8 ±0.7	0.6* ±0.2	-0.5 ±0.2	1.1* ±0.2	3.2 ±0.2	0.4 ±0.3	2.8 ±0.3	0.9* ±0.3
DIDS (0.1 mM)	-1.0 ±0.2	-3.3* ±0.9	-1.3 ±0.2	2.5 ±0.7	-0.1 ±0.3	2.6 ±0.9	3.9 ±0.8	0.2 ±0.6	3.6 ±0.5	0.0 ±0.5
Furosemide (0.1 mM)	-0.6 ±0.2	-1.8 ±0.6	-0.9 ±0.2	1.9 ±0.4	-0.5 ±0.2	2.4 ±0.4	2.9 ±0.6	0.2 ±0.3	2.8 ±0.5	-0.2 ±0.2
Amiloride (0.75 mM) Furosemide (0.1 mM)	-0.7 ±0.2	-2.3 ±0.7	-0.8 ±0.2	0.2* ±0.3	-0.5* ±0.1	0.7* ±0.3	2.5* ±0.5	0.5 ±0.3	2.0 ±0.5	0.5* ±0.4

Values are means±SE. Number of tissue pairs studied with no addition = 17, with 0.1 mM acetazolamide = 15, with 1.0 mM acetazolamide = 6, with 0.75 mM amiloride = 8, with 0.1 mM DIDS = 6, with 0.1 mM furosemide = 6, and with 0.75 mM amiloride plus 0.1 mM furosemide = 7. Separate groups of tissue pairs were studied in 140 mM Na and 21 mM HCO₃⁻-Ringer's solution at CO₂ tensions of 21.6 (pH 7.63) and 69.6 mmHg (pH 7.10). Data (deltas) represent the differences between values at 69.6 mmHg minus values at 21.6 mmHg. Data for addition: none were derived from experiments presented in Table I. * P < 0.05 when compared with addition: none by unpaired t test.

caused qualitatively and quantitatively similar changes in ion flux, as were observed in a bathing solution containing 140 mM sodium (compare Tables IV and V). The effect of amiloride at a PCO₂ of 21.6 mmHg also was similar at 50 and 140 mM sodium: amiloride reduced J_{net}^{Na} 70% and J_{net}^{Cl} 47%, and increased J^R . Nevertheless, when the PCO₂ was raised in 50 mM sodium Ringer's solution, amiloride completely inhibited the increases in J_{ms}^{Na} and J_{net}^{Na} .

Furosemide. To determine whether a cotransport process was involved in the colonic response to CO₂, 0.1 mM furosemide was added to the mucosal bathing solution. At PCO₂ = 21.6 mmHg, furosemide decreased J_{net}^{Na} 55% and J_{net}^{Cl} 67%, respectively (Table III). This effect was additive to that of amiloride: the mucosal addition of both amiloride and furosemide at PCO₂ = 21.6 mmHg resulted in J_{net}^{Na} ($P < 0.05$) and J_{net}^{Cl} ($P < 0.005$) values that were significantly lower than for amiloride alone. In fact, J_{net}^{Na} and J_{net}^{Cl} did not differ from zero under these conditions.

Despite this effect of furosemide at PCO₂ = 21.6 mmHg, a furosemide-inhibitable process did not appear to participate in the colonic response to PCO₂. As shown in Table IV, furosemide did not alter the increments in J_{net}^{Na} or J_{net}^{Cl} caused by increasing bathing solution PCO₂ to 69.6 mmHg. Nevertheless, it was possible that in the presence of amiloride, a role for this process might be revealed. The effect of CO₂, therefore, was measured in tissues exposed to mucosal furosemide in a 21-mM HCO₃ Ringer's solution containing amiloride. As shown in Table IV, the increments in J_{net}^{Na} and J_{net}^{Cl} in response to increasing PCO₂ under these conditions were no different than in tissues exposed to similar PCO₂ changes in the presence of amiloride alone: ΔJ_{net}^{Na} 0.7±0.3 vs. 1.1±0.2 $\mu\text{eq}/\text{cm}^2$ per h, NS; ΔJ_{net}^{Cl} 2.0±0.5 vs. 2.8±0.3 $\mu\text{eq}/\text{cm}^2$ per h, NS).

DIDS. The addition of DIDS to the mucosal bathing solution at a final concentration of 0.1 mM reduced J_{ms}^{Na} , J_{net}^{Na} , and J_{net}^{Cl} at PCO₂ = 21.6 mmHg. A similar reduction in J_{net}^{Cl} did not reach statistical significance. Nevertheless, DIDS at 0.1 mM or at a 10-fold higher concentration did not alter the increments in J_{net}^{Na} or J_{net}^{Cl} induced by increasing the PCO₂ to 69.6 mmHg (data for 0.1 mM DIDS shown in Table IV).

Discussion

In previous studies we reported that physiologic changes in arterial PCO₂ have major effects on electrolyte and water absorption in the rat colon (1-3). The in vivo experimental conditions, however, allowed only tentative conclusions regarding the specificity and site of action of CO₂ (11, 13). The current findings confirm that the action of CO₂ on sodium and chloride absorption is predominant among the acid-base variables, and represents a local effect of CO₂ on the intestinal mucosa. A significant effect of pH was ruled out by the absence of a correlation between bathing solution pH and either J_{net}^{Na} or J_{net}^{Cl} , and by the minimal effect of pH on ion flux (between 7.09 and 7.65) when PCO₂ remained unchanged. The overall magnitude of the CO₂ effect was of interest as well: J_{net}^{Na} and J_{net}^{Cl} increased ~ 0.5 $\mu\text{eq}/\text{cm}^2$ per h per 10-mmHg increase in PCO₂. These increments are equivalent to the 60% increases in sodium and chloride absorption observed in vivo over a similar range of CO₂ tensions (2 $\mu\text{eq}/\text{cm}$ length per h per 10-mmHg change in arterial PCO₂) (2, 3).

The colonic transport pathways affected by CO₂ appear to be active, electroneutral sodium and chloride absorptive processes. The importance of electroneutral absorption was originally suggested during in vivo studies of rat colon by the minimal changes in PD, and the equivalent increases in net sodium and chloride absorption in response to increases in arterial PCO₂ (2, 3, 12). In addition, the rabbit distal colon, which absorbs sodium by conductive, electrogenic pathways (16), did not respond at all to alterations in arterial PCO₂ (17). The nature of this electroneutral process remained uncertain, however, although the in vivo data were most consistent with the participation of Na/H and Cl/HCO₃ ion exchange processes (3, 13). It was presumably these exchangers, located along the luminal membranes of colonic epithelial cells, that responded to the changes in pH_i and intracellular HCO₃ concentrations that mediated the CO₂ effect (11).

In the present study, we examined the role of the various transport pathways by comparing the increments in ion flux caused by CO₂ in the presence and absence of several transport

Table V. Effect of CO₂ and Amiloride on Colonic Electrolyte Transport in Low Sodium Ringer's Solution

Addition	PCO ₂	pH	Isc	PD	G	Na flux			Cl flux			J ^R
						J _{ms}	J _{sm}	J _{net}	J _{ms}	J _{sm}	J _{net}	
	mM		$\mu\text{eq}/\text{cm}^2$ per h	mV	mS/cm ²	$\mu\text{eq}/\text{cm}^2$ per h			$\mu\text{eq}/\text{cm}^2$ per h			
None	21.3	7.64	1.5	8.3	4.6	8.7	1.8	6.9	12.8	7.3	5.5	0.1
	±0.6	±0.01	±0.3	±1.1	±0.6	±2.3	±0.4	±2.1	±1.8	±0.7	±1.3	±1.4
	71.1	7.11	0.7	5.1	3.9	10.5	1.6	8.9	16.3	8.0	8.3	0.1
	±0.3	±0.0	±0.1	±0.8	±0.6	±2.5	±0.3	±2.3	±2.2	±0.5	±2.0	±1.8
<i>P</i> *	<0.001	<0.001	<0.02	<0.001	<0.02	<0.005		<0.005	<0.001	<0.05	<0.01	
Amiloride (0.75 mM)	21.3	7.64	1.2	6.8	4.1 [‡]	3.8 [‡]	1.7	2.1 [§]	10.2 [§]	7.4	2.9 [§]	1.9
	±0.6	±0.01	±0.3	±0.9	±0.6	±0.8	±0.4	±0.6	±0.9	±0.7	±0.5	±0.5
	71.1	7.11	0.5	3.4	3.7	3.9	1.4	2.4	12.9	8.4	4.5	2.8
	±0.3	±0.0	±0.1	±0.6	±0.4	±0.9	±0.3	±0.7	±1.4	±0.7	±1.0	±0.8
<i>P</i> *	<0.001	<0.001		<0.01					<0.01	<0.05	<0.05	

Values are means±SE. Number of tissue pairs studied = 7. Tissue pairs were studied in 50 mM Na, 21 mM HCO₃ Ringer's solution at CO₂ tensions of 21.3 and 71.1 mmHg in the presence and absence of 0.75 mM luminal amiloride. * Paired *t* test. ‡ $P < 0.02$, § $P < 0.05$ when compared with addition: none at PCO₂ = 21.3 mmHg by paired *t* test.

inhibitors. This method allowed us to differentiate between the presence of a transport process and its participation in the colonic response to CO_2 (12). The most dramatic example of such a dissociation was our finding that 0.1 mM furosemide inhibited both sodium and chloride absorption $\sim 55\%$ at $\text{PCO}_2 = 21.6$ mmHg, but did not reduce the increments in absorption caused by increasing the PCO_2 . We interpret this finding to mean that an electroneutral sodium chloride absorptive process inhibited by furosemide, although present in rat colonic epithelial cells and an important pathway for sodium absorption, does not respond to alterations in PCO_2 . It is not clear at present whether furosemide inhibited a sodium chloride absorptive process distinct from Na/H and Cl/HCO₃ ion exchange, or whether these antiporters are sensitive to furosemide (but not CO_2) under certain experimental and physiologic conditions. Indeed, in one recent study using apparently similar methodology, furosemide-sensitive sodium chloride absorption was not found in the rat distal colon (18).

We may compare the effects of furosemide on colonic absorption with those of luminal amiloride. At a PCO_2 of 21.6 mmHg, amiloride also markedly inhibited sodium absorption. In fact, at low CO_2 tensions, the combination of amiloride and furosemide reduced net sodium and chloride absorption to zero. However, whereas CO_2 and furosemide apparently affect separate sodium absorptive pathways, CO_2 and amiloride modulate the same process. We found that the increment in net sodium absorption caused by an increase in PCO_2 was markedly diminished in the presence of amiloride. At luminal bathing solution concentrations reported to inhibit Na/H ion exchange in a variety of epithelia and membrane vesicle preparations (19–22), amiloride reduced the increment in sodium absorption $\sim 60\%$. The presence of amiloride did not affect the increment in chloride absorption, but was associated with a PCO_2 -induced increase in J^{R} that was not observed in its absence. These findings suggest that both Na/H and Cl/HCO₃ ion exchange processes are present in the colonic epithelium and respond to alterations in PCO_2 .

There are two difficulties with this interpretation of the data. First, amiloride did not completely prevent the CO_2 -induced increase in sodium absorption. One possible explanation was that a furosemide-inhibitable sodium absorptive process might have become sensitive to CO_2 (or pH) when Na/H ion exchange was inhibited. We ruled out this possibility by finding that amiloride plus furosemide did not have a greater effect on CO_2 -induced changes in sodium absorption than amiloride alone (Table IV). Another explanation for the incomplete amiloride inhibition was methodological: perhaps the concentration of sodium in the bathing solution (140 mM) interfered with amiloride-inhibition of Na/H ion exchange (19, 22, 23). We examined this possibility by substituting choline chloride for sodium chloride to lower the bathing solution sodium concentration to 50 mM. As a result of this change, amiloride completely prevented the PCO_2 -induced increase in net sodium absorption. This finding appears to confirm the suggestion that the only sodium absorptive process responsive to CO_2 is Na/H ion exchange. It also may account for the apparent absence of amiloride-inhibitable Na/H ion exchange in other studies of rat distal colon in which a similar bathing solution sodium concentration of 140 mM was used (19, 21, 24, 25).

A second difficulty in concluding that both Na/H and Cl/HCO₃ ion exchange are responsive to CO_2 is our finding that

mucosal DIDS at 0.1 and 1 mM did not inhibit CO_2 -induced increments in chloride absorption. DIDS and related disulfonic stilbenes are reported to inhibit anion exchange in intact intestinal epithelia and membrane vesicles (26, 27). In recent studies of the rat proximal and distal colon, however, 2 mM mucosal DIDS and 1 mM 4-acetamido-4'-isothiocyanatostilbene-2,2'-disulfonic acid did not alter $J_{\text{net}}^{\text{Na}}$, $J_{\text{net}}^{\text{Cl}}$, or I_{sc} despite other evidence suggesting that anion exchange was present (18, 28). Similar observations have been made in rabbit ileum (29) and rabbit colon (30). Our results therefore do not absolutely exclude the participation of Cl/HCO₃ ion exchange in the rat distal colon's response to CO_2 . As suggested by others, DIDS may not be a sensitive probe for anion exchange under all experimental conditions.

A role for mucosal carbonic anhydrase activity was tested for in a similar way: by inhibiting the enzyme and stimulating absorption with CO_2 . In fact, acetazolamide markedly reduced the effect of PCO_2 on sodium absorption. This effect of acetazolamide also was observed during the luminal perfusion of the distal colon in intact anesthetized rats (12). The concentration of acetazolamide required to significantly reduce sodium chloride absorption, however, was greater (1 mM) than the concentration required to inhibit the enzyme in broken cell preparations (0.1 mM) (12, 31). Such high concentrations are often needed to inhibit carbonic anhydrase activity in gastrointestinal tissues mounted in Ussing flux chambers (4, 18, 32). This may reflect the moderate water and lipid solubility of acetazolamide and its limited diffusion into tissues (31). Nevertheless, in some tissues, these acetazolamide concentrations may affect ion transport by mechanisms other than carbonic anhydrase inhibition (31). Such nonspecific or toxic effects, however, have not been observed in the rat distal colon (12). The specific effect of carbonic anhydrase inhibition is to alter pHi and as a consequence, electrolyte transport (5, 12, 31, 33). This role for pHi in the colon was suggested in previous studies by the finding that mucosal pHi varied directly with changes in arterial PCO_2 and inversely with changes in sodium absorption (11). In the present study, it is thus likely that acetazolamide blunted the effect of PCO_2 on colonic absorption by altering the pHi response to PCO_2 .

Finally, it was noteworthy that the effect of the plasma HCO₃ concentration on colonic net HCO₃ secretion observed in vivo (3) was not found in the Ussing flux chamber. Only a slightly higher level of J^{R} was found at a bathing solution HCO₃ concentration of 39 mM (pH 7.65) than at 11 mM (pH 7.14). Although J^{R} may not always be a reliable measure of net HCO₃ secretion (4, 29), in our study, J^{R} appeared to reflect HCO₃ transport. For example, J^{R} increased in response to increases in PCO_2 in the presence of either amiloride or acetazolamide, additions that would be expected to cause disproportionate responses in Na/H and Cl/HCO₃ ion exchange. In addition, J^{R} was zero and did not respond to changes in pH in the absence of bathing solution CO_2 and HCO₃ (Hepes buffer). We believe the difference between the in vivo and in vitro findings may be due to the equivalent HCO₃ concentrations in the mucosal and serosal solutions in the chamber, and the differing HCO₃ concentrations in the plasma and luminal perfusate in vivo. The in vivo results thus may have reflected the effect of a transmucosal HCO₃ concentration gradient on intercellular HCO₃ movement. It is also likely that the relatively large chamber volume and unpaired design of the HCO₃ experiments limited our ability to detect small changes in net

HCO₃ secretion. This is especially relevant in light of the recent finding, using a new in vitro technique, that extracellular HCO₃ concentration apparently does modulate active colonic HCO₃ secretion (9, 34). For these reasons, our conclusions regarding the effects of HCO₃ concentration and various inhibitors on J^R and HCO₃ secretion by the colon must be tentative.

In summary, these studies establish a primary role for CO₂ in regulating active sodium chloride absorption by the rat distal colon. The transport pathways stimulated by increasing PCO₂ appear to be Na/H and Cl/HCO₃ ion exchange processes located along the luminal border of colonic epithelial cells. This effect of CO₂ requires mucosal carbonic anhydrase activity to generate the pHi change that mediates the observed changes in electrolyte transport. In general, these findings confirm previous observations, although a separate effect of extracellular HCO₃ concentration on active colonic HCO₃ secretion was not found.

Acknowledgments

The authors appreciate the technical assistance of Peter M. Ingrassia, the helpful advice of Dr. John I. Scheide, and the secretarial assistance of Elaine M. Allard.

This study was supported by the Veterans Administration.

References

1. Feldman, G. M., and A. N. Charney. 1980. Effect of acute metabolic alkalosis and acidosis on intestinal electrolyte transport in vivo. *Am. J. Physiol.* 239:G427-G436.
2. Feldman, G. M., and A. N. Charney. 1982. Effect of acute respiratory alkalosis and acidosis on intestinal electrolyte transport in vivo. *Am. J. Physiol.* 242:G436-G492.
3. Charney, A. N., and L. P. Haskell. 1984. Relative effects of systemic pH, PCO₂, and HCO₃ concentration on colonic ion transport. *Am. J. Physiol.* 246:G159-G165.
4. Binder, H. J., and C. L. Rawlins. 1973. Electrolyte transport across isolated large intestinal mucosa. *Am. J. Physiol.* 225:1232-1239.
5. Phillips, S. F., and P. F. Schmalz. 1970. Bicarbonate secretion by the rat colon: effect of intraluminal chloride and acetazolamide. *Proc. Soc. Exp. Biol. Med.* 135:116-122.
6. Will, P. C., J. L. Lebowitz, and U. Hopfer. 1980. Induction of amiloride-sensitive sodium transport in the rat colon by mineralocorticoids. *Am. J. Physiol.* 238(Renal Fluid Electrolyte Physiol. 7):F261-F268.
7. Yau, W. M., and G. M. Makhlof. 1975. Comparison of transport mechanisms in isolated ascending and descending rat colon. *Am. J. Physiol.* 228:191-195.
8. Foster, E. S., T. W. Zimmerman, J. P. Hayslett, and H. J. Binder. 1983. Corticosteroid alteration of active electrolyte transport in rat distal colon. *Am. J. Physiol.* 245(Gastrointest. Liver Physiol. 8):G668-G675.
9. Jorkasky, D., M. Cox, and G. M. Feldman. 1985. Differential effects of corticosteroids on Na⁺ transport in rat distal colon in vitro. *Am. J. Physiol.* 248(Gastrointest. Liver Physiol. 11):G424-G431.
10. Kurtin, P., and A. N. Charney. 1984. Intestinal ion transport and intracellular pH during acute respiratory alkalosis and acidosis. *Am. J. Physiol.* 247(Gastrointest. Liver Physiol. 10):G24-G31.
11. Wagner, J. D., P. Kurtin, and A. N. Charney. 1985. Effect of systemic acid-base disorders on colonic intracellular pH and ion transport. *Am. J. Physiol.* 249(Gastrointest. Liver Physiol. 12):G39-G47.
12. Charney, A. N., J. D. Wagner, G. J. Birnbaum, and J. N. Johnstone. 1986. Functional role of carbonic anhydrase in intestinal electrolyte transport. *Am. J. Physiol.* 251(Gastrointest. Liver Physiol. 14):G682-G687.
13. Kurtin, P., and A. N. Charney. 1984. Effect of arterial carbon dioxide tension on amiloride-sensitive sodium absorption in the colon. *Am. J. Physiol.* 247(Gastrointest. Liver Physiol. 10):G537-G541.
14. Siggard-Andersen, O. 1974. The Acid-Base Status of the Blood. 4th ed. Williams & Wilkins, Baltimore, MD. 30-34.
15. Snedecor, G. W., and W. G. Cochran. 1967. Statistical Methods. 6th ed. Iowa State University Press, Ames, Iowa.
16. Frizzell, R. A., M. J. Koch, and S. G. Schultz. 1976. Ion transport by rabbit colon. I. Active and passive components. *J. Membr. Biol.* 27:297-316.
17. Charney, A. N., M. Arnold, and N. Johnstone. 1983. Acute respiratory alkalosis and acidosis and rabbit intestinal ion transport in vivo. *Am. J. Physiol.* 244(Gastrointest. Liver Physiol. 7):G145-G150.
18. Binder, H. J., E. S. Foster, M. E. Budinger, and J. P. Hayslett. 1987. Mechanism of electroneutral sodium chloride absorption in distal colon of the rat. *Gastroenterology.* 93:449-455.
19. Benos, D. J. 1982. Amiloride: a molecular probe of sodium transport in tissues and cells. *Am. J. Physiol.* 242(Cell Physiol. 11):C131-C145.
20. Foster, E. S., P. K. Dudeja, and T. A. Brasitus. 1986. Na⁺-H⁺ exchange in rat colonic brush-border membrane vesicles. *Am. J. Physiol.* 250(Gastrointest. Liver Physiol. 13):G781-G787.
21. Kinsella, J. L., and P. S. Aronson. 1981. Amiloride inhibition of the Na⁺-H⁺ exchanger in renal microvillus membrane vesicles. *Am. J. Physiol.* 241(Renal Fluid Electrolyte Physiol. 10):F374-F379.
22. Grinstein, S., and A. Rothstein. 1986. Mechanisms of regulation of the Na⁺/H⁺ exchanger. *J. Membr. Biol.* 90:1-12.
23. Turnheim, K., R. A. Frizzell, and S. G. Schultz. 1978. Interaction between cell sodium and the amiloride-sensitive sodium entry step in rabbit colon. *J. Membr. Biol.* 39:233-256.
24. Budinger, M. D., E. S. Foster, J. P. Hayslett, and H. J. Binder. 1986. Sodium and chloride transport in the large intestine of potassium-loaded rats. *Am. J. Physiol.* 251(Gastrointest. Liver Physiol. 14):G249-G252.
25. Halevy, J., M. E. Budinger, J. P. Hayslett, and H. J. Binder. 1986. Role of aldosterone in the regulation of sodium and chloride transport in the distal colon of sodium-depleted rats. *Gastroenterology.* 91:1227-1233.
26. Hatch, M., R. W. Freel, A. M. Goldner, and D. L. Earnest. 1984. Oxalate and chloride absorption by the rabbit colon: sensitivity to metabolic and anion transport inhibitors. *Gut.* 25:232-237.
27. Knickelbein, R., P. S. Aronson, C. M. Schron, J. Seifter, and J. W. Dobbins. 1985. Sodium and chloride transport across rabbit ileal brush border. II. Evidence for Cl-HCO₃ exchange and mechanism of coupling. *Am. J. Physiol.* 249(Gastrointest. Liver Physiol. 12):G236-G245.
28. Foster, E. S., M. E. Budinger, J. P. Hayslett, and H. J. Binder. 1986. Ion transport in proximal colon of the rat. *J. Clin. Invest.* 77:228-235.
29. Smith, P. L., M. A. Cascairo, and S. K. Sullivan. 1985. Sodium dependence of luminal alkalization by rabbit ileal mucosa. *Am. J. Physiol.* 249(Gastrointest. Liver Physiol. 12):G258-G268.
30. Langridge-Smith, J. E., and M. Field. 1981. Sulfate transport in rabbit ileum: characterization of the serosal border anion exchange process. *J. Membr. Biol.* 63:207-214.
31. Maren, T. H. 1977. Use of inhibitors in physiological studies of carbonic anhydrase. *Am. J. Physiol.* 232(Renal Fluid Electrolyte Physiol. 4):F291-F297.
32. Nellans, H. N., R. A. Frizzell, and S. G. Schultz. 1975. Effect of acetazolamide on sodium and chloride transport by in vitro rabbit ileum. *Am. J. Physiol.* 228: 1808-1814.
33. Jones, B., and D. P. Simpson. 1983. Influence of alterations in acid-base conditions on intracellular pH of intact renal cortex. *Renal Physiol.* 6:28-35.
34. Feldman, G. M., S. F. Berman, and R. L. Stephenson. 1988. Bicarbonate secretion in rat distal colon in vitro: a measurement technique. *Am. J. Physiol.* 254 (Cell Physiol. 23): C383-C390.