

Recombinative events of the T cell antigen receptor delta gene in peripheral T cell lymphomas.

P Kanavaros, ... , M P Lefranc, F Reyes

J Clin Invest. 1991;**87**(2):666-672. <https://doi.org/10.1172/JCI115044>.

Research Article

Recombinative events of the T cell antigen receptor (TCR) delta-chain gene were studied in 37 cases of peripheral T cell lymphoma (PTCL) and related to their clinical presentation and the expression of the alpha beta or gamma delta heterodimers as determined by immunostaining of frozen tissue samples. There were 22 cases of alpha beta, 5 cases of gamma delta, and 10 cases of silent TCR expressing neither the alpha beta nor gamma delta TCR. 5 different probes were used to examine the delta locus. The 22 cases of alpha beta PTCL displayed biallelic and monoallelic deletions; a monoallelic V delta 1 J delta 1 rearrangement was observed in 1 case and a monoallelic germ line configuration in 7 cases. The 5 cases of gamma delta PTCL displayed biallelic rearrangements: the productive rearrangements could be ascribed to V delta 1J delta 1 joining in 3 cases and VJ delta 1 joining in 2 cases according to the combined pattern of DNA hybridization with the appropriate probes and of cell reactivity with the TCR delta-1, delta TCS-1, and anti-V delta 2 monoclonal antibodies. In the VJ delta 1 joining, the rearranged V segments were located between V delta 1 and V delta 2. Interestingly, in the third group of 10 cases of silent PTCL, 5 cases were found to have a TCR [...]

Find the latest version:

<https://jci.me/115044/pdf>



Recombinative Events of the T cell Antigen Receptor δ Gene in Peripheral T cell Lymphomas

Panagiotis Kanavaros,* Jean-Pierre Farcet,** Philippe Gaulard,[§] Corinne Haioun,[‡] Marine Divine,[‡] Jean-Pierre Le Couedic,* Marie Paule Lefranc,^{||} and Félix Reyes**

*Institut Nationale de la Santé et de la Recherche Médicale U 91, [‡]Service d'Hématologie Clinique, [§]Laboratoire d'Anatomopathologie, Hôpital Henri Mondor, 94010 Créteil, France; ^{||}Laboratoire d'Immunogénétique Moléculaire, Unité de Recherche Associée, Centre Nationale de la Recherche Scientifique 1191, Université Montpellier II, Place E. Bataillon, 34095 Montpellier cedex 5 France

Abstract

Recombinative events of the T cell antigen receptor (TCR) δ -chain gene were studied in 37 cases of peripheral T cell lymphoma (PTCL) and related to their clinical presentation and the expression of the $\alpha\beta$ or $\gamma\delta$ heterodimers as determined by immunostaining of frozen tissue samples. There were 22 cases of $\alpha\beta$, 5 cases of $\gamma\delta$, and 10 cases of silent TCR expressing neither the $\alpha\beta$ nor $\gamma\delta$ TCR. 5 different probes were used to examine the δ locus. The 22 cases of $\alpha\beta$ PTCL displayed biallelic and monoallelic deletions; a monoallelic V δ 1 J δ 1 rearrangement was observed in 1 case and a monoallelic germ line configuration in 7 cases. The 5 cases of $\gamma\delta$ PTCL displayed biallelic rearrangements: the productive rearrangements could be ascribed to V δ 1J δ 1 joining in 3 cases and VJ δ 1 joining in 2 cases according to the combined pattern of DNA hybridization with the appropriate probes and of cell reactivity with the TCR δ -1, δ TCS-1, and anti-V δ 2 monoclonal antibodies. In the VJ δ 1 joining, the rearranged V segments were located between V δ 1 and V δ 2. Interestingly, in the third group of 10 cases of silent PTCL, 5 cases were found to have a TCR gene configuration identical to that in the TCR $\alpha\beta$ PTCL, as demonstrated by biallelic δ gene deletion. These 5 cases were CD3 positive. The 5 remaining cases showed a monoallelic δ gene rearrangement with a monoallelic germ line configuration in 4 and a monoallelic deletion in 1. Four of these cases were CD3 negative, which was consistent with an immature genotype the TCR commitent of which could not be ascertained.

Finally, TCR $\gamma\delta$ PTCL consisted of a distinct clinical morphological and molecular entity whereas TCR $\alpha\beta$ and silent PTCL had a similar presentation. (*J. Clin. Invest.* 1991. 87:666-672.) Key words: human T cell • differentiation • V genes

Introduction

The T cell antigen receptor (TCR)¹ is the antigen specific surface molecule characteristic of T cells. Two types of TCR, desig-

Address reprint requests to Dr. J. P. Farcet, INSERM U.91, Hôpital Henri Mondor, 94010 Créteil, France.

Received for publication 18 July 1990 and in revised form 27 September 1990.

1. Abbreviations used in this paper: D, G, R, deletion, germline, rearrangement configuration; PTCL, peripheral T cell lymphoma; TCR, T cell antigen receptor.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/91/02/0666/07 \$2.00

Volume 87, February 1991, 666-672

nated $\alpha\beta$ and $\gamma\delta$, have been described (1-10). Both are heterodimeric molecules expressed in association with the CD3 protein complex (1, 10). The majority of T cells bear TCR $\alpha\beta$ which is responsible for MHC-restricted antigen recognition (1). These T cells include two mutually exclusive subsets expressing either a CD4⁺ CD8⁻ or a CD4⁻ CD8⁺ phenotype. A minority of T cells bear TCR $\gamma\delta$ whose recognition specificity is still under investigation (11). Most $\gamma\delta$ T cells express the CD4⁻ CD8⁻ phenotype (12). The TCR δ locus, which has been recently characterized (13-22), is located within the TCR α locus on chromosome 14 and is flanked by V α segments on its 5' side and by J α segments on its 3' side. To date, one C δ , three J δ , three D δ , and six V δ segments have been reported (13-27). As a model for lymphoid differentiation, leukaemias and lymphomas in humans have provided much information for the elucidation of the developmental hierarchy among TCR genes. The TCR δ gene undergoes rearrangements and/or deletions (28-40) which might constitute the earliest recombinative event in T cell differentiation (30, 32).

In contrast to numerous studies of the TCR δ gene configuration in T cell lymphoblastic malignancies (28-36, 38, 40), few cases of nonlymphoblastic T cell lymphomas have been investigated (37, 40, 41). In this study, using five different probes, we have examined the organization of the TCR δ gene in a large series of peripheral T cell lymphomas (PTCL). The phenotype of these cases, which has been previously reported (42), allowed us to categorize them into three groups expressing either TCR $\alpha\beta$, TCR $\gamma\delta$, or neither. A correlation between the TCR δ gene status and each TCR group has been established.

Methods

Tissue samples. Frozen tissue samples from 37 cases diagnosed as peripheral T cell lymphomas on the basis of both histologic and immunologic features (excluding lymphoblastic lymphomas and mycosis fungoides) were selected for the genotypic study. In the 37 cases, the histologic and phenotypic data as well as the pattern of immunoglobulin and TCR- β gene rearrangements have been previously reported (42). Based on the reactivity of lymphoma cells with anti-CD3, β F₁, and TCR δ -1 monoclonal antibodies, the patients were categorized into three groups: CD3⁺, β F₁⁺, TCR δ -1⁻ (22 cases); CD3⁺, β F₁⁺, TCR δ -1⁺ (5 cases); and CD3[±], β F₁⁻, TCR δ -1⁻ (10 cases), referred to as TCR $\alpha\beta$, TCR $\gamma\delta$, and TCR silent PTCL, respectively.

DNA analysis. DNA extraction was performed as previously reported (43). 10 μ g of high molecular weight DNA was digested with each appropriate restriction enzyme, fractionated in 0.8% agarose gel and blotted onto a nitrocellulose filter. Filters were P³² hybridized with appropriate P³²-labeled probes. After hybridization the filters were washed for 30 min with 2 \times SSC at room temperature, twice for 30 min with 0.5 \times SSC/0.5% SDS at 56°C and finally for 5 min with 0.5 \times SSC at room temperature. Autoradiography was carried out for 2-5 d at -80°C.

DNA probes. Immunoglobulin and TCR- β genes were studied with JH and C β probes, as previously reported (43).

The TCR- δ gene restriction map is shown in Fig. 1. TCR- δ gene rearrangements were analyzed after Bam HI and Hind III digestion using the following probes: (a) J δ S16, specific for the J δ_1 region (15); (b) R21XH, specific for the J δ_2 region (16); (c) TCR V δ_1 , a 240-bp Eco RI-Sac I fragment isolated from the 0-240/38 cDNA clone (13) and specific for the V δ_1 region; (d) pDV2 SPO.S, specific for the V δ_2 region (44); and (e) R1IEE, specific for the C δ region (15).

In the TCR $\gamma\delta$ and TCR silent PTCL, TCR- γ gene rearrangements were analyzed by using the J γ probe pH 60 (45) in Bam HI, Hind III, and Eco RI digests.

Rearrangements and deletions of the TCR δ gene were defined in this study according to previously reported criteria (30). When the germ line band had the same intensity as the control with no additional band, both alleles were considered in germ line configuration (GG); an additional band showed a rearranged allele (RG). When the germ line band was not visible, the δ segment was deleted either on both alleles (DD) or on one allele (RD), depending upon the presence or absence of an additional band. When the germ line band showed a decreased intensity compared with the control, the δ segment was either in a DG or RG configuration depending upon the absence or presence of an additional band. Finally, both alleles could have rearranged (RR).

Results

TCR $\alpha\beta$ PTCL. 22 cases expressed the TCR $\alpha\beta$ and most were ascribed to the diffuse mixed or diffuse large cell histologic subtype (Table I). Patient median age was 48 yr and male/female ratio was 10:1. 20 patients had stage IV disease according to the Ann Arbor classification. Lymph nodes were involved in 17 cases, bone marrow in 11 cases, skin in 8 cases, and liver in 7 cases. The spleen was enlarged in 7 cases. Treatment and survival are shown in Table I.

In 14 cases, biallelic deletions of all TCR δ sequences were detected using the J δ_1 and C δ probes (DD pattern) (Fig. 2). With these probes, monoallelic TCR δ deletions were observed in seven additional cases (DG pattern) (Fig. 2). The deletions involved the entire TCR δ gene in all cases except 4, where the V δ_1 remained in the germ line configuration. Hybridization with the V δ_2 probe was not done in this group. In case 7, a monoallelic J δ_1 rearrangement (for the sake of simplicity, the δ gene VDJ rearrangements will be referred to as VJ rearrangements) with deletion of the germ line band was found (RD

pattern). The joining event involved the V δ_1 segment as demonstrated by the presence of a 12-kb band in Hind III digests using both the J δ_1 and V δ_1 probes (13).

The TCR γ gene was not studied in this group and the immunoglobulin JH and TCR β gene configurations are shown in Table I.

TCR $\gamma\delta$ PTCL. 5 cases expressed the TCR $\gamma\delta$. All had an unusual clinical presentation. Case 27 presented as a lethal midline granuloma with histologic findings consistent with a polymorphic reticulosis evolving into an overt lymphoma as previously reported (46). The 4 other cases were hepatosplenic T cell lymphoma, a new entity among PTCL with clinical histological and phenotypic features which we recently reported in 2 of them (47). The patients (3 males and 1 female) had a median age of 35 yr and presented with hepatosplenomegaly. The liver, bone marrow, and spleen had a sinusoidal/sinusoidal infiltration by monomorphic medium sized cells. The lymph nodes were not involved. Treatment and survival are shown in Table I.

In all 5 cases, biallelic rearrangements (RR pattern) were detected with the J δ_1 probe in Bam HI and Hind III digests (Fig. 3). In cases 23, 24, and 26, a V δ_1 J δ_1 rearrangement was observed using the V δ_1 and J δ_1 probes (Fig. 3). The other allele of these cases as well as the two alleles of cases 25 and 27, displayed a V δ_1 segment in germ line configuration whereas the V δ_2 segment was deleted as shown by hybridization with the V δ_2 probe. Therefore, the J δ_1 rearrangement did not involve either the V δ_1 or the V δ_2 segment; neither was the V δ_3 segment involved in these J δ_1 rearrangements since the C δ probe (which detects V δ_3 rearrangement [48]) revealed only germ line bands in Bam HI digests.

The TCR γ gene had rearranged in the five cases (Fig. 3), while the immunoglobulin JH gene was in germ line configuration and the TCR β gene had rearranged in 2 out of 5 cases (Table II).

TCR silent PTCL. 10 cases expressed neither the $\alpha\beta$ nor $\gamma\delta$ TCR and most were ascribed to the diffuse large cell histologic subtype (Table I). Patient median age was 50 yr and male/female ratio was 2.3:1. All patients had stage IV disease with involvement of lymph nodes in 9 cases, bone marrow in 5 cases, skin in 8 cases, and liver in 4 cases. The spleen was enlarged in 4 cases. Treatment and survival are shown in Table I.




Figure 1. Genomic organization of the human TCR- δ gene. The locations of the relevant Hind III (H) and Bam HI (B) restriction enzyme sites are indicated. The solid bars below the δ gene represent the five probes which were used for hybridization of the Southern blot filters.

Table I. Histology, Phenotype, and Prognosis in 37 Cases of PTCL

Patient No.	Histology*	TCR [‡]	Treatment [§]	Survival	Status
1	DM	$\alpha\beta$	CHOP	8	dead
2	IBL	$\alpha\beta$	ACVBP	2	dead
3	DM	$\alpha\beta$	CVP	2	dead
4	DM	$\alpha\beta$	CHOP	15	dead
5	DM	$\alpha\beta$	ACVBP	35	alive
6	Uncl	$\alpha\beta$	ACVBP	14	dead
7	DM	$\alpha\beta$	CHOP	48	alive
8	ALC	$\alpha\beta$	ACVBP	4	dead
9	DM	$\alpha\beta$	ACVBP	27	dead
10	SL	$\alpha\beta$	CHOP	51	alive
11	DM	$\alpha\beta$	CVP	48	alive
12	DM	$\alpha\beta$	CHOP	49	alive
13	DLC	$\alpha\beta$	CVP	53	alive
14	DLC	$\alpha\beta$	ACVBP	18	dead
15	ALC	$\alpha\beta$	mBACOD	36	alive
16	DM	$\alpha\beta$	CHOP	2	dead
17	DLC	$\alpha\beta$	CHOP	24	alive
18	DLC	$\alpha\beta$	CHOP	follow up not available	not available
19	DLC	$\alpha\beta$	CHOP	2	dead
20	IBL	$\alpha\beta$	mBACOD	5	dead
21	DLC	$\alpha\beta$	CVP	follow up not available	not available
22	DLC	$\alpha\beta$	CHOP	36	alive
23	Uncl	$\gamma\delta$	ACVBP	25	dead
24	Uncl	$\gamma\delta$	ACVBP	44	dead
25	Uncl	$\gamma\delta$	CHOP	36	dead
26	Uncl	$\gamma\delta$	ACVBP	14	alive
27	DM	$\gamma\delta$	CHOP	2	dead
28	DLC	silent	CHOP	6	dead
29	DLC	silent	ACVBP	9	dead
30	DM	silent	CHOP	15	dead
31	DLC	silent	CHOP	6	dead
32	SL	silent	CVP	30	dead
33	DM	silent	ACVBP	36	alive
34	DLC	silent	ACVBP	7	dead
35	DLC	silent	CVP	2	dead
36	ALC	silent	CHOP	1	dead
37	ALC	silent	CHOP	22	dead

* Diffuse mixed (DM); immunoblastic (IBL); unclassifiable (Uncl); anaplastic large cell (ALC); small lymphocytic (SL); and diffuse large cell (DLC).

[‡] TCR expression as determined on tissue section by β F1 and TCR δ -1 monoclonal antibody reactivity using the alkaline antiphosphatase alkaline procedure.

[§] CHOP: cyclophosphamide, adriamycin, oncovin, prednisone, ACVBP: adriamycin, cyclophosphamide, velbe, bleomycine, prednisone. CVP: cyclophosphamide, VM26, prednisone. mBACOD: methotrexate, bleomycin, cyclophosphamide, oncovin, dexamethasone.

^{||} Months.

In 5 cases, TCR δ biallelic deletion was demonstrated by using the V δ 1, J δ 1, J δ 2, and C δ probes (DD pattern) (Fig. 4).

In the 5 other cases, a monoallelic rearrangement (RD or RG pattern) involving either the J δ 1 (cases 29, 35, 37) or the

J δ 2 segment (cases 34, 36) was detected with the J δ 1 and J δ 2 probes (Fig. 4). When the J δ 1 segment was rearranged, the J δ 2 segment was in germ line configuration (Fig. 4). When the J δ 2 segment was rearranged, the J δ 1 segment was deleted. Using the V δ 1 probe in Hind III digests, the V δ 1 segment was shown to be rearranged in a V δ 1J δ 1 joining in 3 cases with a 12-kb band and in a V δ 1 J δ 2 joining in 2 cases with a 7.3-kb band (Fig. 4). In all these 5 cases, the C δ segment was in germ line configuration as shown by the C δ probe.

All 10 cases had rearranged the TCR γ locus as detected by the J γ 1 probe, while the JH gene was in germline configuration in all cases and the C β gene had rearranged in 8 out of 10 cases (Table III).

Discussion

We have analyzed the recombinative events involving the TCR δ locus in 37 cases of PTCL. These were categorized into 3 phenotypic groups according to TCR expression. 22 cases expressed TCR $\alpha\beta$, 5 cases $\gamma\delta$, and 10 cases neither $\alpha\beta$ nor $\gamma\delta$. Using 5 probes specific for the J δ 1, J δ 2, V δ 1, V δ 2, and C δ segments, Southern blot analyses demonstrated deletions and/or rearrangements in all 37 cases of PTCL.

In all cases of TCR $\alpha\beta$ PTCL either biallelic (DD) or monoallelic (DG, RD) deletions were observed. 1 case exhibited a monoallelic V δ 1 J δ 1 rearrangement in addition to the deleted allele. As expected, the TCR $\alpha\beta$ expression implied the deletion of the TCR δ gene locus associated with the TCR α gene rearrangement. The TCR δ locus, nested between the V α and J α regions, is deleted in V α J α joining (23), occasionally sparing the V δ 1 segment which thus remains in germ line configuration (48) as found here in 4 cases. The TCR δ gene configuration in TCR $\alpha\beta$ PTCL was in keeping with the findings of others in T ALL (33, 35) and T cell clones (48, 49) expressing the TCR $\alpha\beta$. During differentiation, whether the TCR δ and α genes rearrange successively in a single pathway or in two different pathways remains controversial (50–52). In our series of β PTCL, most alleles, i.e., 36/44, were deleted, which precludes any conclusion regarding the status of the δ gene before α gene rearrangement. Thus, the deletion could involve a δ gene either in germ line configuration or when rearranged.

The 5 cases of TCR $\gamma\delta$ PTCL had a biallelic TCR δ rearrangement (RR). All 5 cases had a peculiar clinical presentation clearly distinct from TCR $\alpha\beta$ and silent PTCL. 1 case was a midline granuloma evolving into a PTCL (46) and 4 cases were hepatosplenic PTCL, 2 of which we recently reported as a new entity (47). The tropism of malignant $\gamma\delta$ T cells for the sinuses of spleen, bone marrow, and the sinusoids of the liver is remarkable with regard to the preferential localization in the splenic red pulp of their normal counterpart (42). In this report we have extended the study of the genotypic configuration of TCR genes to 5 cases of TCR $\gamma\delta$ PTCL and the V and J segments involved in the TCR δ rearrangement have been characterized. 3 of the 5 cases displayed a productive V δ 1 J δ 1 rearrangement which has been reported as the most frequent pattern in T ALL and T lymphoblastic lymphomas expressing the TCR $\gamma\delta$ (33–35). The δ TCS1⁺/anti-V δ 2⁻ phenotype of the lymphoma cells (47) was in agreement with the genotypic findings and has been reported as that of the major subset of $\gamma\delta$ T cells in normal thymus and spleen (54). The TCR δ productive rearrangements in the other 2 cases also involved the J δ 1 segment. However,





Figure 2. Representative TCR δ gene deletions in TCR $\alpha\beta$ PTCL. Patient number is given below each lane. C, control DNA from granulocytes. Germ line bands are indicated by thin lines near the lane C and their sizes are given in kb. Rearranged bands are shown by arrows. On the top figure, two DG pattern and one DD pattern are shown after J δ 1 hybridization. The same filter was rehybridized with the C β probe in order to demonstrate that an equal amount of DNA was loaded in each lane. The bottom figure illustrates case No. 14 with one allele having deleted J δ 1 and C δ segments while the V δ 1 segment has remained in germ line configuration.

the V segments that joined J δ 1 were neither V δ 1, V δ 2, nor V δ 3 as demonstrated with the appropriate δ probes. In addition, the δ chain encoded by these VJ δ 1 combinations was recognized by the TCR- δ 1 monoclonal antibody, which reacts with all δ chains, and not by the δ TCS-1 and anti-V δ 2 antibodies which react with an epitope encoded by the V δ 1J δ 1 pairing and the V δ 2 segment, respectively (see reference 47). In these 5 cases of $\gamma\delta$ PTCL, the 5 rearranged and nonproductive alleles also involved the V segments located between V δ 1 and V δ 2. Indeed, the V δ 1 segment was in germ line configuration and the V δ 2 segment was deleted, showing that J δ 1 rearranged with V segments located 3' to V δ 1 and 5' to V δ 2. These V segments might belong to the V α pool as reported previously in a $\gamma\delta$ T cell clone (55). Alternatively, other V δ segments embedded in the V α family might rearrange with J δ 1 as suggested in δ TCS1⁻ clones derived from the thymus (56). Taken together, the TCR $\gamma\delta$ PTCL had a genotype and phenotype in marked contrast to those of T cell $\gamma\delta$ clones derived from the peripheral blood of

normal individuals which use preferentially a V δ 2 J δ 1 joining (54, 57-59).

The third group of 10 patients consisted of TCR silent PTCL (Table IV). Their clinical and morphological characteristics were similar to those of TCR $\alpha\beta$ PTCL. 5 of these displayed a TCR genotype identical to that in the group of TCR $\alpha\beta$ PTCL, namely a biallelic TCR δ deletion with a TCR β rearrangement. In addition, the deletion of V δ 1 embedded in the V α pool indicated a V α -J α rearrangement. Therefore, the lymphoma cells seemed to be committed to TCR $\alpha\beta$ expression. In addition, they were CD3⁺ on tissue sections. The β F1⁻ status could be explained in two ways: either the level of β chain expression was below the detection threshold of the immunostaining or nonproductive rearrangements or a translational dysregulation prevented the β chain synthesis. Northern blot analyses were not performed because of insufficient material. The possibility that some of these 5 cases arose from cells initially expressing TCR $\alpha\beta$ cannot be excluded, since a pheno-




Figure 3. Representative TCR δ and γ gene rearrangements in TCR $\gamma\delta$ PTCL. Patient number, control DNA, germ line bands, and rearrangements are indicated as in Fig. 1. Note the additional 12-kb band in Hind III digests, which shows a V δ 1 J δ 1 rearrangement, as demonstrated here by the V δ 1 probe in case 26 and by the J δ 1 probe in case 24.

Table II. Genotype of the TCR $\alpha\beta$ PTCL

Patient No.	JH	C β	V δ_1	J δ_1	C δ
1	G	R	DD	DD	DD
2	G	R	DD	DD	DD
3	G	R	DG	DG	DG
4	G	R	DD	DD	DD
5	G	G	DG	DG	DG
6	G	R	DD	DD	DD
7	G	R	RD	RD	DG
8	G	R	DD	DD	DD
9	G	R	GG	DG	DG
10	G	R	DG	DG	DG
11	G	R	DD	DD	DD
12	G	R	DD	DD	DD
13	G	R	DD	DD	DD
14	R	R	GG	DG	DG
15	G	G	GG	DG	DG
16	G	R	GG	DG	DG
17	G	R	DD	DD	DD
18	G	R	DD	DD	DD
19	G	R	DD	DD	DD
20	G	R	DD	DD	DD
21	G	R	DD	DD	DD
22	G	R	DD	DD	DD

typic drift including the loss of TCR expression can occur along tumor progression while the TCR genotype remains unchanged (47). 5 additional cases of TCR silent PTCL had a monoallelic δ gene rearrangement. 3 of them consisted of a V δ_1 J δ_1 joining and the other 2 of a V δ_1 J δ_2 joining. In these 5 cases unrearranged alleles were either germ line in 4 cases (RG), or deleted in 1 case (RD). 4 cases (1 RD + 3 RG) were CD3⁻ suggesting that the lymphoma cells were immature with respect to TCR differentiation. Their genotype was in contrast to the unequivocal DD and RR genotype of the PTCL expressing the CD3 antigen. Alternatively, the genotype of TCR silent PTCL could be that of a T cell which normally would have died but was rescued by the leukaemic events.

In conclusion, all cases of PTCL displayed a recombinative event in the TCR δ locus. When TCR $\alpha\beta$ was expressed, the δ




Figure 4. Representative TCR δ gene rearrangement in TCR silent PTCL. Patient number, control DNA, germ line bands, and rearrangements are indicated as in Fig. 1. The patient 34 illustrated here had a V δ_1 J δ_2 rearrangement demonstrated by an additional 7.3-kb band in Hind III digests which were hybridized with the V δ_1 and J δ_2 probes.

Table III. Genotype of the TCR $\gamma\delta$ PTCL

Patient No.	JH	C β	J γ_1	V δ_1	V δ_2	J δ_1	J δ_2	C δ
23	G	R	R	RG	DD	RR	GG	GG
						B = >23/10, 5*		
						H = 12/5		
24	G	G	R	RG	DD	RR	GG	GG
						B = >23/12		
						H = 12/10		
25	G	G	R	GG	DD	RR	GG	GG
						B = 15/12, 5		
						H = 7, 6/G		
26	G	G	R	RG	DD	RR	GG	GG
						B = >23/12, 5		
						H = 12/5		
27	G	R	R	GG	DD	RR	GG	GG
						B = 15/12, 5		
						H = 7, 6/G		

B, Bam HI; H, Hind III.

* Numbers correspond to the band size in kb.

gene was deleted on most alleles; when TCR $\gamma\delta$ was expressed, the δ gene was rearranged on all alleles. In these TCR $\gamma\delta$ PTCL, the rearrangements consisted of the J δ_1 segment joined to the V δ_1 segment or yet unsequenced V segments located between V δ_1 and V δ_2 . Interestingly, a distinct subgroup of PTCL expressing neither TCR $\alpha\beta$ nor $\gamma\delta$ encompassed cases with a δ gene configuration identical to that in TCR $\alpha\beta$ PTCL and cases with an "immature" genotype which differed from that in both

Table IV. Genotype of the TCR silent PTCL

Patient No.	JH	C β	J γ_1	V δ_1	J δ_1	J δ_2	C δ
28	G	R	R	DD	DD	DD	DD
29	G	G	R	RG	RG	GG	GG
					B = >23		
					H = 12		
30	G	R	R	DD	DD	DD	DD
31	G	R	R	DD	DD	DD	DD
32	G	R	R	DD	DD	DD	DD
33	G	R	R	DD	DD	DD	DD
34	G	G	R	RG	DG	RG	GG
						B = >23	
						H = 7, 3	
35	G	R	R	RG	RG	GG	GG
					B = >23		
					H = 12		
36	G	R	R	RG	DD	RD	GD
						B = >23	
						H = 7, 3	
37	G	R	R	RG	RG	GG	GG
					B = >23		
					H = 12		

* Patients 28-33: CD3⁺; patients 34-37: CD3⁻.

the TCR $\alpha\beta$ and TCR $\gamma\delta$ PTCL. Most importantly, TCR $\gamma\delta$ PTCL consisted of a clearly defined clinical, morphological, and molecular entity whereas TCR $\alpha\beta$ and silent PTCL had a similar presentation.

Acknowledgments

The authors thank J. M. Massé for illustrations and A. M. Dulac, R. Quintel, and V. Corsetti for secretarial work. They thank T. H. Rabbitts and M. S. Krangel for providing probes.

Supported by Institut Nationale de la Santé et de la Recherche Médicale, Université Paris Val de Marne, and Association de Recherche contre le Cancer.

References

- Kronenberg, M., G. Siu, L. Hood, and N. Shastri. 1986. The molecular genetics of the T-cell antigen receptor and T-cell antigen recognition. *Annu. Rev. Immunol.* 4:529-591.
- Marrack, P., and J. Kappler. 1986. The antigen specific major histocompatibility complex restricted receptor on T-cells. *Adv. Immunol.* 38:1-25.
- Brenner, M. B., J. McLean, D. Dialynas, J. Strominger, J. Smith, F. Owen, J. Seidman, S. Rosen, F. Ip, and M. Krangel. 1986. Identification of a putative second T-cell receptor. *Nature (Lond.)*. 322:145-149.
- Bank, L., R. DePinho, M. Brenner, J. Cassimeris, F. Alt, and L. Chess. 1986. A functional T3 molecule associated with a novel heterodimer on the surface of immature human thymocytes. *Nature (Lond.)*. 322:179-181.
- Weiss, A., M. Newton, and D. Crommie. 1986. Expression of T3 in association with a molecule distinct from the T-cell antigen receptor. *Proc. Natl. Acad. Sci. USA*. 83:6998-7002.
- Borst, J., R. Van de Griend, J. Van Oostveen, S. Ang, C. Melief, J. Seidman, and R. Bolhuis. 1987. A T-cell receptor γ /CD3 complex found on cloned functional lymphocytes. *Nature (Lond.)*. 325:683-688.
- Brenner, M., J. McLean, H. Scheft, J. Riberty, S. Ang, J. Seidman, P. Devlin, and M. Krangel. 1987. Two forms of the T-cell receptor γ protein found on peripheral blood cytotoxic T lymphocytes. *Nature (Lond.)*. 325:689-693.
- Lanier, L. L., N. A. Federspiel, J. J. Ruitenberg, J. H. Phillips, J. P. Allison, D. Littman, and A. Weiss. 1987. The T cell antigen receptor complex expressed on normal peripheral blood CD4⁺CD8⁻ T lymphocytes. A CD3 associated disulphide linked γ chain heterodimer. *J. Exp. Med.* 165:1076-1094.
- Moingeon, P., S. Jitsukawa, F. Faure, F. Troalen, F. Triebel, M. Graziani, F. Forestier, D. Bellet, C. Bohuon, and T. Hercend. 1988. A γ -chain complex forms a functional receptor on cloned human lymphocytes with natural killer-like activity. *Nature (Lond.)*. 325:723-728.
- Brenner, M. B., J. L. Strominger, and M. S. Krangel. 1988. The $\gamma\delta$ T cell receptor. *Adv. Immunol.* 43:133-149.
- Born, W., M. P. Happ, A. Dallas, C. Reardon, R. Kubo, T. Shinnick, P. Brennan, and R. O'Brien. 1990. Recognition of heat shock proteins and $\gamma\delta$ cell function. *Immun. Today* 11:40-48.
- Groh, V., S. Porcelli, M. Fabbri, L. L. Lanier, L. J. Picker, T. Anderson, R. A. Warnke, A. K. Bhana, J. L. Strominger, and M. B. Brenner. 1989. Human lymphocytes bearing T cell receptor $\gamma\delta$ are phenotypically diverse and evenly distributed throughout the lymphoid system. *J. Exp. Med.* 169:1277-1294.
- Hata, S., M. B. Brenner, and M. S. Krangel. 1987. Identification of putative human T cell receptor δ complementary DNA clones. *Science (Wash. DC)*. 238:678-682.
- Loh, E. Y., L. L. Lanier, C. W. Turck, D. R. Littman, M. M. Davis, Y.-H. Chien, and A. Weiss. 1987. Identification and sequence of a fourth human T cell antigen receptor chain. *Nature (Lond.)*. 330:569-572.
- Boehm, T., R. Baer, I. Lavenir, A. Forster, J. J. Waters, E. Nacheva, and T. H. Rabbitts. 1988. The mechanism of chromosomal translocation t(11;14) involving the T-cell receptor C δ locus human chromosome 14q11 and a transcribed region of chromosome 11p15. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:385-391.
- Baer, R., T. Boehm, H. Yssel, H. Spits, and T. H. Rabbitts. 1988. Complex rearrangements within the human J δ -C δ /J α -C α locus and aberrant recombination between J α segments. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:1661-1668.
- Takahara, Y., E. Champagne, H. Griesser, N. Kimura, D. Tkachuk, J. Reimann, A. Okada, F. W. Alt, L. Chess, M. Minden, and T. W. Mak. 1988. Sequence and organization of the human T cell δ chain gene. *Eur. J. Immunol.* 18:283-287.
- Griesser, H., E. Champagne, D. Tkachuk, Y. Takihara, M. Lalande, E. Baillie, M. Minden, and T. W. Mak. 1988. The human T cell receptor α - δ locus: a physical map of the variable, joining and constant region genes. *Eur. J. Immunol.* 18:641-653.
- Isobe, M., G. Russo, F. G. Haluska, and C. M. Croce. 1988. Cloning of the gene encoding the δ subunit of the human T-cell receptor reveals its physical organization within the α -subunit locus and its involvement in chromosome translocations in T-cell malignancy. *Proc. Natl. Acad. Sci. USA*. 85:3933-3945.
- Takahara, Y., D. Tkachuk, E. Michalopoulos, E. Champagne, J. Reimann, M. Minden, and T. W. Mak. 1988. Sequence and organization of the diversity, joining, and constant region genes of the human T-cell δ -chain locus. *Proc. Natl. Acad. Sci. USA*. 85:6097-6110.
- Satyanarayana, K., S. Hata, P. Devlin, M. G. Roncarolo, J. E. DeVries, H. Spits, J. L. Strominger, and M. S. Krangel. 1988. Genomic organization of the human T-cell antigen-receptor α/δ locus. *Proc. Natl. Acad. Sci. USA*. 85:8166-8170.
- Loh, E. Y., S. Cwirra, A. T. Serafini, J. H. Phillips, and L. L. Lanier. 1988. The human T-cell receptor δ chain: genomic organization, diversity, and expression in populations of cells. *Proc. Natl. Acad. Sci. USA*. 85:9714-9728.
- Chien, Y., M. Iwashima, K. B. Kaplan, J. F. Elliott, and M. M. Davis. 1987. A new T-cell receptor gene located within the α locus and expressed early in T-cell differentiation. *Nature (Lond.)*. 327:677-682.
- Chien, Y., M. Iwashima, D. A. Wettstein, K. B. Kaplan, J. F. Elliott, W. Born, and M. M. Davis. 1987. T-cell receptor δ gene rearrangements in early thymocytes. *Nature (Lond.)*. 330:722-727.
- Takahara, Y., E. Champagne, E. Ciccone, L. Moretta, M. Minden, and T. W. Mak. 1989. Organization and orientation of a human T cell receptor δ chain V gene segment that suggests an inversion mechanism is utilized in its rearrangement. *Eur. J. Immunol.* 19:571-574.
- Hata, S., P. Clabby, P. Devlin, H. Spits, J. E. DeVries, and M. S. Krangel. 1989. Diversity and organization of human T cell δ variable gene segments. *J. Exp. Med.* 169:41-57.
- Takahara, Y., J. Reinmann, E. Michalopoulos, E. Ciccone, L. Moretta, and T. W. Mak. 1989. Diversity and structure of human T cell receptor δ chain genes in peripheral blood γ/δ -bearing T lymphocytes. *J. Exp. Med.* 169:393-405.
- Van Dongen, J. M., I. L. M. Wolvers-Tettero, F. Wassenaar, J. Borst, and P. Van den Elsen. 1989. Rearrangement and expression of T-cell receptor δ genes in T-cell acute lymphoblastic leukemias. *Blood*. 74:334-342.
- Dyer, M. J. S. 1989. T-cell receptor δ/α rearrangements in lymphoid neoplasms. *Blood*. 74:1073-1083.
- Foroni, L., M. Laffan, T. Boehm, T. H. Rabbitts, D. Catovsky, and L. Luzzatto. 1989. Rearrangement of the T-cell receptor δ genes in human T-cell leukemias. *Blood*. 73:559-565.
- Biondi, A., E. Champagne, V. Rossi, G. Giudici, A. Cantù-Rajno, G. Masera, A. Mantovan, T. W. Mak, and M. D. Minden. 1989. T-cell receptor δ gene rearrangement in childhood T-cell acute lymphoblastic leukemia. *Blood*. 73:2133-2138.
- Yumura-Yagi, K., J. Hara, N. Terada, S. Ishihara, A. Tawa, Y. Takihara, E. Champagne, M. D. Minden, T. W. Mak, and K. Kawa-Ha. 1989. Analysis of molecular events in leukemic cells arrested at an early stage of T-cell differentiation. *Blood*. 74:2103-2111.
- Loiseau, P., P. Guglielmi, D. Le Paslier, E. Macintyre, A. Gessain, J.-C. Bories, G. Flandrin, Z. Chen, and F. Sigaux. 1989. Rearrangements of the T cell receptor δ gene in T acute lymphoblastic leukemia cells are distinct from those occurring in B lineage acute lymphoblastic leukemia and preferentially involve one V δ gene segment. *J. Immunol.* 142:3305-3311.
- De Villartay, J.-P., A. B. Pullman, R. Andrade, E. Tschachler, O. Colamenici, L. Neckers, D. I. Cohen, and J. Cossman. 1989. γ/δ lineage relationship within a consecutive series of human precursor T-cell neoplasms. *Blood*. 74:2508-2518.
- Griesinger, F., J. M. Greenberg, and J. H. Kersey. 1989. T cell receptor γ and δ rearrangements in hematologic malignancies. Relationship to lymphoid differentiation. *J. Clin. Invest.* 84:506-516.
- Hara, J., S. H. Benedict, E. Champagne, Y. Takihara, T. W. Mak, M. Minden, and E. W. Gelfand. 1988. T cell receptor δ gene rearrangements in acute lymphoblastic leukemia. *J. Clin. Invest.* 82:1974-1982.
- Asou, N., T. Hattori, M. Matsuoka, F. Kawano, and K. Takatsuki. 1989. Rearrangements of T-cell antigen receptor δ chain gene in hematologic neoplasms. *Blood*. 74:2707-2712.
- Raghavachar, A., E. Thiel, T. E. Hansen-Hagge, B. Kranz, and C.-R. Bartram. 1989. Rearrangement of T cell receptor β , γ and δ gene loci in human pre-T cell acute lymphoblastic leukemia. *Leukemia*. 3:413-418.
- Hara, J., K. Yumura-Yagi, A. Tawa, S. Ishihara, M. Murata, N. Terada, Y. Isumi, E. Champagne, Y. Takihara, T. W. Mak, M. Minden, and K. Kawa-Ha. 1989. Molecular analysis of acute undifferentiated leukemia: two distinct subgroups at the DNA and RNA levels. *Blood*. 74:1738-1746.
- Kimura, N., Y. Takihara, T. Akiyoshi, T. Yoshida, K. Oshima, T. Kawara, S. Hisano, M. Kozuru, M. Okumura, and M. Kikuchi. 1989. Rearrangement of T-cell receptor δ chain gene as a marker of lineage and clonality in T-cell lymphoproliferative disorders. *Cancer Res.* 49:4488-4492.
- Tkachuk, D. C., H. Griesser, Y. Takihara, E. Champagne, M. Minden,

- A. C. Feller, K. Lennert, and T. W. Mak. 1988. Rearrangement of T-cell δ locus in lymphoproliferative disorders. *Blood*. 72:353-357.
42. Gaulard, P., P. Bourquelot, P. Kanavaros, C. Haioun, J.-P. LeCouedic, M. Divine, M. Goossens, E.-S. Zafrani, J.-P. Farcet, and F. Reyes. Expression of the α/β and γ/δ T-cell receptors in 57 cases of peripheral T-cell lymphomas. *Am. J. Pathol.* 137:617-623.
43. Henni, T., P. Gaulard, M. Divine, J.-P. Le Couedic, D. Rocha, C. Haioun, Z. Henni, J.-P. Marolleau, Y. Pinaudeau, M. Goossens, J.-P. Farcet, and F. Reyes. 1988. Comparison of genetic probe with immunophenotype analysis in lymphoproliferative disorders: a study of 87 cases. *Blood*. 72:1937-1943.
44. Dariavach, P., and M.-P. Lefranc. 1989. First genomic sequence of the human T-cell receptor V δ 2 gene (TRDV2). *Nucleic Acids Res.* 17:4880-4889.
45. Lefranc, M.-P., and T. H. Rabbitts. 1985. Two tandemly organized human genes encoding the T-cell γ constant-region sequences show multiple rearrangement in different T-cell types. *Nature (Lond.)*. 316:464-466.
46. Gaulard, P., T. Henni, J. P. Marolleau, C. Haioun, M. C. Voisin, M. Divine, M. Goossens, J. P. Farcet, and F. Reyes. 1988. Lethal midline granuloma and lymphomatoid granulomatosis. Evidence for a monoclonal T-cell lymphoproliferative disorder. *Cancer (Phila.)*. 62:705-710.
47. Farcet, J. P., P. Gaulard, J. P. Marolleau, J. P. Le Couedic, T. Henni, M. F. Gourdin, M. Divine, C. Haioun, S. Zafrani, M. Goossens, T. Hercend, and F. Reyes. 1990. Hepatosplenic T cell lymphoma: sinusal/sinusoidal localization of malignant cells expressing the T cell receptor $\gamma\delta$. *Blood*. 75:2213-2220.
48. Hockett, R. D., J.-P. De Villartay, K. Pollock, D. G. Poplack, D. I. Cohen, and S. J. Korsmeyer. 1988. Human T-cell antigen receptor (TCR) δ -chain locus and elements responsible for its deletion are within the TCR α -chain locus. *Proc. Natl. Acad. Sci. USA*. 85:9694-9698.
49. De Villartay, J.-P., D. Lewis, R. Hockett, T. A. Waldmann, S. J. Korsmeyer, and D. I. Cohen. 1987. Deletional rearrangement in the human T-cell receptor α -chain locus. *Proc. Natl. Acad. Sci. USA*. 84:8608-8612.
50. De Villartay, J.-P., R. D. Hockett, D. Coran, S. J. Korsmeyer, and D. I. Cohen. 1988. Deletion of the human T-cell receptor δ -gene by a site-specific recombination. *Nature (Lond.)*. 335:170-174.
51. Winoto, A., and D. Baltimore. 1989. Separate lineages of T cells expressing the $\alpha\beta$ and $\gamma\delta$ receptors. *Nature (Lond.)*. 338:430-432.
52. Takeshita, S., M. Toda, and H. Yamagishi. 1989. Excision products of the T cell receptor gene support a progressive rearrangement model of the $\alpha\delta$ locus. *EMBO (Eur. Mol. Biol. Organ.) J.* 8:3261-3270.
53. Redondo, J. M., S. Hata, C. Brocklehurst, and M. S. Krangel. 1990. A T cell-specific transcriptional enhancer within the human T cell receptor δ locus. *Science (Wash. DC)*. 247:1225-1229.
54. Falini, B., L. Flenghi, S. Pileri, P. Pelicci, M. Fagioli, M. F. Martelli, L. Moretta, and E. Ciccone. 1989. Distribution of T cells bearing different forms of the T cell receptor γ/δ in normal and pathological human tissues. *J. Immunol.* 143:2480-2488.
55. Guglielmi, P., F. Davi, L. D'Auriol, J.-C. Bories, J. Dausset, and A. Bensussan. 1988. Use of a variable α region to create a functional T-cell receptor γ chain. *Proc. Natl. Acad. Sci. USA*. 85:5634-5638.
56. Casorati, G., G. De Libero, A. Lanzavecchia, and N. Migone. 1989. Molecular analysis of human γ/δ + clones from thymus and peripheral blood. *J. Exp. Med.* 70:1521-1535.
57. Hata, S., K. Satyanarayana, P. Devlin, H. Band, J. McLean, J. L. Strominger, M. B. Brenner, and M. S. Krangel. 1988. Extensive junctional diversity of rearranged human T cell receptor γ genes. *Science (Wash. DC)*. 240:1541-1544.
58. Borst, J., A. Wicherink, J. J. M. Van Dongen, E. De Vries, W. M. Comans-Bitter, F. Wassenaar, and P. Van den Elsen. 1989. Non-random expression of T cell receptor γ and δ variable gene segments in functional T lymphocyte clones from human peripheral blood. *Eur. J. Immunol.* 19:1559-1568.
59. Sturm, E., E. Braakman, R. E. Bontrop, P. Chuchana, R. J. Van de Griend, F. Koning, M.-P. Lefranc, and R. L. H. Bolhuis. 1989. Coordinated V γ and V δ gene segment rearrangements in human T cell receptor γ/δ + lymphocytes. *Eur. J. Immunol.* 19:1261-1265.