JCI The Journal of Clinical Investigation

120-kD surface glycoprotein of Pneumocystis carinii is a ligand for surfactant protein A.

P E Zimmerman, ..., J R Paulsrud, W J Martin 2nd

J Clin Invest. 1992;89(1):143-149. https://doi.org/10.1172/JCI115554.

Research Article

Pneumocystis carinii is the most common cause of life-threatening pneumonia in immunocompromised patients. In the current study, surfactant protein A (SP-A), the major nonserum protein constituent of pulmonary surfactant, is demonstrated to bind P. carinii in a specific and saturable manner. SP-A is surface bound and does not appear to be internalized or degraded by the P. carinii organism. Furthermore, SP-A binding to P. carinii is time- and calcium-dependent and is competitively inhibited by mannosyl albumin. In the absence of calcium or the presence of excess mannosyl albumin, SP-A binding to P. carinii is reduced by 95 and 71%, respectively. SP-A avidly binds P. carinii with a Kd of 8 x 10(-9) M and an estimated 8.4 x 10(6) SP-A binding sites per P. carinii organism, as determined from Scatchard plots. SP-A is shown to bind P. carinii in vivo, and a putative binding site for SP-A on P. carinii is demonstrated to be the mannoserich surface membrane glycoprotein gp120. These findings suggest that P. carinii can interact with the phospholipid-rich material in the alveolar spaces by specifically binding a major protein constituent of pulmonary surfactant.

Find the latest version:



120-kD Surface Glycoprotein of *Pneumocystis carinii* Is a Ligand for Surfactant Protein A

P. E. Zimmerman, D. R. Voelker, F. X. McCormack, J. R. Paulsrud, and W. J. Martin II

Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine and Department of Pathology, Indiana University School of Medicine, Indianapolis, Indiana 46202; and Department of Medicine, National Jewish Center for Immunology and Respiratory Medicine, Denver, Colorado 80206

Abstract

Pneumocystis carinii is the most common cause of life-threatening pneumonia in immunocompromised patients. In the current study, surfactant protein A (SP-A), the major nonserum protein constituent of pulmonary surfactant, is demonstrated to bind P. carinii in a specific and saturable manner. SP-A is surface bound and does not appear to be internalized or degraded by the P. carinii organism. Furthermore, SP-A binding to P. carinii is time- and calcium-dependent and is competitively inhibited by mannosyl albumin. In the absence of calcium or the presence of excess mannosyl albumin, SP-A binding to P. carinii is reduced by 95 and 71%, respectively. SP-A avidly binds P. carinii with a K_d of 8×10^{-9} M and an estimated 8.4×10^6 SP-A binding sites per P. carinii organism, as determined from Scatchard plots. SP-A is shown to bind P. carinii in vivo, and a putative binding site for SP-A on P. carinii is demonstrated to be the mannoserich surface membrane glycoprotein gp120. These findings suggest that P. carinii can interact with the phospholipid-rich material in the alveolar spaces by specifically binding a major protein constituent of pulmonary surfactant. (J. Clin. Invest. 1992. 89:143-149.) Key words: binding • lectin • pneumonia • surfactant apoprotein • AIDS

Introduction

The most characteristic morphologic finding in *Pneumocystis* carinii pneumonia is the filling of the alveoli with an abundant network of foamy surfactant-like material that encases the *P. carinii* organisms (1–3). The accumulation of this lipid- and protein-rich material in the alveolar spaces likely contributes to the impairment of gas exchange observed in *P. carinii* pneumonia (4). However, the interaction of *P. carinii* with the components of pulmonary surfactant is poorly understood.

Surfactant protein A (SP-A),¹ a surfactant-associated protein (5), may play a role in surfactant organization and homeostasis. SP-A has been implicated in the adsorption of phospholipid to the alveolar air-liquid interface (6), the in vitro regula-

Address correspondence to W. J. Martin II, M.D., Indiana University School of Medicine, Division of Pulmonary and Critical Care Medicine, 1001 W. 10th St., OPW 425, Indianapolis, IN 46202-28790.

Received for publication 9 January 1991 and in revised form 24 July 1991.

1. Abbreviation used in this paper: SP-A, surfactant protein A.

© The American Society for Clinical Investigation, Inc. 0021-9738/92/01/0143/07 \$2.00 Volume 89, January 1992, 143-149

tion of surfactant secretion (7–9) and the in vitro uptake of phospholipid by alveolar type II cells (10). In addition, SP-A is recognized as a calcium-dependent lectin with a binding specificity for immobilized mannose, fucose, glucose, and galactose (11).

The predominant surface membrane protein on *P. carinii* is a glycoprotein with an estimated molecular mass of 110–120 kD, known as *P. carinii* gp120 (12). Recent studies suggest that gp120 may serve to mediate attachment of *P. carinii* to the alveolar epithelium (13). *P. carinii* gp120 is heavily glycosylated with mannose-containing oligosaccharide chains (14) that could function as ligands for SP-A.

In this study, SP-A is demonstrated to avidly bind *P. carinii* both in vitro and in vivo. This binding is saturable, calcium dependent, and carbohydrate specific. Ligand immunoblotting of size-fractionated *P. carinii* membrane proteins demonstrate that SP-A selectively binds the carbohydrate expressed by the mannose-rich gp120. Finally, SP-A binding to *P. carinii* in vivo is verified by immunoblot analysis of *P. carinii* obtained by lung lavage using polyclonal antibody to rat SP-A.

Methods

Isolation of P. carinii. P. carinii pneumonia was induced in pathogenfree rats by immunosuppression with dexamethasone and transtracheal inoculation of P. carinii organisms as described by Bartlett et al. (15). P. carinii organisms were then harvested by lung lavage as previously described (16, 17). P. carinii were purified from the lavage fluid by a method adapted from Masur and Jones (18). The lavage fluid was centrifuged (600 g for 10 min) to pellet inflammatory and alveolar cells, and the supernatant was saved. The cell pellet was resuspended and cytopreparation smears were made using a cytologic centrifuge (Cytospin II; Shandon Southern Instruments Inc., Sewickley, PA). The cytopreparation smears were stained with Gomori methenamine silver stain (19) and Diff-Quick (Difco Laboratories, Inc., Detroit, MI) stain (20) to confirm the presence of P. carinii organisms. Any suspensions found to contain bacterial, fungal, or inflammatory cell contamination were discarded. The supernatant was centrifuged (2,000 g for 30 min) and the resulting P. carinii pellet was resuspended in HBSS without calcium or magnesium (Gibco Laboratories, Grand Island, NY) plus 0.6 mM EDTA. Examination of this suspension by light microscopy demonstrated P. carinii trophozoites to represent 97-98% of the intact cellular material present. P. carinii trophozoites were quantified by the method of Bartlett et al. (21).

SP-A isolation, purification, and iodination. Surfactant protein A was purified by previously described methods (22). Briefly, surfactant was obtained by lung lavage of rats 4 wk after intratracheal instillation of silica in saline. The surfactant was purified as described by Hawgood et al. (6) and delipidated by extraction with l-butanol. The butanol-insoluble protein precipitate was separated from the lipids by centrifugation (2,000 g for 30 min). The protein was suspended in 5 mM Tris/HCl buffer (pH 7.4) and dialyzed against the same buffer at 4° C for 36-48 h. The protein suspension was centrifuged (150,000 g for 1 h), and the supernatant was collected and then eluted over an affinity

J. Clin. Invest.

column of mannose-Sepharose 6B (23). The SP-A bound to the column in the presence of 2 mM CaCl₂ was then eluted with 2 mM EDTA. The eluate was further purified and residual EDTA removed by gel filtration over a Bio-Gel A 5-m column (Bio-Rad Laboratories, Richmond, CA). Protein content was estimated using the method of Lowry et al. (24).

 125 I-SP-A was prepared as previously described according to the method of Bolton and Hunter (25). Rat SP-A (1–2 mg) was dialyzed at 4°C against a 0.1 M sodium borate buffer (pH 8.5). A 1–1.5-ml aliquot of the SP-A suspension was added directly to an iced Bolton-Hunter Reagent vial (Amersham Corp., Arlington Heights, IL) after the benzene was removed by evaporation under a stream of nitrogen. The reaction mixture was incubated for 30 min on ice with occasional mixing. Free 125 I was removed by dialysis against a 5-mM Tris buffer (pH 7.4). The specific activity of the 125 I-SP-A was $^{31-54}\mu$ Ci/mg protein. The purity of the SP-A and 125 I-SP-A obtained by these methods has previously been demonstrated (22).

Binding of 125 I-SP-A to P. carinii. Isolated P. carinii were incubated 18 h in DME (Whittaker MA Bioproducts, Walkersville, MD) supplemented with 0.6 mg/ml glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin, 0.5 μg/ml amphotericin B, 4 μg/ml gentamicin, 1% BSA, and 5 mM EGTA (pH 7.4). To prepare P. carinii for binding experiments, the P. carinii suspension was centrifuged (2,000 g for 15 min), the supernatant was discarded, and the pellet was washed once with HBSS containing 2 mM CaCl₂ supplemented with 1% BSA (HBSS⁺) and finally resuspended in DME plus 1% BSA to a concentration of 5 × 10⁵ P. carinii organisms/ml. Binding of SP-A to P. carinii trophozoites was quantified using a modification of the method of Proctor et al. (26). Binding assays performed to determine the specific binding of SP-A to P. carinii contained 5×10^4 P. carinii (100 μ l), 60–1,500 ng ¹²⁵I-SP-A (10-30 μl) in sample buffer (5 mM Tris/HCl and 1% BSA, pH 7.4) in the presence and absence of 10 μ g unlabeled SP-A. The reaction suspension was incubated for 30 min in an atmosphere of 90% air and 10% CO₂ at 37°C. The suspension was spun in a microfuge (13,600 g for 2 min). The cell pellet was washed one time with HBSS+, and the pellet was added to a 12 × 75-mm tube for counting. Counts per minute present were determined using a gamma counter, and the amount of ¹²⁵I-SP-A bound was quantified (ng bound = CPM_{pellet}/protein specific activity). To study the effect of time, temperature, calcium, and excess carbohydrate on the binding of SP-A to P. carinii, binding assays were performed using 1×10^6 P. carinii in HBSS⁺, 100 ng ¹²⁵I-SP-A, and increasing concentrations of calcium (0-2 mM), α -methylmannopyranoside (Sigma Chemical Co., St. Louis, MO) (0-750 mM) and mannosyl-BSA $(0-1,000 \mu g)$ with a molar ratio of mannose to albumin of 26:1 (Sigma Chemical Co., St. Louis, MO). The final volume of the mixture was 160 µl. The assay was incubated as above and the amount of 125I-SP-A bound was quantified (ng bound = 100 \times CPM_{pellet}/[CPM_{pellet + supernatant}]).

The ability of anti-SP-A IgG to inhibit binding was also examined in a similar manner using 0, 1.0, and 2.5 μ g of antibody. Polyclonal rabbit anti-rat SP-A IgG was raised in New Zealand white rabbits. The animals were injected with 50-100 µg of purified rat SP-A in Freund's complete adjuvant. A second booster injection of 50-100 µg of rat SP-A in Freund's incomplete adjuvant was given at 14 d. The titer of antibody was assayed 10 d after the booster injection by ELISA using purified rat SP-A as the solid phase. A repeat booster injection was given in cases where the titer was < 1:10,000. Antibody was isolated from immune rabbit serum on Protein-A Sepharose CL-4B (Pharmacia LKB Biotechnology, Inc., Piscataway, NJ) and eluted with 0.1 M acetic acid. The eluted fraction was neutralized immediately in Tris buffer and adsorbed against rat serum bound to Affigel-10 and Affigel-15 columns (Bio-Rad Laboratories). The antibody was exhaustively dialyzed against PBS and stored at -20°C. The purity and specificity of this antibody has previously been demonstrated (9).

To examine whether SP-A is surface bound or internalized, binding assays were performed as above at 37° and at 4°C, a temperature at which internalization is inhibited. A second method to determine whether ¹²⁵I-SP-A was surface bound used release of ¹²⁵I-SP-A from *P*.

carinii by washing the cells with 5 mM Tris/HCl plus 5 mM EGTA (pH 7.4). Since the binding of SP-A to *P. carinii* is calcium dependent, this treatment releases surface-bound but not internalized ¹²⁵I-SP-A.

Finally, autoradiography and TCA precipitation of purified and bound ¹²⁵I-SP-A were used to study whether bound SP-A remains intact or is degraded after binding. *P. carinii* bound by ¹²⁵I-SP-A were solubilized in SDS-PAGE sample buffer (0.125 M Tris, 20% glycerol, 4% SDS, 0.002% bromophenol blue, and 4% 2-mercaptoethanol), and the proteins were size fractionated by SDS-PAGE. The gel was dried and exposed to Kodak X-Omat RP film for 2 h at -70°C. TCA precipitation of purified and *P. carinii*-bound ¹²⁵I-SP-A was done using a TCA final concentration of 26%.

Binding of SP-A to P. carinii membrane proteins. Rat P. carinii organisms obtained from culture (27) were solubilized in 9.5 M urea, 2% (wt/vol) NP-40, 2% ampholines (composed of 1.6% pH range 5-7 and 0.4% pH range 3-10) and 5% 2-mercaptoethanol. The P. carinii proteins were then electrofocused and size fractionated by two-dimensional gel chromatography as described by O'Farrell (28). To demonstrate whether SP-A binds a P. carinii surface protein, the protein in the two-dimensional gel was electrophoretically transferred for 30 min to an Immobilon-P PVDF transfer membrane (Millipore Corp., Bedford, MA) using Towbin buffer (29) supplemented with 0.05% SDS. The membrane was blocked with 3% BSA, 5 mM Tris, 2 mM CaCl₂, 0.5% Triton X-100 pH 7.4 (buffer A) for 2 h. The membrane was blotted with 20 μ g/ml SP-A in buffer A for 2 h at room temperature. After this, all washes and incubations were done at 4°C. After each incubation the membrane was washed five times, 5 min/wash with buffer A. The membrane was immunoblotted with 20 µg/ml polyclonal rabbit antibody to rat SP-A for 2 h. The bound SP-A was visualized using horseradish peroxidase-conjugated goat anti-rabbit IgG antibody developed with 4-chloro-1-naphthol in diethylene glycol (Bio-Rad Laboratories).

Demonstration of in vivo binding of SP-A to P. carinii. To demonstrate whether SP-A is bound to P. carinii organisms in vivo, organisms were collected as before in lavage fluid containing 2 mM CaCl₂. Lung lavage from uninfected animals was obtained and purified in a similar fashion and used as a control. The P. carinii obtained from lung lavage were centrifuged (71,000 g for 90 min) through a 20-60% discontinuous sucrose density gradient. The cell band present at the 20-30% (0.58-0.88 M) sucrose interface was collected and solubilized in SDS-PAGE sample buffer. The P. carinii and the corresponding control fractions were subjected to SDS-PAGE, transferred to nitrocellulose, and then immunoblotted with rabbit polyclonal antibody to rat SP-A. The presence of SP-A was identified using horseradish peroxidase-conjugated goat anti-rabbit IgG antibody directed against the rabbit polyclonal anti-SP-A antibody.

Statistical analysis. Values are expressed as means \pm SEM. Differences between control and experimental data groups were compared using the one-tailed t test or ANOVA with paired comparisons performed using Fisher's LSD method. Statistical significance was accepted for P < 0.01.

Results

¹²⁵I-SP-A binding to P. carinii trophozoites. SP-A binds P. carinii in vitro in a time-dependent manner and achieves apparent equilibrium by 30 min (Fig. 1). At 5 min 34.45 \pm 0.62 ng ¹²⁵I-SP-A/1 × 10⁶ organisms is bound, whereas at the observed times of 30 min and beyond, mean binding of ¹²⁵I-SP-A plateaus at 52.62 \pm 0.56 ng/1 × 10⁶ organisms. Further binding experiments were conducted at 30 min, a time when equilibrium of SP-A binding was established.

The calcium requirement for optimal binding of SP-A to *P. carinii* was examined by repeating the SP-A binding assays in the presence of increasing concentrations of extracellular calcium (Fig. 2). In the absence of calcium only 2.85±0.27 ng of ¹²⁵I-SP-A is bound, whereas in the presence of 2 mM calcium




Figure 1. ¹²⁵I-SP-A binds *P. carinii* in a time-dependent fashion. Binding assays were performed using 1×10^6 *P. carinii* incubated with 100 ng ¹²⁵I-SP-A in HBSS, 2 mM calcium, and 1% BSA for the indicated times. Binding equilibrium is evident at 30 min, with 51.92 ± 1.33 ng ¹²⁵I-SP-A bound/1 \times 10⁶ organisms. Values are means \pm SEM of three experiments performed in duplicate.

59.96±3.05 ng SP-A is bound. Binding appears to plateau beyond calcium concentrations in excess of 1 mM calcium. This is in agreement with previous studies, demonstrating the calcium requirement of SP-A ligand binding (11). Therefore, extracellular calcium appears to also be required for the optimal binding of SP-A to *P. carinii*.

The specificity and ligand concentration dependence of 125 I-SP-A binding to *P. carinii* was demonstrated using increasing amounts of 125 I-SP-A (60–1,500 ng) in the presence and absence of 10 μ g excess unlabeled SP-A (Fig. 3 *A*). Addition of increasing amounts of 125 I-SP-A to the binding assay resulted in an increase in both total and nonspecific binding, with saturation of specific binding evident at concentrations in excess of 1,000 ng 125 I-SP-A added. Scatchard plots (30) using the specific binding data are linear (r = 0.99), indicating a homoge-





Figure 2. Calcium is required for ¹²⁵I-SP-A to bind *P. carinii*. Binding assays were performed as in Fig. 1, using increasing amounts of extracellular calcium. Maximal binding reveals a plateau at calcium concentrations above 1 mM. Values are means±SEM of three experiments performed in triplicate.







Figure 3. Saturation and specific binding of 125 I-SP-A to *P. carinii*. Binding assays were performed using 5×10^4 *P. carinii*, 60–1,500 ng 125 I-SP-A with and without $10~\mu g$ excess unlabeled SP-A. (A) Saturation of specific SP-A binding to *P. carinii* demonstrated in the presence of > 1,000 ng 125 I-SP-A added. Values are means \pm SEM of three experiments performed in triplicate. (B) Scatchard plot of specific binding data. The slope of the line represents a K_d of 8×10^{-9} M for the binding interaction of SP-A and *P. carinii*. The number of SP-A binding sites per organism was calculated from the *x*-intercept to be 8.4×10^6 .

nous population of binding sites for SP-A on *P. carinii* (Fig. 3 *B*). The binding dissociation constant (K_d) is 8×10^{-9} M, with each *P. carinii* trophozoite estimated to have 8.4×10^6 binding sites for SP-A, using an estimated molecular mass for SP-A of 750 kD (31, 32).

P. carinii binding of SP-A occurs at the cell surface and does not appear to require active internalization of the SP-A oligomer. Binding of SP-A to P. carinii occurs both at 37° and at 4°C, a temperature at which membrane vesicle internalization is inhibited (33) (Fig. 4 A). Binding is slightly greater at 4°C (71.17±2.31 ng) compared with 37°C (50.03±2.79 ng). Surface binding is further supported by demonstrating that 92% of ¹²⁵I-SP-A bound at 37°C can be removed from P. carinii by washing the organisms with 5 mM EGTA (Fig. 4 B).

Autoradiography of ¹²⁵I-SP-A bound to *P. carinii* at 37°C reveals that SP-A is not degraded. No SP-A fragments are found upon examination of an autoradiogram prepared from an SDS-PAGE of bound ¹²⁵I-SP-A when compared with puri-






Figure 4. SP-A is bound to the cell surface of *P. carinii*. (A) Binding assays were performed using 100 ng 125 I-SP-A incubated with 1×10^6 *P. carinii* for 30 min in HBSS, 2 mM calcium, and 1% BSA at 37°C and 4°C. Binding is clearly evident at 4°C, a temperature at which receptor internalization inhibited. Values are means±SEM of three experiments performed in triplicate (*P < 0.01). (B) 125 I-SP-A bound to *P. carinii* at 37°C is nearly completely reversed by washing the organisms in 5 mM EGTA. Values are means±SEM of three experiments performed in triplicate (*P < 0.01).

fied 125 I-SP-A (data not shown). Furthermore, there is no difference between the TCA precipitability of purified 125 I-SP-A compared with the 125 I-SP-A incubated with *P. carinii*, 94.1 vs. 94.6%, respectively (P > 0.50). The failure to show a disparity between these two groups provides evidence supporting the absence of formation of small 125 I-SP-A peptide degradation fragments by the *P. carinii* organism.

SP-A binding to *P. carinii* could also be inhibited by polyclonal anti-SP-A IgG (Fig. 5). In the absence of antibody 38.40 ± 3.22 ng ¹²⁵I-SP-A is bound, whereas in the presence of 2.5 μ g anti-SP-A IgG binding is reduced to 9.77 ± 2.05 ng.

To determine whether SP-A recognized carbohydrates expressed by *P. carinii* surface membrane glycoproteins, ¹²⁵I-SP-A binding experiments were conducted in the presence and absence of excess carbohydrate. Because *P. carinii* expresses high mannose-type oligosaccharides on its surface (14) and the specificity of SP-A for mannose has been well described (11), α -methyl-D-mannopyranoside and mannosyl-BSA were





Figure 5. Polyclonal anti-SP-A IgG inhibits SP-A binding to *P. carinii*. ¹²⁵I-SP-A binding assays were performed in the absence and presence of polyclonal anti-SP-A IgG. SP-A binding is inhibited by 75% using 2.5 μ g of antibody. Values are means±SEM of three experiments performed in triplicate (*P < 0.01).

used to determine whether SP-A binding to P. carinii exhibited a specificity for carbohydrate. Mannosyl-BSA effectively inhibits ¹²⁵I-SP-A binding to P. carinii (Fig. 6 A). SP-A binding can be inhibited by 71% in the presence of 2.42 mM mannose when mannose is presented as the multivalent glycoprotein mannosyl-BSA, whereas the monosaccharide α -methyl-Dmannopyranoside (Fig. 6 B) requires concentrations as high as 500 mM to achieve a similar degree of inhibition (73%). Other monosaccharides known to be ligands for SP-A, such as N-acetyl-D-glucosamine also inhibit SP-A binding to P. carinii at high concentrations (data not shown). However, there is a limitation to the use of monosaccharides in studying SP-A binding, because the carbohydrate specificity of SP-A decreases as the monosaccharide concentration increases. The ability of low mannose concentrations, in the form of mannosyl-BSA, to inhibit binding of SP-A to P. carinii suggests that a glycoprotein expressed by P. carinii is a possible ligand for SP-A.

SP-A binding to P. carinii membrane proteins. To demonstrate the SP-A target binding site on P. carinii, SP-A binding to P. carinii membrane proteins was examined by ligand immunoblot analysis. Whole P. carinii proteins were resolved by two-dimensional gel chromatography and stained with Coomassie brilliant blue R-250 (Fig. 7 A). Two-dimensional chromatography of P. carinii proteins reveal a 116-120-kD basic protein, representing the majority of P. carinii surface protein (12) known as gp120.

The immunoreactivity for SP-A present on the ligand immunoblot analysis (Fig. 7 B) demonstrates that SP-A binds the P. carinii 120-kD surface glycoprotein, as well as a smaller protein with an estimated mass of 80 kD. There is no immunoreactivity for SP-A present when immunoblots are performed with omission of the initial SP-A incubation (data not shown). Ligand immunoblots of P. carinii gp120 treated with 50 mM sodium metaperiodate (34), express no immunoreactivity for SP-A (data not shown), demonstrating that oxidation of the carbohydrate residues of gp120 destroys the epitope required for SP-A binding. The inhibition of SP-A binding to whole P. carinii organisms with mannosyl-BSA and the periodate sensi-






Figure 6. ¹²⁵I-SP-A binding to *P. carinii* is inhibited by excess monovalent and multivalent carbohydrates. (A) Binding assays were performed in the presence of increasing amounts of the multivalent ligand mannosyl-BSA (0–1,000 μ g), resulting in an increase in monosaccharide concentration (0–2.42 mM). In the presence of 2.42 mM mannose (1 mg mannosyl-BSA), 71% of ¹²⁵I-SP-A binding is inhibited. There is no inhibition of ¹²⁵I-SP-A binding observed when identical concentrations of the carrier protein, BSA, are used (data not shown). (B) ¹²⁵I-SP-A binding assays were performed in the presence of increasing concentrations of the monosaccharide α -methyl-D-mannopyranoside. At the highest concentration tested ¹²⁵I-SP-A binding is inhibited by 86%. Values are means±SEM of three experiments performed in triplicate (*P < 0.01).

tivity of SP-A binding to *P. carinii* membrane proteins provide strong evidence that SP-A recognizes oligosaccharide moieties of gp120.

In vivo demonstration of SP-A binding to P. carinii. Because SP-A binds P. carinii in vitro, the pivotal issue is whether similar binding occurs in vivo within the alveolar milieu. In vivo SP-A binding to P. carinii was demonstrated by use of immunoblot analysis of P. carinii membrane proteins isolated and purified as described above. The immunoblot (Fig. 8) using polyclonal rabbit antibody to rat SP-A demonstrates the specificity of the antibody to purified SP-A (lane 3). Lanes 1 and 4 represent identical immunoreactivity for SP-A found to be present in the isolate of P. carinii (lane 1) and in P. carinii purified from discontinuous sucrose density gradients (lane 4). There is

no immunoreactivity for SP-A evident in the isolate obtained from pathogen-free rats (lane 2). The absence of immunoreactivity in lane 2 is further evidence that the methods of *P. carinii* isolation from lung lavage does not also co-purify free host SP-A with the *P. carinii* organisms. This experiment provides direct evidence for the in vivo binding of SP-A to *P. carinii*.

Discussion

P. carinii organisms have previously been demonstrated to avidly bind alveolar surfactant (35); however, the mechanism of adherence between surfactant and *P. carinii* has not been defined. This study indicates that SP-A, the major protein constituent of surfactant, binds *P. carinii* in a specific and saturable fashion. Scatchard plots estimate the number of SP-A binding sites per *P. carinii* organism to be $\sim 8.4 \times 10^6$. The linearity of the plots suggest that a homogeneous group of binding sites exist, with a similar affinity for SP-A.

SP-A exists in vivo as a large multivalent oligomeric protein composed of \sim 18 subunits, similar to that of complement component C1q (36). Analysis of the deduced primary amino




Figure 7. SP-A binds P. carinii gp120. (A) Two-dimensional gel chromatography of P. carinii proteins. P. carinii organisms were electrofocused (1, first dimension) using ampholines with a pH range of 3 to 10 and then size fractionated by SDS-PAGE (2, second dimension) and stained with Coomassie brilliant blue. The predominate band seen represents P. carinii gp120. (B) Ligand immunoblot of SP-A binding whole P. carinii proteins. SP-A binds almost exclusively to the protein band corresponding to the mannose-rich cell surface glycoprotein gp120. In addition there is a small amount of immunoreactivity for SP-A present in the 80-kD area representing SP-A binding to a lesser expressed P. carinii glycoprotein.




Figure 8. P. carinii binds SP-A in vivo. P. carinii membrane proteins were prepared and immunoblotted using polyclonal antibody to rat SP-A as described in the methods. Immunoreactivity for SP-A is demonstrated in the same 26-38 kD range in the P. carinii membrane protein preparations (lanes 1 and 4) as is visible in the purified SP-A control lane (lane 3). No immunoreactivity for SP-A is evident in lane 2, representing alveolar lavage proteins prepared from pathogen-free rats in the same fashion as the P. carinii proteins.

acid structure of SP-A reveals that the protein has several discrete structural domains (37–41). Of particular relevance to this study are the amphipathic helix and hydrophobic region, which form a putative phospholipid-binding domain (42), and the globular carboxyl terminus region, which contains a carbohydrate-binding domain (43). The amino acid sequence of the carbohydrate recognition domain shows sequence homology with the carbohydrate-recognition domains of other C-type lectins, specifically the soluble serum lectin, mannose-binding protein, and the hepatic and macrophage mannose receptor (43–45). SP-A has previously been shown to require calcium for binding to immobilized saccharides or to its receptor on isolated alveolar type II cells (22, 46, 47). These data suggest that SP-A binding to *P. carinii* is also a carbohydrate-dependent and calcium-requiring process.

The multiple affinities of SP-A for carbohydrate and phospholipid (42, 48) may enable this protein to simultaneously bind P. carinii and the phospholipid components of surfactant. Therefore, SP-A may facilitate the accumulation of surfactant phospholipids about the organism. Saturation of P. carinii's major antigenic determinant, gp120, by SP-A alone or in combination with adsorbed surfactant may alter the structure of this immunodeterminant such that recognition by host immune surveillance cells could be avoided. Parasite acquisition of host molecules is a recognized means of host immune response avoidance (49). The interaction of P. carinii with SP-A could potentially result in the combined disruption of SP-A's regulatory role of alveolar phospholipid homeostasis and the disorganization of the alveolar phospholipid monolayer, in part explaining the alveolar filling process and hypoxia seen in P. carinii pneumonia (4).

The predominant antigenic determinant on the *P. carinii* cell membrane is the mannose-rich membrane glycoprotein gp120 (34, 50). In this study, SP-A is demonstrated to bind *P.*

carinii gp120, supporting the hypothesis that gp120 is a putative binding site on *P. carinii* for SP-A. The ability of mannosyl-BSA to inhibit binding of SP-A to gp120 and the periodate sensitivity of SP-A binding suggests that the SP-A binding is dependent on intact oligosaccharides present on the gp120 ligand. Furthermore, immunoblot analysis of isolated *P. carinii* membrane proteins demonstrates the presence of SP-A, indicating that SP-A is adherent to *P. carinii* in vivo.

P. carinii gp120 has been demonstrated to play a potential role in the mechanism of adherence of the organism to the alveolar epithelium (13). The binding of the carbohydrate recognition domain of SP-A to P. carinii in the alveolar lining fluid may provide the organism access to the alveolar epithelium via fibronectin-mediated attachment between SP-A's free collagen-like domain and the alveolar epithelium. This is suggested by fibronectin's ability to bind the collagenous domain of the structurally homologous protein complement C1q (51) and fibronectin's ability to mediate attachment of P. carinii to the alveolar epithelium (15).

In summary, this study provides direct evidence that SP-A, the major protein constituent of surfactant, binds *P. carinii*. This binding is specific and saturable, time dependent, carbohydrate specific, and calcium requiring. The binding site on the *P. carinii* organism appears to be the mannose-rich membrane glycoprotein gp120. The binding of SP-A to *P. carinii* in vivo may enhance adsorption of surfactant to the organism, possibly impede host defenses against *P. carinii*, as well as potentiate *P. carinii* adherence to the alveolar epithelium.

Acknowledgments

We thank David Smith for his excellent assistance in the purification and iodination of SP-A and Steve Brady for his time in preparation of the two-dimensional gels and PVDF membranes.

This study was supported by National Institutes of Health grants HL-43524 and HL-46647 (W. J. Martin) and HL-29891 (D. R. Voelker). Dr. Zimmerman is the recipient of the Sunnyside Guild Fellowship in Pulmonary Medicine. Part of this work was performed in the Lord and Taylor Laboratory for Lung Biochemistry and the Anna Perahia Adatto Clinical Research Center, Denver, CO.

References

- Rifkind, D., T. D. Faris, and R. B. Hill. 1966. Pneumocystis carinii pneumonia. Studies on the diagnosis and treatment. Ann. Intern. Med. 65:943-956.
- 2. McNeal, J. E., and T. G. Yaeger. 1960. Observations on a case of *Pneumocystis pneumonia*. Arch. Pathol. 70:397-406.
- 3. Esterly, J. A., and N. E. Warner. 1965. *Pneumocystis carinii* pneumonia: twelve cases in patients with neoplastic lymphoreticular diseases. *Arch. Pathol.* 80:433-441.
- Van Nhieu, J. T., A. M. Vojtek, J. Bernaudin, E. Escudier, and J. Fleury-Feith. 1990. Pulmonary alveolar proteinosis associated with *Pneumocystis car*inii. Chest. 98:801–805.
- 5. King, R. J., and J. A. Clements. 1972. Surface active materials from dog lung. II. Composition and physiological correlations. *Am. J. Physiol.* 223:715-726.
- 6. Hawgood, S., B. J. Benson, J. Schilling, D. Damm, J. Clements, and R. T. White. 1987. Nucleotide and amino acid sequences of pulmonary surfactant protein SP 18 and evidence for cooperation between SP 18 and SP 28-36 in surfactant lipid adsorption. *Proc. Natl. Acad. Sci. USA*. 84:66-70.
- 7. Dobbs, G. L., J. R. Wright, S. Hawgood, R. Gonzalez, K. Venstrom, and J. Nellenbogen. 1987. Pulmonary surfactant and its components inhibit secretion of phosphatidylcholine from cultured rat alveolar type II cells. *Proc. Natl. Acad. Sci. USA*. 84:1010–1014.
- 8. Rice, W. R., G. F. Ross, F. M. Singleton, S. Dingle, and J. A. Whitsett. 1987. Surfactant-associated protein inhibits phospholipid secretion from type II cells. *J. Appl. Physiol.* 63:692–698.

- 9. Kuroki, Y., R. J. Mason, and D. R. Voelker. 1988. Pulmonary surfactant apoprotein A structure and modulation of surfactant secretion by rat alveolar type II cells. *J. Biol. Chem.* 263:3388-3394.
- 10. Wright, J. R., R. E. Wager, S. Hawgood, L. Dobbs, and J. A. Clements. 1987. Surfactant apoprotein $M_\tau=26,000-36,000$ enhances uptake of liposomes by type II cells. *J. Biol. Chem.* 262:2888–2894.
- 11. Haagsman, H. P., S. Hawgood, T. Sargeant, D. Buckley, R. T. White, K. Drickamer, and B. J. Benson. 1987. The major lung surfactant protein, SP 28-36, is a calcium-dependent, carbohydrate-binding protein. *J. Biol. Chem.* 262:13877-13880.
- 12. Radding, J. A., M. Y. K. Armstrong, E. Ullu, and F. F. Richards. 1989. Identification and isolation of a major cell surface glycoprotein of *Pneumocystis carinii*. *Infect. Immun.* 57:2149–2157.
- 13. Pottratz, S. T., J. Paulsrud, and W. J. Martin II. 1990. *Pneumocystis carinii* membrane glycoprotein gp120 is a fibronectin binding protein which mediates Pneumocystis attachment. *Clin. Res.* 38:466a. (Abstr.)
- 14. Nakamura, Y., K. Tanabe, and K. Egawa. 1989. Structure of major surface determinants and DNA diagnosis of *Pneumocystis carinii*. *J. Protozool*. 36:60s-61s.
- 15. Bartlett, M. S., J. A. Fishman, S. F. Queener, M. M. Durkin, M. A. Jay, and J. W. Smith. 1988. New rat model of *Pneumocystis carinii* pneumonia. *J. Clin. Microbiol.* 26:1100-1102.
- 16. Pottratz, S. T., and W. J. Martin II. 1990. Role of fibronectin in *Pneumocystis carinii* attachment to cultured lung cells. *J. Clin. Invest.* 85:351–356.
- 17. Limper, A. H., and W. J. Martin II. 1990. *Pneumocystis carinii:* inhibition of lung cell growth mediated by parasite attachment. *J. Clin. Invest.* 85:391–396.
- 18. Masur, H., and T. C. Jones. 1978. The interaction in vitro of *Pneumocystis carinii* with macrophages and L-cells. *J. Exp. Med.* 147:157-170.
- 19. Brinn, N. T. 1975. Rapid metallic histological staining. *J. Histotechnol.* 6:125–129.
- 20. Cushion, M. T., and P. D. Walzer. 1984. Growth and serial passage of *Pneumocystis carinii* in A549 cell line. *Infect. Immun.* 44:245-251.
- 21. Bartlett, M. S., P. A. Vervanec, and J. W. Smith. 1979. Cultivation of *Pneumocystis carinii* with WI-38 cells. *J. Clin. Microbiol.* 10:796–799.
- 22. Kuroki, Y., R. J. Mason, and D. R. Voelker. 1988. Alveolar type II cells express a high-affinity receptor for pulmonary surfactant protein A. *Proc. Natl. Acad. Sci. USA*. 85:5566-5570.
- 23. Fornstedt, H. P., and J. Porath. 1975. Characterization studies on a new lectin found in seeds of *Vicia ervilia. FEBS (Fed. Eur. Biochem. Soc.) Lett.* 57:187-191.
- 24. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265–275.
- 25. Bolton, A. E., and W. M. Hunter. 1973. The labelling of proteins to high specific radioactivities by conjugation to a ¹²⁵l-containing acylating agent. *Biochem. J.* 133:529-539.
- 26. Proctor, R. A., D. F. Mosher, and P. J. Olbrantz. 1982. Fibronectin binding to Staphylococcus aureus. J. Biol. Chem. 257:14788-14794.
- 27. Durkin, M. M., M. S. Bartlett, S. F. Queener, M. M. Shaw, and J. W. Smith. 1989. A culture method allowing production of relatively pure *Pneumocystis carinii* trophozoites. *J. Protozool.* 36:31s-32s.
- 28. O'Farrell, P. H. 1975. High resolution two-dimensional electrophoresis of proteins. J. Biol. Chem. 250:4007–4021.
- 29. Towbin, H., T. Staehelin, and J. Gordon. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA*. 76:4350–4354.
- 30. Scatchard, G. 1949. The attractions of proteins for small molecules and ions. *Ann. NY Acad. Sci.* 51:660-672.
- 31. King, R. J., D. Simon, and P. M. Horowitz. 1989. Aspects of secondary and quaternary structure of surfactant protein a from canine lung. *Biochim. Biophys. Acta.* 1001:294-301.
 - 32. Haagsman, H. P., R. T. White, J. Schilling, K. Lau, B. J. Benson, J.

- Golden, S. Hawgood, and J. A. Clements. 1989. Studies of the structure of lung surfactant protein SP-A. Am. J. Physiol. 257:L421–L429.
- 33. Stahl, P., P. H. Schlesinger, E. Sigardson, J. S. Rodman, and Y. C. Lee. 1980. Receptor-mediated pinocytosis of mannose glycoconjugates by macrophages: characterization and evidence for receptor recycling. *Cell.* 19:207–215.
- 34. Graves, D. C., S. J. N. McNabb, M. A. Worley, T. D. Downs, and M. H. Ivy. 1986. Analyses of rat *Pneumocystis carinii* antigens recognized by humans and rat antibodies by using western immunoblotting. *Infect. Immun.* 54:96-103.
- 35. Pesanti, E. L. 1987. Phospholipid profile of *Pneumocystis carinii* and its interaction with alveolar type II epithelial cells. *Infect. Immun.* 55:736–741.
- 36. Voss, T., H. Eistetter, and K. P. Schafer. 1988. Macromolecular organization of natural and recombinant lung surfactant protein SP 28-36. Structural homology with the complement factor C1q. J. Mol. Biol. 201:219-227.
- 37. Benson, B., S. Hawgood, J. Schilling, J. Clements, D. Damm, B. Cordell, and R. T. White. 1985. Structure of canine pulmonary surfactant apoprotein: cDNA and complete amino acid sequence. *Proc. Natl. Acad. Sci. USA*. 82:6379–6383.
- 38. White, R. T., D. Damm, J. Miller, K. Spratt, J. Schilling, S. Hawgood, B. Benson, and B. Cordell. 1985. Isolation and characterization of the human pulmonary surfactant apoprotein gene. *Nature (Lond.)*. 317:361–363.
- 39. Floros, J., R. Steinbrink, K. Jacobs, D. Phelps, R. Kriz, M. Recny, L. Sultzman, S. Jones, H. W. Taeusch, H. A. Frank, and E. F. Fritsch. 1986. Isolation and characterization of cDNA clones for the 35-kDa pulmonary surfactant-associated protein. *J. Biol. Chem.* 261:9029–9033.
- 40. Sano, K., J. Fisher, R. J. Mason, Y. Kuroki, J. Schilling, B. Benson, and D. Voelker. 1987. Isolation and sequence of a cDNA clone for the rat pulmonary surfactant-associated protein (PSP-A). *Biochem. Biophys. Res. Commun.* 144:367–374.
- 41. Boggaram, V., K. Qing, and C. R. Mendelson. 1988. The major apoprotein of rabbit pulmonary surfactant. J. Biol. Chem. 263:2939–2947.
- 42. Ross, G. F., R. N. Notter, J. Meuth, and J. A. Whitsett. 1986. Phospholipid binding and biophysical activity of pulmonary surfactant-associated protein (SAP)-35 and its noncollagenous COOH-terminal domains. *J. Biol. Chem.* 261:4283–4291.
- 43. Drickamer, K., M. A. Dordal, and L. Reynolds. 1986. Mannose-binding proteins isolated from rat liver contain carbohydrate-recognition domains linked to collagenous tails. *J. Biol. Chem.* 261:6878–6887.
- 44. Drickamer, K. 1989. Demonstration of carbohydrate-recognition activity in diverse proteins which share a common primary structure motif. *Biochem. Soc. Trans.* 17:13-15.
- 45. Sastry, K., G. A. Herman, L. Day, E. Deignan, G. Bruns, C. C. Morton, and R. A. B. Ezekowitz. 1989. The human mannose-binding protein gene exon structure reveals its evolutionary relationship to human pulmonary surfactant gene and localization to chromosome 10. J. Exp. Med. 170:1175–1189.
- 46. Ryan, R. M., R. E. Morris, W. R. Rice, C. Ciraolo, and J. Whitsett. 1989. Binding and uptake of pulmonary surfactant protein (SP-A) by pulmonary type II epithelial cells. *J. Histochem. Cytochem.* 37:429–440.
- 47. Wright, J. R., J. D. Borchelt, and S. Hawgood. 1989. Lung surfactant apoprotein SP-A (26–36 kDa) binds with high affinity to isolated alveolar type II cells. *Proc. Natl. Acad. Sci. USA*. 86:5410–5414.
- 48. King, R. J., M. C. Carmichael, and P. M. Horowitz. 1983. Reassembly of lipid-protein complexes of pulmonary surfactant. *J. Biol. Chem.* 258:10672–10680.
- 49. Bloom, B. R. 1979. Games parasites play: how parasites evade immune surveillance. *Nature (Lond.)*, 279:21–26.
- 50. Linke, M. J., M. T. Cushion, and P. D. Walzer. 1989. Properties of the major antigens of rat and human *Pneumocystis carinii*. *Infect. Immun*. 57:1547–1555.
- 51. Menzel, J. E., J. S. Smolen, L. Liotta, and K. B. M. Reid. 1982. Interaction of fibronectin with C1q and its collagen-like fragment (CLF). FEBS (Fed. Eur. Biochem. Soc.) Lett. 129:188-192.