

**Keratinocyte growth factor and hepatocyte growth factor/scatter factor are heparin-binding growth factors for alveolar type II cells in fibroblast-conditioned medium**

*J Clin Invest.* 1994;[93\(3\)](#):1347-1347. <https://doi.org/10.1172/JCI116673C1>.

**Correction**

**Find the latest version:**

<https://jci.me/116673C1/pdf>



Panos, R. J., J. S. Rubin, S. A. Aaronson, and R. J. Mason.  
*The Journal of Clinical Investigation*. Volume 92, No. 2, August 1993.

Page 969.

While Dr. Karl G. Csaky was originally cited in the acknowledgments of the article, the authors believe he should be cited as a coauthor since the description and initial application of his technique for the detection of hepatocyte growth factor/scatter factor were first reported in this article. There are no changes to the body of the article.

The author information should read as follows:

Keratinocyte Growth Factor and Hepatocyte Growth Factor/Scatter Factor Are Heparin-binding Growth Factors for Alveolar Type II Cells in Fibroblast-conditioned Medium

Ralph J. Panos,\* Jeffrey S. Rubin,† Karl G. Csaky,‡ Stuart A. Aaronson,‡ and Robert J. Mason§

\*Pulmonary Division, Department of Medicine, Northwestern University Medical School, Chicago, Illinois 60611; †Laboratory of Cellular and Molecular Biology, National Cancer Institute, Bethesda, Maryland 20892; and ‡Pulmonary Division, National Jewish Center for Immunology and Respiratory Medicine, University of Colorado Health Sciences Center, Denver, Colorado 80206.