

Keratinocyte growth factor and hepatocyte growth factor/scatter factor are heparin-binding growth factors for alveolar type II cells in fibroblast-conditioned medium

J Clin Invest. 1994;93(3):1347-1347. <https://doi.org/10.1172/JCI116673C1>.

Correction

Find the latest version:

<https://jci.me/116673C1/pdf>



Correction

Panos, R. J., J. S. Rubin, S. A. Aaronson, and R. J. Mason.
The Journal of Clinical Investigation. Volume 92, No. 2, August 1993.

Page 969.

While Dr. Karl G. Csaky was originally cited in the acknowledgments of the article, the authors believe he should be cited as a coauthor since the description and initial application of his technique for the detection of hepatocyte growth factor/scatter factor were first reported in this article. There are no changes to the body of the article.

The author information should read as follows:

Keratinocyte Growth Factor and Hepatocyte Growth Factor/Scatter Factor Are Heparin-binding Growth Factors for Alveolar Type II Cells in Fibroblast-conditioned Medium

Ralph J. Panos,* Jeffrey S. Rubin,‡ Karl G. Csaky,‡ Stuart A. Aaronson,‡ and Robert J. Mason§

*Pulmonary Division, Department of Medicine, Northwestern University Medical School, Chicago, Illinois 60611; ‡Laboratory of Cellular and Molecular Biology, National Cancer Institute, Bethesda, Maryland 20892; and §Pulmonary Division, National Jewish Center for Immunology and Respiratory Medicine, University of Colorado Health Sciences Center, Denver, Colorado 80206.