

Indian hedgehog couples chondrogenesis to osteogenesis in endochondral bone development

Ung-il Chung, Ernestina Schipani, Andrew P. McMahon, Henry M. Kronenberg

J Clin Invest. 2001;107(3):295-304. <https://doi.org/10.1172/JCI11706>.

Article

Vertebrate skeletogenesis requires a well-coordinated transition from chondrogenesis to osteogenesis. Hypertrophic chondrocytes in the growth plate play a pivotal role in this transition. Parathyroid hormone-related peptide (PTHrP), synthesized in the periarticular growth plate, regulates the site at which hypertrophy occurs. By comparing *PTH/PTHrP receptor*^{-/-}/wild-type (*PPR*^{-/-}/wild-type) chimeric mice with *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric and *Ihh*^{-/-}/wild-type chimeric mice, we provide *in vivo* evidence that *Indian hedgehog* (*Ihh*), synthesized by prehypertrophic and hypertrophic chondrocytes, regulates the site of hypertrophic differentiation by signaling to the periarticular growth plate and also determines the site of bone collar formation in the adjacent perichondrium. By providing crucial local signals from prehypertrophic and hypertrophic chondrocytes to both chondrocytes and preosteoblasts, *Ihh* couples chondrogenesis to osteogenesis in endochondral bone development.

Find the latest version:

<https://jci.me/11706/pdf>



Indian hedgehog couples chondrogenesis to osteogenesis in endochondral bone development

Ung-il Chung,¹ Ernestina Schipani,¹ Andrew P. McMahon,² and Henry M. Kronenberg¹

¹Endocrine Unit, Massachusetts General Hospital, Boston, Massachusetts, USA

²Department of Molecular and Cellular Biology, Harvard University, Cambridge, Massachusetts, USA

Address correspondence to: Henry Kronenberg, Endocrine Unit, Massachusetts General Hospital, 50 Blossom Street, Wellman 503, Boston, Massachusetts 02114, USA.

Phone: (617) 726-3966; Fax: (617) 726-7543; E-mail: kronenberg.henry@mgh.harvard.edu.

Received for publication November 3, 2000, and accepted in revised form December 27, 2000.

Vertebrate skeletogenesis requires a well-coordinated transition from chondrogenesis to osteogenesis. Hypertrophic chondrocytes in the growth plate play a pivotal role in this transition. Parathyroid hormone-related peptide (PTHrP), synthesized in the periartricular growth plate, regulates the site at which hypertrophy occurs. By comparing *PTH/PTHrP receptor*^{-/-}/wild-type (*PPR*^{-/-}/wild-type) chimeric mice with *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric and *Ihh*^{-/-}/wild-type chimeric mice, we provide *in vivo* evidence that *Indian hedgehog* (*Ihh*), synthesized by prehypertrophic and hypertrophic chondrocytes, regulates the site of hypertrophic differentiation by signaling to the periartricular growth plate and also determines the site of bone collar formation in the adjacent perichondrium. By providing crucial local signals from prehypertrophic and hypertrophic chondrocytes to both chondrocytes and preosteoblasts, *Ihh* couples chondrogenesis to osteogenesis in endochondral bone development.

J. Clin. Invest. **107**:295–304 (2001).

Introduction

The skeleton of the developing vertebrate limb forms through endochondral bone development. This process is characterized by replacement of a cartilage mold by bone, during which chondrogenesis and osteogenesis are tightly coordinated in time and space (1, 2). During chondrogenesis, growth plate chondrocytes proliferate in orderly columns, then stop proliferating and become prehypertrophic chondrocytes. These chondrocytes then hypertrophy, produce a characteristic matrix, and die. Hypertrophic chondrocytes in the growth plate play a pivotal role in coordinating chondrogenesis and osteogenesis, as hypertrophic chondrocytes provide a scaffold for subsequent formation of trabecular bone by mineralizing their surrounding matrix, and they may induce a bone collar, the precursor of cortical bone, in the adjacent perichondrium (3).

The switch from a proliferative to a postproliferative state determines the number of chondrocytes in the proliferative versus hypertrophic pools. Parathyroid hormone-related peptide (PTHrP) is involved in this switch. In the growth plate, PTHrP mRNA is expressed by perichondrial cells and proliferating chondrocytes in the periartricular region, whereas the mRNA for its receptor, the *PTH/PTHrP receptor* (*PPR*), is expressed at low levels by proliferating chondrocytes in columns and at higher levels by prehypertrophic chondrocytes (4). Mice homozygous for a null mutation in either the *PTHrP* gene or the *PPR* gene show similar abnormalities in the growth plate; proliferation stops prematurely, yielding a short, irregular layer of proliferating chondrocytes (5–7). A similar phenotype has been reported in human fetuses that carry a null mutation

in the *PPR* gene (Blomstrand chondrodysplasia) (8–11). In contrast, increase in *PTHrP* signaling produces an opposite phenotype, with the lengthening of columns of proliferating chondrocytes (12–14). Together, these results suggest that *PTHrP* regulates the switch from a proliferative to a postproliferative state.

Ihh is one of the vertebrate homologs of the *Drosophila* segment polarity gene, *Hedgehog* (*Hb*) (15–17). *Ihh* mRNA is expressed in prehypertrophic and hypertrophic chondrocytes of the mouse embryo (18). Overexpression of *Ihh* protein through injection of a recombinant retrovirus into embryonic chicken limbs delays hypertrophy of growth plate chondrocytes, as does addition of an active NH₂-terminal fragment of Sonic hedgehog (Shh, a relative of *Ihh* known to mimic *Ihh* actions) to embryonic mouse limbs *in vitro* (6, 19). These gain-of-function phenotypes are associated with an increase in *PTHrP* mRNA expression in the perichondrial region at the ends of long bones. In contrast, the Shh fragment has no effect on either *PTHrP*^{-/-} or *PPR*^{-/-} mouse limbs (6, 19). Further, mice homozygous for a null-mutation in the *Ihh* gene have no detectable *PTHrP* mRNA in their growth plate, and hypertrophic chondrocytes predominate in the *Ihh*^{-/-} growth plate late in fetal development (20). These data suggest that *Ihh* delays the switch from proliferation to hypertrophy of chondrocytes by stimulating *PTHrP* production in the periartricular growth plate. *Ihh* also stimulates chondrocyte proliferation by a mechanism that is independent of *PTHrP* (21).

Studies of the growth plates of chimeric mice illustrate how chondrocyte proliferation and further differentiation are coordinated. *PPR*^{-/-} ES cells were inject-

ed into wild-type blastocysts, which then were developed further in foster mothers. In the growth plate of resultant *PPR*^{-/-}/wild-type chimeric mice, *PPR*^{-/-} chondrocytes ectopically hypertrophy closer to the articular surface, while the switch from proliferation to hypertrophy of wild-type chondrocytes is delayed, leading to abnormally long columns of wild-type proliferating chondrocytes (3). Ectopically hypertrophied *PPR*^{-/-} chondrocytes in *PPR*^{-/-}/wild-type chimeric mice produce Ihh much closer to the periarticular growth plate than wild-type cells, and PTHrP mRNA expression in the periarticular growth plate is upregulated. These data suggest that prehypertrophic and hypertrophic chondrocytes may signal their position through Ihh to the periarticular growth plate, which then responds by acting through PTHrP on columnar proliferating chondrocytes to delay their switch to hypertrophy. However, as prehypertrophic and hypertrophic chondrocytes also express other factors such as bone morphogenetic proteins (Bmps) and Wnts, it is still unclear whether Ihh provides the crucial positional signal.

In addition to its actions on chondrocytes, *Ihh* may control osteogenesis as well. Normally, bone collars are formed in the perichondrium abutting prehypertrophic and hypertrophic chondrocytes. In *PPR*^{-/-}/wild-type chimeric mice, ectopic bone collars are formed in the perichondrium adjacent to ectopically hypertrophied *PPR*^{-/-} chondrocytes (3). *Ihh*^{-/-} mice lack bone collars in their dramatically shortened, deformed limbs (20). When chicken limbs are infected with an Ihh-producing retrovirus, bone collars are induced, despite suppression of hypertrophy of chondrocytes (19). Further, Ihh stimulates osteogenic differentiation of mesenchymal cell lines in vitro, and Shh induces ectopic bone formation in vivo (22, 23). These data suggest that Ihh may be involved in bone collar formation. However, it is still unclear whether Ihh, expressed locally in physiological amounts, determines the site of bone collar formation.

We have hypothesized that Ihh synthesized by prehypertrophic and hypertrophic chondrocytes may be responsible both for signaling the relative position of these cells to the periarticular growth plate and for locally signaling the induction of bone collar formation in the adjacent perichondrium. To test this hypothesis in vivo, we have isolated embryonic stem (ES) cell lines homozygous for a null mutation in both the *PPR* and *Ihh* genes, and generated chimeric mice containing both wild-type and *Ihh*^{-/-}; *PPR*^{-/-} cells. Comparison of *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeras with *PPR*^{-/-}/wild-type chimeras has revealed that, in the absence of *Ihh*, ectopic prehypertrophic/hypertrophic chondrocytes form, but no longer induce ectopic bone collars in the adjacent perichondrium, and their presence no longer induces elongation of the layer of wild-type columnar proliferating chondrocytes nor upregulates PTHrP mRNA expression in the periarticular growth plate. Furthermore, chimeric mice containing both wild-type and *Ihh*^{-/-} cells were also generated. In these mice, bone collar formation occurs in the peri-

chondrium adjacent to clusters of wild-type prehypertrophic/hypertrophic chondrocytes, whereas bone collar formation does not occur in the perichondrium adjacent to clusters of *Ihh*^{-/-} prehypertrophic/hypertrophic chondrocytes. These data suggest that Ihh couples cartilage and bone formation by determining the site at which chondrocytes stop proliferating and further differentiate as well as by locally coordinating this process with the differentiation of adjacent osteoblasts.

Methods

Generation of ES cell lines de novo. To generate ES cell lines homozygous for the *PTH/PTHrP receptor*-null mutation (*PPR*^{-/-}), mice heterozygous for the *PTH/PTHrP receptor*-null mutation (*PPR*^{+/-}) in a C57BL/6-129/SvJ background (6) were mated with each other, and blastocysts were collected. One out of 4 of these blastocysts was expected to be *PPR*^{-/-}. To generate ES cell lines homozygous for the *Ihh*-null mutation (*Ihh*^{-/-}) as well as *PPR*^{-/-}, mice heterozygous for the *Ihh*-null mutation (*Ihh*^{+/-}) in a 129/SvJ background (20) were mated with *PPR*^{+/-} mice in a C57BL6-129/SvJ background. Among pups, *Ihh*^{+/-}; *PPR*^{+/-} mice were selected by Southern blot analysis. Then, these *Ihh*^{+/-}; *PPR*^{+/-} mice were mated with each other, and blastocysts were collected. One of 16 of these blastocysts was expected to be *Ihh*^{-/-}; *PPR*^{-/-}. ES cell lines were established from blastocysts as described previously (24). The animal care was in accordance with the policies of the Massachusetts General Hospital.

Generation of chimeric mice. Chimeras were generated by blastocyst injection as described previously (25). *PPR*^{-/-}, *Ihh*^{-/-}; *PPR*^{-/-}, and *Ihh*^{-/-} ES cells were injected into C57BL/6 blastocysts that carry a β -galactosidase transgene as a marker. To introduce a β -galactosidase transgene into host blastocysts, C57BL/6 male mice carrying a β -galactosidase transgene that was engineered to be expressed ubiquitously (26) were mated with wild-type C57BL/6 female mice. One of two blastocysts was expected to carry the transgene. To produce chimeras with various degrees of ES cell contributions, the number of ES cells injected into the blastocyst cavity was varied from five to 15. At least two independently established ES cell lines of each genotype yielded an identical phenotype. All the ES cell lines gave germline transmission. Resultant chimeric mice were studied at E14.5, 15.5, 16.5, 17.5, and 18.5, and mostly the data from d17.5 embryos were presented. All observations in the results were verified in at least five different chimeric mice. The degree of chimerism was estimated by staining for β -galactosidase activity of the skin and the periarticular growth plate. In *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric (and also *Ihh*^{-/-}/wild-type chimeric) mice, the entire length of bone starts becoming shorter than that of wild-type littermates as the degree of contributions from mutant cells goes over approximately 50%. Therefore, when the length of the layer of columnar proliferating chondrocytes in *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric mice was compared with that of wild-type littermates, chimeric mice with less than 50% contributions from mutant cells were used.

Southern blot analysis. Southern blot analysis was performed as described previously (20). To detect disrupted alleles of the *PPR* gene, 10 µl (~10 µg) of genomic DNA were digested with SacI, Southern blotted, and probed with a 1.0-kb SacI-KpnI fragment containing a part of the first intron. The sizes of genomic DNA fragments expected from the normal and disrupted alleles are 5.6 and 4.9 kb, respectively. To detect disrupted alleles of the *Ihh* gene, 10 µl (~10 µg) of genomic DNA were digested with NcoI, Southern blotted, and probed with a 1.2-kb KpnI-HindIII fragment containing a part of the 5'-end of the gene. The sizes of genomic DNA fragments expected from the normal and disrupted alleles are 7.6 and 8.4 kb, respectively. To detect β-galactosidase transgene alleles, 10 µl (~10 µg) of genomic DNA were digested with StuI, Southern blotted, and probed with a 140-bp fragment containing a 5'-flanking region of the transgene (26). The sizes of DNA fragments expected from the normal and the mutant alleles are ~7 and ~12 kb, respectively.

Histological analysis. Chimeras were sacrificed at various ages, dissected, and fixed in 4% paraformaldehyde/PBS at 4°C for 4 hours. For detection of β-galactosidase activity, tissues were stained with X-Gal (5-bromo-4-chloro-3-indolyl β-D-galactoside) as described previously (27). Subsequently, they were processed, embedded in paraffin, and cut. Sections were stained with hematoxylin and eosin (H&E) or Nuclear Fast Red (Fluka, Neu-Ulm, Switzerland) for morphological study. For detection of mineralization, sections were stained with 1% silver nitrate (von Kossa method); mineral is stained black.

Analysis of BrdU incorporation. Pregnant mice were injected intraperitoneally with 50 µg BrdU per gram body weight 1 hour before sacrifice. Embryos were treated to generate sections as already described here. BrdU was detected by BrdU Staining Kit (Zymed Laboratories Inc., South San Francisco, California, USA) according to the manufacturer's instruction. All the BrdU-positive and -negative nuclei of columnar proliferating chondrocytes were counted. At least three sections were counted for each of two wild-type and two *PPR*^{−/−}/wild-type chimeric littermate embryos at E17.5.

In situ hybridization. Tissues were fixed in 4% paraformaldehyde/PBS overnight at 4°C, processed, embedded in paraffin, and cut. In situ hybridization was performed as described previously (4) by using complementary ³⁵S-labeled riboprobes for mouse *type X collagen* (gift from B. Olsen, Harvard Medical School, Boston, Massachusetts, USA), mouse *Ptc1* (R. Johnson, Stanford University, Stanford, California, USA), mouse *Ihh* (17), mouse *PTHrP* (J. Wysolmerski, Yale University, New Haven, Connecticut, USA), mouse *Id1* (R. Benezra, Memorial Sloan-Kettering, New York, New York, USA), mouse *ostecalcin*, mouse *Cbfα1* (G. Karsenty, Baylor College of Medicine, Houston, Texas, USA), mouse *Bmp2* through 7 (E. Robertson, Harvard University, Cambridge, Massachusetts, USA), human *Follistatin* (Y. Sidis and H. Keutmann, Massachusetts General Hospital, Boston, Massachusetts, USA), mouse

Noggin (R. Harland, University of California at Berkeley, Berkeley, California, USA), mouse *Gremlin* (A. Zuniga, University of Utrecht, Utrecht, The Netherlands), mouse *Dan* (ICRFc100D11145; Resource Center of the German Human Genome Project at the Max-Planck-Institute for Molecular Genetics, Berlin, Germany), and mouse *Chordin* (S. Ang, IGBMC, Strasbourg, France).

Generation of transgenic mice. Mice expressing a constitutively active human *PPR* transgene under the control of the rat *α1 (II) collagen* promoter (*caPPR*) were generated and identified, as described previously (13). To evaluate chondrocyte-specific expression of the transgene, RT-PCR was performed as described previously (13). In brief, total RNA was extracted from sternal cartilage and from calvariae of newborn transgenic mice, and reverse-transcribed into a single-stranded DNA with a reverse primer G2, specific to the human *PPR* gene. PCR was then performed using the same reverse primer G2 and the forward primer 2F. The PCR products were analyzed by Southern blot using the ³²P-labeled oligonucleotides H40 and 2FA, specific to the human *PPR* gene. Chondrocyte-specific expression of the transgene was also confirmed by *in situ* hybridization with an antisense probe for the NH₂-terminal extracellular domain portion of the human *PPR* gene, which is significantly different from that of the mouse *PPR* gene in nucleotide sequence and does not recognize the mouse *PPR* mRNA (28).

To rescue *PTHrP*^{−/−} mice, the *caPPR* transgene was introduced into *PTHrP*^{−/−} mice by appropriate matings, and these mice were mated together to generate *caPPR*⁺; *PTHrP*^{−/−} mice.

Results

Isolation of *Ihh*^{−/−}; *PPR*^{−/−} ES cells and generation of chimeric mice. ES cell lines homozygous for a null-mutation in both the *Ihh* and *PTH/PTHrP receptor* genes were generated de novo from inner cell masses of blastocysts derived from mating *Ihh*^{−/−}; *PPR*^{−/−} mice with each other. Of 20 ES cell lines established, two double homozygous mutants were isolated. In addition, ES cell lines with other genetic combinations such as wild-type, *Ihh*^{+/−}, *Ihh*^{−/−}, *PPR*^{+/−}, and *PPR*^{−/−} cell lines were also isolated. To distinguish visually between cells descended from injected ES cells and cells descended from host blastocysts, one copy of a β-galactosidase transgene engineered to be expressed widely in mouse tissues (26) was introduced into host blastocysts by appropriate matings. All cells in the growth plate from the mice carrying one copy of the β-galactosidase transgene alone exhibited β-galactosidase activity and normal morphology (data not shown). In mice generated by injecting wild-type ES cells into wild-type blastocysts expressing one copy of the β-galactosidase transgene, cells derived from ES cells and cells derived from host blastocysts behaved indistinguishably, as expected (data not shown).

***Ihh*^{−/−}; *PPR*^{−/−} chondrocytes undergo ectopic hypertrophy in the absence of ectopic *Ihh* signaling.** The wild-type fetal growth plate consists of three major layers of chondrocytes; that is, the proliferating, prehypertrophic, and




Figure 1

Ectopic differentiation of growth plate chondrocytes deficient in *PThrP* signaling. (a-c) Sections of the tibiae from d17.5 wild-type (a), *PThrP*^{-/-} (b), and *PPR*^{-/-} (c) embryos were stained with H&E. The growth plates of *PThrP*^{-/-} mice and *PPR*^{-/-} mice have a shorter layer of columnar proliferating chondrocytes. In *PThrP*^{-/-} mice, they represent less than 30% of wild-type (b, bracket), and in *PPR*^{-/-} mice, they are almost completely absent (c, arrowhead). (d-g) H&E staining and *in situ* hybridization with a mouse *type X collagen* antisense probe of the sections of the tibiae from d17.5 *PPR*^{-/-}/wild-type chimera (d and e, respectively) and *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimera (f and g, respectively) embryos. In the absence of *PThrP* signaling, mutant cells ectopically hypertrophy when they move from the layer of periarticular proliferating chondrocytes into the layer of columnar proliferating chondrocytes (arrowheads). Staining for β -galactosidase activity of both chimeric growth plates shows that all the mutant chondrocytes undergo ectopic hypertrophy, whereas wild-type cells do not (data not shown). Horizontal bar = 100 μ m.

hypertrophic layers. The proliferating layer contains two distinct types of chondrocytes. Proliferating chondrocytes close to the articular surface are round, randomly scattered (Figure 1a), and express *PThrP* mRNA. We refer to this population as periarticular proliferating chondrocytes. Proliferating chondrocytes immediately below periarticular proliferating chondrocytes are flat, form longitudinal columns (Figure 1a), and weakly express *PPR* mRNA. We refer to this population as columnar proliferating chondrocytes. As previously shown, the growth plates of *PThrP*^{-/-} mice and *PPR*^{-/-} mice have a shortened layer of proliferating chondrocytes (Figure 1, b and c). Columnar proliferating chondrocytes are mainly affected by loss of *PThrP* signaling (in *PThrP*^{-/-} mice, they represent less than 30% of wild-type, and in *PPR*^{-/-} mice, they are almost completely absent), whereas the periarticular proliferating chondrocytes appear relatively intact (Figure 1, b and c). In the growth plate of chimeric mice containing *PPR*^{-/-} and wild-type cells, *PPR*^{-/-} chondrocytes ectopically adopt the hypertrophic phenotype, as they progress from the layer of periarticular proliferating chondrocytes to the layer of columnar proliferating chondrocytes (Figure 1, d and e). Thus, *PThrP* signaling directly prevents the switch from proliferation to hypertrophy

of columnar proliferating chondrocytes. In the growth plate of chimeric mice containing *Ihh*^{-/-}; *PPR*^{-/-} and wild-type cells, mutant chondrocytes similarly hypertrophy ectopically in the layer of columnar proliferating chondrocytes (Figure 1, f and g). Staining for β -galactosidase activity of both chimeric growth plates shows that all the mutant chondrocytes undergo ectopic hypertrophy, whereas wild-type cells do not (3) (data not shown).

Ihh signals the relative position of prehypertrophic and hypertrophic chondrocytes to the periarticular growth plate. Previous studies have suggested that prehypertrophic and hypertrophic chondrocytes may signal their position to the periarticular growth plate, which then delays the switch from proliferation to hypertrophy of columnar proliferating chondrocytes through *PThrP* signaling. As a result, a certain distance between the periarticular growth plate and the layers of prehypertrophic and hypertrophic chondrocytes is maintained. This distance affects bone length by determining the number of chondrocytes in proliferative versus hypertrophic pools, as hypertrophic chondrocytes contribute to longitudinal bone growth and proliferating chondrocytes serve as a reservoir for future hypertrophic chondrocytes. *Ihh* may signal the relative position of prehypertrophic and hypertrophic chondrocytes to the peri-

ticular growth plate (3). To test this hypothesis in vivo, we have compared *PPR*^{-/-}/wild-type chimeric mice and *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric mice.

In the wild-type growth plate, *Ihh* mRNA is expressed in prehypertrophic and hypertrophic chondrocytes (18) (Figure 2a). In *PPR*^{-/-}/wild-type chimeric mice, mutant ectopic prehypertrophic/hypertrophic chondrocytes also express *Ihh* mRNA (Figure 2b, asterisk), whereas in *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric mice, mutant prehypertrophic/hypertrophic chondrocytes do not express *Ihh* mRNA, as expected (Figure 2c). Patched1 (*Ptc1*) is a receptor for Hedgehog (*Hh*) proteins (29, 30). Because its transcription is upregulated by *Hh* signaling, *Ptc1* mRNA expression is a marker for actions of *Hh* proteins (31). In the wild-type growth plate, *Ptc1* mRNA is expressed most strongly in columnar proliferating chondrocytes close to *Ihh*-expressing prehypertrophic/hypertrophic chondrocytes, with its expression decreasing toward the end of bone (3) (Figure 2d). In the growth plate of *PPR*^{-/-}/wild-type chimeric mice, however, proliferating chondrocytes surrounding ectopic prehypertrophic/hypertrophic chondrocytes overexpress *Ptc1* mRNA, providing evidence for ectopic action of *Ihh* (Figure 2e, bracket). In contrast, in the growth plate of *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric mice, there is no overexpression of *Ptc1* mRNA in proliferating chondrocytes surrounding ectopic prehypertrophic/hypertrophic chondrocytes, and the expression pattern of *Ptc1* mRNA looks similar to that in the wild-type growth plate, except that mutant ectopic hypertrophic chondrocytes do not express *Ptc1* mRNA (Figure 2f). Thus, although

Ihh^{-/-}; *PPR*^{-/-} chondrocytes do ectopically hypertrophy, they do not induce the *Ihh* signaling pathway ectopically, unlike *PPR*^{-/-} chondrocytes.

The growth plate of *PPR*^{-/-}/wild-type chimeric mice exhibits elongation of the layer of wild-type columnar proliferating chondrocytes, compared with the wild-type growth plate (3) (Figure 3, a and b). The proliferation rate of columnar proliferating chondrocytes of the *PPR*^{-/-}/wild-type chimeric growth plate is not significantly different from that of columnar proliferating chondrocytes of the wild-type growth plate (percentage of BrdU-positive nuclei: 11.0 ± 0.5% vs. 11.8 ± 0.6%, respectively). Given that *Ihh* is ectopically expressed, this ectopic *Ihh* signaling may upregulate PTHrP expression, which, in turn, might delay hypertrophic differentiation of wild-type columnar proliferating chondrocytes. In the growth plate of *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric mice, the length of the layer of wild-type columnar proliferating chondrocytes remains the same as the length of the layer of columnar proliferating chondrocytes in the wild-type growth plate, despite the presence of ectopic prehypertrophic/hypertrophic chondrocytes (Figure 3, c and d). Thus, ectopically synthesized *Ihh* is required to trigger the lengthening of columns of wild-type proliferating chondrocytes.

In the wild-type growth plate, PTHrP mRNA is expressed weakly in periarticular proliferating chondrocytes and adjacent perichondrial cells at E17.5 day (Figure 3e). If the position of prehypertrophic and hypertrophic chondrocytes is sensed by the periarticular growth plate, which then upregulates PTHrP mRNA expression to slow down the switch from proliferation

Figure 2

Ihh signaling by ectopically hypertrophied *PPR*^{-/-} and *Ihh*^{-/-}; *PPR*^{-/-} chondrocytes. (a–c) *In situ* hybridization of the sections of the tibiae from d17.5 embryos with a mouse *Ihh* antisense probe. In the wild-type growth plate (a), *Ihh* mRNA is expressed in prehypertrophic and hypertrophic chondrocytes. In the chimeric growth plates, ectopically differentiated *PPR*^{-/-} chondrocytes also express *Ihh* mRNA (b, asterisk), whereas ectopically differentiated *Ihh*^{-/-}; *PPR*^{-/-} cells do not (c). (d–f) *In situ* hybridization of the sections of the tibiae from d17.5 embryos with a mouse *Ptc1* antisense probe. *Ptc1* is a transcriptional target of *Ihh* signaling. In the wild-type growth plate (d), *Ptc1* mRNA is expressed most strongly in columnar proliferating chondrocytes adjacent to prehypertrophic chondrocytes with the expression decreasing toward the end of bone. *Ptc1* mRNA is also expressed in the perichondrium and the primary spongiosa. In the *PPR*^{-/-}/wild-type chimeric growth plate (e), periarticular and columnar proliferating chondrocytes surrounding ectopic prehypertrophic/hypertrophic chondrocytes overexpress *Ptc1* mRNA (bracket), reflecting ectopic actions of *Ihh*. In contrast, in the *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric growth plate (f), there is no ectopic expression of *Ptc1* mRNA despite the presence of ectopic prehypertrophic/hypertrophic chondrocytes. Left, bright field; right, dark field. Horizontal bar = 100 μ m.





Figure 3

to hypertrophy of columnar proliferating chondrocytes, PTHrP mRNA expression is expected to increase when prehypertrophic and hypertrophic chondrocytes are placed closer than normal to the periarticular growth plate. In the growth plate of *PPR*^{-/-} mice, there are almost no columnar proliferating chondrocytes, and therefore the layer of periarticular proliferating chondrocytes is placed immediately next to the layers of prehypertrophic and hypertrophic chondrocytes. Consistent with the hypothesis just described, PTHrP mRNA expression is strongly upregulated in the *PPR*^{-/-} periarticular growth plate (Figure 3f, bracket). In the growth plate of *PPR*^{-/-}/wild-type chimeric mice, ectopic prehypertrophic/hypertrophic chondrocytes are also placed next to the layer of periarticular proliferating chondrocytes, but fewer in number compared to the *PPR*^{-/-} growth plate. Therefore, PTHrP mRNA expression is expected to increase, but to a lesser extent than in the *PPR*^{-/-} growth plate. PTHrP mRNA in the growth plate of *PPR*^{-/-}/wild-type chimeric mice confirms this expectation (Figure 3g, bracket). In contrast, in the growth plate of *Ihh*^{-/-};*PPR*^{-/-}/wild-type chimeric mice, the presence of ectopic prehypertrophic/hypertrophic chondrocytes does not lead to upregulation of PTHrP mRNA expression in the periarticular growth plate (Figure 3h).

Taken together, these data strongly suggest that Ihh, synthesized by prehypertrophic and hypertrophic chondrocytes, signals the relative position of these cells to the periarticular growth plate, which then responds by upregulating PTHrP mRNA expression to delay the switch to hypertrophy of columnar proliferating chondrocytes.

Ihh determines the location of bone collar formation. Bone collars are precursors of the cortical regions of long bones. Normally, bone collars form in the perichondrium abutting prehypertrophic and hypertrophic chondrocytes (Figure 4a, arrowhead). Bone collars in *PTHrP*^{-/-} mice and *PPR*^{-/-} mice are also formed in the perichondrium abutting prehypertrophic and hypertrophic chondrocytes, despite much shorter layers of columnar proliferating chondrocytes (5, 6). As reported previously (3), in the growth plate of *PPR*^{-/-}/wild-type chimeric mice, ectopic bone collars are formed near clusters of ectopic prehypertrophic/hypertrophic chondrocytes (Figure 4b, brackets). These ectopic bone collars express typical markers for osteoblasts, such as type I collagen, alkaline phosphatase, *Cbfa1*, bone sialoprotein (data not shown), osteopontin (Figure 5b, bracket), and osteocalcin (Figure 5e, bracket) mRNAs. These results suggest that hypertrophy of chondrocytes and differentiation of osteoblasts are coupled. To demonstrate that prehypertrophic and

hypertrophic chondrocytes induce bone collar formation, we took advantage of the rescue of the *PTHrP*^{-/-} mice by a *constitutively active PPR* (*caPPR*) transgene (13). Ribs, with long arrays of hypertrophic chondrocytes and associated bone collar, were examined to determine the effect of manipulating hypertrophic differentiation of chondrocytes. Expression of type X collagen mRNA (Figure 4, e–g), and mineralization (Figure 4, h–j) were analyzed. Normally, rib cartilage close to the sternum does not hypertrophy prenatally, and no bone collar forms (Figure 4, e and h). In *PTHrP*^{-/-} mice, however, chondrocytes in the same region prematurely hypertrophy and at the same time acquire bone collars in the adjacent perichondrium (Figure 4, f and i). When a *caPPR* gene, which is engineered to be expressed only in chondrocytes under the control of *type II collagen* promoter (13), is introduced by appropriate matings into *PTHrP*^{-/-} mice, ectopic hypertrophy of rib cartilage is reversed, in association with the disappearance of ectopic bone collars (Figure 4, g and j). Similar findings are observed in the tibia (data

not shown). Because the transgene is expressed in chondrocytes, but not in osteoblasts (as assessed by PCR in these mice, data not shown), these data strongly suggest that prehypertrophic and hypertrophic chondrocytes induce bone collars in the adjacent perichondrium.

As noted earlier here, Ihh may provide the local signal whereby prehypertrophic and hypertrophic chondrocytes induce bone collar formation. In the growth plate of *Ihh*^{-/-}; *PPR*^{-/-}/wild-type chimeric mice, although mutant cells ectopically hypertrophy, they fail to induce ectopic bone collars in the adjacent perichondrium (Figure 4, c and d), and no ectopic expression of osteopontin and osteocalcin mRNAs was observed in this region (Figure 5, c and f). Furthermore, chimeric mice containing wild-type and *Ihh*^{-/-} cells were generated. In the growth plate of *Ihh*^{-/-}/wild-type chimeric mice, mutant cells do not ectopically hypertrophy, but instead hypertrophy in synchrony with wild-type cells. The architecture of the growth plate of *Ihh*^{-/-}/wild-type chimeric mice is well preserved despite its smaller size, probably due to Ihh secret-

Figure 4





Figure 5

Expression of osteoblastic markers in the chimeric growth plates. (a–c) *In situ* hybridization of sections of the tibiae from d17.5 embryos with a mouse *osteopontin* antisense probe. In the wild-type growth plate (a), osteopontin mRNA is expressed in osteoblasts as well as in mature chondrocytes, and its expression pattern overlaps the locations of bone collars and cartilaginous mineralization. In the *PPR*^{−/−}/wild-type chimera growth plate (b), there is ectopic expression of osteopontin mRNA in ectopic bone collars (bracket). Ectopic hypertrophic chondrocytes also express osteopontin mRNA (asterisk). In contrast, in the *Ihh*^{−/−};*PTH/PPR*^{−/−}/wild-type chimera (c), there is no ectopic expression of osteopontin mRNA in the perichondrium, whereas ectopic hypertrophic chondrocytes still express osteopontin mRNA (asterisk). Arrowheads denote where the ectopic prehypertrophic layer starts. (d–f) *In situ* hybridization of sections of the tibiae from d17.5 embryos with a mouse *osteocalcin* antisense probe. In the wild-type growth plate (d), osteocalcin mRNA is expressed in mature osteoblasts. In the *PPR*^{−/−}/wild-type chimera (e), there is ectopic expression of osteocalcin mRNA in ectopic bone collars (bracket), whereas in the *Ihh*^{−/−};*PPR*^{−/−}/wild-type chimera (f), there is no ectopic expression of osteocalcin mRNA in the perichondrium. Arrowheads denote where the ectopic prehypertrophic layer starts. Horizontal bar = 100 μ m.



ed from wild-type cells. In these mice, bone collars do not form in the perichondrium adjacent to clusters of *Ihh*^{−/−} prehypertrophic/hypertrophic chondrocytes (Figure 4k), whereas bone collars do form in the perichondrium adjacent to clusters of wild-type prehypertrophic/hypertrophic chondrocytes (Figure 4l, brackets). These findings provide *in vivo* evidence that Ihh, secreted by prehypertrophic and hypertrophic chondrocytes, is locally required for induction of bone collar formation.

Bmp signaling alone is not sufficient for induction of bone collar formation. Because *decapentaplegic* (*Dpp*), a homolog of *Bmps*, works downstream of *Hh* signaling in *Drosophila* in some settings (32), and *Bmps* have been implicated in bone formation (33), it is possible that ectopic Ihh action in the growth plate of *PPR*^{−/−}/wild-type chimeric mice may work through *Bmp* signaling. Expression patterns of *Bmp2*, *4*, *5*, *6*, and *7* mRNAs were analyzed by *in situ* hybridization. In the growth plate of *PPR*^{−/−}/wild-type chimeric mice, expression patterns of *Bmp 4*, *5*, and *7* mRNAs do not change significantly, compared with those of wild-type mice (data not shown), whereas ectopic hypertrophic chondrocytes express *Bmp2* and *Bmp6* mRNAs (Figure 6, b and e, respectively), just as wild-type hypertrophic chondrocytes do (Figure 6, a and d, respectively). In the growth plate of *Ihh*^{−/−};*PPR*^{−/−}/wild-type chimeric mice, ectopic hypertrophic chondrocytes also express *Bmp2* and *Bmp6* mRNAs (Figure 6, c and f, respectively), but they do not induce ectopic bone collar formation in the adjacent perichondrium. There is no significant difference in mRNA expression patterns of *Bmp* inhibitors such as *Noggin*, *Gremlin*, *Dan*, *Chordin*, and *Follistatin* between the growth plates of *Ihh*^{−/−};*PPR*^{−/−}/wild-


type chimera and *PPR*^{−/−}/wild-type chimera mice (data not shown). Further, mRNA expression of *Id1*, one of the transcriptional target genes of *Bmp* signaling, is similarly upregulated in the growth plate of both chimeras (data not shown). These data suggest that *Bmp* signaling does not mediate the actions of Ihh on the periarticular growth plate. They also suggest that *Bmp* signaling alone cannot account for induction of bone collar formation in the perichondrium by adjacent prehypertrophic/hypertrophic chondrocytes.

Discussion

Ihh couples chondrogenesis and osteogenesis in endochondral bone development. Previous data suggested that Ihh and PTHrP participate in a negative feedback loop that regulates the rate of chondrocyte differentiation. However, the model was based on the experiments involving supraphysiological amounts of PTHrP and Ihh. The current study uses a chimeric system to demonstrate the role of Ihh in chondrogenesis in a more physiological setting. By comparing chimeric growth plates that differ genetically only by the presence or absence of the *Ihh* gene, we define the mechanism used by ectopic prehypertrophic/hypertrophic cells to extend the columns of normal proliferating chondrocytes. In the *PPR*^{−/−}/wild-type chimeric growth plates, but not in the *Ihh*^{−/−};*PPR*^{−/−}/wild-type chimeric growth plates, ectopic Ihh regulates the amount of PTHrP synthesized in the periarticular region. When Ihh is synthesized nearer to the articular end of the growth plate, PTHrP synthesis is increased. In this way, Ihh signals the relative position of prehypertrophic and hypertrophic chondrocytes to the periartic-

Figure 6

Bmp signaling alone is not sufficient for induction of bone collars. (a–c) *In situ* hybridization of sections of the tibiae from d17.5 embryos with a mouse *Bmp2* antisense probe. In the wild-type growth plate (a), *Bmp2* mRNA is expressed in hypertrophic chondrocytes and the perichondrium, whereas both in the *PPR*^{−/−}/wild-type chimeric growth plate (b) and in the *Ihh*^{−/−}; *PPR*^{−/−}/wild-type chimeric growth plate (c), ectopic hypertrophic mutant cells also express *Bmp2* mRNA (arrowheads). (d–f) *In situ* hybridization of sections of the tibiae from d17.5 embryos with a mouse *Bmp6* antisense probe. In the wild-type growth plate (d), *Bmp6* mRNA is expressed in hypertrophic chondrocytes, whereas both in the *PPR*^{−/−}/wild-type chimeric growth plate (e) and in the *Ihh*^{−/−}; *PPR*^{−/−}/wild-type chimeric growth plate (f), ectopic hypertrophic mutant cells also express *Bmp6* mRNA (arrowheads). Horizontal bar = 100 μ m.



ular growth plate. PTHrP then delays the transition from proliferation to hypertrophy of growth plate chondrocytes. Thus, through PTHrP, Ihh controls the site at which hypertrophy occurs (Figure 7).


Ihh also determines the site of formation of the bone collar, which will become cortical bone. Previous work showed that *Ihh*^{−/−} mice had no bone collar and that overexpression of Ihh by retroviral infection induced bone collar formation. However, it was unclear whether Ihh is locally required for this process *in vivo*, as the growth plates of *Ihh*^{−/−} mice had a dramatically altered structure, and overexpression by retroviral infection involved supraphysiological amounts of Ihh. The current study shows that Ihh synthesized by prehypertrophic and hypertrophic chondrocytes is locally required for the induction of mature osteoblasts in the

adjacent perichondrium in the growth plate with relatively normal architecture. The current study also shows that Bmp2 and Bmp6 synthesized by the same cells are not sufficient for this induction. Our *in vivo* data suggest that Ihh may induce differentiation of mature osteoblasts in the adjacent perichondrium, perhaps in synergy with Bmp2 and Bmp6 (Figure 7).

It is noteworthy that Ihh converts only a select group of cells into mature osteoblasts. Perichondrial cells that will eventually become mature osteoblasts upon exposure to Ihh are a distinct population of cells, in that they express low levels of alkaline phosphatase, type I collagen and *Cbfa1* mRNAs (4) (data not shown). Therefore, they are probably already committed to some extent toward the osteoblastic lineage and probably are susceptible to the induction of full maturation

Figure 7

Interactions of Ihh and PTHrP signaling pathways in the fetal growth plate. This figure shows a schema of the fetal growth plate of the mouse long bone. The end of the bone is at the top. As proliferating chondrocytes differentiate into prehypertrophic chondrocytes and then hypertrophic chondrocytes, they express Ihh. Ihh directly or indirectly stimulates PTHrP synthesis in the periarticular growth plate including the perichondrium and periarticular proliferating chondrocytes (long solid arrow). In this way, Ihh signals the relative position of prehypertrophic and hypertrophic chondrocytes to the periarticular growth plate. Ihh also has a positive effect on chondrocyte proliferation (short solid arrow). PTHrP then acts directly on columnar proliferating chondrocytes to delay their differentiation into prehypertrophic and hypertrophic chondrocytes, which synthesize Ihh (solid t-bar). Thus, through PTHrP, Ihh controls the site at which hypertrophy occurs. Ihh also acts on perichondrial cells to induce mature osteoblasts, which form a bone collar (white arrow).



by *Hh* signaling. Further study is required to clarify what signaling pathway initially commits perichondrial cells to the osteoblast lineage.

Involvement of Hh signaling in other modes of bone development. Although *Ihh* is required for the differentiation of bone collar osteoblasts, the role of *Ihh* in the differentiation of osteoblasts in other settings is less certain. These settings include endochondral osteoblasts of the primary spongiosa, which lay down a bone matrix on top of a cartilaginous matrix, and osteoblasts of intramembranous bones, such as the bones of the vault of the skull. The primary spongiosa expresses *Ptc1* mRNA, which is a marker for *Hh* signaling. The source of *Hh* protein for this signaling might be *Ihh* secreted by prehypertrophic and hypertrophic chondrocytes. *Hh* proteins usually do not travel very far owing to the sequestering action of *Ptc1* in responding cells. However, because hypertrophic chondrocytes do not express *Ptc1*, *Ihh* may be able to reach the primary spongiosa. *Ihh*^{-/-} mice do have vascular invasion in the center of mineralized cartilage of the humerus, although osteoblasts with their associated matrix are not clearly present. However, mRNAs for proteins such as type I collagen are detectable, suggesting that early osteoblasts may be present. The severe abnormality in cartilage development and neonatal lethality of *Ihh*^{-/-} mice make it difficult to use these mice to determine the possible role of *Ihh* in the primary spongiosa.

Intramembranous bones such as the parietal bone do form in *Ihh*^{-/-} mice. Although they are not normal, they do contain osteoblasts capable of ossification, presumably through an *Ihh*-independent pathway. Either another *Hh* molecule (*Shh* or *Dhh*) is active in these bones or, alternatively, osteoblasts in these bones do not require *Hh* signaling. From this perspective, the role of *Ihh* in bone collar formation emphasizes a special situation in which chondrocyte differentiation and osteoblast differentiation may need careful coordination. *Ihh* provides this coordination.

Acknowledgments

We thank E. Li for technical support in ES cell isolation, and the members of Endocrine Unit, Massachusetts General Hospital, for helpful discussion. This work was funded by a grant (to H. Kronenberg and A.P. McMahon) from NIH (P01 DK56246). U. Chung at the time of this research was a Medical Foundation and Charles A. King Trust Fellow.

1. Erlebacher, A., Filvaroff, E.H., Gitelman, S.E., and Deryck, R. 1995. Toward a molecular understanding of skeletal development. *Cell*. **80**:371-378.
2. Marks, S.C., and Hersey, D.C. 1996. The structure and development of bone. In *Principles of bone biology*. J.P. Bilezikian, L.G. Raisz, and G.A. Rodan, editors. Academic Press. San Diego, California, USA. 3-14.
3. Chung, U.I., Lanske, B., Lee, K., Li, E., and Kronenberg, H. 1998. The parathyroid hormone/parathyroid hormone-related peptide receptor coordinates endochondral bone development by directly controlling chondrocyte differentiation. *Proc. Natl. Acad. Sci. USA*. **95**:13030-13035.
4. Lee, K., Deeds, J.D., and Segre, G.V. 1995. Expression of parathyroid hormone-related peptide and its receptor messenger ribonucleic acids during fetal development of rats. *Endocrinology*. **136**:453-463.
5. Karaplis, A.C., et al. 1994. Lethal skeletal dysplasia from targeted disruption of the parathyroid hormone-related peptide gene. *Genes Dev.* **8**:277-289.
6. Lanske, B., et al. 1996. PTH/PTHrP receptor in early development and Indian hedgehog-regulated bone growth. *Science*. **273**:663-666.
7. Amizuka, N., Warshawsky, H., Henderson, J.E., Goltzman, D., and Karaplis, A.C. 1994. Parathyroid hormone-related peptide-depleted mice show abnormal epiphyseal cartilage development and altered endochondral bone formation. *J. Cell Biol.* **126**:1611-1623.
8. Jobert, A.S., et al. 1998. Absence of functional receptors for parathyroid hormone and parathyroid hormone-related peptide in Blomstrand chondrodyplasia. *J. Clin. Invest.* **102**:34-40.
9. Zhang, P., Jobert, A.S., Couvineau, A., and Silve, C. 1998. A homozygous inactivating mutation in the parathyroid hormone/parathyroid hormone-related peptide receptor causing Blomstrand chondrodyplasia. *J. Clin. Endocrinol. Metab.* **83**:3365-3368.
10. Karaplis, A.C., et al. 1998. Inactivating mutation in the human parathyroid hormone receptor type 1 gene in Blomstrand chondrodyplasia. *Endocrinology*. **139**:5255-5258.
11. Karperien, M., et al. 1999. A frame-shift mutation in the type I parathyroid hormone (PTH)/PTH-related peptide receptor causing Blomstrand lethal osteochondrodyplasia. *J. Clin. Endocrinol. Metab.* **84**:3713-3720.
12. Weir, E.C., et al. 1996. Targeted overexpression of parathyroid hormone-related peptide in chondrocytes causes chondrodyplasia and delayed endochondral bone formation. *Proc. Natl. Acad. Sci. USA*. **93**:10240-10245.
13. Schipani, E., et al. 1997. Targeted expression of constitutively active receptors for parathyroid hormone and parathyroid hormone-related peptide delays endochondral bone formation and rescues mice that lack parathyroid hormone-related peptide. *Proc. Natl. Acad. Sci. USA*. **94**:13689-13694.
14. Schipani, E., Kruse, K., and Juppner, H. 1995. A constitutively active mutant PTH/PTHrP receptor in Jansen-type metaphyseal chondrodyplasia. *Science*. **268**:98-100.
15. Krauss, S., Concorde, J.P., and Ingham, P.W. 1993. A functionally conserved homolog of the *Drosophila* segment polarity gene *hh* is expressed in tissues with polarizing activity in zebrafish embryos. *Cell*. **75**:1431-1444.
16. Riddle, R.D., Johnson, R.L., Laufer, E., and Tabin, C. 1993. Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell*. **75**:1401-1416.
17. Echelard, Y., et al. 1993. Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell*. **75**:1417-1430.
18. Bitgood, M.J., and McMahon, A.P. 1995. Hedgehog and Bmp genes are coexpressed at many diverse sites of cell-cell interaction in the mouse embryo. *Dev. Biol.* **172**:126-138.
19. Vortkamp, A., et al. 1996. Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science*. **273**:613-622.
20. St-Jacques, B., Hammerschmidt, M., and McMahon, A.P. 1999. Indian hedgehog signaling regulates proliferation and differentiation of chondrocytes and is essential for bone formation. *Genes Dev.* **13**:2072-2086.
21. Karp, S.J., et al. 2000. Indian hedgehog coordinates endochondral bone growth and morphogenesis via parathyroid hormone-related protein-dependent and -independent pathways. *Development*. **127**:543-548.
22. Kinto, N., et al. 1997. Fibroblasts expressing Sonic hedgehog induce osteoblast differentiation and ectopic bone formation. *FEBS Lett.* **404**:319-323.
23. Nakamura, T., et al. 1997. Induction of osteogenic differentiation by hedgehog proteins. *Biochem. Biophys. Res. Commun.* **237**:465-469.
24. Robertson, E.J. 1987. Embryo-derived stem cell lines. In *Teratocarcinoma and embryonic stem cells*. E.J. Robertson, editor. IRL Press. Oxford, United Kingdom. 71-112.
25. Bradley, A. 1987. Production and analysis of chimaeric mice. In *Teratocarcinoma and embryonic stem cells*. E.J. Robertson, editor. IRL Press. Oxford, United Kingdom. 113-152.
26. Zambrowicz, B.P., et al. 1997. Disruption of overlapping transcripts in the ROSA beta geo 26 gene trap strain leads to widespread expression of betagalactosidase in mouse embryos and hematopoietic cells. *Proc. Natl. Acad. Sci. USA*. **94**:3789-3794.
27. Rossert, J., Eberspaecher, H., and de Crombrugghe, B. 1995. Separate cis-acting DNA elements of the mouse pro-alpha 1(I) collagen promoter direct expression of reporter genes to different type I collagen-producing cells in transgenic mice. *J. Cell Biol.* **129**:1421-1432.
28. Schipani, E., et al. 1993. Identical complementary deoxyribonucleic acids encode a human renal and bone parathyroid hormone (PTH)/PTH-related peptide receptor. *Endocrinology*. **132**:2157-2165.
29. Stone, D.M., et al. 1996. The tumour-suppressor gene patched encodes a candidate receptor for Sonic hedgehog. *Nature*. **384**:129-134.
30. Marigo, V., Davey, R.A., Zuo, Y., Cunningham, J.M., and Tabin, C.J. 1996. Biochemical evidence that patched is the Hedgehog receptor. *Nature*. **384**:176-179.
31. Goodrich, L.V., Johnson, R.L., Milenkovic, L., McMahon, J.A., and Scott, M.P. 1996. Conservation of the hedgehog/patched signaling pathway from flies to mice: induction of a mouse patched gene by Hedgehog. *Genes Dev.* **10**:301-312.
32. Ingham, P.W. 1998. Transducing hedgehog: the story so far. *EMBO J.* **17**:3505-3511.
33. Wozney, J.M., et al. 1988. Novel regulators of bone formation: molecular clones and activities. *Science*. **242**:1528-1534.