JCI The Journal of Clinical Investigation

Effect of glucagon on intracellular pH regulation in isolated rat hepatocyte couplets.

D Alvaro, P Della Guardia, A Bini, A Gigliozzi, S Furfaro, T La Rosa, C Piat, L Capocaccia

J Clin Invest. 1995;96(2):665-675. https://doi.org/10.1172/JCI118109.

Research Article

To elucidate mechanisms of glucagon-induced bicarbonate-rich choleresis, we investigated the effect of glucagon on ion transport processes involved in the regulation of intracellular pH (pHi) in isolated rat hepatocyte couplets. It was found that glucagon (200 nM), without influencing resting pHi, significantly stimulates the CI-/HCO3- exchange activity. The effect of glucagon was associated with a sevenfold increase in cAMP levels in rat hepatocytes. The activity of the CI-/HCO3- exchanger was also stimulated by DBcAMP + forskolin. The effect of glucagon on the CI-/HCO3- exchange was individually blocked by two specific and selective inhibitors of protein kinase A, Rp-cAMPs (10 microM) and H-89 (30 microM), the latter having no influence on the glucagon-induced cAMP accumulation in isolated rat hepatocytes. The CI-channel blocker, NPPB (10 microM), showed no effect on either the basal or the glucagon-stimulated CI-/HCO3- exchange; however, this effect was achieved through a significant inhibition of the glucagon-stimulated cAMP accumulation in rat hepatocytes. Colchicine pretreatment inhibited the basal as well as the glucagon-stimulated CI-/HCO3- exchange activity. The Na+/H+ exchanger was unaffected by glucagon either at basal pHi or at acid pHi values. In contrast, glucagon, at basal pHi, stimulated the Na(+)-HCO3- symport. The main findings of this study indicate that [...]

Find the latest version:



Effect of Glucagon on Intracellular pH Regulation in Isolated Rat Hepatocyte Couplets

Domenico Alvaro, Paola Della Guardia, Adriano Bini, Alessandro Gigliozzi, Sonia Furfaro, Tiziana La Rosa, Claudia Piat, and Livio Capocaccia

II Department of Gastroenterology, University of Rome "La Sapienza," Rome, Italy

Abstract

To elucidate mechanisms of glucagon-induced bicarbonaterich choleresis, we investigated the effect of glucagon on ion transport processes involved in the regulation of intracellular pH (pH_i) in isolated rat hepatocyte couplets. It was found that glucagon (200 nM), without influencing resting pH_i, significantly stimulates the CI-/HCO3 exchange activity. The effect of glucagon was associated with a sevenfold increase in cAMP levels in rat hepatocytes. The activity of the Cl-/HCO3 exchanger was also stimulated by DBcAMP + forskolin. The effect of glucagon on the Cl⁻/HCO₃ exchange was individually blocked by two specific and selective inhibitors of protein kinase A, Rp-cAMPS (10 μ M) and H-89 (30 μ M), the latter having no influence on the glucagon-induced cAMP accumulation in isolated rat hepatocytes. The Cl - channel blocker, NPPB (10 µM), showed no effect on either the basal or the glucagon-stimulated Cl⁻/ HCO₃ exchange. In contrast, the protein kinase C agonist, PMA (10 μ M), completely blocked the glucagon stimulation of the Cl-/HCO3 exchange; however, this effect was achieved through a significant inhibition of the glucagonstimulated cAMP accumulation in rat hepatocytes. Colchicine pretreatment inhibited the basal as well as the glucagon-stimulated Cl⁻/HCO₃ exchange activity. The Na⁺/H⁺ exchanger was unaffected by glucagon either at basal pHi or at acid pH_i values. In contrast, glucagon, at basal pH_i, stimulated the Na+-HCO₃ symport. The main findings of this study indicate that glucagon, through the cAMP-dependent protein kinase A pathway, stimulates the activity of the Cl⁻/HCO₃ exchanger in isolated rat hepatocyte couplets, a mechanism which could account for the in vivo induced bicarbonate-rich choleresis. (J. Clin. Invest. 1995. 96:665-675.) Key words: glucagon • hepatocytes • Cl⁻/ HCO₃ exchanger • cAMP • protein kinase A

Introduction

Glucagon, in the liver, exerts a number of metabolic effects such as inhibition of glycogen synthesis and stimulation of gluconeogenesis and glycogenolysis.

Part of this work was presented at the American Gastroenterological Association Meeting in New Orleans on 15–18 May 1994 and was published in abstract form (1994. *Gastroenterology*. 106:858A).

Address correspondence to Domenico Alvaro, MD, II Department of Gastroenterology, University of Rome "La Sapienza," via Valsolda 45/i, 00141 Rome, Italy. Phone: 39-6-4453319; FAX: 39-6-4440806.

Received for publication 13 September 1994 and accepted in revised form 29 March 1995.

Moreover, glucagon, at physiological doses, induces a bicarbonate-rich choleresis in humans (1) as well as in many different animal species, including the guinea pig (2), dog (3), and rat (4). Mechanisms of glucagon choleresis are still unknown. It has been established recently (2) that glucagon choleresis: (a) is canalicular in origin as it is associated with an increased clearance of erythritol and is not influenced by bile ductule proliferation; (b) is mediated by the intracellular messenger cAMP but not by inositol phosphates nor by prostaglandin release; and (c) requires an intact microtubular system (colchicine sensitivity).

Bile salt-independent bile flow is the result of active solute transport followed by osmotic water flow in bile canaliculi (5). Bicarbonate together with glutathione represents the main solute responsible for this fraction of bile flow (5). The ion transport processes underlying basal or hormonal-stimulated bicarbonate secretion in the hepatocyte are currently under investigation. Recent progress has been made in studies exploring intracellular. pH (pH_i) regulation in isolated hepatocytes (6-8). These studies identify the presence of an electroneutral Na⁺-independent Cl⁻/ HCO₃ exchanger (7, 8) which functions as a counterpoint to the acid extruding systems (Na+/H+ exchanger and Na+-HCO₃ symport) (6) and which has been detected recently in the hepatocyte apical domain of human liver (9). The Cl⁻/ HCO₃ exchanger is present in all the bicarbonate-secreting epithelia (10-19). In biliary epithelium for example, this exchanger, functionally coupled with apically located Cl - channels, is stimulated by secretin, a hormone which induces a ductular bicarbonate-rich choleresis (18, 19). The role of the Cl⁻/HCO₃ exchanger in the generation of bile salt-independent bile flow has been supported recently by studies in the isolated perfused rat liver, where stimulation of this exchanger by intracellular alkalinization is followed by a bicarbonate-rich choleresis (20).

In an attempt to elucidate the mechanisms of glucagon choleresis, this study focuses on the effect of glucagon on the H⁺/HCO₃ transport processes involved in pH_i regulation in isolated rat hepatocyte couplets (IRHC).¹ The main data emerging from this study demonstrate that glucagon, through the cAMP-dependent protein kinase A pathway, stimulates the activity of the Cl⁻/HCO₃ exchanger, a mechanism which could account for bicarbonate-rich choleresis induced by the hormone. Glucagon induces an upregulation of pH_i regulatory mechanisms operating in the presence of bicarbonate since, besides the Cl⁻/HCO₃ exchanger (acid loader), the Na⁺-HCO₃ symporter

J. Clin. Invest.

[©] The American Society for Clinical Investigation, Inc. 0021-9738/95/08/0665/11 \$2.00 Volume 96, August 1995, 665-675

^{1.} Abbreviations used in this paper: BCECF-AM, 2,7-bis(carboxyethyl)-5(6)-carboxyl-fluorescein-acetomethylester; β_i , intrinsic buffering power; DBcAMP, N^6 ,2'-O-dibutyryladenosine-3'-5'-cyclic monophosphate; DIDS, 4,4,diisothiocyano-2,2' disulfonic acid stilbene; H-89, N-[2-(p-bromocinnamyl-amino)ethyl]-5-isoquinolinesulfonamide HCl; IRHC, isolated rat hepatocyte couplets; L-15, Liebowitz-15; NPPB, 5'-nitro-2'-(3-phenylpropyl-amino)-benzoate; Rp-cAMPS, Rp-adenosine-3',5'-monophosphothioate.

(acid extruder) was also stimulated by the hormone. We have also shown that glucagon stimulation of the Cl⁻/HCO₃⁻ exchanger does not involve 5'-nitro-2'-(3-phenylpropyl-amino)-benzoate (NPPB)-sensitive Cl⁻ channels, is partially blocked by the microtubular inhibitor colchicine, and is completely inhibited by the protein kinase C agonist, PMA, which, however, significantly decreases the cAMP accumulation induced by the hormone.

Methods

Materials. Glucagon, collagenase A, bovine serum albumin, EDTA, penicillin/streptomycin, heparin, Hepes, D(+)glucose, insulin, soybean trypsin inhibitor (type I-s), amiloride, DMSO, deoxyribonuclease (DN-25), nigericin, 4,4, diisothiocyano-2,2' disulfonic acid stilbene (DIDS), Na⁺-gluconate, K⁺-gluconate, hemicalcium gluconate, N⁶, 2'-O-dibutyryladenosine-3'-5'-cyclic monophosphate (DBcAMP), forskolin, PMA, and 3-isobutyl-1-methyl-xanthine (IBMX) were purchased from Sigma Chemical Co. (St. Louis, MO). 2,7,bis(carboxyethyl)-5(6)carboxy-fluorescein-acetomethylester (BCECF-AM) was obtained from Molecular Probes, Inc. (Eugene, OR). Liebowitz 15 (L-15), MEM, α-MEM, L-glutamine, gentamicin, and fetal calf serum were obtained from Gibco Laboratories (Grand Island, NY). NPPB was provided by Prof. R. Greger (Freiburg, Germany). Rp-adenosine-3',5'-monophosphothioate (Rp-cAMPS) was purchased from BioLog (Bremen, Germany). N-[2-(p-bromocinnamyl-amino)ethyl]-5-isoquinolinesulfonamide HCl (H-89) was purchased from Calbiochem-Novabiochem (San Diego, CA). cAMP[125I] assay system was purchased from Amersham International (Little Chalfont, United Kingdom).

Isolation of rat hepatocyte couplets. Male Wistar rats (F. Morini, Reggio Emilia, Italy), weighing 240–260 grams, were housed in temperature- and light-controlled rooms and allowed free access to water and laboratory chow. Animals received humane care and the study protocol was in compliance with our institution's ethical guidelines.

IRHC were prepared by a modification (21) of Seglen's procedure (22). The animals were anesthetized with pentobarbital (50 mg/kg intraperitoneally). The portal vein was cannulated by a 16-gauge cannula, and the liver was perfused in situ for 7 min with Ca2+, Mg2+-free Hanks' medium (NaCl 120 mM, KCl 5.0 mM, NaHCO₃ 25 mM, KH₂PO₄ 0.66 mM, Na₂HPO₄ 0.33 mM, 0.1% D-glucose) and then with Ca²⁺, Mg²⁺ containing Hanks' buffer supplemented with 0.05% collagenase A plus 0.8 U trypsin inhibitor/U of collagenase tryptic activity. When the capsula began to detach, the liver was removed from the carcass, grasped at the hilum by forceps, and the hepatocytes were combed into L-15 medium at 0°C. Cell suspension was then sequentially filtered through 80- and 45-μm mesh nylon monofilament screens and centrifuged at 50 g. Supernatant was discarded, and the cells were washed three times with cold L-15 medium. Cell suspension was plated on small cover slip fragments (5 × 5 mm), layered on culture plastic wells (Corning Inc., Corning, NY) incubated at 37°C in an air-equilibrated incubator. In experiments performed in the absence of HCO₃, cells were plated in L-15 medium containing penicillin/streptomycin (100,000 U/100 μ g/liter), L-glutamine (2 mM), FCS (10%), and insulin (0.1 μ M). In experiments performed in KRB, cells were plated in α -MEM medium containing the same additive as L-15, but 25 mM NaHCO₃ was used to replace 25 mM Hepes. Medium was changed after 2 h and experiments were performed between 2 and 6 h after plating.

The yield of our preparation was $1-1.9\times10^7$ hepatocytes per gram of liver, with a viability (Trypan blue exclusion) ranging from 75 to 88% (n=73 cell preparations). Couplets represent $20\pm5\%$ of the cell preparation, with a viability constantly > 90%. Cell viability was not significantly decreased by preincubation with Cl⁻-free solutions, DIDS (0.5 mM), amiloride (1 mM), NPPB $(10 \mu\text{M})$, colchicine or β -lumicolchicine (10^{-5} M) , Rp-cAMPS $(10 \mu\text{M})$, or H-89 $(30 \mu\text{M})$.

 pH_i determination. pH_i of IRHC was measured using a microfluorimetric single-cell method (SPEX-AR-CM-micro system; Spex Industries, Edison, NJ) and BCECF-AM as a fluorescent pH_i indicator (18, 23, 24). Cells on glass coverslips were loaded with BCECF-AM (12 μm) for 10 min, washed for 10 min in a BCECF-free medium, and transferred into a thermostated perfusion chamber placed on the stage of a Nikon Diaphot inverted microscope. The perfusion media were constantly warmed at 37°C and gassed both by direct bubbling and by an artificial lung. For the experiments performed in HCO3-enriched media, gas-impermeable tubing was used. Coverslips were scanned under DIC optics and an isolated couplet with a well developed canalicular pole was selected. Couplets with membrane blebs were avoided. Signal recording was obtained from one of the two hepatocytes constituting the couplet by using a pin hole device (Nikon) inserted in the emission light pathway. The microscope was connected with a rotating chopper mirror able to rapidly alternate the light generated by a 150 W xenon lamp between two excitation beams (490 and 440 nm). Light was attenuated to 0.1% by heat reflectors and neutral density filters (Omega Optics, Eugene, OR). The emitted light was read at 530 nm, integrated by the computer with 500-ms acquisitions every 1,500 ms and displayed in real time. All measurements for each excitation wavelength were corrected for background fluorescence obtained from dye-free cells. Fluorescence intensity exceeded background autofluorescence by at least 30-fold. The 490/440 fluorescence intensity ratio data were converted to pH_i values by using the nigericin (12 μ m) calibration curve technique (6-8, 18, 23-25). Over the pH range of 6.4-7.7, fluorescence varied in a linear fashion with pHout.

Total and intrinsic intracellular buffering power. The intrinsic buffering power (β_i) was determined at different pH_i as described (7, 18, 24, 26-30), by exposing the cells to 25 mM NH₄Cl in Hepes, Na⁺free buffered solutions and then decreasing the NH₄Cl concentration by 5 or 10 mM for each step to 0 mM (n = 10). The β_i was then calculated from the midpoint change in pH_i at each step. β_i values were then plotted versus pH_i using a best-fit program (Enzfitter). β_i/pH_i curves were also evaluated in IRHC perfused with 200 nM glucagon (n = 8). As in other studies (7, 18, 24) a single exponential function provides a better description of the known β_i/pH_i relationships. β_i changed from 10.4 mM/pH U at pH_i 7.5 to 25 mM/pH U at pH_i 7 and to 66 mM/ pH U at pH_i 6.5 and was not significantly influenced by glucagon at each pH_i value. We found values of β_i in IRHC which were very close to those measured in subconfluent monolayer of rat hepatocytes (7). For each experimental protocol, β_i was obtained from the relative β_i pH_i curves as estimated above, for that pH_i at which δ pH_i/ δ t was measured. H⁺ fluxes were then calculated by the formula: H⁺ fluxes = β_i $\times \delta pH_i/min$. In this formula, β_{tot} substituted for β_i when experiments were performed in the presence of bicarbonate. β_{tot} was calculated from the formula $\beta_{tot} = \beta_i + 2.303 \times [HCO_3^-]_i$ where intracellular [HCO₃] is derived from the Henderson-Hasselbach equation.

cAMP levels in isolated rat hepatocytes. cAMP was measured in isolated rat hepatocytes cultured in α -MEM and incubated with glucagon, agonists, or inhibitors under the same conditions as those for pH_i measurement in IRHC. cAMP was extracted from cells and measured by using a kit (Amersham International) composed by AmprepTM SAX minicolumn and cAMP[125 I] assay system and by following instructions included in the kit.

Solutions. The compositions of solutions used in the study have been previously detailed (7, 8, 18, 24). Glucagon, made up as a concentrated solution in the appropriate perifusion buffer containing 1% (wt/vol) bovine serum albumin, was infused (1:50 vol/vol dilution) into the perifusion fluid, at a rate calculated to produce the required final concentration. NPPB, forskolin, and PMA were dissolved in DMSO and then diluted (1:1,000, vol/vol) in the perifusion solution to the required final concentration.

Statistical analysis. Data are presented as arithmetic means \pm standard deviations. Statistical analysis was conducted using the paired or unpaired Student's t test as appropriate or ANOVA when three groups were compared.

Results

Experiments in bicarbonate-free media. In nominally bicarbonate-free, Hepes-buffered media, IRHC maintained a basal pH_i

Table I. Recovery from an Acute Acid Load in the Presence (KRB) or Absence (Hepes) of Bicarbonate

Condition	Basal pH _i	Nadir pH _i	Recovery rates	JH ⁺	
			pH U/min	mM/min	
Hepes					
Controls $(n = 7)$	7.09 ± 0.07	6.70±0.09	0.145 ± 0.042	6.35 ± 2.50	
Glucagon	7.11±0.08	6.71±0.15	0.130 ± 0.020	5.58±1.89	
KRB + amiloride					
Controls $(n = 8)$	7.18±0.10	6.69 ± 0.07	0.172 ± 0.032	9.63 ± 3.20	
Glucagon	7.15±0.07	6.70±0.08	0.165 ± 0.042	8.94±3.72	
KRB, Cl ⁻ depletion + amiloride					
Controls $(n = 8)$	7.40 ± 0.07	7.12±0.02	0.063 ± 0.016	3.28 ± 0.82	
Glucagon	7.43 ± 0.06	7.13 ± 0.02	0.086±0.018*	4.49±1.20*	

Hepes, IRHC perfused with nominally bicarbonate-free Hepes-buffered media were submitted to a 20 mM NH₄Cl acid load. The recovery rate from this acid load is a measure of the Na⁺/H⁺ exchanger activity. When two consecutive NH₄Cl pulses were performed in the same cell, the second during superfusion with 200 nM glucagon, no difference in the recovery rates were found. KRB + amiloride, IRHC perfused with KRB were submitted to two consecutive 20 mM NH₄Cl acid loads. At the moment of NH₄Cl withdrawal, 1 mM amiloride was administered to exclude the Na⁺/H⁺ exchanger. The recovery from the NH₄Cl-amiloride acid pulse is a measure of the Na⁺-HCO₃ symport activity. When two consecutive NH₄Cl-amiloride pulses were performed in the same cell, the second in the presence of 200 nM glucagon, no difference in the recovery rates was found. KRB, Cl⁻ depletion + amiloride, to measure the activity of the Na⁺-HCO₃ symport at pH₁ values close to basal pH₁ of IRHC, we used a protocol where Cl⁻-depleted IRHC were submitted to a 15 mM NH₄-amiloride acid load. In these experimental conditions nadir acidification was 7.12±0.02. Glucagon administered in correspondence of a second NH₄-amiloride acid load significantly increased the recovery rate with respect to the basal values (first control NH₄-amiloride acid load). Findings indicate that glucagon stimulates the Na⁺-HCO₃ symport at pH₁ values close to the basal pH₁ of IRHC but not at acid pH₁ values. Data are means±SD. * P < 0.03 vs. control values.

of 7.09 ± 0.05 (n=13). Basal pH_i was unmodified by 10 min of exposure to 200 nM glucagon (n = 13). In the absence of bicarbonate, pH_i regulation in hepatocytes was mediated by the activity of the Na⁺/H⁺ exchanger (6, 23). The lack of a glucagon effect on basal pHi, in bicarbonate-free media, suggests that this hormone does not influence the activity of the Na⁺/ H+ exchanger. However, we directly evaluated whether glucagon may influence the activity of this exchanger, assessed by the effect of amiloride on basal pH_i and the recovery from acid load. Amiloride (1 mM), a specific inhibitor of the Na⁺/H⁺ exchange, promoted a 0.14±0.04 pH U acidification, the cell recovering to the basal pH_i after amiloride withdrawal. When amiloride exposure/withdrawal was repeated in the same cell but during superfusion with 200 nM glucagon, the same acidification was found (-0.15 ± 0.04 pH U; n=6). The activity of the Na⁺/H⁺ exchanger was then measured by evaluating the rate of pH_i recovery from an acute acid load (20 mM NH₄Cl⁻) in Hepes. The recovery from this acid load is driven by the activity of the Na⁺/H⁺ exchanger (6, 23). When two consecutive NH₄Cl⁻ pulses were applied, no significant difference in the maximal rate (at nadir pH_i) of pH_i recovery was found (n = 7). To evaluate the effect of glucagon, when the recovery from the first pulse was completed, the cell was superfused with 200 nM glucagon, and after 4-6 min a second NH₄Cl⁻ pulse was performed. The rate of recovery at nadir pH_i was compared with those measured after the first control pulse. During glucagon exposure, both the degree of acidification after NH₄Cl⁻ withdrawal (nadir pH_i = 6.71 ± 0.15) and the maximal rate of recovery $(0.130\pm0.020 \text{ pH U/min}; \text{JH}^+ = 5.58\pm1.89 \text{ mM/min})$ min) were similar to control values (n = 7; Table I). These findings further confirm that glucagon has no effect on the activity of the Na⁺/H⁺ exchanger, evaluated at either basal pH_i (amiloride effect on basal pH_i) or at acid pH_i values (recovery from an acute acid load).

Experiments in bicarbonate-containing media. The basal pH_i of IRHC cultured and perfused with media supplemented with

25 mM HCO $_3^-$ (7.20±0.07, n=72), was higher (P<0.001) than in the nominal absence of HCO $_3^-$, indicating that a HCO $_3^-$ loading mechanism is active. Exposure for 10 min to 200 nM glucagon (n=10), 500 nM glucagon (n=5), 200 nM glucagon + 100 μ M IBMX (n=5) showed no detectable changes in basal pH_i. Basal pH_i also remained unchanged when glucagon was administered in cells perifused in HCO $_3^-$ -containing media, with amiloride to inhibit the Na $^+$ /H $^+$ exchanger (n=6), thus excluding the involvement of this acid extruder as a mechanism of pH_i maintenance during glucagon exposure.

Effect of glucagon and DBcAMP + forskolin on Cl⁻/ HCO₃ exchange (Table II, Fig. 1). The activity of the Cl⁻/ HCO₃ exchanger was measured in IRHC, cultured and perfused with HCO3-enriched media, by using the method of acute external Cl⁻ removal/readmission (7, 8, 18, 24, 27–30). Acute Cl⁻ removal (equimolar substitution with gluconate) produces an intracellular alkalinization due to the exit of internal Cl in exchange with external HCO₃, pH_i recovering to baseline when external Cl⁻ is readmitted to drive the exchanger in the opposite direction. The net pH_i increase promoted by Cl⁻ removal (δ pH_i) and the rate of both pH_i increase after Cl⁻ removal and pH_i recovery after C1⁻ readmission were the parameters measured to evaluate the activity of the Cl⁻/HCO₃ exchanger. The effect of Cl⁻ removal was completely abolished by 0.5 mM DIDS pretreatment (40-60 min of preincubation; n = 6) indicating that the increase in pH_i induced by acute Cl⁻ removal depends on the transport of HCO₃ across the cell membrane. When two consecutive C1 removal/readmission maneuvers were performed no significant difference was found in the three measured parameters (n = 10). To evaluate the effect of glucagon, a second Cl removal/readmission maneuver was performed during superfusion with the hormone (200 nM; n = 10) and findings compared with those obtained after the first (control) maneuver. During glucagon perfusion, the net pH_i increase (0.28±0.04 pH U) and the maximal rate of alkalinization $(0.145\pm0.025 \text{ pH U/min}; H^+ \text{ flux} = 7.78\pm1.58 \text{ mM/min}) \text{ in-}$

Table II. Effect of Glucagon and DBcAMP + Forskolin on Chloride/Bicarbonate Exchange Activity

			Cl ⁻ removal		Cl ⁻ readmission	
	Basal pH _i	$\delta \mathrm{pH_{i}}$	pH U/min	H ⁺ flux	pH U/min	H ⁺ flux
		pH U		mM/min		mM/min
Controls $(n = 10)$	7.18±0.06	0.20 ± 0.07	0.063±0.013	3.18±0.73	0.096±0.042	6.51±3.01
Glucagon	7.18±0.07	$0.28\pm0.04*$	0.145±0.025*	7.78±1.58*	$0.145\pm0.048^{\ddagger}$	10.72±4.92‡
Controls $(n = 13)$	7.22±0.07	0.19 ± 0.07	0.065±0.030	3.60±1.96	0.105±0.028	7.20±2.02
DBcAMP + forskolin	7.20 ± 0.08	0.26±0.07*	0.127±0.040*	6.58±2.13*	0.160±0.070*	11.10±5.17*
DBcAMP + forskolin	7.20 ± 0.08	0.26±0.07*	0.127±0.040*	6.58±2.13*	0.160±0.070*	11.1

Two consecutive Cl^- removal/readmission maneuvers were performed, the second during superfusion with glucagon (200 nM, n = 10) or DBcAMP (100 μ M) + forskolin (10 μ M) (n = 13). The net pH_i increase (δ pH_i) and the maximal rate of pH_i increase after acute Cl⁻ removal as well as the maximal rate of pH_i recovery after Cl⁻ readmission were compared between the first (control) and the second (glucagon or DBcAMP + forskolin) acute Cl⁻ removal/readmission maneuver. Data are means \pm SD. * P < 0.02 vs. control values. \pm P < 0.05 vs. control values.

duced by Cl⁻ removal were significantly (P < 0.02) higher then control values (Table II, Fig. 1). Glucagon also increased (P < 0.05) the rate of pH_i recovery after Cl⁻ readmission (Table II, Fig. 1). This demonstrates that glucagon stimulates the activity of the Cl⁻/HCO₃⁻ exchanger.

To assess whether glucagon stimulation of the Cl⁻/HCO₃ exchange could be mediated by cAMP (31), we tried to reproduce the effect of the hormone by using a mixture composed




Figure 1. Effect of glucagon or DBcAMP + forskolin on pH_i changes promoted by acute chloride removal and readmission. IRHC were perfused with bicarbonate containing solutions (KRB). Two consecutive Cl⁻ removal/readmission (equimolar substitution with gluconate) maneuvers were then performed, the second during superfusion with 200 nM glucagon (upper tracing, n=10), or DBcAMP (100 μ M) + forskolin (10 μ M) (lower tracing, n=13). The net pH_i increase (δ pH_i) and the maximal rate of pH_i increase after acute Cl⁻ removal as well as the maximal rate of pH_i recovery after Cl⁻ readmission were significantly higher during exposure to glucagon or DBcAMP + forskolin with respect to control values (first acute Cl⁻ removal/readmission maneuver).

of a membrane permeant cAMP analogue (DBcAMP, $100~\mu M$) and a stimulator of adenyl cyclase (forskolin, $10~\mu M$). During superfusion with DBcAMP + forskolin, the net pH_i increase (0.26±0.07 pH U) and the maximal rate of alkalinization (0.127±0.040 pH U/min; H⁺ flux = $6.58\pm2.13~\text{mM/min}$) after Cl⁻ removal and the rate of pH_i recovery after Cl⁻ readmission (0.160±0.070 pH U/min; H⁺ flux = $11.10\pm5.17~\text{mM/min}$) were higher (P < 0.02) than corresponding control values (first maneuver, Table II, Fig. 1; n = 13). This indicates that the Cl⁻/HCO₃ exchanger is stimulated in conditions capable of increasing intracellular cAMP.

Effect of protein kinase A inhibitors (Rp-cAMPS, H-89) on glucagon stimulation of the Cl⁻/HCO₃ exchanger activity (Table III, Fig. 2). Cells were preincubated (30 min) and perfused with 10 μ M Rp-cAMPS (n=10) or with 30 μ M H-89 (n=10). Two consecutive Cl⁻ removal/readmission maneuvers were then performed, the second during superfusion with 200 nM glucagon. The effect of protein kinase A inhibitors on the basal (first Cl⁻ removal/readmission maneuver) and on the glucagon-stimulated (second maneuver) Cl⁻/HCO₃ exchange activity was evaluated in comparison (ANOVA) with control experiments performed in the absence of inhibitors, but with glucagon superfusion during the second Cl⁻ removal/readmission maneuver (Table III, Fig. 2).

IRHC pretreated and perfused with Rp-cAMPS or with H-89 showed a basal activity of the Cl⁻/HCO₃ exchanger (first Cl removal/readmission maneuver) similar to controls (n = 12). In fact, the net pH_i increase and the rate of alkalinization after Cl removal as well as the rate of pH_i recovery after Cl readmission were unchanged either by Rp-cAMPS or by H-89. In contrast, both Rp-cAMPS and H-89 significantly inhibited the glucagon-stimulated Cl⁻/HCO₃ exchange activity. In fact, when glucagon was administered (second Cl- removal/readmission maneuver), the net pH_i increase and the rate of alkalinization after Cl removal as well as the rate of pH_i recovery after Cl- readmission were significantly lower in IRHC pretreated and perfused with Rp-cAMPS (P < 0.03) or with H-89 (P < 0.02) in comparison with the corresponding parameters measured in control IRHC. Glucagon increased by 106% the basal rate of alkalinization induced by Cl⁻ removal (P < 0.01) and by 66% the basal rate of pH_i recovery after Cl⁻ readmission (P < 0.02) in control IRHC. In contrast, glucagon increased the basal rate of alkalinization induced by Cl⁻ removal by only 36% (P < 0.05) in Rp-cAMPS-treated IRHC while no significant increase was found in H-89-treated cells. The rate of pH_i recovery after Cl - readmission was unchanged by glucagon in either Rp-cAMPS- or H-89-treated cells.

Table III. Effect of Protein Kinase A Inhibitors (Rp-cAMPS, H-89) on the Basal and Glucagon-stimulated Chloride/Bicarbonate Exchange Activity

			Cl- removal		Cl ⁻ readmission	
	Basal pH _i	δ p H_i	pH U/min	H ⁺ flux	pH U/min	H ⁺ flux
		pH U		mM/min		mM/min
Controls $(n = 12)$	7.22±0.06	0.19±0.07	0.067±0.034	3.68±1.85	0.091±0.043	6.36±3.34
Glucagon	7.21 ± 0.05	0.26±0.05*	0.138±0.045* [‡]	7.49±2.44* [‡]	0.151±0.046* [‡]	10.83±3.39* [‡]
Rp-cAMPS $(n = 10)$	7.21±0.10	0.22±0.06	0.069±0.018	3.85±0.93	0.100±0.024	7.57±2.11
Rp-cAMPS + glucagon	7.22 ± 0.06	0.24±0.04	0.094±0.031§	5.20±1.62 [§]	0.105 ± 0.034	7.62 ± 2.00
H-89 $(n = 10)$	7.22±0.06	0.20±0.04	0.073±0.024	4.05±1.30	0.094±0.031	6.63±2.55
H-89 + glucagon	7.20 ± 0.10	0.23 ± 0.06	0.080 ± 0.025	4.39 ± 1.36	0.109 ± 0.037	7.42 ± 2.73

Two consecutive Cl⁻ removal/readmission maneuvers were performed, the second during superfusion with 200 nM glucagon, in IRHC pretreated (30 min) and perfused with 10 μ M Rp-cAMPS (n=10) or with 30 μ M H-89 (n=10). Data were compared (ANOVA) with respect to control untreated IRHC (n=12). The net pH_i increase (δ pH_i) and the maximal rate of pH_i increase after acute Cl⁻ removal as well as the maximal rate of pH_i recovery after Cl⁻ readmission measured during the first maneuver (basal activity of the Cl⁻/HCO₃⁻ exchanger) were not influenced by Rp-cAMPS nor by H-89. The same parameters measured in the presence of glucagon (second maneuver) were significantly higher in untreated IRHC with respect to IRHC treated with Rp-cAMPS or with H-89. Data are means±SD. * P < 0.02 vs. control basal values. P < 0.03 vs. Rp-cAMPS + glucagon or H-89 + glucagon.

When two consecutive $C1^-$ removal/readmission maneuvers were performed in IRHC pretreated and perfused with RpcAMPS (n = 6) or with H-89 (n = 6) but without glucagon administration, no significant differences were found in the three measured parameters between the first and the second maneuver.

Effect of the Cl⁻ channel blocker, NPPB, on the glucagon stimulation of the Cl⁻/HCO₃ exchanger activity (Table IV. Fig. 3). Since the activity of Cl⁻/HCO₃ exchangers may be regulated either by a direct effect of second messengers on the transporter or, indirectly, by modifying the rate of Cl- flux across Cl - channels in the cell membrane as described in other epithelia (11-15, 19, 32, 33), the effect of NPPB, a specific and potent Cl⁻ channel blocker (34), on the glucagon response in IRHC was assessed. NPPB alone showed no effect on the activity of the Cl⁻/HCO₃ exchange. In fact, when two consecutive Cl⁻ removal/readmission maneuvers were performed, the second in the presence of 10 μ M NPPB, no differences in the pH_i changes induced by Cl⁻ removal/readmission were found (n = 7, Table IV). To evaluate the effect of NPPB on the glucagon stimulation of the Cl⁻/HCO₃ exchange, after the first control maneuver, cells were superfused with 10 μ M NPPB and 2 min later with NPPB + glucagon (n = 7; Table IV, Fig. 3). Even in the presence of NPPB, glucagon stimulates (P < 0.02) the net pH_i increase after Cl^- removal (0.27±0.05 pH U), the rate of alkalinization after Cl removal (0.157±0.052 pH U/ min; H^+ flux = 9.34±3.16 mM/min) and the rate of pH_i recovery after Cl⁻ readmission (0.197±0.089 pH U/min; H⁺ flux = 13.40 ± 6.59 mM/min) with respect to the control values (first maneuver; Table IV, Fig. 3). These findings indicate that NPPB does not influence the glucagon stimulation of the Cl-/ HCO₃ exchanger, suggesting that the hormonal effect is not mediated by NPPB-sensitive Cl - channels.

Effect of PMA on the glucagon stimulation of Cl^-/HCO_3^- exchanger (Table IV, Fig. 3). To evaluate the role of protein kinase C on the basal and glucagon-stimulated Cl^-/HCO_3^- exchange, we investigated the effect of the agonist PMA on the pH_i changes promoted by Cl^- removal/readmission in the ab-

sence of or during glucagon exposure. PMA alone showed no effect on the activity of the Cl-/HCO3 exchanger. In fact, when two consecutive C1- removal/readmission maneuvers were performed, the second in the presence of 10 μ M PMA, no significant differences in the pH_i changes induced by Cl removal/readmission were found (n = 7, Table IV). To evaluate the effect of PMA on the glucagon stimulation of the Cl⁻/ HCO₃ exchange, at the end of the first control maneuver, cells were superfused with 10 μ M PMA and 4 min later with PMA + glucagon. A second Cl⁻ removal/readmission maneuver was then performed and findings compared with the first control maneuver (n = 13; Table IV, Fig. 3). During superfusion with glucagon + PMA, the net pH_i increase after Cl⁻ removal (0.22±0.05 pH U), the rate of alkalinization after Cl⁻ removal $(0.070\pm0.034 \text{ pH U/min}; H^+ \text{ flux} = 3.98\pm1.02 \text{ mM/min})$ and the rate of pH_i recovery after Cl⁻ readmission (0.130±0.063 pH U/min; H⁺ flux = 7.05 ± 3.17 mM/min) were similar to control values (first control maneuver; Table IV; Fig. 3). These findings indicate that PMA blocks the glucagon stimulation of the activity of the Cl⁻/HCO₃ exchanger.

Effect of colchicine on the glucagon stimulation of the Cl⁻/ HCO₃ exchanger activity (Table IV, Fig. 4). To evaluate the role of the microtubular system on the basal and glucagonstimulated Cl⁻/HCO₃ exchange activity, IRHC were pretreated with the microtubular inhibitor colchicine (10^{-5} M) or with its inactive analogue, β -lumicolchicine (10⁻⁵ M), for 4-5 h. Two consecutive Cl removal/readmission maneuvers were then performed, the second during superfusion with 200 nM glucagon. Colchicine pretreatment (n = 10) significantly (P < 0.03)decreased the net pH_i increase (0.12±0.04 pH U) and the rate of alkalinization after Cl⁻ removal (0.040±0.018 pH U/min; H^+ flux = 2.20±0.77 mM/min) as well as the rate of pH_i recovery after C1⁻ readmission (0.054±0.014 pH U/min; H⁺ flux = 3.36 ± 0.92 mM/min) in comparison with β -lumicolchicine-pretreated cells (n = 10; Table IV, Fig. 4). Moreover, during glucagon exposure in colchicine-pretreated cells, the net pH_i increase (0.22±0.08 pH U) and the rate of alkalinization after Cl⁻ removal (0.107±0.035 pH U/min; H⁺ flux




Figure 2. Effect of protein kinase A inhibitors (Rp-cAMPS, H-89) on pH_i changes promoted by acute chloride removal and readmission in the absence or presence of glucagon. Two consecutive Cl⁻ removal/readmission maneuvers were performed, the second during superfusion with 200 nM glucagon, in IRHC pretreated (30 min) and perfused with 10 μ M Rp-cAMPS (n=10, middle tracing) or with 30 μ M H-89 (n=10, lower tracing). Data were compared with respect to control untreated IRHC (n=12, upper tracing). The net pH_i increase (δ pH_i) and the maximal rate of pH_i increase after acute Cl⁻ removal as well as the maximal rate of pH_i recovery after Cl⁻ readmission measured during the first maneuver (basal activity of the Cl⁻/HCO₃ exchanger) were not influenced by Rp-cAMPS nor by H-89. The same parameters measured in the presence of glucagon (second maneuver) were significantly higher in untreated IRHC with respect to IRHC treated with Rp-cAMPS or with H-89.

= 5.92 ± 1.91 mM/min) as well as the rate of pH_i recovery after Cl⁻ readmission (0.120 ±0.034 pH U/min; H⁺ flux = 7.14 ± 2.99 mM/min) were significantly (P < 0.02) lower in comparison with those measured during glucagon exposure in β -lumicolchicine-treated cells. However, colchicine-pretreated cells still responded to the hormone, since both the net pH_i increase or the rate of alkalinization after Cl⁻ removal and the rate of pH_i recovery after Cl⁻ readmission were significantly (P < 0.05) increased by glucagon exposure (second Cl⁻ removal/readmission maneuver) in comparison with control values (first

Cl⁻ removal/readmission maneuver; Table IV, Fig. 4). These findings indicate that the inhibition of microtubular function decreased both the basal and glucagon-stimulated Cl⁻/HCO₃⁻ exchanger activity.

Effect of glucagon on the activity of the Na⁺-HCO₃ symport (Table I, Fig. 5). The activity of the Na⁺-HCO₃ symport was measured by evaluating the rate of recovery from 20 mM NH₄Cl pulse in cells cultured and perfused with HCO3enriched media. This recovery is driven by two acid extruder mechanisms, the Na⁺/H⁺ exchanger and the Na⁺-HCO₃⁻ symport (6). To exclude the influence of the Na⁺/H⁺ exchanger, when NH₄Cl was withdrawn, cells were perfused with 1 mM amiloride. When two consecutive NH₄Cl-amiloride pulses were performed (n = 7), no differences in the degree of acidification (nadir pH_i) and in the maximal rate of pH_i recovery from the acid load were found. To investigate the effect of glucagon, a second NH₄Cl-amiloride pulse was performed during glucagon exposure, and the maximal rate of pH; recovery was compared with the first control pulse (n = 8; Table I, Fig. 5). Both the degree of acidification (nadir pH_i = 6.70 ± 0.08) and the maximal rate of recovery from NH₄Cl⁻-amiloride pulse in the presence of glucagon were similar to the values measured in the absence of glucagon (nadir pH_i = 6.69 ± 0.07), indicating that the hormone has no effect on the activity of the Na⁺-HCO₃⁻ symport measured at acid pH_i values.

The Na⁺-HCO₃ symport activity and the effect of glucagon were also evaluated at pH_i values approaching the basal pH_i of IRHC and in conditions where Na⁺/H⁺ and Cl⁻/HCO₃ exchangers were blocked. For this purpose, cells were preincubated (40 min) and perfused with Cl--free media (equimolar substitution with gluconate). In these experimental conditions, IRHC showed a higher (P < 0.01) basal pH_i (7.40 ± 0.07) in comparison with cells incubated and perfused with media containing normal Cl - concentration, due to the blockade of the acid loader Cl⁻/HCO₃ exchanger. When Cl⁻-depleted IRHC were submitted to a 15 mM NH₄ acid pulse and amiloride was administered at the moment of NH₄ withdrawal, nadir acidification (7.12±0.02) was close to the basal pH_i of IRHC in medium containing normal Cl⁻ concentration (7.19±0.07). If H₂-DIDS (0.5 mM) was administered together with amiloride at the moment of NH_4^+ withdrawal (n = 4), the recovery from the acid load was completely abolished, indicating that the entrance of HCO₃ is responsible for this pH_i recovery.

Two consecutive 15 mM NH₄⁺ pulses were performed, the second during glucagon exposure and, at the moment of NH₄⁺ withdrawal, amiloride was superfused to exclude the influence of the Na⁺/H⁺ exchanger (n = 8; Table I, Fig. 5). In these experimental conditions, glucagon significantly (P < 0.03) stimulated the maximal rate of pH_i recovery (0.086 ± 0.018 pH U/min; JH⁺ = 4.49 ± 1.20 mM/min) from the NH₄⁺ acid load (nadir pH_i = 7.13 ± 0.02) with respect to control values (first pulse: nadir pH_i = 7.12 ± 0.02 ; recovery rate = 0.063 ± 0.016 pH U/min; JH⁺ = 3.28 ± 0.82 mM/min). This indicates that glucagon stimulates the activity of Na⁺-HCO₃⁻ symport at basal pH_i.

cAMP levels in isolated rat hepatocytes (Fig. 6). Fig. 6 shows the cAMP levels in isolated rat hepatocytes, incubated with glucagon, agonists, or inhibitors, under the same conditions as those for the pH_i measurement in IRHC. Exposure to 200 nM glucagon for 5 min increases cAMP levels sevenfold (from 0.98 ± 0.35 to 6.8 ± 2.4 pmol/ 10^6 cells; P < 0.001). Pretreatment with the protein kinase A inhibitor H-89 (30 μ M) for 40

Table IV. Effect of NPPB, PMA, or Colchicine on Basal and Glucagon-stimulated Chloride/Bicarbonate Exchanger Activity

		Cl ⁻ removal			Cl ⁻ readmission	
	Basal pH _i	δpH_i	pH U/min	H+ flux	pH U/min	H ⁺ flux
	-	pH U		mM/min		mM/min
Controls $(n = 7)$ NPPB	7.22±0.05	0.21±0.05	0.062±0.039	3.36±1.17	0.115±0.039	6.92±2.22
	7.21±0.06	0.20±0.06	0.060±0.032	3.16±1.06	0.103±0.069	6.22±2.01
Controls $(n = 7)$	7.21±0.04	0.19±0.04	0.067±0.030	3.36±1.07	0.120±0.035	7.22±2.42
Glucagon + NPPB	7.20±0.04	0.27±0.05*	0.157±0.052*	9.34±3.16*	0.197±0.089*	13.40±6.59*
Controls $(n = 7)$ PMA	7.21±0.09	0.22±0.06	0.058±0.018	3.48±1.07	0.102±0.050	6.14±2.86
	7.20±0.08	0.21±0.07	0.060±0.034	3.58±1.10	0.109±0.059	6.34±2.87
Controls $(n = 13)$	7.19±0.05	0.20±0.04	0.061±0.020	3.28±0.87	0.110±0.050	6.64±3.06
Glucagon + PMA	7.20±0.05	0.22±0.05	0.070±0.034	3.98±1.02	0.130±0.063	7.05±3.17
β -Lumicolchicine controls ($n = 10$)	7.20±0.06	0.21±0.04	0.068±0.018	3.48±0.97	0.110±0.050	6.79±3.22
β -Lumicolchicine + glucagon	7.21±0.08	0.29±0.07*	0.147±0.035*	8.02±1.91*	0.160±0.054 [‡]	11.87±5.42‡
Colchicine controls (n = 10)	7.23±0.04	0.12±0.04 [§]	0.040±0.018 [§]	2.20±0.77 [§]	$0.054\pm0.014^{\$}$	3.36±0.92 [§]
Colchicine + glucagon	7.21±0.05	0.22±0.08*	0.107±0.035*	5.92±1.91*	$0.120\pm0.034^{\ddagger }$	7.14±2.99 [‡]

To evaluate the effect of NPPB or PMA on the basal and glucagon-stimulated Cl^-/HCO_3^- exchanger activity, two consecutive Cl^- removal/readmission maneuvers were performed, the second during superfusion with NPPB alone ($10 \mu M$, n=7), NPPB ($10 \mu M$) + 200 nM glucagon (n=7), PMA alone ($10 \mu M$, n=7), or PMA ($10 \mu M$) + 200 nM glucagon (n=13). The net pH_i increase (δpH_i) and the maximal rate of pH_i increase after acute Cl^- removal as well as the maximal rate of pH_i recovery after Cl^- readmission were compared between the first (control) and the second (agonist or inhibitor with or without glucagon) acute Cl^- removal/readmission maneuver. To evaluate the effect of colchicine on the basal and glucagon-stimulated Cl^-/HCO_3^- exchanger activity, two consecutive Cl^- removal/readmission maneuvers were performed in cells pretreated with colchicine ($10^{-5} M$, 4-5 h), the second during superfusion with glucagon (200 nM, n=10). Findings were compared with β -lumicolchicine-pretreated cells (n=10). Data are means $\pm SD$. * P<0.02 vs. control values. * P<0.03 vs. β -lumicolchicine control values. * P<0.03 vs. β -lumicolchicine glucagon values.

min did not significantly influence basal cAMP levels nor the cAMP increase induced by exposure to 200 nM glucagon for 5 min. Nor did the Cl⁻ channel blocker, NPPB (10 μ M; 10 min) significantly change the basal or glucagon-stimulated cAMP levels. The protein kinase C agonist, PMA (10 μ M, 5 min of incubation), did not significantly change the basal cAMP levels but cells incubated with PMA for 4 min before exposure to 200 nM glucagon (5 min) exhibited 55% lower cAMP levels (3.06±1.03 pmol/10⁶ cells) with respect to glucagon-stimulated controls (6.8 \pm 2.4 pmol/10⁶ cells; P < 0.01). Colchicine pretreatment (10⁻⁵ M; 5 h) failed to significantly influence the accumulation of cAMP induced by 200 nM glucagon with respect to β -lumicolchicine-treated cells. cAMP levels were not determined in cells treated with Rp-cAMPS or with DBcAMP because these two compounds interfere with the cAMP assay system.

Discussion

Ion transport processes involved in pH_i regulation have been defined previously in rat hepatocyte subconfluent monolayers (6-8). In our study, pH_i measurement was performed in polarized IRHC, using a single cell microfluorimetric equipment. This method has the advantage of recording the signal from a polarized cell system, which represents the functional secretory unit in the liver (5), rather than averaging the signal from a whole cell population, containing either polarized or nonpolarized cells. Furthermore, viability of IRHC was considerably higher than the whole cell population and was less variable between different cell preparations. Consistent with the above,

the activities of transport processes involved in pH_i regulation measured in IRHC were higher than those reported in subconfluent rat hepatocyte monolayers (6-8).

Effect of glucagon on the Cl⁻/HCO₃ exchanger activity. In all the cell types studied (7, 8, 11-15, 18, 24, 27-30), the Cl⁻/HCO₃ exchanger has been shown to work as an acid loader driven by the Cl⁻ gradient. As an acid loader, this exchanger functions as a counterpoint to the acid extruder systems in the pH_i regulation on hepatocytes (6-8). The Cl⁻/HCO₃ exchanger was first identified in canalicular rat liver plasma membrane vesicles (35), then in monolayer culture of isolated hepatocytes (7) and recently it has been detected immunohistochemically in the hepatocyte apical domain of human liver (9). The two acid extruders, Na+-HCO3 symport and Na+/H+ exchanger, in addition to isolated rat hepatocytes (6), have also been identified in rat hepatocyte basolateral membrane vesicles (36, 37). It could thus be assumed that in the polarized IRHC, the Cl⁻/HCO₃ exchanger is located on the apical side and the two acid extruders are located on the basolateral side. Electrophysiological studies on IRHC have demonstrated that the resistances of tight junctions that seal the canalicular lumen from the bathing media are comparable with those of leaky epithelia and allow rapid ion equilibration between these two compartments (38). This implies that the behavior of the Cl⁻/HCO₃ in response to Cl - removal/readmission from the bathing media should be the same whether the exchanger is restricted to the canalicular domain of IRHC or not.

In the isolated perfused rat liver, stimulation of Cl⁻/HCO₃ exchanger by intracellular alkalinization induces a bicarbonate-rich choleresis, thereby suggesting the involvement of




Figure 3. Effect of NPPB or PMA on glucagon-stimulated pH_i changes promoted by acute chloride removal and readmission. (Upper tracing) After the first control Cl - removal/readmission (equimolar substitution with gluconate) maneuver, IRHC was perfused with the chloride channel blocker NPPB (10 μ M) and later with NPPB + 200 nM glucagon. A second Cl⁻ removal/readmission maneuver was then performed (n = 7). During superfusion with glucagon + NPPB, the net pH_i increase and the rate of alkalinization after Cl removal as well as the rate of pH_i recovery after Cl⁻ readmission were significantly higher than control values. (Lower tracing) After the first control Cl removal/readmission maneuver, IRHC was perfused with the protein kinase C agonist, PMA (10 μ M), and later with PMA + 200 nM glucagon. A second Cl^- removal/readmission maneuver was then performed (n = 13). During superfusion with glucagon + PMA, the net pH_i increase and the rate of alkalinization after Cl - removal as well as the rate of pH_i recovery after Cl - readmission were similar with respect to control values.

this exchanger in bicarbonate excretion in bile and in the generation of bile salt-independent canalicular bile flow (20). We demonstrated that glucagon stimulates the activity of the Cl⁻/HCO₃⁻ exchanger, a mechanism which could sustain the bicarbonate-rich choleresis induced by this hormone. Glucagon increased the rate of Cl⁻/HCO₃⁻ exchange by 100% when measured during Cl⁻ removal or by 50% after Cl⁻ readmission. In comparison, the excretion rates of bicarbonate in bile were increased by 34% in the rat (4) and by 71% in the guinea pig (2).

A number of studies have shown previously that glucagon, after stimulation of adenylate cyclase activity, increased the intracellular level of cAMP in isolated rat hepatocytes (31), in the isolated perfused rat (39), and in guinea pig liver (2). By extrapolating measurements of the entire preparation of isolated rat hepatocytes, the concentration of glucagon used to stimulate the Cl⁻/HCO₃ exchange in IRHC induces an almost maximal increase of cAMP levels. That cAMP is the intracellular messenger mediating the glucagon-induced bicarbonate biliary excretion was shown previously in the isolated perfused guinea pig liver (2), where a bicarbonate-rich choleresis was induced




Figure 4. Effect of colchicine on pH_i changes promoted by acute chloride removal and readmission in basal conditions and during glucagon exposure. IRHC were pretreated with colchicine (4–5 h, 10^{-5} M, n=10) or with its inactive analogue (β -lumicolchicine, n=10) in bicarbonate-containing media. Two consecutive Cl⁻ removal/readmission (equimolar substitution with gluconate) maneuvers were then performed, the second during superfusion with 200 nM glucagon. The net pH_i increase (δ pH_i) and the maximal rate of pH_i increase after acute Cl⁻ removal as well as the maximal rate of pH_i recovery after Cl⁻ readmission were significantly lower in colchicine-than in β -lumicolchicine—treated IRHC, either in basal conditions (first maneuver) or during glucagon exposure.

by glucagon but not by a glucagon analogue, TH-glucagon, which did not increase the intracellular level of cAMP. In addition, in the perfused rat liver, DBcAMP induced a bicarbonaterich choleresis (40). Our finding that DBcAMP + forskolin completely mimics the stimulatory effect of glucagon on Cl⁻/ HCO₃ exchange strongly suggests that this hormone enhances the bicarbonate excretion in bile via cAMP-mediated stimulation of an apically located (9) Cl⁻/HCO₃ exchanger. This is in keeping with studies concerning a number of different bicarbonate-secreting epithelia, where the intracellular messenger cAMP appears to stimulate the HCO₃ excretion mediated by the Cl⁻/HCO₃ exchanger (10-13, 17, 18). We then investigated the involvement of protein kinase A in the glucagon stimulation of the Cl⁻/HCO₃ exchanger activity using two different protein kinase A inhibitors, Rp-cAMPS and H-89. Rp-cAMPS is a distereomer of an analogue of natural cAMP which binds to the regulatory subunit of both type I and type II cAMPdependent protein kinase without dissociating the catalytic subunit (41). Rp-cAMPS is also a poor substrate for several phosphodiesterases (42). In rat hepatocytes, Rp-cAMPS inhibited glucagon-induced glycogenolysis, a cAMP-mediated effect (43). H-89, a recently synthesized isoquinolimesulfonamide, exhibited selective and potent inhibition on protein kinase A (44). Cells were preincubated and then perfused with RpcAMPS or with H-89, as indicated by previous studies, to




Figure 5. Effect of glucagon on the rate of pH_i recovery from an acute acid load in normal or chloride-depleted IRHC. (Upper tracing) In the presence of bicarbonate (KRB), two consecutive 20 mM NH₄Cl pulses were performed, the second during exposure to 200 nM glucagon (n = 8). At the moment of NH₄Cl withdrawal, amiloride was superfused to exclude the influence of the Na⁺/H⁺ exchanger. The degree of acidification and the rate of pH_i recovery after NH₄Cl withdrawal were not changed by glucagon. (Lower tracing) Cl⁻ depletion was performed by preincubation (40 min) and perfusion with Cl⁻-free media (equimolar substitution with gluconate). Two consecutive 15 mM NH₄ pulses were performed, the second during exposure to 200 nM glucagon (n = 8). At the moment of NH₄ withdrawal, amiloride was superfused to exclude the influence of the Na⁺/H⁺ exchanger. Glucagon significantly increased the rate of pH_i recovery from the acute acid load with respect to control values (first pulse).

achieve the maximal inhibitory effect on protein kinase A (43, 44). We demonstrated that the stimulatory effect of glucagon on Cl⁻/HCO₃⁻ exchange was almost completely blocked by pretreatment and perfusion with Rp-cAMPS or with H-89. Both these inhibitors had no impact on basal pH_i, nor on the basal activity of the Cl⁻/HCO₃⁻ exchanger and, at least with regard to H-89, did not decrease the cAMP accumulation induced by glucagon in isolated hepatocytes. Thus, our findings indicate that protein kinase A is directly involved in the glucagon stimulation of the Cl⁻/HCO₃⁻ exchange.

Epithelia that secrete HCO₃ via the Cl⁻/HCO₃ exchanger may regulate this secretory process by directly affecting the exchanger or, indirectly, by modifying the activity of Cl⁻ channels. In the presence of an in-to-out directed electrochemical Cl⁻ gradient, the activation of apically located Cl⁻ channels increases the luminal Cl⁻ concentration, thus favoring the gradient for the Cl⁻/HCO₃ exchange. To investigate whether the glucagon stimulation of the Cl⁻/HCO₃ exchange activity is a primary event or occurs secondary to Cl⁻ channel activation, the effect of NPPB, a potent and specific Cl⁻ channel blocker, was studied (34). We demonstrated that NPPB failed to influence either basal or glucagon-stimulated Cl⁻/HCO₃ exchange. This finding excludes that NPPB-sensitive Cl⁻ channels are




Figure 6. cAMP levels in isolated rat hepatocytes. Cells plated in 25 mM bicarbonate-containing media (KRB) were incubated with glucagon, agonists, or inhibitors, under the same conditions as those for the pH_i measurement in IRHC. Cell exposed for 5 min to increased concentration (from 1 nM to 1 μ M) of glucagon (GLUC.) exhibited a progressive accumulation of cAMP, with a sevenfold enhancement of basal cAMP levels obtained at 200 nM glucagon. Pretreatment with the protein kinase A inhibitor H-89 (30 μ M) for 40 min did not significantly influence basal cAMP levels nor cAMP increase induced by exposure to 200 nM glucagon for 5 min. The Cl⁻ channel blocker, NPPB (10 μ M; 10-min incubation), did not significantly change the basal nor the glucagon-stimulated cAMP levels. The protein kinase C agonist, PMA (10 μ M, 5-min incubation), did not significantly change the basal cAMP levels, but cells incubated with PMA for 4 min before exposure to 200 nM glucagon (5 min) exhibited 55% lower cAMP levels (3.06±1.03 pmol/10⁶ cells) with respect to 200 nM glucagon-stimulated controls $(6.8\pm2.4 \text{ pmol}/10^6 \text{ cells}; P < 0.01)$. Colchicine (COLCHI.) pretreatment (10⁻⁵ M; 5 h) failed to significantly influence the accumulation of cAMP induced by 200 nM glucagon with respect to β -lumicolchicine $(\beta$ -LUMICOLCHI.) - treated cells. *P < 0.01 vs. 200 nM glucagon. Data are means \pm SD from duplicate measurements in n = 6 cell preparations. Statistical analysis was conducted using ANOVA.

involved in glucagon stimulation of the exchanger. Different Cl⁻ channels could, however, be involved also in light of a recent description, in rat liver canalicular plasma membranes, of two types (30- and 90-pS conductance) of Cl⁻ channels with NPPB-insensitive conductance and kinetics (45).

In different cell types, several ion transport systems regulated by protein kinase A show inverse sensitivity to protein kinase C (46, 47). In the isolated perfused rat liver, protein kinase C agonists, phorbol esters, inhibit DBcAMP-stimulated bile flow (48) and, in isolated rat hepatocytes, they inhibit the HCO₃- and DBcAMP-stimulated Cl⁻/HCO₃ exchanger activity (8). This prompted us to explore the role of protein kinase C in the glucagon stimulation of the Cl⁻/HCO₃ exchanger. We demonstrated that the phorbol ester, PMA, completely blocked

the effect of glucagon on Cl⁻/HCO₃⁻ exchanger activity but, however, this effect was achieved through a marked inhibition of the glucagon-induced cAMP accumulation in isolated hepatocytes. Although an additional effect at the level of the Cl⁻/HCO₃⁻ exchanger cannot be excluded, our findings are consistent with a counterregulatory effect of protein kinase C, or PMA itself, at the level of glucagon receptors or adenylate cyclase. This is in keeping with previous studies showing that phorbol esters cause a dose-dependent inhibition of the glucagon-stimulated adenylate cyclase in plasma membranes isolated from rat hepatocytes, probably through protein kinase C activation and modification of the guanine nucleotide regulatory protein system (49).

The microtubular system is thought to be involved in many different transport processes in the liver as well as in the generation of bile flow (5). As far as glucagon choleresis is concerned, this is blocked by the microtubular inhibitor colchicine, which did not interfere with the metabolic effects or cAMP accumulation induced by the hormone (2). We found that both basal and glucagon-stimulated Cl⁻/HCO₃ exchange activities were inhibited by colchicine in comparison with β -lumicolchicine. However, in colchicine-treated cells, a glucagon stimulation of Cl⁻/HCO₃ exchanger is still present, although at a lower level than in β -lumicolchicine controls. The effect of colchicine occurred without influencing the glucagon-induced cAMP accumulation in isolated hepatocytes. These findings could be explained by the fact that the number of Cl⁻/HCO₃ units in the canalicular membranes was decreased by colchicine through a blockade of a microtubular-dependent targeting of vesicles containing the exchanger to the canalicular plasma membranes. Such regulatory mechanisms of the Cl⁻/HCO₃ exchanger have been proposed (8) after observations that colchicine inhibits the stimulation of the Cl⁻/HCO₃ exchanger induced by incubation of isolated hepatocytes in HCO₃-containing media.

Effect of glucagon on pH_i regulation in IRHC. Despite the increased bicarbonate excretion via Cl⁻/HCO₃ activation, glucagon showed no effect on basal pH_i of IRHC. The maintenance of basal pH_i, despite the increased bicarbonate excretion via Cl⁻/HCO₃ exchange stimulation, indicates that a compensatory mechanism in the cell is, in parallel, activated by glucagon. We investigated whether glucagon stimulation of the acid loading mechanism (Cl⁻/HCO₃ exchanger) is balanced by an increased activity of acid extruders. We have excluded an effect of glucagon on the Na⁺/H⁺ exchanger both when the activity of this acid extruder was measured at basal pHi (amiloride exposure withdrawal) and after an acute acid load. The other acid extruding system, Na+-HCO₃ symport, was not stimulated by glucagon at acid pH_i, as evaluated by the recovery rate from an acute acid load in KRB (20 mM NH₄Cl) and in the presence of amiloride (nadir pH = 6.7). However, since the stimulatory effect of glucagon on the Cl⁻/HCO₃ was demonstrated at basal pH_i (Cl⁻ removal) or at higher pH_i values (Cl⁻ readmission), we tried to use a protocol whereby the activity of the Na+-HCO₃ symport and the effect of glucagon could be measured at pHi values close to basal pHi of IRHC. Indeed, when we measured the recovery rate from a 15 mM NH₄ acid load in Cl^- -depleted and amiloride-perfused IRHC (nadir pH_i = 7.13) a significant stimulation by glucagon was observed. It was demonstrated previously, in Cl--depleted and amiloride-perfused hepatocyte monolayers, that the recovery from an acute acid load is driven by the Na+-HCO₃ symport (6), which is the major acid extruding mechanism in the presence of bicarbonate.




Figure 7. Proposed mechanisms of glucagon stimulation of the Cl $^-/HCO_3^-$ exchanger activity in IRHC. The figure shows the two acid extruders (Na $^+-H^+$ exchanger and Na $^+-HCO_3^-$ symport) located on the basolateral side and the acid loader (Cl $^-/HCO_3^-$ exchanger) on the canalicular side of one of the two hepatocytes forming a couplet. BC, bile canaliculus. After binding to receptor (R), glucagon stimulates adenylate cyclase (AC) and enhances cAMP levels which, through protein kinase A (PKA), increases the activity of the apically located Cl $^-/HCO_3^-$ exchanger. Despite the glucagon-stimulated bicarbonate excretion, resting pH_i of hepatocytes remains unchanged because of the parallel increase of bicarbonate loading through the stimulation of Na $^+$ -HCO $_3^-$ symport (dotted-line arrow). The Na $^+/H^+$ exchanger was unaffected by glucagon.

We further observed how, in these experimental conditions, the pH_i recovery from acid load is completely abolished by H₂-DIDS, evidence that HCO₃ entrance is involved. Thus, our findings indicate that glucagon stimulates bicarbonate loading throughout the Na⁺-HCO₃ symport but that this takes place only at basal pHi and not at acid pHi values. A pH dependence of the Na+-HCO3 symport and of its regulation, demonstrated recently in basolateral membrane vesicles isolated from rabbit renal cortex (50), could explain this finding. On the other hand, the glucacon stimulation of this symport occurred in Cl⁻-depleted cells, a condition where the Cl⁻/HCO₃ exchanger is almost entirely blocked. This indicates that HCO₃ entrance is stimulated by the hormone independently and not as a consequence of increased bicarbonate extrusion via Cl⁻/HCO₃ exchanger. The Na+-HCO3 symport could be stimulated subsequent to the increased Na+ gradient induced by glucagon stimulation of (Na+-K+)-ATPase, an effect already demonstrated in isolated hepatocytes (51).

The conclusions of our study are outlined in Fig. 7. Glucagon enhances cAMP levels which in turn, through protein kinase A, stimulate the Cl⁻/HCO₃ exchanger. This mechanism could account for in vivo induced bicarbonate-rich choleresis. Glucagon induces an upregulation of the pH_i regulatory mechanisms operating in the presence of bicarbonate since the activity of the bicarbonate loading system, Na⁺-HCO₃ symport, is also stimulated, thus avoiding changes of resting pH_i. We have also shown that glucagon stimulation of the Cl⁻/HCO₃ exchange is not mediated by NPPB-sensitive chloride channels, requires an intact microtubule system, and is counterregulated by the protein kinase C agonist, PMA, which blocks the cAMP accumulation induced by the hormone.

From a physiologic point of view, molecular mechanisms of glucagon-induced bicarbonate excretion should mainly operate in the fasting state in relationship to the raised serum level of the hormone. In these conditions, the bicarbonate excretion stimulated by the hormone may well play an important role as an osmotic driving force for canalicular bile formation since bile salt biliary secretion is at the minimal rate.

Acknowledgments

The authors thank Prof. S. Adamo and Dr. F. Folli for helpful discussions and Mrs. Joan Crowley for manuscript preparation.

References

- 1. Forker, E. L. 1977. Mechanisms of hepatic bile formation. Annu. Rev. Physiol. 39:323-347.
- 2. Lenzen, R., V. J. Hruby, and N. Tavoloni. 1990. Mechanism of glucagon choleresis in guinea pigs. Am. J. Physiol. 259:G736-G744.
- 3. Khedis, A., M. Dumont, and S. Erlinger. 1974. Influence of glucagon on canalicular bile production in the dog. *Biomedicine (Paris)*. 21:176-181.
- 4. Thomsen, O. O., and J. A. Larsen. 1981. The effect of glucagon, dibutyryl cyclic AMP, and insulin on bile production in the intact rat and the perfused rat liver. *Acta Physiol. Scand.* 111:23-30.
- Nathanson, H. M., and J. L. Boyer. 1991. Mechanisms and regulation of bile secretion. *Hepatology*. 14:551-566.
 Gleeson, D., N. D. Smith, and J. L. Boyer. 1989. Bicarbonate-dependent
- 6. Gleeson, D., N. D. Smith, and J. L. Boyer. 1989. Bicarbonate-dependent and -independent intracellular pH regulatory mechanisms in rat hepatocytes. Evidence for Na⁺:HCO₃⁻ cotransport. J. Clin. Invest. 84:312-321.
- 7. Benedetti, A., M. Strazzabosco, J. C. Corasanti, P. Haddad, J. Graf, and J. L. Boyer. 1991. Cl⁻/HCO₃ exchanger in isolated rat hepatocytes: role in regulation of intracellular pH. *Am. J. Physiol.* 261:G512–G522.
- 8. Benedetti, A., M. Strazzabosco, O. C. Ng., and J. L. Boyer. 1994. Regulation of activity and apical targeting of the Cl⁻/HCO₃ exchanger in rat hepatocytes. *Proc. Natl. Acad. Sci. USA*. 91:792-796.
- 9. Martines Anso, E., J. E. Castillo, J. Diez, J. F. Medina, and J. Prieto. 1994. Immunohistochemical detection of chloride/bicarbonate anion exchangers in human liver. *Hepatology*. 19:1400-1406.
- 10. Flemström, G., J. R. Heylings, and A. Garner. 1982. Gastric and duodenal HCO₃ transport in vitro: effects of hormones and local transmitters. *Am. J. Physiol.* 242:G100-G110.
- 11. Case, R. M., and B. E. Argent. 1986. Bicarbonate secretion by pancreatic duct cells: mechanism and control. *In* The Exocrine Pancreas: Biology, Pathobiology, and Diseases. V. L. W. Go, J. D. Gardner, F. P. Brooks, E. Lebenthal, E. P. Di Magno, and G. A. Scheele, editors. Raven Press, New York. 213–243.
- 12. Gray, M. A., J. R. Greenwell, and B. E. Argent. 1988. Secretin-regulated chloride channel on the apical plasma membrane of pancreatic duct cells. *J. Membr. Biol.* 105:131-142.
- 13. Novak, I., and R. Greger. 1988. Properties of the luminal membrane of isolated rat pancreatic ducts. Effect of cyclic AMP and blockers of chloride transport. *Pfügers Archiv.* 411:546-553.
- 14. Stuenkel, E. L., T. E. Machen, and J. A. Williams. 1988. pH regulatory mechanisms in rat pancreatic ductal cells. *Am. J. Physiol.* 254:G925-G930.
- 15. Stetson, D. L., R. Beauwens, J. Palmisano, P. P. Mitchell, and P. R. Steinmetz. 1985. A double-membrane model for urinary bicarbonate secretion. *Am. J. Physiol.* 249:F546-F552.
- 16. Sundaram, U., R. Knikelbein, and J. W. Dobbins. 1991. pH regulation in ileum: Na⁺-H⁺ and Cl⁻/HCO₃ exchange in isolated crypt and villus cells. *Am. J. Physiol.* 260:G440-G449.
- 17. Schuster, V. L. 1985. Cyclic adenosine monophosphate-stimulated bicarbonate secretion in rabbit cortical collecting tubules. *J. Clin. Invest.* 75:2056–2064.
- 18. Alvaro, D., W. K. Cho, A. Mennone, and J. L. Boyer. 1993. Effect of secretin on intracellular pH regulation in isolated rat bile duct epithelial cells. *J. Clin. Invest.* 92:1314-1325.
- 19. Fitz, J. G., S. Basavappa, J. M. McGill, O. Mehlus, and J. A. Cohn. 1993. Regulation of membrane chloride current in rat bile duct epithelial cells. *J. Clin. Invest.* 91:319-328.
- 20. Bruck, R. A., A. Benedetti, M. Strazzabosco, and J. L. Boyer. 1993. Intracellular alkalinization stimulates bile flow and vesicular-mediated exocytosis in IPRL. Am. J. Physiol. 28:G347-G353.
- 21. Gautam, A., O. C. Ng, and J. L. Boyer. 1987. Isolated rat hepatocyte couplets in short-term culture: structural characteristics and plasma membrane reorganization. *Hepatology*, 7:216-223.
- 22. Seglen, P. O. 1976. Preparation of isolated rat liver cells. *Methods Cell Biol.* 13:29-84.
- 23. Strazzabosco, M., S. Sakisaka, T. Hayakawa, and J. L. Boyer. 1991. Effect of UDCA on intracellular and biliary pH in isolated rat hepatocyte couplets and perfused livers. Am. J. Physiol. 260:G58-G69.
- 24. Alvaro, D., 'A. Mennone, and J. L. Boyer. 1993. Effect of ursodeoxycholic acid on intracellular pH regulation in isolated rat bile duct epithelial cells. *Am. J. Physiol.* 265:G783-G791.
- 25. Thomas, J. A., R. N. Buchsbaum, A. Zimniack, and E. Racker. 1979. Intracellular pH measurements in Erlich ascites tumor cells utilizing spectroscopic probes generated in situ. *Biochemistry*. 18:2210-2218.
- 26. Wenzl, E., and T. Machen. 1989. Intracellular pH dependence of buffer capacity and anion exchange in the parietal cell. Am. J. Physiol. 257:G741-G747.

- 27. Weintraub, W. H., and T. E. Machen. 1989. pH regulation in hepatoma cells: roles for Na⁺-H⁺ and Cl⁻-HCO₃ exchange and for Na⁺-HCO₃ symport. *Am. J. Physiol.* 257:G317-G327.
- 28. Boyarsky, G., M. B. Ganz, R. B. Sterzel, and W. F. Boron. 1988. pH regulation in single glomerular mesangial cells. I. Acid extrusion in absence and presence of HCO₃⁻. Am. J. Physiol. 255:C844-C856.
- 29. Boyarsky, G., M. B. Ganz, R. B. Sterzel, and W. F. Boron. 1988. pH regulation in single glomerular mesangial cells. II. Na⁺-dependent and -independent Cl⁻/HCO₁ exchangers. Am. J. Physiol. 255:C857—C869.
- 30. Roos, A., and W. F. Boron. 1981. Intracellular pH. Physiol. Rev. 61:296-432.
- 31. Corvera, S., J. Huerta-Bahena, J. T. Pelton, V. J. Hruby, D. Trivedi, and J. A. Garcia-Sainz. 1984. Metabolic effects and cyclic AMP levels produced by glucagon, (1-N-Trinitrophenylhistidine,12-homoerginine) glucagon and forskolin in isolated rat hepatocytes. *Biochim. Biophys. Acta.* 804:434-441.
- 32. Gray, M. A., J. R. Greenwell, and B. E. Argent. 1988. Secretin-regulated chloride channel on the apical plasma membrane of pancreatic duct cells. *J. Membr. Biol.* 105:131-142.
- 33. Argent, B. E., S. Arkle, M. E. J. Cullen, and R. Green. 1986. Morphological, biochemical and secretory studies on rat pancreatic ducts maintained in tissue culture. Q. J. Exp. Physiol. 71:633-648.
- 34. Wangermann, P., M. Wittner, A. Di Stefano, H. C. Henglert, H. J. Lang, E. Schlatter, and R. Greger. 1986. Cl⁻-channel blockers in the thick ascending limb of the loop of Henle. Structure activity relationship. *Pflügers Arch.* 407(Suppl.):S128-S141.
- 35. Meier, P. J., R. G. Knickelbein, R. H. Moseley, J. W. Dobbins, and J. L. Boyer. 1985. Evidence for carrier-mediated chloride/bicarbonate exchange in canalicular rat liver plasma membrane vesicles. *J. Clin. Invest.* 75:1256-1263.
- 36. Arias, I. M., and M. Forgac. 1984. The sinusoidal domain of the plasma membrane of rat hepatocytes contains an amiloride-sensitive Na⁺/H⁺ antiport. *J. Biol. Chem.* 259:5406-5408.
- 37. Renner, E. L., J. R. Lake, B. F. Scharschmidt, B. Zimmerli, and P. J. Meier. 1988. Rat hepatocytes exhibit basolateral Na⁺/HCO₃⁻ cotransport. *J. Clin. Invest.* 83:1225–1235.
- 38. Graf, J., R. M. Henderson, B. Krumpholz, and J. L. Boyer. 1987. Cell membrane and transepithelial voltages and resistances in isolated rat hepatocytes couplets. *J. Membr. Biol.* 95:241-254.
- 39. Schworer, C., and G. E. Mortimore. 1979. Glucagon-induced autophagy and proteolysis in rat liver: mediation by selective deprivation of intracellular amino acids. *Proc. Natl. Acad. Sci. USA*. 76:3169-3173.
- 40. Hayakawa, T., R. Bruck, O. C. Ng, and J. L. Boyer. 1990. DBcAMP stimulates vesicle transport and HRP excretion in isolated rat liver. Am. J. Physiol. 259:G727—G735
- 41. Rothermel, J. D., and L. H. P. Botelho. 1988. A mechanistic and kinetic analysis of the interaction of the diastereomers of adenosine 3',5'-(cyclic) phosphorothioate. *Biochem. J.* 251:757-762.
- 42. Jarvest, R. L., G. Lowe, J. Baraniak, and W. J. Stec. 1982. A stereochemical investigation of the hydrolysis of cyclic AMP and the (Sp)- and (Rp)-diastereoisomers of adenosine cyclic 3':5'-phosphorothioate by bovine heart and baker's-yeast cyclic AMP phosphodiesterases. *Biochem. J.* 203:461-470.
- 43. Rothermel, J. D., B. Jastorff, and L. H. P. Botelho. 1984. Inhibition of glucagon-induced glycogenolysis in isolated rat hepatocytes by the Rp diastereomer of adenosine 3',5'-(cyclic) phosphorothioate. *J. Biol. Chem.* 259:8151–8155.
- 44. Chijiwa, T., A. Mishima, M. Hagiwara, M. Sano, K. Hayashi, T. Inoue, K. Naito, T. Toshioka, and H. Hidaka. 1990. Inhibition of forskolin-induced neurite outgrowth and protein phosphorylation by a newly synthesized selective inhibitor of cyclic AMP-dependent protein kinase, N-[2-(p-bromocinnamyl-amino)ethyl]-5-isoquinolinesulfonamide (H-89), of PC12D pheochromocytoma cells. J. Biol. Chem. 265:5267-5272.
- 45. Sellinger, M., S. A. Weinman, R. M. Henderson, A. Zweifach, J. L. Boyer, and J. Graf. 1992. Anion channels in rat liver canalicular plasma membranes reconstituted into planar lipid bilayers. Am. J. Physiol. 262:G1027-G1032.
- Reuss, L. 1989. Ion transport across gallbladder epithelium. Physiol. Rev. 69:503-545.
- 47. Krapf, R., and R. J. Alpern. 1993. Cell pH and transepithelial H/HCO₃ transport in the renal proximal tubule. *J. Membr. Biol.* 131:1-10.
- 48. Corasanti, J. G., N. D. Smith, E. G. Gordon, and J. L. Boyer. 1989. Protein kinase C agonists inhibit bile secretion independently of effects on the microcirculation in the isolated perfused rat liver. Henotology. 10:8-13
- microcirculation in the isolated perfused rat liver. *Hepatology*. 10:8-13.
 49. Heyworth, C. M., S. P. Wilson, D. J. Gawler, and M. D. Houslay. 1985. The phorbol ester TPA prevents the expression of both glucagon desensitisation and the glucagon-mediated block of insulin stimulation of the peripheral plasma membrane cyclic AMP phosphodiesterase in rat hepatocytes. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 187:196-200.
- 50. Soleimani, M., G. A. Lesoine, J. A. Bergman, and T. D. McKinney. 1991. A pH modifier site regulates activity of the Na⁺:HCO₃⁻ cotransporter in basolateral membranes of kidney proximal tubules. *J. Clin. Invest.* 88:1135–1140.
- 51. Fehlmann, M., and P. Freychet. 1981. Insulin and glucacon stimulation of (Na⁺-K⁺)-ATPase transport activity in isolated rat hepatocytes. *J. Biol. Chem.* 256:7449-7453.