

Dissecting stepwise mutational impairment of megakaryopoiesis in a model of Down syndrome–associated leukemia

Edward J. Evans Jr., James DeGregori

J Clin Invest. 2022;132(14):e161659. <https://doi.org/10.1172/JCI161659>.

Commentary

Individuals with Down syndrome (DS) have more than 100-fold increased risk of acute megakaryoblastic leukemia (AMKL), but its pathogenesis is poorly understood. In this issue of the *JCI*, Arkoun et al. engineered stepwise DS-AMKL–associated mutations in *GATA1*, *MPL*, and *SMC3* in human induced pluripotent stem cell (iPSC) clones from individuals with DS to dissect how each mutation affects gene expression control and megakaryocytic differentiation. The authors showed that the mutations cooperatively promote progression from transient myeloproliferative disorder to DS-AMKL. This study highlights the importance of mutation order and context in the perturbations of transcriptional and differentiation pathways involved in the evolution of hematologic malignancies, which will be critical for the development of preventative and therapeutic interventions.

Find the latest version:

<https://jci.me/161659/pdf>



Dissecting stepwise mutational impairment of megakaryopoiesis in a model of Down syndrome-associated leukemia

Edward J. Evans Jr.^{1,2} and James DeGregori^{1,2}

¹Department of Biochemistry and Molecular Genetics and ²Linda Crnic Institute for Down Syndrome, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA.

Individuals with Down syndrome (DS) have more than 100-fold increased risk of acute megakaryoblastic leukemia (AMKL), but its pathogenesis is poorly understood. In this issue of the *JCI*, Arkoun et al. engineered stepwise DS-AMKL-associated mutations in *GATA1*, *MPL*, and *SMC3* in human induced pluripotent stem cell (iPSC) clones from individuals with DS to dissect how each mutation affects gene expression control and megakaryocytic differentiation. The authors showed that the mutations cooperatively promote progression from transient myeloproliferative disorder to DS-AMKL. This study highlights the importance of mutation order and context in the perturbations of transcriptional and differentiation pathways involved in the evolution of hematologic malignancies, which will be critical for the development of preventative and therapeutic interventions.

Acute megakaryoblastic leukemia of Down syndrome

Down syndrome (DS) is the result of trisomy of chromosome 21 (T21). Children with DS have a heightened susceptibility to many hematologic malignancies, especially leukemia (1, 2). Notably, 5% to 10% of children with DS under four years of age are diagnosed with the preleukemic state of transient myeloproliferative disorder (TMD), which is invariably driven by *GATA1* mutations generating a protein with a truncated N-terminus (*GATA1s*). Fortunately, in most of these children, the TMD and the associated *GATA1s* muta-

tion spontaneously resolve with age (3–5). However, 20% to 30% of these patients with TMD acquire additional mutations and progress to acute megakaryoblastic leukemia (DS-AMKL). The additional mutations occur in cohesin family genes, the CTCF gene encoding a key regulator of chromatin architecture, genes encoding members of the polycomb repressive complex-2 (PRC2), including *EZH2*, and components of cytokine signaling, such as *JAK2* or *MPL* (6–8). To help understand the evolution of TMD to DS-AMKL, it is necessary to develop human-relevant models that can recapitulate the DS-AMKL

mutations and allow examination of progressive perturbations of megakaryocytic differentiation and other disease phenotypes. In this issue of the *JCI*, Arkoun and colleagues accomplish this objective using a stepwise technique to introduce *GATA1*, *MPL*, and *SMC3* mutants into induced pluripotent stem cells (iPSCs) from humans with or without DS (9). The researchers uncovered the individual contributions of each variant and how they could cooperate with T21 to lead to DS-AMKL.

The authors generated 20 different disomic and trisomic iPSC clones harboring a combination of *GATA1s*, *MPL*^{W515K}, and heterozygous loss of *SMC3* (*SMC3*^{+/-}) using CRISPR/Cas9 technology for stepwise gene editing and validated these changes with functional assays. *MPL* is a transmembrane receptor for thrombopoietin, which is necessary for megakaryocyte maturation into platelets. The intracellular domain mediates signaling through interactions with *JAK2*. Multiple *MPL* gain-of-function amino acid substitutions at position 515 result in myeloproliferative disorders through thrombopoietin-independent activation of the *JAK/STAT* pathway (10). Interestingly, W515K/L mutations have also been seen in both AMKL from people with T21 and in leukemias from euploid individuals (D21) with somatic acquisition of an additional chromosome 21, potentially contributing to altered megakaryocytic differentiation (7, 11). *MPL* mutations in the context of T21 and *Gata1s* are sufficient to induce megakaryocytic leukemia in mice (12). Additionally, the authors hypothesized that haploinsufficiency of the cohesin gene *SMC3* through heterozygous inactivation would alter chromatin accessibility of *GATA1s* binding and consequently alter the transcriptional control of megakaryocytic differentiation. Given that these mutations individually contribute to disrupting the myeloid lineage, the stepwise iPSC models

► **Related Article:** <https://doi.org/10.1172/JCI156290>

Conflict of interest: JD serves as a member of the American Association of Cancer Research (AACR) Special Conferences Committee and the AACR Cancer Evolution Working Group Steering Committee and serves as an Executive Subcommittee member for the AACR Aging, Stress, and Cancer Task Force; is a co-chair and on the Steering Committee for the National Cancer Institute (NCI) and the National Institute on Aging (NIA) Onco-Aging Consortium; and is president of the International Society for Evolution, Ecology, and Cancer (ISEEC). JD also serves as associate editor at Molecular Cancer Research (AACR), editor-in-chief for Aging and Cancer (Wiley); associate editor at Evolution, Medicine, and Public Health (Oxford); and associate editor at Aging Cell (Wiley). JD is on the External Advisory Board at the University of Alabama O'Neal Comprehensive Cancer Center and is an External Advisory Board member at Moffitt Cancer Center and Mitotherapy. JD has a patent titled, "Use of tyrosine kinase inhibitor in cancer treatment" (US patent no. US10076520B2).

Copyright: © 2022, Evans et al. This is an open access article published under the terms of the Creative Commons Attribution 4.0 International License.

Reference information: *J Clin Invest.* 2022;132(14):e161659. <https://doi.org/10.1172/JCI161659>.




Figure 1. Progressive disruption in megakaryopoiesis in a model of DS-AMKL. The *GATA1s* mutation in hematopoietic progenitors induced in T21 iPSCs impairs megakaryopoiesis and leads to inefficient platelet generation and a TMD-like state, at least in part mediated by reduced chromatin accessibility and expression at NFE2 target genes. Subsequent inactivation of the cohesin complex component *SMC3* (*SMC3*^{+/-}) further reduces chromatin accessibility and expression of NFE2 target genes, leading to impaired platelet formation and the progression of a more AMKL-like state.

that the authors constructed enabled the authors to parse how each mutation interacted with the others to result in the DS and DS-AMKL phenotypes.

Cooperativity between mutants

There are two major strengths of the Arkoun et al. study. First, the authors started with hematopoietic progenitors derived from human iPSC cells generated from individuals with T21 or D21, contrasting with many previous studies using mouse models. Second, as mutations were generated in a stepwise fashion, the effect of individual mutations on cellular phenotypes could be compared with that of those in combination, providing insight as to how these mutations cooperate to lead to DS-AMKL. Using RNA-Seq for gene expression, ATAC-Seq for chromatin accessibility, and staining of α -granules and lysosomes for assessment of megakaryocytic maturation, Arkoun et al. showed phenotypic transformations at each step and how T21 enabled these changes. The authors showed that *GATA1s* with the addition of *SMC3*^{+/-} increased MK progenitors, synergistically prevented proplatelet (PPT) formation through a disrupted demarcation membrane system, and mimicked gene expression signatures previously reported for the progression from TMD to DS-AMKL. These observations are consistent with a differentiation block, enabling the expansion of progenitors through self-renewal, a hallmark of many forms of leuke-

mia. Notably, the *SMC3*^{+/-} mutation had opposite effects on platelet maturation and NFE2 inhibition, depending on whether or not it was preceded by the *GATA1s* mutation. Thus, the order in which mutations were acquired affected phenotypes controlling progression to DS-AMKL. Through RNA-Seq and ATAC-Seq, the authors showed that *GATA1s* hindered the NFE2 transcriptional pathway through reduced chromatin accessibility, with the subsequent *SMC3*^{+/-} mutation enhancing these effects (Figure 1). Importantly, ectopic expression of NFE2 partially restored megakaryopoiesis. NFE2 is a member of the Cap'N'Collar transcription factor family that regulates megakaryocytic maturation and platelet production and is important for oxidative stress response (13). Given that people with DS have increased reactive oxidative species and defective mitochondria (14), inhibition of the cytoprotective NFE2-regulated transcriptional program upon mutation of *GATA1s* (causing TMD) could provide additional pressure for further disease evolution. The mechanism by which cohesin mutations contribute to leukemogenesis has been elusive, and the Arkoun et al. study now provides a model through altered transcriptional control of key differentiation pathways. Future investigations will hopefully flesh out how cohesin disruption interacts with *GATA1s* mutation to further impair chromatin accessibility at these genes and thus promote TMD to AMKL progression.

The influence of the T21 context

The fact that *GATA1s*-driven TMD is apparently unique to individuals with DS (7) and that the prevalence of AMKL is substantially higher in people with DS highlights the importance of T21 in the onset of these hematologic malignancies. AMKL is much rarer in children without DS, and these cases of AMKL rarely exhibit the *GATA1s* mutation (7, 8). Notably, in some of these non-DS-AMKLs, somatic gain of chromosome 21 is observed. Other differences are apparent, including the frequency of mutations in cohesin complex components, occurring in about half of DS-AMKL, but less commonly present in non-DS-AMKL (approximately 10%) (7). Patients with non-DS-AMKL often exhibit chromosomal rearrangements that generate fusion proteins not seen in DS-AMKL. One could speculate that these alterations contribute to the impaired megakaryocytic differentiation otherwise imparted by *GATA1s* mutation in DS-AMKL. Intriguingly, instances of non-DS-AMKL that lack these fusion proteins frequently exhibit mutations in *GATA1* and are more likely to have cohesin mutations (15). Thus, DS-AMKL and non-DS-AMKL may represent convergent evolution to a similar disease through distinct mutational routes.

A recent study from Wagenblast and colleagues, leveraging CRISPR to engineer DS-AMKL mutations in human CD34⁺ fetal hematopoietic progenitors from individuals with and without DS, demonstrated that

inducing GATA1s in T21 (but not D21) progenitors leads to a TMD-like disease, even though GATA1s with a mutated cohesin gene can result in AMKL in both D21 and T21 contexts (16). T21 hematopoietic progenitors exhibit reduced clonogenic capacity relative to D21 progenitors, and GATA1s mutation restores clonogenicity selectively in the T21 background, suggesting that GATA1s mutation only enhances progenitor fitness in the T21 context (16). Of course, since subsequent mutations need to happen in the *GATA1s*-mutated background, the expanded GATA1s TMD clone (the target for the next mutation) promoted by the T21 context greatly increases the odds of progression to AMKL. Importantly, the Wagenblast et al. study also established that *GATA1s* mutation only engenders the preleukemic state in fetal or newborn hematopoietic progenitors, providing an explanation for why the window for DS-AMKL genesis is limited to very young children.

Arkoun et al. also recognized the importance of comparing isogenic disomic clones with mutations analogous to the trisomic models. Transcriptome analysis showed enrichment of the *MYC* and *MYB* proliferation pathways in *GATA1s/MPL^{W515K}/SMC3^{-/-}* (GMS) combined mutant clones in isogenic T21 clones relative to their D21 counterparts, consistent with greater proliferative and clonogenic potential when these mutations were engineered in the T21 background. Accordingly, the authors demonstrated that a *MYC* inhibitor caused greater reductions in the megakaryoblast proliferation in a T21-GMS context relative to D21-GMS, providing a targetable pathway for DS-AMKL and highlighting the value of investigating multistep carcinogenesis in different genotypic contexts.

The need for multiple models and next steps

It is worth emphasizing that Arkoun et al. constructed their model to specifically investigate megakaryocytic differentiation and proliferation in the DS context. While noting the consistency of their results with other published models of DS-AMKL, particular deviations serve as reminders of the limitations and the difficulties of fully recapitulating phenotypes observed in DS-AMKL. As noted by the authors, their iPSC model did not generate a leukemic state amenable to serial transplantation,

which is commonly used to showcase self-renewal, a hallmark of blood cancers. Additionally, while the *in vitro* model provides a powerful tool for molecular dissection, such a system lacks the extrinsic microenvironment present *in vivo*. Given that people with DS have increased inflammation (17), including hyperactive interferon signaling (18), and the tight interplay between inflammation and megakaryocyte and platelet maturation and function (19), *in vivo* studies of TMD and AMKL, whether in mouse models or by using human samples, could provide additional insight into DS-AMKL evolution.

Like many good studies, Arkoun et al. generates numerous intriguing questions and future directions. (a) Are there distinctions in megakaryocytic phenotypes in DS-AMKL with different cohesin gene alterations, and how do they interact with GATA1s and T21? (b) What are the cell-extrinsic factors that could influence the deleterious impact of AMKL mutations on megakaryocytic maturation, and are those factors T21-dependent? (c) How would additional mutations in the PRC2 complex, such as in *EZH2*, influence the transcriptional control of megakaryocytic differentiation and its alteration during AMKL development? (d) And perhaps most importantly, can an enhanced understanding of TMD and DS-AMKL evolution elucidate interventions to reduce progression (particularly at the TMD stage) or, better, to eliminate the disease? Hopefully, questions like these can be addressed in the future to better understand perturbations of megakaryopoiesis in people with DS and highlight therapeutic options to mitigate the prevalence and burden of DS-AMKL. With advancements in technology and pervasive use of multiomics approaches, we have the tools and opportunity to reduce the impact of TMD and AMKL on kids with DS.

Acknowledgments

This work was generously supported by the Blumenthal Fellowship from the Linda Crnic Institute for Down Syndrome (to EJE) and by R01HD103828 from the NIH/National Institute of Child Health and Human Development (NICHD) INCLUDE Project (to JD). We thank Joaquin Espinosa and Kelly Sullivan of the Linda Crnic Institute for insightful comments.

Address correspondence to: James DeGroot, University of Colorado Anschutz Medical Campus, Mail Stop 8101, 12801 East 17th Ave., Aurora, Colorado 80045, USA. Phone: 303.724.3230; Email: james.degroot@cuanschutz.edu.

- Hasle H, et al. Risks of leukaemia and solid tumours in individuals with Down's syndrome. *Lancet*. 2000;355(9199):165-169.
- Zeller B, et al. Acute leukaemia in children with Down syndrome: a population-based Nordic study. *Br J Haematol*. 2005;128(6):797-804.
- Crispino JD. GATA1 in normal and malignant hematopoiesis. *Semin Cell Dev Biol*. 2005;16(1):137-147.
- Roberts I, et al. GATA1-mutant clones are frequent and often unsuspected in babies with Down syndrome: identification of a population at risk of leukemia. *Blood*. 2013;122(24):3908-3917.
- McNulty M, Crispino JD. Acute megakaryocytic leukemia. *Cold Spring Harb Perspect Med*. 2020;10(2):a034884.
- Hussein K, et al. MPLW515L mutation in acute megakaryoblastic leukaemia. *Leukemia*. 2009;23(5):852-855.
- Yoshida K, et al. The landscape of somatic mutations in Down syndrome-related myeloid disorders. *Nat Genet*. 2013;45(11):1293-1299.
- Labuhn M, et al. Mechanisms of progression of myeloid preleukemia to transformed myeloid leukemia in children with down syndrome. *Cancer Cell*. 2019;36(2):123-138.
- Arkoun B, et al. Stepwise GATA1 and SMC3 mutations alter megakaryocyte differentiation in a Down syndrome leukemia model. *J Clin Invest*. 2022;132(14):e156290.
- Defour JP, et al. Oncogenic activation of MPL/thrombopoietin receptor by 17 mutations at W515: implications for myeloproliferative neoplasms. *Leukemia*. 2016;30(5):1214-1216.
- Pardanani AD, et al. MPL515 mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients. *Blood*. 2006;108(10):3472-3476.
- Malinge S, et al. Increased dosage of the chromosome 21 ortholog *Dyrk1a* promotes megakaryoblastic leukemia in a murine model of Down syndrome. *J Clin Invest*. 2012;122(3):948-962.
- Motohashi H, et al. NF-E2 domination over Nrf2 promotes ROS accumulation and megakaryocytic maturation. *Blood*. 2010;115(3):677-686.
- Bordi M, et al. mTOR hyperactivation in Down Syndrome underlies deficits in autophagy induction, autophagosome formation, and mitophagy. *Cell Death Dis*. 2019;10(8):1-17.
- De Rooij JDE, et al. Pediatric non-Down syndrome acute megakaryoblastic leukemia is characterized by distinct genomic subsets with varying outcomes. *Nat Genet*. 2017;49(3):451-456.
- Wagenblast E, et al. Mapping the cellular origin and early evolution of leukemia in Down syndrome. *Science*. 2021;373(6551):eabf6202.
- Huggard D, et al. Immune dysregulation in children with Down Syndrome. *Front Pediatr*. 2020;8:73.
- Sullivan KD et al. Trisomy 21 consistently activates the interferon response. *Elife*. 2016;5:1-28.
- Margraf A, Zarbock A. Platelets in inflammation and resolution. *J Immunol*. 2019;203(9):2357-2367.