







	Uncomplicated CDI (n=39)	Recurrent CDI within 90 days (n=12)	Death within 90 days (n=5)	P value
Female Sex	20 (51.3%)	7 (58.3%)	4 (80%)	0.14
Age in a year, Mean (IQR)	56.5 (46-67)	59.3 (52-71.5)	72.4 (63-80.5)	0.02
Race/ethnicity:				0.07
White/Caucasian	30 (76.9%)	10 (83.3%)	3 (60%)	0.05
Black/African- American	7 (17.9%)	2 (16.7%)	2 (40%)	0.03
Hispanic/Latino	1 (2.6%)	0 (0%)	0 (0%)	0.51
Asian/Other	1 (2.6%)	0 (0%)	0 (0%)	0.51
Peak WBC, Mean (IQR)	12.6 (6.1-16.3)	12.2 (3.0-19.2)	14.1 (6.2-22.6)	0.82
Charlson Comorbidity Index, Mean (IQR)	4.3 (2-6)	4.8 (2.5-7)	8 (6-10)	0.001


Supplemental Table 1: Basic patient characteristics.




Supplemental Figure 1.











Supplemental Figure 2.




Supplemental Figure 3.



Supplemental Figure 4.




Supplemental Figure 5.




Supplemental Figure 6.


A




B




Supplemental Figure 7.



Supplemental Figure 8.




Supplemental Figure 9.




Supplemental Figure 10.


A




B




Supplemental Figure 11.



B



C



Supplemental Figure 12.