

FoxO1 expression in osteoblasts regulates glucose homeostasis through regulation of osteocalcin in mice

Marie-Therese Rached, Aruna Kode, Barbara C. Silva, Dae Young Jung, Susan Gray, Helena Ong, Ji-Hye Paik, Ronald A. DePinho, Jason K. Kim, Gerard Karsenty, Stavroula Kousteni

J Clin Invest. 2010;120(3):932-932. <https://doi.org/10.1172/JCI39901C1>.

Corrigendum

Metabolism

Original citation: *J. Clin. Invest.* 2010;120(1):357–368. doi:10.1172/JCI39901. Citation for this corrigendum: *J. Clin. Invest.* 2010;120(3):932. doi:10.1172/JCI39901C1. The legend for Figure 5H was incorrect. The correct text appears below. (H) Changes in uncarboxylated or undercarboxylated Ocn in serum of WT and *Foxo1ob*^{–/–} mice; n = 5 mice/group. The authors regret the errors.

Find the latest version:

<https://jci.me/39901C1/pdf>





Table 3

Summary of original and new experiments with Substance P and intracellular calcium

	I	II	III	I + II	II + III
	Original experiment with suspect SubP	Original experiments with good SubP	New experiments	Published data	Corrected data
Resting $[Ca^{2+}]_i$	72–105 nM	70–120nM	85–140 nM	70–120nM	70–140 nM
<i>n</i> , subjects; <i>n</i> cells	1; 6	4; 33	4; 25	5; 39	8; 58
<i>n</i> (%) cells responding to SubP	6/6 (100%) ^A	25/33 (76%)	22/25 (88%)	31/39 (79%)	47/58 (81%)
$[Ca^{2+}]_i$ response to SubP (peak value, nM)	134 ± 34 ^A	140 ± 32	125 ± 18	139 ± 33	133 ± 35
<i>n</i> cells responding to carbachol (1 or 10 μM)	6/6 (100%)	33/33 (100%)	25/25 (100%)	39/39 (100%)	58/58 (100%)
$[Ca^{2+}]_i$ response to Carb 1 μM (peak value, nM)	194 ± 10	186 ± 17	Not done	187 ± 19	186 ± 17
$[Ca^{2+}]_i$ response to Carb 10 μM (peak value, nM)	256 ± 20	252 ± 15	202 ± 35	253 ± 17	231 ± 36
PAS positivity in SubP responsive cells	2/6 (33%)	7/25 (28%)	Not done	9/31 (29%)	7/25 (28%)
PAS positivity in SubP nonresponsive cells	NA	6/8 (75%)	Not done	6/8 (75%)	6/8 (75%)
% cells responding to SubP (10 μM) in presence of atropine	Not done	Not done	22/22 (100%)	Not done	22/22 (100%)

^APublished data generated using possibly contaminated Substance P.

Corrigendum

FoxO1 expression in osteoblasts regulates glucose homeostasis through regulation of osteocalcin in mice

Marie-Therese Rached, Aruna Kode, Barbara C. Silva, Dae Young Jung, Susan Gray, Helena Ong, Ji-Hye Paik, Ronald A. DePinho, Jason K. Kim, Gerard Karsenty, Stavroula Kousteni

Original citation: *J Clin Invest.* 2010;120(1):357–368. doi:10.1172/JCI39901.

Citation for this corrigendum: *J Clin Invest.* 2010;120(3):932. doi:10.1172/JCI39901C1.

The legend for Figure 5H was incorrect. The correct text appears below.

(H) Changes in uncarboxylated or undercarboxylated Ocn in serum of WT and *Foxo1^{ob-/-}* mice; *n* = 5 mice/group.

The authors regret the error.

Erratum

Urea-induced ROS generation causes insulin resistance in mice with chronic renal failure

Maria D'Apolito, Xueliang Du, Haihong Zong, Alessandra Catucci, Luigi Maiuri, Tiziana Trivisano, Massimo Pettoello-Mantovani, Angelo Campanozzi, Valeria Raia, Jeffrey E. Pessin, Michael Brownlee, and Ida Giardino

Original citation: *J Clin Invest.* 2010;120(1):203–213. doi:10.1172/JCI37672.

Citation for this erratum: *J Clin Invest.* 2010;120(3):932. doi:10.1172/JCI37672E1.

During the preparation of the manuscript, the urea infusion rate was incorrectly given. The correct sentence containing the infusion rate appears below.

The rats were allowed to recover from the surgery for 5 days and then were either infused with isotonic PBS or urea (100 mg/kg/h) for 48 hours using a microdialysis pump.

The JCI regrets the error.