Supplemental figure legends


Suppl Figure 1. Expression and purification of ecto-LRP4. (A) Schematic diagrams of full length LRP4 and ecto-LRP4. (B) Sliver staining of wash-through and purified ecto-LRP4.


Suppl Figure 2. LRP4 sera did not recognize AChR α protein. Lysates of HEK293 cells expressing AChR α were subjected to western blot by anti-AChR antibody mAb35 and by anti-LRP4 sera.


Suppl Figure 3. No effect of motor neuron-specific LRP4 mutation on muscle weakness induced by LRP4 immunization. (A, B) Similar reduction in body weight and muscle strength between LRP4-injected *HB9-LRP4*^{+/+} mice (control) and *HB9-LRP4*^{-/-} mice. (C, D) Reduced CMAP amplitudes at 20 Hz (C) or at 40 Hz (D) in LRP4-injected *HB9-LRP4*^{+/+} mice (control) and *HB9-LRP4*^{-/-} mice. (E) Reduced CMAP amplitudes of the 10th stimulation at different stimulation frequencies. (F) Representative mEPP traces. Traces on the right were enlarged from those underlined on the left. (G, H) Similar reduction of mEPP amplitude and frequency in the two genotypes after LRP4 immunization. Control, n = 3; *HB9-LRP4*^{-/-} mice, n = 5; LRP4-injected *HB9-LRP4*^{-/-} mice, n = 5. 3-4 muscle fibers were recorded from each mouse.


Suppl Figure 4. Immunization of rabbits with ecto-LRP4. (A) Detection of anti-LRP4 antibodies in sera of ecto-LRP4-injected rabbits. Sera were collected one week after first boost and subjected to ELISA. n = 3. (B) Images of representative control and LRP4-injected rabbits. (C) Reduced body weight of LRP4-injected rabbits.

Suppl Figure 5. IgG purification from LRP4-injected rabbit sera. (A) Schematic diagram of IgG purification procedures. (B, C) Analysis of purified rabbit IgGs on 8% non-denaturing PAGE (B) and 12% denaturing SDS-PAGE (C). Fractions surrounding 25th fraction were pooled for injection.










Control

LRP4



В Α Size Exclusion Chromatography Sera collection 150 Saturated ammonium sulfate precipitation 100 Dissolution in PBS Dialysis in PBS 50-Concentration by centri-YM30 column Chromatography on sephacryl S-200 column OD280 quantification Analysis by PAGE



40th Fraction

Albumin

lgG(H)

lgG(L)