

## $\alpha$ ENaC: leading the charge

Young S. Oh, ... , Sunil Saxena, David G. Warnock

*J Clin Invest.* 1999;104(7):849-850. <https://doi.org/10.1172/JCI8378>.

### Commentary

The amiloride-sensitive epithelial Na<sup>+</sup> channel (ENaC) is expressed in the apical membranes of many Na<sup>+</sup>-absorptive epithelia, such as renal tubules, distal colon, and lungs. The  $\alpha$  subunit of ENaC ( $\alpha$ ENaC), and subsequently the  $\beta$  and  $\gamma$  subunits, were identified by expression cloning (1, 2). The latter subunits potentiate amiloride-sensitive Na<sup>+</sup> currents more than 100-fold when coinjected with  $\alpha$ ENaC into *Xenopus* oocytes, but neither the  $\beta$  nor the  $\gamma$  subunit — alone or coexpressed — is sufficient to generate appreciable Na<sup>+</sup> currents in this system in the absence of  $\alpha$ ENaC. Thus, this subunit appears to be required for the assembly or function of the ENaC complex. The 3 known ENaC subunits are homologous to each other, having about 35% amino acid identity and conserved membrane topology. Site-directed mutagenesis has defined a short segment (the P region depicted in Figure 1) that precedes the second transmembrane domain of all 3 ENaC subunits and is involved in ion permeation and sensitivity to amiloride. Several naturally occurring  $\alpha$ ENaC splice variants have been identified, but their physiological significance has not been characterized (Figure 1) (3). Clinical disorders due to malfunction of the ENaC complex are well described (4). Liddle's syndrome, an autosomal dominant form of volume-expanded low-renin hypertension, is caused by gain-of-function mutations in the COOH-termini of the  $\beta$  or  $\gamma$  ENaC subunits that constitutively [...]

Find the latest version:

<https://jci.me/8378/pdf>



Young S. Oh,<sup>1,2,3</sup> Sunil Saxena,<sup>1,3</sup> and David G. Warnock<sup>1,3</sup>

<sup>1</sup>Department of Medicine,

<sup>2</sup>Department of Neurobiology, and

<sup>3</sup>Nephrology Research and Training Center, University of Alabama–Birmingham, Birmingham, Alabama 35294, USA

Address correspondence to: David G. Warnock, Division of Nephrology, University of Alabama–Birmingham, UAB Station, Birmingham, Alabama 35294-0007, USA. Phone: (205) 934-3585; Fax: (205) 934-1879; E-mail: [dwarnock@nrtc.dom.uab.edu](mailto:dwarnock@nrtc.dom.uab.edu).

The amiloride-sensitive epithelial Na<sup>+</sup> channel (ENaC) is expressed in the apical membranes of many Na<sup>+</sup>-absorptive epithelia, such as renal tubules, distal colon, and lungs. The  $\alpha$  subunit of ENaC ( $\alpha$ ENaC), and subsequently the  $\beta$  and  $\gamma$  subunits, were identified by expression cloning (1, 2). The latter subunits potentiate amiloride-sensitive Na<sup>+</sup> currents more than 100-fold when coinjected with  $\alpha$ ENaC into *Xenopus* oocytes, but neither the  $\beta$  nor the  $\gamma$  subunit — alone or coexpressed — is sufficient to generate appreciable Na<sup>+</sup> currents in this system in the absence of  $\alpha$ ENaC. Thus, this subunit appears to be required for the assembly or function of the ENaC complex. The 3 known ENaC subunits are homologous to each other, having about 35% amino acid identity and conserved membrane topology. Site-directed mutagenesis has defined a short segment (the P region depicted in Figure 1) that precedes the second transmembrane domain of all 3 ENaC subunits

and is involved in ion permeation and sensitivity to amiloride. Several naturally occurring  $\alpha$ ENaC splice variants have been identified, but their physiological significance has not been characterized (Figure 1) (3).


Clinical disorders due to malfunction of the ENaC complex are well described (4). Liddle's syndrome, an autosomal dominant form of volume-expanded low-renin hypertension, is caused by gain-of-function mutations in the COOH-termini of the  $\beta$  or  $\gamma$  ENaC subunits that constitutively activate ENaC activity (5). In contrast, loss-of-function mutations in the  $\alpha$  and  $\gamma$  ENaC subunits have been found in autosomal recessive pseudohypoaldosteronism type I (PHA-I) with salt wasting, hyperkalemia, and metabolic acidosis (6, 7).

The critical role of  $\alpha$ ENaC in vivo has been demonstrated by  $\alpha$ ENaC gene knockouts in mice; these animals expire with severe respiratory insufficiency (8). In contrast, patients with PHA-I do not

experience such respiratory disturbances but do exhibit abnormal renal Na<sup>+</sup> handling. This discrepancy between the clinical phenotype and the knockout mouse models might be explained by physiological differences between the species, such as the existence of a protective mechanism that is either unique to the human lung or not expressed in the mouse lung. Alternatively, the  $\alpha$ ENaC knockout animal might prove to be an inadequate model for human disease if the human mutant alleles retain some biological function.


In support of the latter class of explanation, Bonny et al. (9) now provide evidence that an  $\alpha$ ENaC truncation mutant promotes the channel activity of ENaC complexes containing normal  $\beta$  and  $\gamma$  subunits. It appears that the previous failure to detect functional  $\beta\gamma$  ENaC channels was primarily due to delayed expression of these channels in oocytes and to the relatively low levels of the resulting amiloride-sensitive Na<sup>+</sup> currents. Moreover, a truncated  $\alpha$ ENaC, equivalent to the R508X mutation that has been described in human PHA-I cases, can assemble with  $\beta$  and  $\gamma$  subunits to produce higher amiloride-sensitive Na<sup>+</sup> currents than the  $\beta\gamma$  channels alone. Although these current levels are only about 10% of those found with wild-type  $\alpha\beta\gamma$  ENaC, they could be sufficient, if expressed in vivo, to prevent development of the perinatal pulmonary phenotype in PHA-I patients. Bonny et al. have further proposed that the truncated  $\alpha$ ENaC, which retains the NH<sub>2</sub>-terminus and the extracellular cysteine-rich domain, may contain a sorting signal that promotes the assembly and proper intracellular trafficking of the  $\beta\gamma$  channels to the cell membrane.

Whether  $\alpha$ ENaC is necessary for expression of ENaC activity in cells remains a key question, but the formation of functional ENaC  $\beta\gamma$  complexes could account for the difficulty of detecting  $\alpha$ ENaC in some tissues (10).



**Figure 1**

$\alpha$ ENaC-associated splice variants and mutations identified in human (h), rat (r), and chicken (c). The  $\alpha$ ENaC contains 2 transmembrane domains (TMs), 2 cysteine-rich domains (CRDs), a leucine-rich domain (LRD), and a pore-forming region (P) just before TM2. The human  $\alpha$ ENaC2+59 contains an additional 59 amino acids at the NH<sub>2</sub>-terminus (14). Three human  $\alpha$ ENaC splice variants (h $\alpha$ ENaC C229x, h $\alpha$ ENaC-19, and h $\alpha$ ENaC+22) cause loss of function in *Xenopus* oocytes (3). The h $\alpha$ ENaC+Alu contains an Alu fragment at the CRD2 (Y.S. Oh and D.G. Warnock, unpublished data). Two  $\alpha$ ENaC splice variants ( $\alpha$ ENaCax and  $\alpha$ ENaCb) were identified in rat taste tissue (15), and  $\alpha$ ENaCx was identified in chicken cochlea (16); these splice variants, as well as h $\alpha$ ENaC+22 and h $\alpha$ ENaC+Alu, use the same splice site in the CRD2 exon (3). Four human  $\alpha$ ENaC mutations (G37S, I68x, C133Y, and R508x) have been identified in PHA-I individuals (6, 7). Here, “x” denotes a prematurely truncated protein, whether the new stop codon is generated by point mutation or frameshift.



**Figure 2**

Possible subunit arrangement of the ion-conducting pore in  $\alpha_x\beta\gamma$  channels and  $\beta\gamma$  channels. A possible ENaC structure consisting of 2  $\alpha$ , 1  $\beta$ , and 1  $\gamma$  subunit is shown as the wild-type channel. A truncated PHA-I mutation, R508x, was created in rat  $\alpha$ ENaC at the equivalent position (L535x), and  $\alpha_{L535x}$ , together with rat  $\beta$  and  $\gamma$  subunits, was expressed in *Xenopus* oocytes (9). Coexpression of rat  $\beta$  and  $\gamma$  ENaC subunits also generated amiloride-sensitive  $\text{Na}^+$  currents in oocytes. Interestingly,  $\alpha_{L535x}\beta\gamma$  and  $\beta\gamma$  channels displayed very similar macroscopic characteristics, such as reduced amiloride affinity and reduced  $\text{Li}^+$  selectivity, suggesting that the conductive pores of these channels are composed of P and TM2 domains of  $\beta$  and  $\gamma$  subunits, respectively.

The special role of  $\alpha$ ENaC in channel expression is confirmed by the work of Masilamani et al., also in this issue of the *JCI* (11). For the first time, these authors have used subunit-specific antibodies to investigate the in vivo regulation of ENaC expression in the renal collecting tubule, and, in particular, the physiologically relevant modulation of ENaC by aldosterone and dietary salt balance. Previous studies of ENaC expression have utilized heterologous expression systems, but the intrinsic limitations of this approach and the importance of examining protein expression and localization in vivo are now evident. Masilamani et al. demonstrate the apical localization of the assembled ENaC complex in response to aldosterone. They also show that  $\alpha$ ENaC is crucial for assembly and targeting of the channel complex to the apical membrane. There is also a provocative migration shift of the  $\gamma$  subunit on gel analysis, consistent with some sort of post-translational processing, perhaps by the CAP-1 serine protease? (12).

Beyond the evident importance of clarifying the pathophysiology of PHA-

I, the significance of both of these papers (9, 11) lies in the insights they offer into normal cell physiology. The analysis of this inborn error lets us appreciate the special role of  $\alpha$ ENaC in "leading the charge" of ionic current flow across the apical membrane. Archibald Gerrod would be pleased but not surprised (13).

1. Lingueglia, E., Voilley, N., Waldmann, R., Lazdunski, M., and Barbry, P. 1993. Expression cloning of an epithelial amiloride-sensitive  $\text{Na}^+$  channel. *FEBS Lett.* **318**:95-99.
2. Canessa, C.M., Horisberger, J.-D., and Rossier, B.C. 1993. Epithelial sodium channel related to proteins involved in neurodegeneration. *Nature.* **361**:467-470.
3. Tucker, J.K., et al. 1998. Cloning and functional studies of splice variants of the  $\alpha$ -subunit of the amiloride-sensitive  $\text{Na}^+$  channel. *Am. J. Physiol.* **274**:C1081-C1089.
4. Scheinman, S.J., Guay-Woodford, L.M., Thakker, R.V., and Warnock, D.G. 1999. Genetic disorders of renal electrolyte transport. *N. Engl. J. Med.* **340**:1177-1187.
5. Warnock, D.G. 1996. Polymorphism in the beta subunit and  $\text{Na}^+$  transport. *J. Am. Soc. Nephrol.* **7**:2490-2494.
6. Grunder, S., et al. 1997. A mutation causing pseudohypoaldosteronism type 1 identifies a conserved glycine that is involved in the gating of the epithelial sodium channel. *EMBO J.* **16**:899-907.
7. Chang, S.S., et al. 1996. Mutations in subunits of the epithelial sodium channel cause salt wasting

- with hyperkalaemic acidosis, pseudohypoaldosteronism type 1. *Nat. Genet.* **12**:248-253.
8. Hummler, E., et al. 1996. Early death due to defective neonatal lung liquid clearance in  $\alpha$ ENaC-deficient mice. *Nat. Genet.* **12**:325-328.
9. Bonny, O., et al. 1999. Functional expression of a pseudohypoaldosteronism type I mutated epithelial  $\text{Na}^+$  channel lacking the pore-forming region of its  $\alpha$  subunit. *J. Clin. Invest.* **104**:967-974.
10. Kopp, U.C., et al. 1998. Amiloride-sensitive  $\text{Na}^+$  channels in pelvic uroepithelium involved in renal sensory receptor activation. *Am. J. Physiol.* **275**:R1780-R1792.
11. Masilamani, S., Kim, G.-H., Mitchell, C., Wade, J.B., and Knepper, M.A. 1999. Aldosterone-mediated regulation of ENaC  $\alpha$ ,  $\beta$  and  $\gamma$  subunit proteins in rat kidney. *J. Clin. Invest.* **104**:R19-R23.
12. Coles, J.A., Orkand, R.K., Yamate, C.L., and Tsacopoulos, M. 1986. Free concentrations of Na, K, and Cl in the retina of the honeybee drone: Stimulus-induced redistribution and homeostasis. *Ann. NY Acad. Sci.* **481**:303-317.
13. Bearn, A.G. 1993. *Archibald Gerrod and the individuality of man*. Clarendon Press, Oxford, United Kingdom. 227 pp.
14. Thomas, C., Auerbach, S., Stokes, J.B., and Volk, K.A. 1998. 5' heterogeneity in epithelial sodium channel  $\alpha$ -subunit mRNA leads to distinct NH<sub>2</sub>-terminal variant proteins. *Am. J. Physiol.* **274**:C1312-C1323.
15. Li, X.-J., Xu, R.-H., Guggino, W.B., and Snyder, S.H. 1995. Alternatively spliced forms of the  $\alpha$  subunit of the epithelial sodium channel: distinct sites for amiloride binding and channel pore. *Mol. Pharmacol.* **47**:1133-1140.
16. Killick, R., and Richardson, G. 1997. Isolation of chicken alpha ENaC splice variants from a cochlear cDNA library. *Biochim. Biophys. Acta.* **1350**:3-37.