

Isolated polycystic liver disease genes define effectors of polycystin-1 function

Whitney Besse, ... , Vicente E. Torres, Stefan Somlo

J Clin Invest. 2017;127(9):3558-3558. <https://doi.org/10.1172/JCI96729>.

Corrigendum

Original citation: *J Clin Invest.* 2017;127(5):1772–1785. <https://doi.org/10.1172/JCI90129> Citation for this corrigendum: *J Clin Invest.* 2017;127(9):3558. <https://doi.org/10.1172/JCI96729> In the original article, the RT-PCR primer sequences listed in Methods were incorrectly labeled as Pkd1. The correct primer sequences for Pkd1 are in the revised paragraph below. Quantitative PCR and reverse transcription PCR. RNA was isolated from cultured cells using Trizol Reagent (Invitrogen). cDNA was reverse transcribed from RNA using reagents from New England Biolabs. Primers for Pkd1 quantitative PCR (forward, GCTACAGGGCATCCTGGTG; reverse, GGCTGTCAGCGAGAGCTTGAA) were designed using NCBI's primer-designing tool (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). Quantitative PCR was done by Bio-Rad CFX Connect Real-Time PCR Detection System. Primers for Xbp1 RT-PCR have been published previously (1). The authors regret the error.

Find the latest version:

<https://jci.me/96729/pdf>



Corrigendum

Isolated polycystic liver disease genes define effectors of polycystin-1 function

Whitney Besse, Ke Dong, Jungmin Choi, Sohan Punia, Sorin V. Fedeles, Murim Choi, Anna-Rachel Gallagher, Emily B. Huang, Ashima Gulati, James Knight, Shrikant Mane, Esa Tahvanainen, Pia Tahvanainen, Simone Sanna-Cherchi, Richard P. Lifton, Terry Watnick, York P. Pei, Vicente E. Torres, and Stefan Somlo

Original citation: *J Clin Invest.* 2017;127(5):1772–1785. <https://doi.org/10.1172/JCI90129>.

Citation for this corrigendum: *J Clin Invest.* 2017;127(9):3558. <https://doi.org/10.1172/JCI96729>.

In the original article, the RT-PCR primer sequences listed in Methods were incorrectly labeled as *Pkd1*. The correct primer sequences for *Pkd1* are in the revised paragraph below.

Quantitative PCR and reverse transcription PCR. RNA was isolated from cultured cells using Trizol Reagent (Invitrogen). cDNA was reverse transcribed from RNA using reagents from New England Biolabs. Primers for *Pkd1* quantitative PCR (forward, GCTA-CAGGGCATCCTGGTG; reverse, GGCTGTCAGCGAGAGCTTGAA) were designed using NCBI's primer-designing tool (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). Quantitative PCR was done by Bio-Rad CFX Connect Real-Time PCR Detection System. Primers for *Xbp1* RT-PCR have been published previously (1).

1. Iwakoshi NN, Lee AH, Vallabhajosyula P, Otipoby KL, Rajewsky K, and Glimcher LH. Plasma cell differentiation and the unfolded protein response intersect at the transcription factor XBP-1. *Nat Immunol.* 2003;4(4):321–329.

The authors regret the error.